1

How the Central American Seaway and an ancient northern passage affected flatfish diversification

5
6 Lisa Byrne[1], François Chapleau[1], and Stéphane Aris-Brosou[*,1,2]
7
8 [1]*Department of Biology, University of Ottawa, Ottawa, ON, CANADA*
9 [2]*Department of Mathematics & Statistics, University of Ottawa, Ottawa, ON, CANADA*
10
11 [*]**Corresponding author: E-mail:** *sarisbro@uottawa.ca*
12

Abstract

While the natural history of flatfish has been debated for decades, the mode of diversification of this biologically and economically important group has never been elucidated. To address this question, we assembled the largest molecular data set to date, covering > 300 species (out of *ca*. 800 extant), from 13 of the 14 known families over nine genes, and employed relaxed molecular clocks to uncover their patterns of diversification. As the fossil record of flatfish is contentious, we used sister species distributed on both sides of the American continent to calibrate clock models based on the closure of the Central American Seaway (CAS), and on their current species range. We show that flatfish diversified in two bouts, as species that are today distributed around the Equator diverged during the closure of CAS, while those with a northern range diverged after this, hereby suggesting the existence of a post-CAS closure dispersal for these northern species, most likely along a trans-Arctic northern route, a hypothesis fully compatible with paleogeographic reconstructions.

Keywords: Pleuronectiformes, Bayesian dating, trans-Arctic migration, vicariance, Central American Seaway, Isthmus of Panama

The Pleuronectiformes, or flatfishes, are a speciose group of ray-finned fish, containing

14 families and over 800 known species (Munroe 2015). Flatfish begin life in the pelagic

zone, but undergo a larval metamorphosis in which one eye, either left or right,

depending on the species, migrates to the other side of the cranium. The adult fish then

adopts a benthic lifestyle. Flatfish have asymmetric, laterally-compressed bodies, and

have lost their swim bladders during transformation. With eyes facing upwards, flatfish

are also capable of protruding them. This singular morphology long puzzled taxonomists

(Norman 1934), and the phylogeny of this group remains poorly resolved.

At the highest taxonomic level, flatfishes are generally considered to be monophyletic,

based on both morphological (Chapleau 1993), and molecular evidence (Berendzen et al.

2002; Pardo et al. 2005; Azevedo et al. 2008; Betancur-R et al. 2013; Harrington et al.

2016). All these studies also support the monophyletic status of most families within the

order, to the exception of the Paralichthyidae. As all molecular studies to date have

essentially focused on the monophyletic status of the order, they were based on as many

representative species of each order. As a result, intra-ordinal relationships, among

genera and even families, are still debated. It can therefore be expected that taking

advantage of both species- and gene-rich evidence, while incorporating paleontological

and/or geological data in the framework of molecular clocks, should help clarify not only

the phylogenetic status of this family (dos Reis et al. 2015), but also – and more critically

here – their evolutionary dynamics.

However, very few flatfish fossils are known (Chanet 1997; Friedman 2012), and

placed with confidence on the flatfish evolutionary tree (Parham et al. 2012; Campbell et

al. 2014a; Harrington et al. 2016). As a result, calibrating a molecular clock becomes

challenging. On the other hand, a dense species sampling may enable us to take

advantage of a singular feature of flatfishes: some extant species are found both in the

Pacific and Atlantic oceans. Furthermore, the existence of geminate species pairs of

flatfishes, where sister taxa have one member in each ocean, suggests a speciation event

pre-dating the formation of the Isthmus of Panama, which occurred approximately 12 to 3

million years ago [MYA] (Haug and Tiedemann 1998; O'Dea et al. 2016). Also possible

is the role played by the opening of the Bering Strait around 5.5-5.4 MYA (Gladenkov et

al. 2002), which would have permitted the trans-Arctic migration of ancestral populations

from one ocean to the other, as documented in marine invertebrates (*e.g.*, Durham and

MacNeil 1967; Reid 1990) or vertebrates (*e.g.*, Grant 1986), but to date, never in flatfish.

Our driving hypothesis is then that the sole information relative to the formation of the

Isthmus of Panama can be used to calibrate molecular clocks, and allows us to unravel

the timing of flatfish evolution, as how rapidly they diversified remains an unsolved

question. We show here that the diversification of flatfish in the seas surrounding the

Americas followed a complex pattern, related to not only the closure of the Isthmus of

Panama, but also to a warming event that opened up a trans-Arctic northern route.

69   Results

To test how the evolutionary dynamics of flatfish were affected by a major geological

event, the closure of the Central American Seaway (CAS), we reconstructed dated

Bayesian phylogenetic trees from a large data matrix under four relaxed molecular clock

models, each one of them being based on a different calibration scheme (Fig. 1). Under

the first model, no calibration priors were placed on internal nodes. This initial tree, with

the rogue sequences removed (data on GitHub: see Methods), was used to identify pairs

of sister taxa that are split between the two oceans, with one species in the Atlantic and

the other in the Pacific. This led us to single out twelve pairs of such species. These sister

species happened not to be evenly distributed on the estimated phylogeny (Fig. 2, top),

but they are the only geminate species included in GenBank (as of August 2016). On

each pair, we placed calibration date priors corresponding to the closure of the CAS (the

'ALL' model). An examination of the posterior distributions of their divergence times

suggests that some species have a very narrow speciation window, where all the mass of

the posterior distribution is between 5-3 MYA, while others have a wider distribution

(Fig. 3A). Closer inspection of these distributions further reveals that most of the species

with narrow posterior distributions have a northern range (Fig. 2, and 3A, in blue), while

those with the wider posterior distributions have a "southern" distribution, closer to the

Isthmus of Panama (Fig. 2, and 3A, in red). To further assess this observation, we first

went back to the original clock model, removed all priors initially placed on sister taxa

(the 'NONE' model), and were able to validate that even in this case: northern and

southern species showed, to one exception each (*Hippoglossus hippoglossus* and *H.*

stenolepis in the north, and *Poecilopsetta natalensis* and *P. hawaiiensis* in the south; Fig.

2), shifted posterior distributions (Fig. 3B). The former pair was actually not estimated as

being sister species in any of the four clock models (Fig. 1), while *P. natalensis* and *P.*

hawaiiensis, although inhabiting the Western Indian and Eastern Pacific oceans, occupy

ranges that do not extend to the coasts of the Americas as with the other identified sister

taxa. Models with priors placed only on northern (the 'NORTH' model: Fig. 3C) or

southern (the 'SOUTH' model: Fig. 3D) species also showed a similar temporal shift.

This shift suggested that southern species diverged early, before the complete closing of

the CAS, while northern species diverged later, at or possibly after the isthmus was

completed. Averaging these posterior distributions for the northern and southern species,

to the exception of the two outliers noted above, showed these results more clearly (Fig.

3E-H). To assess the robustness of this result to the inclusion of the *P. natalensis* and *P.*

hawaiiensis sister species, we conducted an additional set of analyses removing all prior

information from this pair. The ALL and SOUTH models, which are the only ones where

this prior occurs, show that the general pattern that we found above holds: southern

species diverged early, before the complete closing of the CAS, and northern species

diverged later, at or possibly after the isthmus was completed (Fig. S1).

All these results were obtained assuming that Psettodidae is an appropriate outgroup

for these analyses. However, the position of Psettodidae in flatfishes is still debated

(Betancur-R. and Orti 2014; Campbell, et al. 2014a; Campbell, et al. 2014b; Harrington,

et al. 2016), so that their inclusion might affect our results. To assess this possibility, we

reran all four models without Psettodidae, letting the molecular clock root the tree (Aris-

Brosou and Rodrigue, 2012). Figure S2 shows the exact same results as above (Fig. 3), so

that our dating results are also robust to the inclusion, or not, of Psettodidae.

In an attempt to tease out these models and their predictions about the exact timing of

divergence between northern and southern species, we assessed model fit by means of the

modified Akaike's Information Criterion (AICM; Baele et al. 2013), which accounts for

uncertainty in the MCMC sampling (Raftery et al. 2007). Even if model ranking based on

this measure is known to be unstable (Baele et al. 2013), it is clear that the models with

priors only on the northern or on the southern species perform significantly (> 200 AIC

units) more poorly than the two other models, which may be difficult to tease apart

(Table 1). The predictions of the best models suggest that *H. hippoglossus* and *H.*

stenolepis, both northern species, consistently diverged before the complete closure of the

seaway, in tandem with the average southern species, and that the average northern

species diverged in tandem with *P. natalensis* and *P. hawaiiensis* (Fig. 3E-F).

One intriguing result from these analyses is that the northern sister species seem to

come from a single clade (Fig. 2), which would suggest that it is possible to identify the

direction of the trans-Arctic migration. For this, we retrieved the current species range of

all flatfish analyzed here (Fig. S3), and indicated their average latitude on the

phylogenetic trees (Fig. S4). This northern clade, highly supported (posterior probability

> 0.9; see asterisk in Fig. S4), is mostly comprised of species currently found in the

Pacific (*e.g.*, *Reinhardtius evermanni*), suggesting that the northern sister species that we

identified originated from the Pacific, and that their trans-Arctic migration was from the

Pacific into the Atlantic in a northern direction.

Discussion

Our results show that flatfish underwent a first speciation at the time when the CAS

closed, which led to the species that, today, have an equatorial range (Fig. 2), as the

formation of the Isthmus of Panama resulted in a barrier to gene flow leading to their

speciation. Our results also show that species that today have a northern range (Fig. 2)

either emerged at the closure of the seaway, or after its closure. These timing estimates

imply that this second bout of speciation was not caused by gene flow impeded by the

closure of the seaway, but demand an interpretation involving trans-Arctic gene flow,

through a northern route, before being interrupted, most likely this time by a climatic

event. Critically, these conclusions are robust to the inclusion of outgroup species

(Psettodidae), of outliers species (*P. natalensis* and *P. hawaiiensis*), and are consistently

found across the four clock models implemented here, including the one (NONE) relying

on no other fossil information than the 47.8-42.1 MYA ancestor to this order (Chanet

1997; Friedman 2012) – hereby indicating that our data are highly informative with

respect to the timing of the divergence of these sister species (and rejecting our initial

hypothesis) following their trans-CAS and tans-Arctic migrations.

Trans-Arctic migrations from the Pacific to the Atlantic are not new, and have been

described in both invertebrates (*e.g.*, Durham and MacNeil 1967) and vertebrates (*e.g.*,

Grant 1986), but never using a relaxed molecular clock relying on as little information as

a single fossil and a large multigene data set covering an entire order (Pleuronectiformes).

While seminal work was usually based on paleontological (Durham and MacNeil 1967),

or on morphological evidence (Reid 1990), subsequent work started using allozymes and

other proteins (Grant 1986; Zaslavskaya et al. 1992), followed by genetic evidence,

mostly at the population level using mitochondrial DNA (Rawson and Harper 2009).

However, most of the recent work that relies on molecular clocks either makes strong

assumptions on substitution rates (Rawson and Harper 2009; Liu et al. 2011), uses

divergences that are vicariance-motivated but not supported by any fossil information

(McCusker and Bentzen 2010), or posit in which direction the trans-Arctic migration

took place (Dodson et al. 2007) to infer trans-Arctic migrations. Our work is the first to

use sister species to demonstrate the existence and elucidate the timing of both the trans-

CAS and the trans-Arctic events, while also infering in which direction the latter took

place, without making too many assumptions.

Strikingly, the geological evidence is directly in line with our date estimates. The

fossil record suggests that the first aquatic connection between the Pacific and Arctic

(and Atlantic) oceans through the Bering Strait occurred approximately 5.5-5.4 MYA

(Gladenkov et al. 2002) due to a rise in sea levels, linked to tectonic activity

(Marincovich and Gladenkov 2001). This would have permitted the trans-Arctic

migration of populations ancestral to today's northern species from one ocean to the other

through this ancient "northern passage." This global warming event, between the late

Miocene to early Pleiocene, was then followed by a significant period of cooling during

the Pleiocene into the Pleistocene (Zachos et al. 2001), leading to periods of repeated

glaciations and a subsequent ice age. These cold events would have resulted in the

closure of this ancient "northern passage," hereby stopping the trans-Arctic gene flow

between the two oceans, and leading to the more recent speciation of the northern taxa.

For approximately a million years after the first opening of the Bering Strait, water

flowed through the strait in a southern direction, from the Arctic to the Pacific ocean,

until the formation of the Isthmus of Panama occurred close to the equator (Berta 2012).

With the formation of the Isthmus, and hence the closing of the CAS, the ocean currents

reversed due to a change in global ocean circulation (Haug and Tiedemann 1998; De

Schepper et al. 2015), and have since flowed in a northern direction through the Bering

Strait, from the Pacific to the Arctic (Marincovich 2000). This again is absolutely

consistent with our results (Fig. S4), with dispersal or migration from the Pacific into the

Atlantic in a northern direction through this strait, which is known as the trans-Arctic

interchange (Vermeij 1991). Fossil data also show that the Bering land bridge has been

exposed and submerged on multiple occasions since the Pleistocene (Gladenkov and

Gladenkov 2004). These openings and closings of the Bering Strait could have provided a

mechanism for divergence and the evolution of sister taxa (Taylor and Dodson 1994;

Väinölä 2003).

Our results also have implication at the family level of flatfish. Further significant

global cooling during the Pleistocene resulted in major glaciation events (Zachos et al.

2008) that could be responsible for creating barriers that isolated populations. All of the

remaining sister taxa in our analysis, who have divergence estimates of less than 2 MYA

in our study belong to the family Pleuronectidae (*sensu* Cooper and Chapleau 1998). The

Pleuronectidae are the predominant flatfish family found in cold and temperate seas of

the northern hemisphere (Norman 1934; Cooper and Chapleau 1998). There are far more

Pleuronectidae species in the Pacific Ocean, most of them endemic to the north Pacific

Ocean off of North America and Asia in the region extending from the Bering Strait to

the gulf of California (Norman 1934). None of the arctic species are restricted solely to

arctic waters (Munroe 2015). Munroe also noted that Cooper (in an unpublished

manuscript) identified areas of endemism among the current distribution of the

Pleuronectidae. It has been shown that during the trans-Arctic interchange, there was a

far higher number of species (up to eight times higher) migrating to the Atlantic than to

the Pacific (Vermeij 1991). Fossil and phylogenetic evidence suggest the Pacific Ocean

as the origin for diversification of the Pleuronectidae (Munroe 2015), and our

phylogenetic results are highly congruent with this hypothesis.

It is possible that the outliers, the Atlantic and Pacific halibut (*H. hippoglossus* and *H.*

stenolepis, respectively), diverged during one of the first openings of the Bering Strait.

The estimated dates from the molecular clock analysis are approximately 5.5 MYA, in

accordance with the hypothesized dates for the first aquatic connection (Marincovich and

Gladenkov 2001). The remaining sister taxa have a younger age estimate of ~1-2 MYA,

corresponding with global cooling during the Pleistocene and repeated glaciations

(Zachos et al. 2001), which likely formed more barriers to genetic flow. In the second

pair of outliers, *P. hawaiiensis* inhabits waters of the Eastern Central Pacific to the

Hawaiian Islands, while *P. natalensis* inhabits coastal waters of Eastern Africa (Fig. 2).

Due to the far reaching range of *P. natalensis*, and the relatively younger age estimates

for divergence (1-2 MYA), these results beg for future research into other vicariant

hypotheses, or dispersal routes, as these two members of *Poecilopsetta* diverged long

after the Isthmus had closed. One candidate would be the Indo-Pacific barrier, which

formed an almost continuous land bridge between Australia and Asia, and could have

hence limited dispersal between the Pacific and the Indian oceans, hereby promoting

speciation (Gaither et al. 2011). This barrier formed during the Pleistocene glacial cycles

(2.58-0.01 MYA; Voris 2000), which is, again, fully consistent with our date estimates

(Fig. 3, S1).

Based on the most extensive multigene sequence alignment available to date across

all flatfish species, we have showed here that the evolutionary dynamics of sister species

that are distributed across the two oceans strongly supported the existence of two bouts of

speciation: one triggered by the closure of the CAS 12-3 MYA, and a second one due to

the closure of an ancient northern passage 5-0.01 MYA. Our work adds to a growing

body of evidence pointing not only to the role of geological processes in shaping biotic

turnovers (Bacon et al. 2015), but also on the consequences of climate change that can

either trigger speciation events (Reid 1990), or promote faunal interchanges whose

ecological and economic impacts are difficult to predict (Wisz et al. 2015).

Materials and Methods

Data retrieval and alignment

Prior to retrieving sequence data, GenBank was surveyed to identify all the genes

belonging to species of Pleuronectiform families (as of August 2016), based on

GenBank's taxonomy browser. DNA sequences for a total of 332 flatfish species (out of

over 800 species in the order) were identified and downloaded for five nuclear genes

(KIAA1239, MYH6, RIPK4, RAG1, SH3PX3), and four mitochondrial genes (12S, 16S,

COX1 and CYTB). These represented all the taxa having at least one of these six gene

sequences in GenBank; see Table S1 for the corresponding accession numbers. Diversity

was richly sampled as species from 13 of the 14 families in the order Pleuronectiformes

were included in our catchment. In line with current consensus in flatfish systematics, the

family Psettodidae was chosen as the outgroup to all other taxa (Chapleau 1993).

These sequences were aligned using MUSCLE ver. 3.8.31 (Edgar 2004) on a gene-

by-gene basis. Each alignment was visually inspected with AliView ver. 1.18 (Larsson

2014), and was manually edited where necessary. In particular, large indels (> 10 bp)

were removed prior to all phylogenetic analyses. The 5' and 3' ends of sequences were

also trimmed. The aligned sequences were then concatenated into a single alignment

using a custom R script.

Data pre-processing

To gauge the phylogenetic content of our data set, we performed a first series of

molecular clock analyses on the concatenated data matrix, with all the nine gene

sequences obtained above (12S, 16S, COI, Cyt-b, KIAA1239, MYH6, RIPK4, SH3PX3

and RAG1), and all the 332 taxa representing all families of flatfish. To account for rate

heterogeneity along this alignment, partitions were first identified with PartitionFinder 2

(Lanfear 2012), selecting the best-fit model of nucleotide substitution based on Akaike's

Information Criterion (AIC). This optimal partition scheme was then employed in a first

Bayesian phylogenetic analysis, conducted with BEAST ver. 1.8.3 (Drummond and

Rambaut 2007). This program implements a Markov chain Monte Carlo (MCMC)

sampler, which co-estimates both tree topology and divergence times. As determined by

PartionFinder, a GTR+I+Γ model was applied to each data partition. These models of

evolution across partitions were assumed to be independent (*unlinked*, in BEAST

parlance), while both clock model and tree model partitions were shared (*linked*) by all

partitions. An uncorrelated relaxed clock was employed with a lognormal distribution

prior on rates, and a Yule speciation prior (Drummond et al. 2006). Due to the paucity of

reliably placed fossils on the flatfish tree, a unique calibration point was placed on the

most recent common ancestor (MRCA) of the ingroup, as a mean-one exponential prior,

with an offset of 40 million years reflecting the age of 47.8-42.1 MYA (Chanet 1997;

Friedman 2012). To stabilize the estimate, a lognormal ln(0, 1.5) prior with a 40 MYA

offset was also placed on the root of the tree (root height). Two separate MCMC samplers

were run, each for 10,000,000 generations. Trees were sampled every 5,000 generations,

and convergence was checked visually using Tracer ver. 1.6 (available at:

http://tree.bio.ed.ac.uk/software/ tracer/). Tree log files from each run were combined in

LogCombiner, after conservatively removing 10% of each run as burn-in. The resulting

maximum a posteriori (MAP) tree was generated with TreeAnnotator (Drummond and

Rambaut 2007).

As the topology of this resulting MAP tree was unconventional, we suspected the

presence of rogue taxa, *i.e.* species evolving either much faster or much slower than the

majority, which can contribute negatively to consensus tree support (Aberer et al. 2013).

Rogue taxa were identified using the RogueNaRok (Aberer et al. 2013) webserver

(http://rnr.h-its.org). The consensus trees from the preliminary analysis using 10,000

iterations were used as the tree set for rogue taxon analysis. A total of 22 rogues were

identified and pruned from the analysis, leaving 310 species.

Molecular Dating

To assess the impact of the closure of the CAS on flatfish evolutionary dynamics, a

second set of partitioned relaxed molecular clock analyses was performed (without the

rogue sequences). The timing of the closure of the CAS is estimated to have occurred

between 12 and 3 MYA (Duque-Caro 1990; Coates et al. 1992; Haug and Tiedemann

1998; O'Dea et al. 2016), and we used this time window as a prior to inform the relaxed

molecular clock-based phylogenetic reconstructions. The analyses were performed on the

same concatenated data set, with the same single fossil calibration, but we also placed a

lognormal prior ln(3,1.5), that has most of its mass on the 12-3 MYA window, on the

MRCA of each pair of sister taxa (the 'ALL' model). From the initial BEAST analyses,

twelve pairs of taxa were selected based on two criteria: (i) being sister species on that

initial tree, (ii) with one species distributed in the Pacific and one in the Atlantic ocean

(Fig. 2). Again, two independent MCMC samplers were run each for 100 million

iterations, with samples taken every 5000 step.

Because these pairs of sister species show a contrasted geographic distribution,

having either a southern (equatorial) or a northern range (Fig. 2), two additional sets of

analyses were performed. In a first set, calibration priors (ln(3,1.5), as *per* above) were

placed only sister taxa that had a geographic range in the southern hemisphere (the

'SOUTH' model), while in a second set, identical calibration priors were placed only on

sister species with a northern range (the 'NORTH' model). Finally, a set of analyses was

performed using no sister taxa calibrations at all (the 'NONE' model). For each analysis,

results from the two MCMC runs were combined using LogCombiner after removing an

even more conservative burn-in of 50%. The final MAP tree was generated with

TreeAnnotator.

In an attempt to rank these different models (priors on all sister taxa; only on southern

taxa; only on southern taxa; no "CAS" priors), the AICM was computed for each model

(Baele et al. 2013). These computations were performed in Tracer for each of the four

different calibration models, based on 100 replicates.

317   *Species distribution data*

In order to map the distribution of each species of flatfish, we resorted to the Global

Biodiversity Information Facility (GBIF: https://www.gbif.org/) database, accessed with

the R library rgbif (Chamberlain 2017). Up to 1,000 records were retrieved for each

species, and plotted on a geographic map based on GBIF observational records. Where

needed, species locations were summarized by mean latitude and longitude (Fig. S4).

Approximate attribution to a particular ocean was based on species-specific mean

longitudes (Atlantic: between 20˚ and 120˚; Indian: 120˚ and -100˚; Pacific otherwise).

325   *Availability of Computer Code and Data*

326   Accession numbers, sequence alignments, BEAST models (as xml files), estimated

327   phylogenetic trees (maximum *a posteriori* trees from BEAST), and the R scripts used to

328   plot the figures in this study are available from https://github.com/sarisbro.

Acknowledgments

This work was supported by the University of Ottawa (LB), and the Natural Sciences
Research Council of Canada (FC, SAB). We are grateful to Jonathan Dench for
discussions and comments, and to Compute Canada and Ontario's Center for Advanced
Computing for providing us access to their resource.

References

Aberer AJ, Krompass D, Stamatakis A. 2013. Pruning rogue taxa improves phylogenetic
accuracy: An efficient algorithm and webservice. *Syst Biol*. 62:162–166.

Aris-Brosou S, Rodrigue N. 2012. The essentials of computational molecular evolution.
in Evolutionary Genomics: statistical and computational methods. Anisimova M,
editor. Methods in Molecular Biology (Methods and Protocols), vol 855. Totowa
(NJ): Humana Press.

Azevedo MFC, Oliveira C, Pardo BG, Martínez P, Foresti F. 2008. Phylogenetic analysis
of the order Pleuronectiformes (Teleostei) based on sequences of 12S and 16S
mitochondrial genes. *Genet Mol* Biol. 31:284–292.

Bacon CD, Silvestro D, Jaramillo C, Smith BT, Chakrabarty P, Antonelli A. 2015.
Biological evidence supports an early and complex emergence of the Isthmus of
Panama. *Proc Natl Acad Sci U S A*. 19112:6110–6115.

Baele G, Li WLS, Drummond AJ, Suchard MA, Lemey P. 2013. Accurate model
selection of relaxed molecular clocks in Bayesian phylogenetics. *Mol Biol Evol*.
30:239–243.

Berendzen PB, Dimmick WW, McEachran JD. 2002. Phylogenetic relationships of
Pleuronectiformes based on molecular evidence. *Copeia* 2002:642–652.

Berta A. 2012. Return to the Sea: The life and evolutionary times of marine mammals.
Berkeley (CA): University of California Press.

Betancur-R R, Ortí G. 2014. Molecular evidence for the monophyly of flatfishes
(Carangimorpharia: Pleuronectiformes). *Mol Phylogenet Evol*. 73: 18–22.

Betancur-R R, Li C, Munroe TA, Ballesteros JA, Ortí G. 2013. Addressing gene tree
discordance and non-stationarity to resolve a multi-locus phylogeny of the flatfishes

359 (Teleostei: Pleuronectiformes). *Syst Biol*. 62:763–785.

360 Campbell J, López JA, Satoh TP, Chen WJ, Miya M. 2014a. Mitochondrial genomic

361   investigation of flatfish monophyly. *Gene* 551:176-182.

362 Campbell MA, Chen WJ, López JA. 2014b. Molecular data do not provide unambiguous

363   support for the monophyly of flatfishes (Pleuronectiformes): a reply to Betancur-R

364   and Ortí. *Mol Phylogenet Evol*. 75:149–53.

365 Chamberlain S. 2017. rgbif: Interface to the Global 'Biodiversity' Information Facility

366   API. R package version 0.9.9. https://CRAN.R-project.org/package=rgbif

367 Chanet B. 1997. A cladistic reappraisal of the fossil flatfishes record: consequences on

368   the phylogeny of the Pleuronectiformes (Osteichthyes: Teleostei). *Ann des Sci Nat*

369   *Zool Paris*. 18:105–117.

370 Chapleau F. 1993. Pleuronectiform relationships : a cladistic reassessment. *Bull Mar Sci*.

371   52:516–540.

372 Coates AG, Jackson JBC, Collins LS, Cronin TM, Dowsett J, Bybell LM, Jung P,

373   Obando JA. 1992. Closure of the Isthmus of Panama : The near-shore marine record

374   of Costa Rica and western Panama. *Geol Soc Am Bull*. 104:814–828.

375 Cooper JA, Chapleau F. 1998. Monophyly and intrarelationships of the family

376   Pleuronectidae (Pleuronectiformes), with a revised classification. *Fish Bull*. 96:686–

377   726.

378 De Schepper S, Schreck M, Beck KM, Matthiessen J, Fahl K, Mangerud G. 2015. Early

379   Pliocene onset of modern Nordic Seas circulation related to ocean gateway changes.

380   *Nat Commun*. 6:8659.

381 Dodson JJ, Tremblay S, Colombani F, Carscadden JE, Lecomte F. 2007. Trans-Arctic

382   dispersals and the evolution of a circumpolar marine fish species complex, the

383   capelin (*Mallotus villosus*). *Mol Ecol*. 16:5030–5043.

384 dos Reis M, Donoghue PCJ, Yang Z. 2015. Bayesian molecular clock dating of species

385   divergences in the genomics era. *Nat Rev Genet*. 17:71–80.

386 Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. 2006. Relaxed phylogenetics and

387   dating with confidence. *PLoS Biol*. 4:699–710.

388 Drummond AJ, Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by sampling

389   trees. *BMC Evol Biol*. 7:214.

Duque-Caro H. 1990. Neogene stratigraphy, paleoceanography and paleobiology in

northwest South America and the evolu- tion of the Panama Seaway. Palaeogeogr.

Palaeoclimatol. *Palaeoecol*. 77:203–234.

Durham JW, MacNeil FS. 1967. Cenozoic migrations of marine invertebrates through the

Bering Strait region. In: Hopkins DM, editor. The Bering land bridge. Stanford (CA),

Stanford University Press. p. 326–349.

Edgar R. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res*. 32(5):1792–1797.

Friedman M. 2012. Osteology of †*Heteronectes chaneti* (Acanthomorpha,

Pleuronectiformes), an Eocene stem flatfish, with a discussion of flatfish sister-

group relationships. *J Vert Paleontol.* 32:735-756.

Gaither MR, Bowen BW, Bordenave TR, Rocha LA, Newman SJ, Gomez JA, van

Herwerden L, Craig MT. 2011. Phylogeography of the reef fish *Cephalopholis*

argus (Epinephelidae) indicates Pleistocene isolation across the indo-pacific barrier

with contemporary overlap in the coral triangle. *BMC Evol Biol*. 11:189.

Gladenkov AY, Oleinik AE, Marincovich L, Barinov KB. 2002. A refined age for the

earliest opening of Bering Strait. *Palaeogeogr Palaeoclimatol Palaeoecol*. 183:321–

328.

Gladenkov AY. 2004. Onset of connections between the Pacific and Arctic Oceans

through the Bering Strait in the Neogene. *Stratigr Geol Correl*. 12:175–187.

Grant WS. 1986. Biochemical genetic divergence between Atlantic, *Clupea harengus*,

and Pacific, *C. pallasi*, Herring. *Copeia* 1986:714–719.

Harrington RC, Faircloth BC, Eytan RI, Smith WL, Near TJ, Alfaro ME, Friedman M.

2016. Phylogenomic analysis of carangimorph fishes reveals flatfish asymmetry

arose in a blink of the evolutionary eye. *BMC Evol Biol*. 16:224.

Haug GH, Tiedemann R. 1998. Effect of the formation of the Isthmus of Panama on

Atlantic Ocean thermohaline circulation. *Nature* 393:673–676.

Larsson A. 2014. AliView: A fast and lightweight alignment viewer and editor for large

datasets. *Bioinformatics* 30:3276–3278.

Liu JX, Tatarenkov A, Beacham TD, Gorbachev V, Wildes S, Avise JC. 2011. Effects of

Pleistocene climatic fluctuations on the phylogeographic and demographic histories

of Pacific herring (*Clupea pallasii*). *Mol Ecol*. 20:3879–3893.

Marincovich L, Gladenkov AY. 2001. New evidence for the age of Bering Strait. *Quat*

Sci Rev. 20:329–335.

Marincovich L. 2000. Central American paleogeography controlled Pliocene Arctic

Ocean molluscan migrations. *Geology* 28:551–554.

McCusker MR, Bentzen P. 2010. Phylogeography of 3 North Atlantic Wolffish species

(*Anarhichas* spp.) with Phylogenetic Relationships within the Family

Anarhichadidae. J Heredity. 101:591–601.

Munroe TA. 2015. Systematic Diversity of the Flatfishes. In: Gibson RN, Nash RDM,

Geffen AJ, van der Veer HW, editor. Flatfishes: biology and exploitation.

Chichester, West Sussex: John Wiley & Sons. p. 13–44.

Norman JR. 1934. A systematic monograph of the flatfishes (Heterosomata). London:

Printed by order of the British Museum.

O'Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-Moreno SA, Cione AL, Collins

LS, de Queiroz A, Farris DW, Norris RD, et al. 2016. Formation of the Isthmus of

Panama. *Sci Adv*. 2:e1600883–e1600883.

Pardo BG, Machordom A, Foresti F, Porto-Foresti F, Azevedo MFC, Bañon R, Sánchez

L, Martínez P. 2005. Phylogenetic analysis of flatfish (Order Pleuronectiformes)

based on mitochondrial 16S rDNA sequences. *Sci Mar*. 69:531–543.

Parham JF, Donoghue PCJ, Bell CJ, Calway TD, Head JJ, Holroyd PA, Inoue JG, Irmis

RB, Joyce WG, Ksepka DT, et al. 2012. Best practices for justifying fossil

calibrations. *Syst Biol*. 61:346–359.

Raftery AE, Newton MA, Satagopan JM, Krivitsky PN. 2007. Estimating the integrated

likelihood via posterior simulation using the Harmonic Mean Identity. *Bayesian Stat.*

8:1–45.

Rawson PD, Harper FM. 2009. Colonization of the northwest Atlantic by the blue mussel,

Mytilus trossulus postdates the last glacial maximum. *Mar Biol*. 156:1857–1868.

Reid DG. 1990. Trans-arctic migration and speciation induced by climatic change: the

biogeography of *Littorina* (Mollusca: Gastropoda). *Bull Mar Sci*. 47:35–49.

Taylor EB, Dodson JJ. 1994. A molecular analysis biogeography within (genus

Osmevus) of relationships and a species complex of Holarctic fish. *Mol Ecol.* 3:235–

452  248.

453 Väinölä R. 2003. Repeated trans-Arctic invasions in littoral bivalves: Molecular

454  zoogeography of the *Macoma balthica* complex. *Mar Biol*. 143:935–946.

455 Vermeij GJ. 1991. Anatomy of an invasion: the trans-Arctic interchange. *Paleobiology*

456  17:281–307.

457 Voris HK. 2000. Maps of Pleistocene sea levels in Southeast Asia: Shorelines, river

458  systems and time durations. *J Biogeogr*. 27: 1153-1167.

459 Wisz MS, Broennimann O, Grønkjær P, Møller PR, Olsen SM, Swingedouw D,

460  Hedeholm RB, Nielsen EE, Guisan A, Pellissier L. 2015. Arctic warming will

461  promote Atlantic–Pacific fish interchange. *Nat Climate Change*. 5:261–265.

462 Zachos J, Pagani M, Sloan L, Thomas E, Billups K. 2001. Trends, rhythms, and

463  aberrations in global climate 65 Ma to present. *Science* 292:686–693.

464 Zachos JC, Dickens GR, Zeebe RE. 2008. An early Cenozoic perspective on greenhouse

465  warming and carbon-cycle dynamics. *Nature* 451:279–283.

466 Zaslavskaya NI, Sergievsky SO, Tatarenkov AN. 1992. Allozyme similarity of Atlantic

467  and Pacific species of *Littorina* (Gastropoda: Littorinidae). J Moll Stud. 58:377–384.

470   **Table 1**. AICM values for the phylogenetic analyses using four different calibration

471   schemes. Models are ranked from the best AICM model (top) to the worst model. SE:

472   standard error.

Calibration Model	AICM value	SE
ALL	374,082.199	+/-1.579
NONE	374,110.488	+/-3.789
SOUTH	374,367.749	+/-0.696
NORTH	374,746.628	+/-3.789

Figure captions

FIG. 1. Phylogenetic trees reconstructed based on relaxed clock models. Four models were employed, representing different specifications of prior distributions set on sister taxa. (**A**) No priors were set on sister taxa. (**B**) Priors were set on all pairs of taxa. (**C**) Priors were set only on sister taxa with a current northern range. (**D**) Priors were set only on sister taxa with a current southern range. Horizontal scale is in million years ago (MYA). The closure of the CAS (12-3 MYA) is shown between vertical gray broken lines.

FIG. 2. Distribution of the geminate species used in this study. Geographic distribution of the twelve pairs of sister species of flatfish used to calibrate the relaxed molecular clock models. Original data come from GBIF (www.gbif.org; accessed Nov 9, 2017). The top panel shows the phylogenetic distribution of these species based on the clock model including prior ages on all twelve pairs of species; scale bar: 5 MYA.

FIG. 3. Posterior densities of divergence times for sister taxa used as calibration points in the relaxed molecular clock analyses, under four different calibration schemes. NONE: no priors were placed on sister taxa; ALL: priors were placed on all pairs of sister taxa; NORTH: priors were placed only on pairs of sister taxa with a northern distribution; SOUTH: priors were placed only on pairs of sister taxa with a southern distribution. Top row (**A-D**) shows all 17 distributions, while bottom row (**E-H**) shows range-averaged distributions (solid) to the exception of outlier pairs (dashed lines). Densities are color-coded for species with northern (blue) and southern (red) range. Dashed vertical gray lines indicate the closure of the CAS (21-3 MYA).

9:L. aspera   8:L.proboscidea

9:L. limanda

10:H. platessoides
12:G. cynoglossus
11:H. hippoglossus

11:H. stenolepis

7:P. quadrituberculatus

7:P. platessa

1:H. stomata

1:H. oblonga
8:L. ferruginea
3:C. arctifrons
5:S. micrurum
6:C. chittendeni
4:C. minutus

2:P. hawaiiensis

5:S. maculiferum

3:C. platophrys

6:C. querna

4:C. darwini

2:P. natalensis

Northern

Southern

12:*Glyptocephalus stelleri/Glyptocephalus cynoglossus*
11:*Hippoglossus stenolepis/Hippoglossus hippoglossus*
10:*Hippoglossoides dubius/Hippoglossoides platessoides*
9:*Limanda aspera/Limanda limanda*
8:*Limanda ferruginea/Limanda proboscidea*
7:*Pleuronectes platessa/Pleuronectes quadrituberculatus*
6:*Cyclopsetta chittendeni/Cyclopsetta querna*
5:*Syacium maculiferum/Syacium micrurum*
4:*Citharichthys darwini/Citharichthys minutus*
3:*Citharichthys arctifrons/Citharichthys platophrys*
2:*Poecilopsetta hawaiiensis/Poecilopsetta natalensis*
1:*Hippoglossina oblonga/Hipoglossina stomata*