1   **Nonlinear averaging of thermal experience predicts population growth rates in a thermally**
2   **variable environment**
3
4
5   Joey R. Bernhardt[1*], Jennifer M. Sunday[1,2], Patrick L. Thompson[1], and Mary I. O'Connor[1]
6
7

8   [1]Department of Zoology
9   Biodiversity Research Centre
10 University of British Columbia
11 Vancouver, BC, V6T 1Z4, Canada
12
13 [2] School of Marine and Environmental Affairs
14 University of Washington
15 Box 355685
16 Seattle, WA 98195-5685
17
18 ***Corresponding author:** joey.bernhardt@biodiversity.ubc.ca

19
20 **Short running title:** Population growth under thermal variability
21

22 **Keywords:** thermal variability, population growth, phytoplankton, Jensen's inequality, scale
23 transition theory

24
25 **Article type:** Letter
26 **Abstract:** 149 words
27 **Main text:** 5013 words
28 **Number of references:** 30
29 **Number of figures:** 3
30
31 **Author contributions:** JRB conceived and designed the study, along with help from JMS, PLT
32 and MIO. JRB carried out the experiments, analyzed the data and wrote the first draft of the
33 manuscript. All authors contributed to writing the manuscript. All authors gave final approval for
34 publication.
35
36 **Data accessibility:** All data will be made available on Dryad should the manuscript be accepted.
37
Competing interests: We have no competing interests.

Abstract

Incorporating biological effects of environmental variability into predictions of species'

performance is a persistent challenge that limits our ability to project across temporal scales in

ecology. Here we test the hypothesis that population growth rates in fluctuating thermal

environments can be predicted quantitatively from the curvature of thermal performance curves

(TPCs) using scale transition theory. Using experimental populations of the green alga

Tetraselmis tetrahele grown over a range of temperatures in constant and fluctuating thermal

regimes, we show that population growth rates in fluctuating thermal regimes can be accurately

and quantitatively predicted based on performance in constant thermal regimes. We extrapolate

from these results to predict how population growth and critical temperatures for persistence of

89 phytoplankton species assessed in constant conditions would differ in naturally variable

thermal environments. Together these results demonstrate a key improvement upon our ability to

predict population dynamics in the context of global change.

Introduction

Organisms live in variable environments and their performance reflects their responses to

environmental variation over time and space. Demographic rates and outcomes that integrate

temporal environmental variation may differ substantially from what might be predicted based

on short-term physiological responses to constant, non-varying environments. This difference

complicates projections of population performance based on physiological assays under constant

conditions (Estay *et al.* 2014), prompting calls for ecologists to explicitly incorporate

environmental variation into predictions and models of population and species' performance in

the field (Dowd *et al.* 2015). Because environmental variability differs across regions and the

lifespans of organisms, an approach that allows quantitative scaling from thermal reaction norms

of population growth generated under constant laboratory conditions to population performance

in a variable thermal regime may be particularly useful for understanding patterns of abundance

and distribution, and species' responses to climate change.

Biological responses to environmental variation depend on the shape of the relationship between

performance and an environmental gradient (Ruel & Ayres 1999; Drake 2005; Lawson *et al.*

2015). When performance, *P*, changes nonlinearly with environmental conditions, mean

performance in a variable environment $\overline{P(E)}$ does not necessarily equal performance at the mean

environmental condition $P(\bar{E})$. This fact, captured by the well-known mathematical rule Jensen's

inequality, leads to clear predictions about how environmental variability should affect

performance over time. In the context of temperature, the relationship between organismal or

population performance and temperature, called a thermal performance curve (TPC) (Fig. 1A), is

almost always nonlinear (Schulte 2015) so the often implicitly assumed linear relationship

between environment and population growth in demographic models of environmental

variability is inadequate to describe population dynamics over temperature gradients (Lande *et*

al. 2003). The potential ecological and evolutionary effects of Jensen's inequality have been

shown in several recent studies (Martin & Huey 2008; Estay *et al.* 2014; Vasseur *et al.* 2014;

Denny 2017). Yet, ecologists struggle to incorporate thermal variability when making

predictions about the effects of temperature on growth, abundance, and distributions of species in

nature, often assuming that species' thermal experiences are well represented by the mean

temperature of their environment.

Principles of non-linear averaging have been formalized in scale transition theory (Chesson *et al.*

2005), which builds on Jensen's inequality, and makes predictions for how non-linear dynamics

change over spatial and temporal scales. Scale transition theory is an approach developed for

metacommunity models that allows for quantitative predictions for how variation at one scale

emerges at another when mediated by a nonlinear relationship such as a TPC. While Jensen's

inequality provides qualitative predictions about how the shape of non-linear TPCs will influence

the outcome of performance in the face of environmental variation, scale transition theory allows

for quantitative predictions of these effects. Here, we use scale transition theory in the context of

relating estimated performance over time-averaged scales to performance at short time scales

generated under static conditions. Specifically, the expected performance under variable

conditions, $\overline{P(T)}$, can be approximated by

$$\overline{P(T)} \approx P(\bar{T}) + \frac{1}{2}P''(\bar{T})\sigma^2{}_T \tag{1}$$

where $P''(\bar{T})$ is the second derivative of the TPC at an average temperature, $\bar{T}$ and σ_T is the

standard deviation of temperature experienced by the organism (Chesson *et al.* 2005; Estay *et al.*

2014; Dowd *et al.* 2015). Equation 1 holds whenever $P(T)$ can be approximated accurately by its

second order Taylor expansion, and is likely to be most accurate when σ_T is small relative to the

thermal breadth. Equation 1 shows that where the relationship between performance, *P,* and

temperature is an accelerating function with a positive second derivative (Fig. 1B), *P* will

increase with increasing temporal variance (σ^2) in temperature due to Jensen's inequality.

Similarly, *P* will decrease with increasing variance where the TPC is decelerating (i.e. negative

second derivative, Fig. 1B). Where the TPC is linear, temperature variability should have no

effect on P because the effects at cold temperatures counterbalance the effects of warm

temperatures when averaged over time. By quantifying the curvature at a given point on a

thermal performance curve, scale transition theory provides a way to mathematically predict the

strength and direction of effects of temperature fluctuations around that temperature on

performance.

The typical shape of TPCs (Fig. 1A) (Dell *et al.* 2011), with a decelerating phase at higher

temperatures within a thermal niche and an accelerating phase at lower temperatures, suggests

negative effects of thermal variation at high temperatures and benefits at low temperatures.

Current estimates of the consequences of thermal variability for population-level performance

such as growth rate have assumed a certain shape to the curve (i.e. a Gaussian rise and a

parabolic fall; Vasseur *et al.* 2014) , thus forcing certain outcomes of variability. Yet, new

evidence suggests that the shape and skew of the performance curve can vary substantially

among species, phenotypes, or contexts (Thomas *et al.* 2012, 2017), and this is likely to have

important implications for the effect of environmental variation on performance. Empirical

evidence for how variability affects population growth rates to date has been based on tests at

only two temperatures (Bozinovic *et al.* 2011). This restricted sampling of the TPC has

precluded testing of quantitative predictions based on the curvature of the TPC.

Here we asked: Does population growth in a thermally variable environment reflect the effects of

non-linear averaging over the TPC? To answer this question, we experimentally tested whether

population growth in thermally variable environments can be predicted by scale transition

theory. We experimentally quantified a TPC for population growth of a green alga, *Tetraselmis*

tetrahele, in a constant thermal regime, and then used this TPC and Equation 1 to predict how

population growth differs in a fluctuating temperature regime. Following from Equation 1, we

predicted that increased temperature variability over the range of cold temperatures to the left of

the inflection point of the TPC (i.e. in the accelerating portion of the curve, where the second

derivative of the TPC is positive; Fig. 1B) would lead to higher growth rates relative to constant

conditions, and that temperature variability in the range of warm temperatures to the right of the

inflection point would be detrimental (i.e. where the second derivative of the TPC is negative;

Fig. 2A, dashed curve).

Then, drawing on a global dataset of empirical TPCs for phytoplankton population growth rates,

which vary in shape (skew and position) as well as geographic origin, we use scale transition

theory (Equation 1) to estimate *in situ* population growth rates, given levels of *in situ*

environmental variation at each species' isolation location. We estimate the extent to which

predicted growth rates at each isolation location differ when they are predicted by scale transition

theory, as compared to when they are predicted based on mean temperatures only. By including a

range of phytoplankton TPC shapes from a global distribution, we explore the consequences of

considering thermal variability in projections of population growth rates, with implications for

patterns of abundance and distribution.

Methods

Experimental quantification of TPCs in constant and fluctuating environments

We experimentally quantified TPCs in constant and varying thermal environments for

Tetraselmis tetrahele, a globally distributed coastal marine phytoplankton species. The cultured

strain used here was obtained from the Canadian Centre for the Culture of Microorganisms

(UW414), and was originally isolated off the coast of Vancouver Island, British Columbia,

Canada. The *T. tetrahele* was maintained in laboratory culture in ESAW medium (Enriched

Seawater, Artificial Water, Harrison *et al.* 1980) at 16°C on a 16:8 light:dark cycle under nutrient

and light saturated conditions before the start of the experiments.

We initiated twenty replicate experimental populations of *T. tetrahele* in 30 mL glass test tubes

containing 20 mL of 10uM nitrate ESAW medium at a density of ~ 700 cells/mL under constant

temperature conditions at 0°C, 5°C, 10°C, 15°C, 20°C, 24°C, 27°C, 29°C, and 32°C, and under

fluctuating temperature conditions with the same mean temperatures as the constant conditions,

but fluctuating ± 5°C; i.e. 0°C – 10°C, 5°C – 15°C, 10°C – 20°C, 15°C – 25°C, 19°C – 29°C and

22°C – 32°C. We created fluctuating temperature treatments by programming temperature-

controlled incubators (Panasonic MR 154) to switch between the low and high temperature once

per day (i.e. approximately 11.5 hours at the high temperature and approximately 11.5 hours at

the low temperature, with 30 minutes of transition time in between each temperature cycle). To

avoid experimental artifacts associated with confounding the temperature cycles with daily light

cycles, we grew all experimental populations under 24-hour continuous light, at saturating light

intensities of 150 umol/m^2/s (see Fig. S1 in Supporting Information). The source population was

acclimated to 24-hour continuous light for four months prior to the experiment. We sampled

each of the twenty replicate populations, with four replicates sampled destructively at each of

five time points over the period corresponding to the exponential growth phase at each

175 temperature. The intervals between sampling periods depended on the temperature, such that for

176 the warmer temperatures (>15°C) sampling time points were condensed (up to 3 samples per

177 day) to capture exponential growth, while at the colder temperatures (<=15°C) sampling

178 intervals were spread out over longer time periods (i.e. one to five days between sampling points

179 at 0°C). Population abundances and cell biovolumes (using area-by-diameter estimation, ABD)

180 were measured from 250 uL samples using a FlowCAM (flow rate = 0.3 ml/min; FlowCAM VS

181 Series, Fluid Imaging Technologies).

183 *Statistical estimation of performance curves*

184 We estimated the intrinsic rate of population growth, *r*, during the exponential growth phase as:

186 
$$N(t) = N(0)e^{rt}$$
(2),

188 where *N(t)* is the number of individuals at time *t,* using non-linear least squares regression with

189 the *nls* function in R (R Core Team 2017).

191 For constant and variable conditions, we fit a thermal performance curve for growth rate

192 (Thomas *et al.* 2012),

193 
$$r(T) = ae^{bT}\left[1 - \left(\frac{T-z}{\frac{w}{2}}\right)^2\right]$$
(3)

195 where growth rate, *r,* is a function of temperature, *T*, *a* and *b* are parameters from the Eppley

196 curve (Eppley 1972) that describe maximum observed population growth rates as a function of

temperature, z determines the location of the maximum of the quadratic portion of the function

and w is the range over which the growth rate is positive (i.e. the thermal breadth). We estimated

these parameter values using maximum likelihood estimation with the *mle2* function in the

bblme package (Bolker 2017). Importantly, this TPC equation does not force the curve to take on

any particular shape, such that estimated curves can be either fully decelerating throughout the

entire range of temperatures over which growth is positive, or can have any combination of

accelerating and decelerating portions. We favored this TPC equation over others because of this

flexibility, because it is parameterized with biologically meaningful parameters, and because has

been used previously in studies of phytoplankton TPCs (Thomas *et al.* 2012, 2016). For

comparison, we fitted several other functional forms to the experimental dataset (e.g. a Gaussian

x Gompertz function, as in Martin & Huey 2008) but did not find any better fits.

To generate estimates of uncertainty in critical temperatures of the TPC (e.g. T_{opt}), we

determined confidence intervals around fitted thermal performance curves using parametric

bootstrapping, following the approach outlined in Equation S.3 in (Thomas *et al.* 2012). We then

used the TPC fitted under constant conditions and its upper and lower confidence intervals to

make predictions under variable conditions. For the thermal performance curve fitted to the nine

experimental temperatures under constant conditions, we simulated nine new performance values

drawn from a normal distribution such that the mean of the distribution corresponded to the

value of Equation 3 at each of the original experiment temperatures, given the estimated

parameters and the standard deviation $= \sigma$, where σ was calculated by accounting for uncertainty

in the estimation of the original maximum likelihood standard deviation, $\hat{\sigma}$, using:

$$\hat{\sigma}^2 = \sigma^2 \sqrt{\frac{n-1}{X}} \tag{4}$$

where n is the number of data points ($n = 9$), and X is a random number drawn from the chi-

squared distribution having ($n - 1$) degrees of freedom (Gelman & Hill 2007). We then fit

Equation 3 to these nine simulated performance estimates using maximum likelihood estimation

and repeated this process 10000 times to generate 10000 new thermal performance curves. We

calculated 95% confidence intervals as the range between the 2.5[th] and 97.5[th] quantiles. We used

these 95% confidence intervals to calculate the upper and lower bound of the predicted

population growth rate under fluctuating thermal conditions using Equation 1 (Fig. 2A, dashed

band). We repeated the same curve fitting and bootstrapping process for the fluctuating

temperature regimes, where the TPC (Equation 3) was fit to the mean temperature in each

treatment.

We quantified four critical temperatures of the TPC: the optimal temperature for population

growth, T_{opt}, the minimum temperature for positive population growth, $T_{r>0,min}$, the maximum

temperature for positive population growth, $T_{r>0,max}$, and thermal niche breadth under constant

conditions, w. We quantified the analogs of these critical temperatures under fluctuating

conditions and refer to them as the minimum mean and maximum mean temperatures for

positive population growth under fluctuating conditions, $\bar{T}_{r>0,min}$ and $\bar{T}_{r>0,max}$ respectively, the

mean temperature for optimal growth under fluctuating conditions, $\bar{T}_{opt}$, and thermal niche

breadth, $\bar{w}$. If the $T_{r>0,min}$ from the estimated curve was below -1.8°C, we set $T_{r>0,min}$ to be -

1.8°C because we assumed that *Tetraselmis tetrahele* cannot maintain positive population

growth below the freezing point of seawater. We then calculated the range of temperatures over

which population growth rate is positive as the difference between $T_{r>0,min}$ and

$T_{r>0,max}$. Following Equation 1, we expected that increased temperature variability should

increase and decrease population growth in the accelerating and decelerating portions of the TPC

respectively, and that the shift between positive and negative effects of temperature variability

should occur at the inflection of the constant-temperature TPC. We predicted that increased

variability should push the lower and upper limits of the TPC under variable conditions ($\bar{T}_{r>0,min}$

and $\bar{T}_{r>0,max}$) to lower temperatures than $T_{r>0,min}$ and $T_{r>0,max}$. We expected the maximum

growth rate, r_{max}, to decrease under variable temperature conditions relative to constant

conditions because T_{opt} is always in a decelerating portion of the constant TPC curve. Finally,

we expected the thermal breadth under fluctuating conditions, $\bar{w}$, to also decrease under

fluctuating conditions if $T_{r>0,min}$ is close to freezing, thus preventing $\bar{T}_{r>0,min}$ from shifting to

lower temperatures to compensate for decreased $\bar{T}_{r>0,max}$.

Applying scale transition theory to estimate in situ *phytoplankton population growth rates*

We used scale transition theory (Equation 1) to estimate population growth rates in thermally

variable environments for a set of diverse phytoplankton species. We estimated TPCs for 89

species based on published phytoplankton growth rates (Thomas *et al.* 2016) measured in the lab

at arrays of constant temperatures using the fitting approach described in the lab experiment

above. We restricted our analysis of the global phytoplankton dataset to marine species that met

the following conditions: 1) there were experimental population growth rate estimates from at

least five temperatures (n = 89 phytoplankton strains), 2) estimated $T_{r>0,max}$ and $T_{r>0,min}$ were

within 5°C of the highest and lowest measurement temperatures respectively, 3) growth rates at

two or more temperatures above and below T_{opt} were reported, and 5) fitted curves had an R^2 of

at least 0.85, to ensure the estimated parameters describing the shape of the curve accurately

captured the curvature of the TPCs. In addition to these curve shape criteria, all the curves in the

dataset met the quality control criteria outlined in Thomas *et al.* 2016 (namely that the

experimental populations were grown with adequate light and nutrient supply, and non-stressful

pH and salinities).

For each of these 89 species, we used historical reconstructed sea surface temperature data to

characterize thermal regimes at isolation locations reported in the original studies. We estimated

thermal regimes (mean and variance) for each phytoplankton isolation location (latitude and

longitude) reported in Thomas *et al.* 2016. For each species' isolation location, we extracted

monthly sea surface temperatures (SST) from NOAA's Extended Reconstructed Sea Surface

Temperature dataset, ERSSTv3 (Smith *et al.* 2008). This dataset has a spatial resolution of 0.5-

degree (50 km). We calculated the mean and standard deviation of monthly sea surface

temperatures from 1986 – 2005. We used monthly temperature data because these data reflect

seasonal variation that is central to understanding patterns of long-term population persistence.

Statistical estimation of TPCs

We produced two TPCs for each phytoplankton species – one generated assuming temperatures

remain constant through time (akin to constant lab conditions), and one that incorporates natural

patterns of thermal variability from their habitat. For the 'constant' scenario, we fitted Equation 3

to each species' population growth rate dataset using the methods described above, and estimated

a growth rate, r, for the species at the isolation location using Equation 3, where T = mean SST at

the isolation location. For the 'variable' scenario, we estimated growth rates using the mean and

the variance of SST at each isolation location, following Equation 1, using $P(\bar{T})$ equal to the

value of Equation 3 fitted to the experimental data, and σ_T as the standard deviation of SST. For

each of these two scenarios, we compared three attributes of TPCs: T_{opt}, $T_{r>0,min}$ and $T_{r>0,max}$.

Given that differences in TPCs based on constant vs variable thermal environments depend on

curve shape and temperature variance (Equation 1), we also explored how the discrepancies in

estimated critical temperatures and population growth rates at average *in situ* temperatures

depend on the shape of the TPC observed in constant lab conditions. We examined the effects of

curve attributes including the skew, using a curve skewness metric developed by Thomas et al.

(2016) (Supplementary Equation 5 in Thomas *et al.* 2016), which standardizes the absolute

skewness of the curve by the niche width, *w*, using OLS regression. All analyses were conducted

in R (version 3.4.1, R Core Team 2017).

Results

1) Does population growth in a thermally variable environment reflect the effects of non-linear

averaging over the TPC?

Population growth in fluctuating conditions differed from that in constant thermal conditions

over the thermal gradient, and the differences were predicted quantitatively by scale transition

theory (95% CI of the growth rate estimates under fluctuating conditions overlapped with

predicted growth rates from Equation 1; points and dashed band in Fig. 2B). Consistent with

expectations based on nonlinear averaging of temporal variation in temperature, experimental

populations of *T. tetrahele* had higher population growth rates under fluctuating temperature

conditions compared to constant conditions over accelerating portions of the TPC (i.e. at low

mean temperatures; 5°C and 10°C), but lower growth rates under fluctuating temperature

conditions compared to constant conditions over decelerating portions of the TPC (i.e. at mean

temperatures above 10°C, including 15°C, 24°C and 27°C and 29°C) (Fig. 2B). Notably,

population growth was lower under fluctuating conditions relative to constant conditions at

24°C, which is close to T_{opt} in this population of *T. tetrahele* (Fig. 2B). Populations had negative

growth rates at 32°C. The shift between positive effects of temperature fluctuations on

population growth at low temperatures and negative effects of fluctuations at warmer

temperatures aligned with the inflection point of the constant-temperature TPC (16.96 °C, 95%

CI: 16.71°C, 17.12°C), providing strong empirical support for scale transition theory in

predicting population growth in thermally variable environments.

Thermal variation altered estimated parameter values and key features of the thermal

performance curve. The maximum exponential growth rate ($\bar{r}_{max}$) was lower under variable

conditions than constant conditions, $\bar{r}_{max}$ =1.20 day^{-1} (95% CI: 1.07 day^{-1}, 1.33 day^{-1}) vs r_{max} =

1.62 day^{-1} (95% CI: 1.52 day^{-1}, 1.71 day^{-1}) (Fig. 2A, B). Estimated mean optimal temperatures

for growth rate were lower under variable conditions: $\bar{T}_{opt}$ = 21.40°C (95% CI 20.37°C,

22.23°C) vs. T_{opt} = 24.56°C (95% CI: 24.27°C, 24.86°C). Estimated $\bar{r}_{max}$ and $\bar{T}_{opt}$ were

quantitatively consistent with theoretical predictions from Equation 1 (i.e. had 95% CI

overlapping the predicted values from Equation 1; Fig. 2B). Maximum mean temperatures for

positive growth rates were lower under variable conditions $\bar{T}_{r>0,max}$ = 29.96°C (95% CI:

28.59°C, 32.61°C) vs $T_{r>0,max}$ = 31.91°C (95% CI: 31.73°C, 32.11°C). The range of

temperatures associated with positive growth rates, accounting for the freezing point of seawater,

i.e. $T_{r>0,max} - T_{r>0,min}$ was 33.67°C (95% CI: 33.09°C, 33.91°C) under constant conditions

and 31.52°C (95% CI: 30.09°C, 33.54°C) under variable conditions. The estimated thermal

breadth, w, was also lower under variable conditions: $\bar{w}$ = 36.68°C, 95% CI: 13.87°C, 66.29°C,

vs w = 36.51°C, 95% CI: 32.87°C, 42.51°C). Maximum mean temperatures for positive growth,

the range of temperatures associated with positive growth rates and w were all lower under

variable conditions than constant conditions, but were not statistically distinguishable (i.e. had

overlapping 95% CI).

2) How different are predicted TPCs in variable natural environments from predictions based on

TPCs generated under constant conditions?

When we estimated the TPCs of the 89 phytoplankton species for constant and varying

temperature regimes, we found that for the 90% of species that show negative skew, $\bar{T}_{opt}$ in

variable environments is lower than T_{opt} in constant environments (Fig. 3C), while for the

remaining 10% of species which show a positive skew, thermal variability is expected to

increase $\bar{T}_{opt}$ relative to T_{opt}. The magnitude of the difference between $\bar{T}_{opt}$ and T_{opt} increased

with increasing standard deviation of sea surface temperature, and is well explained by curve

skew (positive slope = 62.92, 95% CI: 51.72, 74.12) and standard deviation of sea surface

temperature (negative slope = -0.22, 95% CI: -0.28, -0.15) (Adjusted R^2 = 0.61, $F_{2,106}$ = 85.85, p

< 0.001) (Fig. 3C). Phytoplankton growth rate estimates that include the effects of thermal

variability, $\bar{r}$, differ from those that do not account for *in situ* thermal variability, r, (Fig. 3D).

Generally, predicted growth rates under variable conditions are lower than growth rates under

constant conditions (i.e. $\bar{r} - r < 0$, data points below the line y = 0 in Fig. 3D), however for

some species living in regions with thermal regimes typically colder than the species' T_{opt},

growth rates in these environments can exceed those in constant conditions ($\bar{r} > r$); an example

of such a species and its TPC is shown in Fig. 3F. Importantly, the differences between r and $\bar{r}$

are greatest for species whose isolation locations have mean temperatures that are close to their

T_{opt}. Predicted upper thermal limits are almost always lower under variable conditions

($\bar{T}_{r>0,max} < T_{r>0,max}$) (Fig. 3E), and the difference between $\bar{T}_{r>0,max}$ and $T_{r>0,max}$ increases

with increasing skewness (positive slope = 38.03, 95% CI: 30.13, 45.92) and standard deviation

of SST (negative slope = -0.42, 95% CI: -0.46, -0.37,) (Adjusted $R^2 = 0.79$, $F_{2,106} = 207.8$, $p <$

0.001). Thermal breadth, $\bar{w}$, is consistently estimated to be lower under variable conditions, and

the difference between w and $\bar{w}$ increases with SST standard deviation, but not curve skewness

(Adjusted $R^2 = 0.50$, $F_{2,106} = 55.53$, $p < 0.001$).

Discussion

Thermal performance curves are widely used to predict patterns of species abundance and

distribution (Deutsch *et al.* 2008; Sunday *et al.* 2012; Thomas *et al.* 2012). And yet, these curves

are often used with the assumption that performance under constant lab conditions is analogous

to performance under naturally variable climate regimes. We have shown experimentally that

realized TPCs in variable environments differ from those in constant environments as predicted

qualitatively by Jensen's inequality (Ruel & Ayres 1999; Denny 2017) and quantitatively match

predictions based on scale transition theory (Chesson *et al.* 2005). Not only do variable vs.

constant condition estimated population growth rates differ for any given mean temperature, but

so do the key parameters and critical temperatures of the TPC. This result aligns with previous

theoretical predictions (Estay *et al.* 2014; Dowd *et al.* 2015) that simply extrapolating critical

temperatures from constant condition lab experiments to naturally variable environments is

likely to introduce substantial error into projections of population performance, often over-

predicting species performance in the field. In the application of scale transition theory to 89

species' TPCs and known *in situ* thermal variability, we find that estimated critical temperatures

in realistically varying environments deviate substantially from observed critical temperatures

under constant conditions. Failing to account for variability can result in greatly over- or

underestimating population performance in natural populations. This has important implications

for how we use TPCs to make predictions about species distributions under current and future

climates.

Consistent with expectations based on Jensen's inequality and scale transition theory, our

experiment showed that temperature fluctuations increased population growth rate relative to

constant conditions over the accelerating portions of the TPC, decreased growth rates over the

decelerating portions of the curve, and the shift between positive and negative effects of

fluctuations matched the inflection point of the constant-temperature TPC (Fig. 1, Fig. 2).

Fluctuating thermal conditions changed several aspects of the thermal performance curve,

including the mean temperature of optimal population growth ($\bar{T}_{opt}$) and the maximum growth

rate attainable ($\bar{r}_{max}$) – effectively shifting the TPC toward lower temperatures and lower

population growth rates overall. However, this does not necessarily mean that species'

population growth performances are altogether lower in variable environments: for populations

living at mean temperatures in the accelerating part of the TPC, our results suggest that they will

benefit from environmental variation. Indeed, the *T. tetrahele* isolate used here was collected at a

location where temperatures are far colder than its $\bar{T}_{opt}$, in the accelerating portion of the

negatively skewed TPC (at mean temperature = 6.92°C, Smith *et al.* 2008).

When we applied scale transition theory to estimate the growth rates of globally distributed

phytoplankton species, we found that the effect of variability on predicted phytoplankton thermal

performance depended strongly on the shape and skew of the TPC and the degree of thermal

variability in the oceans from which the phytoplankton originated. Previous approaches have, in

the absence of more complete datasets, assumed a certain shape to the curve (i.e. a Gaussian rise

and a parabolic fall; Vasseur *et al.* 2014), thus forcing certain outcomes of variability. Here we

used a model that does not prescribe any particular shape (i.e. allows for fully decelerating

curves, or curves with accelerating portions, or any combination of decelerating and accelerating

portions), thus enabling a more complete exploration of the potential effects of temperature

variability on population performance. Importantly, whether the TPC is positively- or negatively-

skewed determined the direction of the shift between T_{opt} and $\bar{T}_{opt}$. The majority of the curves in

the dataset were negatively skewed, and in these cases variability shifted $\bar{T}_{opt}$ to colder

temperatures. Negatively skewed TPCs are widely observed across ectothermic taxa (Huey &

Kingsolver 1989; Dell *et al.* 2011), suggesting that the direction of the effects of thermal

variability observed in our experiment may be general across many ectothermic taxa and

ecosystems. Because the shape and skew of the TPC determine performance in variable

environments, the mechanisms that determine the shape of the thermal performance curve can

have an important influence on the outcome of thermal variability on population persistence.

More studies of the diversity of TPC shapes among species will elucidate the extent to which

environmental variability increases or decreases performance optima relative to constant lab

conditions.

Environmental variation may not always be detrimental for population growth rates, in contrast

to what is commonly assumed (Lande *et al.* 2003), and may play an important role in setting

species range edges, where populations live close to the margins of their thermal limits.

Temperature fluctuations may enhance the probability of population persistence when

environmental temperatures fall within the accelerating portion of the TPC. In contrast,

temperature fluctuations will decrease performance at the warm end of the TPC. Consistent with

the argument that 'suboptimal' is optimal (Martin & Huey 2008), we show both experimentally

(Fig. 2B) and theoretically (using empirical TPCs and *in situ* temperatures; Fig. 3D) that

population growth rates are often lower under variable thermal conditions relative to constant

ones, and this negative effect of temperature variation is greatest for populations living close to

their thermal optima.

Importantly, several recent advances in our understanding of species responses to global change

have relied on the shapes of TPCs generated under constant conditions to predict species range

shifts and other responses to climate change (e.g. Deutsch *et al.* 2008; Thomas *et al.* 2012). Here

we show that natural levels of environmental variability can introduce differences in critical

mean temperatures for population persistence of up to 3°C, which is on par with the magnitude

of predicted temperature changes over the next 100 years (Frölicher & Paynter 2015). Other

work has compared acute thermal physiological limits (e.g. CTmax, CTmin) to environmental

temperatures at range limits to assess relative sensitivities of range edges to warming (e.g.

Sunday *et al.* 2012), yet the underlying nonlinear thermal performance curve expected for these

ectotherms suggests that variability at warm range edges will have a stronger effect on

population persistence than variability at cold range edges. Specifically, our findings suggest

that approaches based on direct applications of lab-determined critical temperatures may under-

predict range edges at boundaries defined by cold temperatures, and over-predict range edges at

boundaries defined by warm temperatures.

While here we assumed a normal distribution of temperatures around the temporal average, this

is a simplification of real thermal regimes, which likely show more complex patterns variability

and temporal autocorrelation, which can further modify the effects of variability on populations

(Gonzalez & Holt 2002). Resource supply is also likely to covary with temperature, potentially

altering the outcomes of thermal variability on population growth rate. Importantly, organisms

may be able to acclimatize or adapt to fluctuating conditions over longer time scales (Angilletta

2009), with the potential to alter the shape and limits of the TPC, so the global predictions we

make here based on scale transition theory should be viewed as null models which do not

incorporate biological responses to environmental variability, and should be tested empirically

(Estay *et al.* 2014; Dowd *et al.* 2015). Nevertheless, even in the simplest scenario of

environmental variability, we predict significant changes in realized mean TPC critical

temperatures.

Understanding population responses to temperature now and into the future involves

understanding biological responses to changes in average temperatures in the context of

environmental variation. Omitting the effects of environmental variation from population and

species distribution models may thus limit our ability to predict species' responses to changes in

mean temperatures, even if variability patterns remain unchanged. We show that the effects of

environmental variation can be predicted based on the shape of the functional relationship

between population growth and the environment, adding another tool to the kit for forecasting

species' responses to the environment in a changing world.

Acknowledgements: We thank M. Thomas for sharing code used for thermal performance curve

fitting, and C. Harley and A. Gehman for helpful comments on previous versions of the

manuscript.

Funding: Funding was provided by Vanier Canada Graduate Scholarship (J.R.B), the

Biodiversity Research Centre (J.M.S), a Killam Postdoctoral Fellowship (P.L.T.), and the

Natural Sciences and Engineering Research Council (M.I.O).

References

Angilletta, M.J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis. Oxford*

University Press.

Bolker, B. (2017). bbmle: Tools for General Maximum Likelihood Estimation.

Bozinovic, F., Bastías, D.A., Boher, F., Clavijo-Baquet, S., Estay, S.A. & Angilletta, M.J.

(2011). The mean and variance of environmental temperature interact to determine

physiological tolerance and fitness. *Physiol. Biochem. Zool.*, 84, 543–52.

Chesson, P., Donahue, M.J., Melbourne, B. & Sears, A.L.W. (2005). Scale Transition Theory for

Understanding Mechanisms in Metacommunities. In: *Metacommunities: spatial dynamics*

and ecological communities. pp. 279–306.

Dell, A.I., Pawar, S. & Savage, V.M. (2011). Systematic variation in the temperature dependence

of physiological and ecological traits. *Proc. Natl. Acad. Sci.*, 108, 10591–10596.

Denny, M. (2017). The fallacy of the average: on the ubiquity, utility and continuing novelty of

Jensen's inequality. *J. Exp. Biol.*, 220, 139–146.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., *et al.*

(2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl.*

Acad. Sci., 105, 6668–6672.

Dowd, W.W., King, F.A. & Denny, M.W. (2015). Thermal variation, thermal extremes and the

physiological performance of individuals. *J. Exp. Biol.*, 218.

Drake, J.M. (2005). Population effects of increased climate variation. *Proc. R. Soc. London B*

Biol. Sci., 272.

Eppley, R.W. (1972). Temperature and phytoplankton growth in the sea. *Fish. Bull*, 70, 1063–

1085.

Estay, S.A., Lima, M. & Bozinovic, F. (2014). The role of temperature variability on insect

performance and population dynamics in a warming world. *Oikos*, 123, 131–140.

Frölicher, T.L. & Paynter, D.J. (2015). Extending the relationship between global warming and

cumulative carbon emissions to multi-millennial timescales. *Environ. Res. Lett.*, 10, 75002.

Gelman, A. & Hill, J. (2007). Data Analysis Using Regression and Multilevel. In: *Cambridge*

University Press. p. 625.

Gonzalez, A. & Holt, R.D. (2002). The inflationary effects of environmental fluctuations in

source-sink systems. *Proc. Natl. Acad. Sci. U. S. A.*, 99, 14872–7.

Harrison, P.J., Waters, R.E. & Taylor, F.J.R. (1980). A broad spectrum artificial sea water

medium for coastal and open ocean phytoplankton. *J. Phycol.*, 16, 28–35.

Huey, R.B. & Kingsolver, J.G. (1989). Evolution of thermal sensitivity of ectotherm

performance. *Trends Ecol. Evol.*, 4, 131–135.

Lande, R., Engen, S. & Saether, B.-E. (2003). *Stochastic Population Dynamics in Ecology and*

Conservation. Oxford University Press.

Lawson, C.R., Vindenes, Y., Bailey, L. & Pol, M. (2015). Environmental variation and

population responses to global change. *Ecol. Lett.*, 18, 724–736.

Martin, T.L. & Huey, R.B. (2008). Why 'suboptimal' is optimal: Jensen's inequality and

ectotherm thermal preferences. *Am. Nat.*, 171, E102-18.

R Core Team. (2017). R: A Language and Environment for Statistical Computing.

Ruel, J.J. & Ayres, M.P. (1999). Jensen's inequality predicts effects of environmental variation.

Trends Ecol. Evol.

Schulte, P.M. (2015). The effects of temperature on aerobic metabolism: towards a mechanistic

understanding of the responses of ectotherms to a changing environment. *J. Exp. Biol.*, 218,

1856–66.

Smith, T.M., Reynolds, R.W., Peterson, T.C., Lawrimore, J., Smith, T.M., Reynolds, R.W., *et al.*

(2008). Improvements to NOAA's Historical Merged Land–Ocean Surface Temperature

Analysis (1880–2006). *J. Clim.*, 21, 2283–2296.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2012). Thermal tolerance and the global redistribution

of animals. *Nat. Clim. Chang.*, 2, 686.

Thomas, M.K., Aranguren-Gassis, M., Kremer, C.T., Gould, M.R., Anderson, K., Klausmeier,

C.A., *et al.* (2017). Temperature–nutrient interactions exacerbate sensitivity to warming in

phytoplankton. *Glob. Chang. Biol.*, 23, 3269–3280.

Thomas, M.K., Kremer, C.T., Klausmeier, C.A. & Litchman, E. (2012). A global pattern of

thermal adaptation in marine phytoplankton. *Science (80).*, 338, 1085–1088.

Thomas, M.K., Kremer, C.T. & Litchman, E. (2016). Environment and evolutionary history

determine the global biogeography of phytoplankton temperature traits. *Glob. Ecol.*

Biogeogr., 25, 75–86.

Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S., *et al.*

(2014). Increased temperature variation poses a greater risk to species than climate

warming. *Proc. R. Soc. London B Biol. Sci.*, 281.

 Figures

Figure 1. A) A thermal performance curve and its key features (T_{min}, T_{max}, T_{opt}, thermal breadth,

w). This negatively-skewed curve shows an exponential increase typical of processes following

an Arrhenius function, with an accelerating region to the left of the inflection point (grey vertical

line), followed by a decelerating region to the right of the inflection point. Notice that the

accelerating region corresponds to the region with a positive second derivative (panel B). Figure

adapted from Dowd *et al.* 2015, but parameterized with *T. tetrahele* data from this study.

Figure 2. Thermal performance curves for *T. tetrahele* populations growing under constant and

variable temperature conditions. A) Exponential growth rates under constant temperature

conditions; blue line is fitted thermal performance curve and blue shading corresponds to 95% CI

generated from parametric bootstrapping. The dashed curve represents predicted growth rate

under thermally variable conditions ($\sigma_T = 5°C$) based on scale transition theory (Equation 1).

Points and error bars are observed mean growth rates and 95% CI. T_{opt} and r_{max} and their 95%

CI are shown in blue shaded boxes. B) Exponential growth rates under thermally fluctuating

conditions (±5°C); dashed curve is predicted based on panel A and Equation 1, green line is the

fitted thermal performance curve, green shading corresponds to 95% CI generated from

parametric bootstrapping. Green points and error bars are observed mean growth rates in the

fluctuating temperature regime and 95% CI. T_{opt} and r_{max} in constant conditions and their 95%

CI are shown in blue shaded boxes, $\bar{r}_{max}$ and $\bar{T}_{opt}$ in variable conditions and their 95% CI are

shown in the green shaded boxes. Predictions from Equation 1 and associated 95% CI are shown

in black dashed lines.

Figure 3. Curve skew and environmental variability predict differences between performance

under constant and variable conditions. A) Diagram showing predicted differences in key

features of a thermal performance curve under constant (black line) and variable (grey line)

environmental conditions. The distance labeled "C", which refers to the difference between

$\bar{T}_{opt}$ and T_{opt}, corresponds to the data plotted in panel C of this figure, while the distance labeled

"D", i.e. $\bar{r} - r$, refers to the predicted difference between growth rates calculated from a TPC

based on constant conditions vs. a TPC based on variable *in situ* conditions, plotted in panel D,

and the distance labeled "E" refers to the difference between $\bar{T}_{r>0,max}$ and $T_{r>0,max}$ and

corresponds to the data plotted in panel E of this figure. B) Map of all phytoplankton isolation

locations used in the analysis. The color of the ocean and the points corresponds to standard

deviation of monthly sea surface temperatures over the time period 1986-2005 (Smith *et al.*

2008). C) The difference between predicted increase $\bar{\bar{T}}_{opt}$ and T_{opt} generated under constant lab

conditions; color coding as in B. D) Predicted differences in phytoplankton growth rates that do

not incorporate *in situ* temperature variation (*r*) vs. predicted growth rates based on Equation 1.

Blue circles highlight two species shown for illustrative purposes in panels F and G; color coding

as in B. E) The difference between $\bar{\bar{T}}_{r>0,max}$ and $T_{r>0,max}$; color coding as in B. F) Example

TPC for *Thalassiosia pseudonana* which shows a negative skew. Mean monthly temperature at

the isolation location for this species is shown in the vertical grey dashed line, T_{opt} in the vertical

black line, and the monthly SD is shown in the grey shaded region. G) Example TPC for a

species with a positive skew TPC, *Detonula confervacea*; vertical lines and shading as in F.