Evolution of sexually-transferred steroids in *Anopheles* mosquitoes

Emilie Pondeville[1,2*], Nicolas Puchot[1,3], Michael Lang[4], Floriane Cherrier[1], Francis

Schaffner[5,6], Chantal Dauphin-Villemant[7], Emmanuel Bischoff[1,8], Catherine Bourgouin[1,3*]

Affiliations

[1]CNRS, URA 3012, Institut Pasteur, 75724 Paris, Cedex 15, France.

[2]MRC-University of Glasgow Centre for Virus Research, G61 1QH, Glasgow, Scotland.

[3]Functional Genetics of Infectious Diseases, Institut Pasteur, 75724 Paris, Cedex 15, France.

[4]Institut Jacques Monod, CNRS, UMR7592, Université Paris Diderot, Sorbonne Paris Cité,

Paris, France.

[5]Francis Schaffner Consultancy, Riehen, Switzerland.

[6]Swiss National Centre for Vector Entomology, Institute of Parasitology, Vetsuisse Faculty,

University of Zürich, Zürich, Switzerland.

[7]Université Pierre et Marie Curie-Paris 6, CNRS, 7 quai Saint Bernard, 75005 Paris, France.

[8]Institut Pasteur, Unit of Genetics and Genomics of Insect Vectors, 75724 Paris, Cedex 15,
France.
18

[*]Correspondence to: emilie.pondeville@glasgow.ac.uk; catherine.bourgouin@pasteur.fr

Abstract

Human malaria, which remains a major public health problem, is transmitted by a subset of

Anopheles mosquitoes belonging to only three out of eight subgenera: *Anopheles*, *Cellia* and

Nyssorhynchus. Unlike almost every other insect species, it was shown that males of some

Anopheles species produce and transfer steroid hormones to females during copulation and

that this transfer mediates reproductive changes. Steroids are consequently seen as a potential

target for malaria vector control. Here, we analysed the evolution of sexually-transferred

steroids and their effects on female reproductive traits across *Anopheles* by using a set of 16

mosquito species (5 *Anopheles*, 8 *Cellia*, and 3 *Nyssorhynchus*), including malaria vector and

non-vector species. We show that male steroid production and transfer are specific to the

Cellia subgenus and that there is no correlation between mating-induced effects in females

and sexually-transferred steroids. In the light of our results, male steroid production, transfer

and post-mating effects in females do not correlate with their ability to transmit human

malaria, which overturns the suggestion from previous studies and suggests that manipulation

of steroid-response pathways in the field should be considered with caution in order to benefit

malaria vector control strategies.

 Main text

Introduction

Anopheles mosquitoes are mostly known for their ability to transmit to mammals

malaria caused by *Plasmodium* parasites. Among 472 species named in this genus, 41 have

been classified as dominant vector species (DVS) of human malaria [1-3]. The *Anopheles*

genus is further subdivided into 8 subgenera of which 3 (*Anopheles*, *Cellia*, and

Nyssorhynchus) contains all known DVS of human malaria [3]. Despite a significant decrease

of malaria incidence in the last 15 years due to vector control and improvement of

chemoprevention, diagnostic testing and treatment, human malaria remains a world-wide

burden with more than 200 million cases reported and an estimated 445 000 deaths in 2016

[4]. With the increase of insecticide resistance in malaria mosquitoes [4,5], new vector control

strategies to limit and possibly eliminate malaria transmission are being developed such as

replacement of wild malaria-susceptible mosquito populations by resistant ones and/or

decrease of vector populations by manipulating mosquito reproduction [6-12]. Unlike

vertebrates, it was considered that insect adult males do not produce significant amounts of

steroid hormones until it was shown that males of *Anopheles gambiae*, the main vector of

human malaria in Africa, produce and transfer high quantities of 20-hydroxyecdysone (20E)

to females during copulation [13]. While ovarian steroid synthesis occurs in mosquito females

upon blood meal triggering egg development [13-18], sexual transfer of steroids by males

likely represents a nuptial gift that affects female reproduction in the malaria vector.

Consistent with this, sexually-transferred steroids were shown to induce refractoriness to

further copulation and to stimulate oviposition in *An. gambiae* females [19]. A study further

reported that mating-induced phenotypes in *Anopheles* females would only occur in species

whose males produce and transfer steroids to females during mating and that would be

specific to dominant human malaria vectors. Since steroids contribute to promote oogenesis and possibly favouring *Plasmodium* survival, the authors concluded that sexual transfer of steroids is likely to have shape anopheline ability to transmit malaria to humans [20]. Steroids became as such a promising target to manipulate mosquito female reproduction with the aim to reduce malaria vector populations specifically. Consequently, field use of 20E receptor agonists has been recently proposed to reduce malaria transmission [21]. In the present work, we investigated the evolutionary history of male steroid production using a large set of mosquito species belonging to the three *Anopheles* subgenera (*Anopheles*, *Cellia*, *Nyssorhynchus*) that contain all known DVS of human malaria. We also investigated the post-mating effects on female reproductive traits whether the cognate males produce or not steroids. The main conclusions of our work are that i) production of steroids in male mosquitoes is restricted to the *Cellia* subgenus, and ii) the effect of mating on female reproductive potential vary across species of the 3 *Anopheles* subgenera and this is not correlated to male steroid production.

Results

Male steroid production is specific to the *Cellia* subgenus

We first measured steroid titers in sexually mature virgin males from 19 different mosquito species. These mosquito species were selected within two *Culicidae* subfamilies, *Anophelinae* and *Culicinae*. Within the *Anophelinae* subfamily, 16 species distributed all over the world were chosen to cover 3 different subgenera (*i.e. Cellia*, *Nyssorhynchus* and *Anopheles*) of the *Anopheles* genus (Figure 1A). In the *Culicinae* subfamily, *Aedes aegypti*, *Aedes albopictus* and *Culex pipiens* that are vectors of arboviruses were also analysed. As shown on Figure 1B (right panel), male 20E production occurs only within the *Cellia* subgenus (*Anophelinae*) and is absent in the *Culicinae*. Interestingly, *Anopheles*

quadriannulatus whose males produce similar levels of steroids compared to *Anopheles*

stephensi is not a malaria vector unlike *An. stephensi* and other *Cellia* species investigated in

the present study. Indeed, its refractoriness or low susceptibility to the human malaria parasite

has been experimentally determined [22,23], confirming epidemiological data [24].

Conversely, production of 20E was undetected in male mosquitoes from the two other

Anopheles subgenera, *Anopheles* and *Nyssorhynchus* of which all tested members are

registered as DVS and/or experimentally shown to be highly susceptible to *Plasmodium*

falciparum [25,26].

To uncover the evolutionary history of male steroid production in mosquitoes, we

further consolidated the phylogenetic relationships of the species we analysed. To this effect,

we used DNA sequences of partial regions of the coding sequence of mitochondrial genes

(*COI*, *COII, ND5 and CYTB*) and nuclear genes (*g6pd* and *white* as well as ribosomal

subunits *18S* and *28S*) from the 19 mosquito species plus *Chagasia bathana* (*Anophelinae*

subfamily, *Chagasia* genera) as outgroup for *Anopheles* mosquitoes [27]. Phylogenetic

analysis of the data set was performed by Bayesian inference (Figure 1B left panel) and by

maximum likelihood (Supplementary Fig. 1). Phylogenetic relationships inferred from these

analyses resulted in different topologies mainly at the subgenus level and with varying node

support values. In both analyses, the members of each subgenus formed monophyletic groups

and the branching orders within the subgenus *Cellia* was identical. Our results are in

agreement with ones obtained by Sallum *et al.* [28] with the *Cellia* clade being the outgroup

of the subgenera *Anopheles and Nyssorhynchus* in the Bayesian approach while in the

maximum likelihood approach, *Nyssorhynchus* is the outgroup of *Anopheles* and *Cellia*, also

in agreement with recently published phylogenies [1,29,30]. The *Anopheles* subgenus has also

been found by others to be a basal lineage to *Cellia* and *Nyssorhynchus* [31]. Thus,

relationships between the different subgenera *Cellia*, *Nyssorhynchus* and *Anopheles*, are still

not entirely resolved. The difficulty in determining the relationships among these subgenera

might be due to limits and bias in gene and taxon sampling, but also to different methods used

[3,27]. This could also be due to the fact that the radiation of *Anopheles* and *Cellia* species

happened roughly at the same time [29,31].

From the Bayesian phylogenetic analysis and using fossil data, we obtained species

divergence estimates, which largely conform to recent phylogenies [29,31,32]

(Supplementary Table 1). The age of the last common ancestor of *Anopheles* genus is about

84.1 Ma (112.7-55.8, 95% confidence interval) and the most ancestral node within the *Cellia*

subgenus is dated to 69.2 Ma (93.0-45.6, 95% confidence interval) (Figure 1B,

Supplementary Table 1). As males from all *Cellia* species tested so far have the ability to

produce 20E, according to the parsimony law steroid production by mosquito males probably

originated once in the early *Cellia* lineage, at about 84.1-69.2 Ma. (112.7-45.6, 95%

confidence interval), *i.e.* during the late Cretaceous. Thus, steroid production by mosquito

males is most probably a shared derived character from the last common ancestor of *Cellia*

mosquitoes and represents as such a synapomorphy of this subgenus. *An. stephensi* males

transfer steroid upon mating to females (Supplementary Fig. 2) as do *An. gambiae* males [13]

and males of other *Cellia* species such as *Anopheles arabiensis* and *Anopheles dirus* [20].

This strongly suggests that transfer of steroids to females during mating is part of the "male

20E production" synapomorphy of *Cellia* mosquitoes. Our geographical mapping of nearly all

contemporary mosquito species belonging to the *Cellia* subgenus (224 species) against the

ones of *Anopheles* subgenus (184 species) (Figure 2) suggests that the common ancestor of

the *Cellia* subgenus diverged after separation of South America and Africa in agreement with

previous observations [3,33]. This biogeographic calibration is consistent with divergence

times obtained from our phylogenetic analysis placing the origin of steroid production by

males of the *Cellia* subgenus around 84.1-69.2 Ma (112.7-45.6, 95% highest posterior density

interval), *i.e.* after the separation of South America and Africa, which started at least 100 Ma

ago with no land bridge for about 80 to 50 Ma [34,35].

Mating-induced effects do not correlate with sexually-transferred steroids

To get a deeper understanding of the effect(s) of sexually-transferred steroids in *Cellia*

female mosquitoes, we investigated the influence of mating on two female reproductive traits

which are known to be regulated by ovarian 20E. On the one hand, steroid hormones trigger

ovarian follicle detachment from the germarium in *An. stephensi* and *Ae. aegypti* [14,36], and

ovarian steroids produced upon blood feeding stimulate vitellogenesis and egg development,

on the other hand [14,16-18,37]. We therefore analysed these two phenotypes in virgin and

mated females from 12 *Anopheles* species (8 *Cellia*, 3 *Anopheles* and 1 *Nyssorhynchus*). As

expected, mating induces the separation of the secondary ovarian follicle from the germarium

in *An. stephensi* (Figure 3, see also confocal pictures Supplementary Fig. 3), a *Cellia* species

whose males produce and transfer steroids during mating. Mating induces as well the

separation of the secondary follicle in *Anopheles minimus* and *Anopheles merus*, although to a

lesser extent. However, no secondary follicle detachment was observed after mating in *An.*

dirus nor detachment of the third follicle for the *Cellia* species whose secondary follicle is

already fully detached in virgin females (*Anopheles farauti*, *An. gambiae*, *An. arabiensis* and

An. quadriannulatus). Among the species whose males do not produce steroids, mating also

induces secondary follicle detachment in some species (*Anopheles atroparvus* and *Anopheles*

freeborni), but not in others, even in species showing partial detachment of the secondary

follicle in virgin females (*Anopheles quadrimaculatus* and *Anopheles albimanus*). Similarly,

mating increases the number of developed eggs in some mosquito species while not in others

and this is not correlated to male steroid production (Figure 4 and Table 1). As an example,

mating has an effect on egg development in *An. albimanus* (*Nyssorhynchus*), whose males do

not produce steroids, but not in *An. gambiae* nor in *An. arabiensis* (*Cellia*). Intriguingly, those

results are in discrepancy with previous studies reporting that mating increases egg

development in *An. gambiae* and *An. arabiensis* [20,38] but not in *An. albimanus* [20,39].

These differences across studies are not due to the origin of the blood used to feed females

(animal or human) as we obtained the same results with *An. albimanus* and *An. gambiae* fed

on mouse or human blood (Supplementary Fig. 4). It is likely that some variations can be

observed between strains of a single *Anopheles* species as already described for some *Aedes*

species [40,41]. It cannot be excluded that male steroids transferred upon mating benefit

reproduction of some *Cellia* species but only under certain environmental conditions as

different ecological pressures such as nutrition resources can favour or not the maintenance

and importance of nuptial gifts, as shown for *Ae. aegypti* [42-44].

Importantly, across the species investigated, mating induces detachment of the

secondary follicle from the germarium in some species of which a low proportion of virgin

females has the secondary follicle already detached as well as in some species of which a high

proportion of virgin females has a detached follicle. Similarly, the occurrence of mating-

induced effects on egg development is not linked to the ability of virgin females to develop a

low or high number of eggs after blood feeding. Moreover, among species within the *Cellia*

subgenus, the occurrence or absence of mating-induced phenotypes in females are not linked

to the different quantities of steroids produced by the cognate males (Figure 1 and Table 1)

and likely transferred to females during mating as determined for *An. gambiae*, *An. stephensi*,

An. arabiensis and An. dirus (Supplementary Fig. 2, [13,20]). Indeed, as depicted in Table 1,

mating triggers both secondary follicle detachment and a rise in the number of developed eggs

in *An. stephensi*, or only an increase in egg development in *An. dirus*, while this does not hold

for *An. gambiae* and *An. arabiensis*, two species whose males produce and transfer higher

amounts of steroids than *An. stephensi* and *An. dirus*. Overall, our analysis demonstrates that

post-mating responses increasing fecundity in females indeed exist in *Cellia* species, some of

which are likely mediated by male steroids. Importantly, they also occur in mosquito species

whose males do not produce and transfer steroids such as in species belonging to the

Anopheles and *Nyssorhynchus* subgenera (Table 1). Recently, male 20E has been shown to

induce refractoriness to further mating in *An. gambiae* mosquitoes [19]. Similarly to results

obtained for the two post-mating responses analysed in this study, female monoandry

(insemination by a single male) occurs in species whose males produce steroids but also in

ones whose males do not [45-48]. While few data are available for mosquitoes, it is well

known that post-mating responses are quite conserved among insects. However, the rapid

evolution of insect reproductive systems often results in species-specific genes and signalling

pathways that ultimately trigger similar post-mating changes in different insect species [49].

For instance, *Ae. aegypti* males transfer to females Juvenile Hormone (JH) and *Drosophila*

flies transfer male accessory gland peptides such as Sex Peptide upon copulation to trigger

physiological and behavioural changes in mated females [50,51]. Likewise, males from *Cellia*

species transfer steroid hormones while males from *Anopheles* and *Nyssorhynchus* species are

likely to transfer other and not yet identified molecule(s) to achieve similar effects. As JH is

transferred to female during mating in *Ae. aegypti* mosquitoes but also in the Lepidoptera

Heliothis virescens [52], sexual transfer of steroid hormones in other mosquito subgenera,

genera or even insect orders not yet tested cannot even be excluded.

Conclusions

A deep understanding of the selective forces driving reproductive strategy diversity

and their functional consequences are critical for designing strategies for management of

insect pests. It was previously suggested that post-mating responses in *Anopheles* mosquitoes

only exist in species whose males produce and transfer steroids to females and that would be

specific to dominant human malaria vectors [20]. Analysing the evolution of steroid

production by male mosquitoes from a larger set of mosquito species reveals that this

physiological trait is a synapomorphy of the *Cellia* subgenus. Importantly, there is no

correlation between the evolution of sexually-transferred 20E and malaria transmission to

humans (Table 1). Consistent with this, while we show that male steroid production and

subsequent transfer to females is likely to have evolved only once in the common ancestor of

Cellia species, phylogenetic analyses on malaria mosquitoes support a convergent evolution

with independent acquisitions of vectorial capacities in *Anopheles* mosquitoes [53-55].

Furthermore, we demonstrate that mating-induced phenotypes are variable among species and

possibly even among strains or under different environmental conditions. These differences

are independent of males ability to produce and transfer steroids to females and are not

correlated to malaria vectorial capacity. Apart from follicle detachment, increase in egg

development and induction of refractoriness to mating, numerous other functions of steroids

have been described in adult insects [19,20,56-61]. Thus, sexually-transferred steroids could

mediate different functions with more or less direct benefit for the reproduction of *Cellia*

females, due to the rapid evolution of reproductive systems between species. The recent

interest in sexually-transferred steroids in anopheline mosquitoes has led to propose targeting

20E pathways with 20E receptor agonists to manipulate mosquito female reproduction as a

mean to reduce vector populations and malaria transmission [21,62]. In the light of our

results, manipulating sexual-transfer of steroids in mosquitoes or using 20E receptor agonists

in the field to manipulate *Anopheles* mosquito population and malaria transmission should be

considered with caution in order to benefit malaria vector control strategies and also because

it risks to affect other untargeted non-malaria vector species.

It remains open as what were the evolutionary forces that have initially promoted the

acquisition and radiation of this presumably costly male steroid production and sexual gift to

females in *Cellia* mosquitoes around 84.1-69.2 Ma. At this time, two main paleogeological

events that had impact on environmental conditions may have led to pressures driving the

evolution of steroid production and transfer by males in *Cellia* species: i) the Gondwana

break up at 100 Ma with separation of South America and Africa [63]; ii) the Cretaceous-

Paleogene extinction event at 65.5 Ma [64,65]. Because such geographical isolation and

environmental stresses are believed to drive traits contributing to animal species survival, it is

likely that transfer of steroids by males to females have favoured *Cellia* species populations at

this critical time.

Methods

Mosquito species and rearing

Ecdysteroid production by sexually mature males was analysed in 19 mosquito species.

Sixteen (16) species belong to the subfamily *Anophelinae* (*An. arabiensis*, *An. dirus*, *An.*

farauti, *An. gambiae* form M, *An. merus*, *An. minimus*, *An. quadriannulatus, An. stephensi*,

An. albimanus, *An. albitarsis*, *An. aquasalis*, *An. atroparvus*, *An. freeborni*, *An. plumbeus, An.*

pseudopunctipennis and *An. quadrimaculatus*) and 3 species belong to the subfamily

Culicinae (*Ae. aegypti, Ae. albopictus* and *Cx pipiens*). *An. gambiae* form M, now called *An.*

coluzzii (N'Gousso strain), *An. stephensi* (Sda 500) were permanently reared at Institut

Pasteur (France). *An. albimanus* STECLA (MRA-126), *An. arabiensis* DONGOLA (males

generously prepared by T. Bukhari and MRA-856), *An. quadrimaculatus* ORLANDO (MRA-

139), *An. minimus* MINIMUS1 (MRA-729), *An. dirus* WRAIR2 (MRA-700), *An. farauti*

FAR1 (MRA-489), *An. freeborni* F1 (MRA-130), *An. atroparvus* EBRO (MRA-493), *An.*

quadriannulatus SANGWE (MRA-1155) and *An. merus* OPHANSI (MRA-803), were

obtained through BEI Resources, NIAID, NIH, and contributed by M.Q Benedict (MRA-

126,-856,-139,-729,-700,-489,-130,-493), C. Aranda (MRA-493), W. Takken (MRA-1155)

and R. Mahraj (MRA-803). *Ae. aegypti* (Liverpool strain) and *Ae. albopictus* (Ho Chi Min

Ville, Vietnam) were a generous gift from A-B. Failloux (Institut Pasteur, France). Eggs from

Cx. pipiens (anautogenous strain) were kindly provided by M. Weill (ISEM, Montpellier,

France). Mosquito larvae were reared at 27°C in deionized water supplemented with minerals

and fed on TetraMin Baby-E fish food from the day of hatching to the fourth larval instar

supplemented with pieces of cat food. Male and female adults were maintained at 27°C, under

68% relative humidity and a 12/12h light/dark cycle, and provided free access to a 10%

wt/vol sucrose solution for the first 5 days post-emergence (PE). Female mosquitoes (first

gonotrophic cycle) were allowed to feed for 30 min on the blood of an anesthetized mouse or

rabbit depending on mosquito species preference (see Supplementary Table 2). *An. plumbeus*

L4 larvae and pupae were harvested from a natural pond in Switzerland and reared up to the

adult stage at the Institut Pasteur. Mature males of *An. albitarsis* and *An. aquasalis* were

kindly prepared and provided by D. Valle and L. Moreira (Fiocruz, Brazil) and *An.*

pseudopunctipennis by F. Lardeux (IRD, Bolivia).

Mosquito sampling, ecdysteroid extraction and quantification

To measure ecysteroid titers in virgin adult males and females, males and females were

separated on the day of adult emergence, transferred individually 5 days later in methanol and

stored at -20°C until ecdysteroid extraction. For the transfer experiments, *An. stephensi* males

and females were separated on the day of adult emergence for 5 days to allow male sexual

maturation. Mating experiments and sampling were performed as described in [13]. Total

ecdysteroids from individual whole mosquitoes were extracted with methanol and re-

dissolved in enzyme immuno assay (EIA) buffer. Empty tubes were treated similarly in

parallel to be used as a negative control (referred as extraction blank). Ecdysteroids were

quantified by EIA, with 20-hydroxyecdysone-2-succinate coupled to peroxydase as a tracer

(dilution 1:100,000) and the L2 antiserum (gift from M. De Reggi (Marseille, France);

dilution 1:100,000). Calibration curves were generated with ecdysone (E; 3,6 - 500 pg/tube)

diluted in EIA buffer, and titers were expressed as E equivalents. Under these conditions,

detection limit is 2 pg E equivalents. All measurements were performed in duplicate and the

results are expressed as mean values ±SEM of several (n=20 at least) independent samples

and have been repeated on two independent cohorts of mosquitoes. Samples at or above the

highest value of the calibration curve were diluted and quantify again. The intra- and inter-

assay variation coefficients were 3,9% and 5,6%, respectively. For steroid titers in whole

males from different species, data were subjected to statistical analysis using Kruskall-Wallis

test for nonparametric data followed by Dunn's post-test (control group: extraction blank).

The indicated p values are those obtained with Dunn's test. For the transfer experiment in *An.*

stephensi, results were subjected to statistical analysis using Mann-Whitney test.

Taxon sampling and DNA sequencing for phylogenetic analysis

We chose 20 *Culicidae* species for phylogenetic and comparative analysis. We selected 16

species of the genus *Anopheles (Anophelinae* subfamily*)* of which *An. arabiensis*, *An. dirus*,

An. farauti, *An. gambiae*, *An. merus*, *An. minimus*, *An. stephensi*, *An. quadriannulatus*, *An.*

atroparvus, *An. freeborni*, *An. plumbeus*, *An. pseudopunctipennis*, *An. quadrimaculatus*, *An.*

albimanus, *An. albitarsis*, and *An. aquasalis*. As outgroups, we chose *Chagasia bathana*

(*Anophelinae* subfamily, *Chagasia* genus), and 3 mosquito species belonging to the subfamily

Culicinae with *Ae. aegypti*, *Ae. albopictus* (*Aedes* genus) and *Cx. pipiens* (*Culex* genus).

Sequence data were generated for *An. aquasalis*, *An. atroparvus*, *An. merus* and *An.*

plumbeus. Genomic DNA was obtained from single individuals using the DNeasy Blood and

tissue kit (QIAGEN). A common set of molecular markers were chosen based on the

availability of sequence data for *Ch. bathana*. Partial genomic regions of four nuclear genes

(*g6pd, white, 18S and 28S*) and four mitochondrial genes (*COI*, *COII*, *ND5* and *CYTB*) were

amplified by PCR with gene-specific or degenerate primers (sequences in Supplementary

Table 3). For PCR amplifications, we used 0.4 µM oligonucleotides, 1 U GoTaq® DNA

Polymerase (Promega) per 35 µl reaction volume, 2 mM $MgCl_2$, and 200 µM dNTP and

reactions were carried out using standard thermocycle conditions. PCR products were purified

and Sanger-sequenced with gene-specific primers or with T7, SP6 universal primers at

Cogenics (www.cogenics.com, Beckman Coulter, GenBank Accession Numbers in

Supplementary Table 4). Sequence data of the remaining species were obtained from

GenBank and VectorBase (Supplementary Table 4). Sequences were examined and aligned

with Geneious 6.1.3 (Biomatters). Our dataset was not complete, we did not find or generate

sequence data for 10 / 160 (8 genes X 20 species) gene specific sequences and among the rest

15 / 150 of sequences were only partially covering the locus. The missing sequence data

represent 5.6% of the total dataset and was annotated as a "?" (missing value) in the

alignments for phylogenetic analysis. Introns were removed from *g6pd* and *white* sequences

and the extremities of all protein coding sequences were trimmed to be in codon frame.

Alignments for protein coding genes were re-aligned with the Geneious translation alignment

program. In addition to gene specific alignments a concatenated dataset was generated in the

following order: *COI-COII-ND5-CYTB-18S-28S-g6pd-white*. The number of informative sites

was calculated using MEGA4 [66].

Phylogenetic analysis

We used DNA sequences of partial regions of the coding sequence of mitochondrial genes

(*COI*, *COII*, *ND5 and CYTB*) and nuclear ones (*18S*, *28S*, *g6pd* and *white*) from the 19

mosquito species plus *Chagasia bathana*. The combined dataset had 4398 positions, including

2602 constant positions, 1776 variable positions and 1356 parsimony informative positions.

Phylogenetic analysis of the concatenated, five-partition data set was performed by maximum

likelihood in PhyML [67] and by Bayesian inference in BEAST [68]. For all analyses, partition specific models of nucleotide substitution were selected using the Akaike Information Criterion as calculated in jModelTest 2.1.3 [69,70]. Maximum likelihood inference was done on the concatenated dataset in PhyML [67] using a GTR+I+G model of nucleotide substitution. Node support was determined by performing 100 bootstrap replicates. Bayesian phylogenetic analysis was performed with BEAST v1.7.5 [68] on the concatenated data set using five partitions with unlinked models of nucleotide substitution (Supplementary Table 5). The partitions corresponded to a single partition for mitochondrial sequences (*COI*, *COII, ND5* and *CYTB*) and four more partitions for the genes *18S*, *28S*, *g6pd*, and *white*. Mitochondrial genes were combined into one partition because they are closely linked in the mitochondrial genome and largely evolve as a single unit with little to no recombination [71]. We used a common strict clock model and a yule birth process as tree prior. While the Bayesian approach places the species of the subgenus *Nyssorhynchus* as sister group of the *Anopheles* subgenus species, they formed the outgroup of the *Anopheles* and *Cellia* lineages in the maximum likelihood approach. Overall, the Bayesian phylogeny revealed high posterior probabilities for each node (>0.9) while the maximum likelihood analysis lacked strong statistical support at the nodes that separate the lineages of the three *Anopheles* subgenera. For species divergence time estimates, we therefore used the Bayesian phylogenetic analysis. We assigned calibration fossil ages to set priors for most recent common ancestors (MRCA). We used *Cx. winchesteri* (33,9-55,8 Ma) [72] (see also http://mosquito-taxonomic-inventory.info/category/fossil-culicidae/fossil-culicidae) to approximate the age of the most recent node shared by *Cx. pipiens* – and *Ae. aegypti*/ *Ae. albopictus*; and *An. dominicanus* (33,9-40,4 Ma) [73,74] to estimate the age of the most recent node shared by *Anopheles* species. MRCA priors that incorporate fossil calibration dates were assumed to follow an exponential distribution, in the two above mentioned cases

with an offset of 33.9 Ma and a mean according to the mean ages of the fossils. Based on a recently published phylogeny of Kamali *et al.* [53], the root age of the *Culicidae* was set to 147 Ma and the prior was assumed to be normally distributed with a standard deviation of 20 Ma. The Markov-Chain Monte-Carlo (MCMC) run was performed with a chain length of 10^8 generations and was recorded every 1000 generations. Estimates were computed with Tracer version 1.5 (http://tree.bio.ed.ac.uk/software/tracer/) and MCMC output analysis was done using TreeAnnotator [68]. The first 2000 sampled trees were discarded as the burn-in. The phylogeny was visualized and annotated with Figtree version 1.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Cellia and *Anopheles* species distribution mapping

Total numbers of mosquito species belonging to the *Cellia* or *Anopheles* subgenera per country were taken from the Walter Reed Biosystematics Unit (WRBU, http://www.wrbu.org/). Data were further represented on maps created with R [75] using the "sp" package [76].

Analysis of the secondary follicle detachment in virgin and mated non-blood fed females

Males and females were separated upon emergence to keep females virgin. A portion of females were put in a cage with males to obtain mated females. After 7 days, female ovaries were dissected, and the spermatheca of mated females checked for presence of spermatozoa. Ovaries from each female were then mounted on a slide in mounting media and observation of the secondary follicle detachment was performed using a transmitted light microscope. 30 to 60 females were analysed per species and per condition. Data were subjected to Chi-square test. For *An. stephensi*, ovaries from either virgin or mated females were fixed with 4% paraformaldehyde, washed 3 times with PBS-Tween 20 0.05% (PBS-T) and stained with

DAPI. After 4 washes with PBS-T, ovaries were mounted on a slide in mounting media.

Pictures were taken using a SP5 Leica confocal microscope.

Analysis of egg development in virgin and mated blood fed females

Virgin and mated females were prepared as described above. A portion of females were put in

a cage with males to obtain mated females. On 7 days PE, females were allowed to blood feed

on anesthetized mouse or rabbit (Supplementary Table 2). *An. gambiae* and *An. albimanus*

females were also fed with fresh human blood (ICAReB Platform, Center for Translational

Research, Institut Pasteur, Paris, France) for comparison with mouse-fed mosquitoes. Ovaries

were dissected 48 hours after blood feeding, and the total number of eggs in virgin and mated

females was counted. Mated status was verified by observing spermatozoa in the spermatheca

and only females with a filled spermatheca were taken into account. 30 to 60 females were

dissected per species and per condition. Data were subjected to Mann-Whitney non-

parametric test.

References

1. Harbach RE, Kitching IJ (2016) The phylogeny of Anophelinae revisited: inferences about
406   the origin and classification of *Anopheles* (Diptera: Culicidae). Zoologica Scripta 45:
407   34-47.
408   2. Hay SI, Sinka ME, Okara RM, Kabaria CW, Mbithi PM, et al. (2010) Developing global
409   maps of the dominant *Anopheles* vectors of human malaria. PLoS Med 7: e1000209.
410   3. Harbach RE (2013) The Phylogeny and Classification of Anopheles. In: Manguin S, editor.
411   Anopheles mosquitoes - New insights into malaria vectors: InTech. pp. 3-55.
412   4. WHO (2017) World Malaria Report 2017 Geneva: World Health Organization. 160 p.
413   5. Ranson H, Lissenden N (2016) Insecticide Resistance in African *Anopheles* Mosquitoes: A
414   Worsening Situation that Needs Urgent Action to Maintain Malaria Control. Trends
415   Parasitol 32: 187-196.
416   6. Alphey L (2014) Genetic control of mosquitoes. Annu Rev Entomol 59: 205-224.
417   7. Gantz VM, Jasinskiene N, Tatarenkova O, Fazekas A, Macias VM, et al. (2015) Highly
418   efficient Cas9-mediated gene drive for population modification of the malaria vector
419   mosquito Anopheles stephensi. Proc Natl Acad Sci U S A 112: E6736-6743.
420   8. Hammond A, Galizi R, Kyrou K, Simoni A, Siniscalchi C, et al. (2016) A CRISPR-Cas9
421   gene drive system targeting female reproduction in the malaria mosquito vector
422   Anopheles gambiae. Nat Biotechnol 34: 78-83.

9. McLean KJ, Jacobs-Lorena M (2016) Genetic Control Of Malaria Mosquitoes. Trends
Parasitol 32: 174-176.
10. Macias VM, Ohm JR, Rasgon JL (2017) Gene Drive for Mosquito Control: Where Did It
Come from and Where Are We Headed? Int J Environ Res Public Health 14.
11. Beaghton A, Hammond A, Nolan T, Crisanti A, Godfray HC, et al. (2017) Requirements
for Driving Antipathogen Effector Genes into Populations of Disease Vectors by
Homing. Genetics 205: 1587-1596.
12. Lambert B, North A, Burt A, Godfray HCJ (2018) The use of driving endonuclease genes
to suppress mosquito vectors of malaria in temporally variable environments. Malar J
17: 154.
13. Pondeville E, Maria A, Jacques JC, Bourgouin C, Dauphin-Villemant C (2008) *Anopheles*
gambiae males produce and transfer the vitellogenic steroid hormone 20-
hydroxyecdysone to females during mating. Proc Natl Acad Sci U S A 105: 19631-
19636.
14. Redfern CPF (1982) 20-hydroxyecdysone and ovarian development in *Anopheles*
stephensi. J Insect Physiol 28: 97-109.
15. Hagedorn HH, O'Connor JD, Fuchs MS, Sage B, Schlaeger DA, et al. (1975) The ovary as
a source of alpha-ecdysone in an adult mosquito. Proc Natl Acad Sci U S A 72: 3255-
3259.
16. Lu YH, Hagedorn HH (1986) Egg development in the mosquito *Anopheles albimanus*. Int
J Invertebr Repr Dev 9: 79-94.
17. Raikhel AS, Brown MR, Belles X (2005) Hormonal control of reproductive processes. In:
Gilbert LI, Iatrou K, Gill SS, editors. Comprehensive Molecular Insect Science.
Elsevier, Oxford, UK. pp. 433-491.
18. Swevers L, Raikhel AS, Sappington TW, Shirk P, Iatrou K (2005) Vitellogenesis and
post-vitellogenic maturation of the insect ovarian follicle. In: Gilbert LI, Iatrou K, Gill
SS, editors. Comprehensive Molecular Insect Science. Elsevier, Oxford, UK. pp. 87-
155.
19. Gabrieli P, Kakani EG, Mitchell SN, Mameli E, Want EJ, et al. (2014) Sexual transfer of
the steroid hormone 20E induces the postmating switch in *Anopheles gambiae*. Proc
Natl Acad Sci U S A 111: 16353-16358.
20. Mitchell SN, Kakani EG, South A, Howell PI, Waterhouse RM, et al. (2015) Mosquito
biology. Evolution of sexual traits influencing vectorial capacity in anopheline
mosquitoes. Science 347: 985-988.
21. Childs LM, Cai FY, Kakani EG, Mitchell SN, Paton D, et al. (2016) Disrupting Mosquito
Reproduction and Parasite Development for Malaria Control. PLoS Pathog 12:
e1006060.
22. Takken W, Eling W, Hooghof J, Dekker T, Hunt R, et al. (1999) Susceptibility of
Anopheles quadriannulatus Theobald (Diptera: Culicidae) to *Plasmodium falciparum*.
Trans R Soc Trop Med Hyg 93: 578-580.
23. Habtewold T, Povelones M, Blagborough AM, Christophides GK (2008) Transmission
blocking immunity in the malaria non-vector mosquito *Anopheles quadriannulatus*
species A. PLoS Pathog 4: e1000070.
24. Sinka ME, Bangs MJ, Manguin S, Coetzee M, Mbogo CM, et al. (2010) The dominant
Anopheles vectors of human malaria in Africa, Europe and the Middle East:
occurrence data, distribution maps and bionomic precis. Parasit Vectors 3: 117.
25. Sinka ME, Bangs MJ, Manguin S, Rubio-Palis Y, Chareonviriyaphap T, et al. (2012) A
global map of dominant malaria vectors. Parasit Vectors 5: 69.

26. Schaffner F, Thiery I, Kaufmann C, Zettor A, Lengeler C, et al. (2012) *Anopheles*
plumbeus (Diptera: Culicidae) in Europe: a mere nuisance mosquito or potential
malaria vector? Malar J 11: 393.
27. Harbach RE (2007) The Culicidae (Diptera): a review of taxonomy, classification and
phylogeny. Zootaxa 1668: 591-638.
28. Sallum MAM, Schultz TR, Foster PG, Aronstein K, Wirtz RA, et al. (2002) Phylogeny of
Anophelinae (Diptera: Culicidae) based on nuclear ribosomal and mitochondrial DNA
sequences. Syst Entomol 27: 361-382.
29. Neafsey DE, Waterhouse RM, Abai MR, Aganezov SS, Alekseyev MA, et al. (2015)
Mosquito genomics. Highly evolvable malaria vectors: the genomes of 16 Anopheles
mosquitoes. Science 347: 1258522.
30. Wei Y, Cheng B, Zhu G, Shen D, Liang J, et al. (2017) Comparative physical genome
mapping of malaria vectors *Anopheles sinensis* and *Anopheles gambiae*. Malar J 16:
235.
31. Freitas LA, Russo CA, Voloch CM, Mutaquiha OC, Marques LP, et al. (2015)
Diversification of the Genus *Anopheles* and a Neotropical Clade from the Late
Cretaceous. PLoS One 10: e0134462.
32. Moreno M, Marinotti O, Krzywinski J, Tadei WP, James AA, et al. (2010) Complete
mtDNA genomes of *Anopheles darlingi* and an approach to anopheline divergence
time. Malar J 9: 127.
33. Krzywinski J, Grushko OG, Besansky NJ (2006) Analysis of the complete mitochondrial
DNA from *Anopheles funestus*: an improved dipteran mitochondrial genome
annotation and a temporal dimension of mosquito evolution. Mol Phylogenet Evol 39:
417-423.
34. Scotese CR (2004) A continental drift flipbook. J Geol 112: 729-741.
35. de Oliveira BF, Molina CE, Marroig G (2009) Paleogeography of the South Atlantic: a
Route for Primates and Rodents into the NewWorld? In: Garber PA, Bicca-Marques
JC, Strier KB, Estrada A, Heymann EW, editors. Developments in Primatology:
Progress and Prospects: Springer New York pp. 55-68.
36. Beckemeyer EF, Lea AO (1980) Induction of follicle separation in the mosquito by
physiological amounts of ecdysterone. Science 209: 819-821.
37. Clements AN (1992) The biology of mosquitoes. Chapman & Hall. pp. 509.
38. Baldini F, Gabrieli P, South A, Valim C, Mancini F, et al. (2013) The interaction between
a sexually transferred steroid hormone and a female protein regulates oogenesis in the
malaria mosquito Anopheles gambiae. PLoS Biol 11: e1001695.
39. Lounibos LP (1994) Variable egg development among *Anopheles* (*Nyssorhynchus*):
control by mating? Physiol Entomol 19: 51-57.
40. O'Meara GF, Evans DC (1977) Autogeny in saltmarsh mosquitoes induced by a substance
from the male accessory gland. Nature 267: 342-344.
41. Gillett JD (1956) Genetic differences affecting egg-laying in the mosquito *Aedes*
(Stegomyia) *aegypti* (Linnaeus). Ann Trop Med Parasitol 50: 362-374.
42. Gwynne DT, Simmons LW (1990) Nature 346: 172-174.
43. Gwynne DT (2008) Sexual conflict over nuptial gifts in insects. Annu Rev Entomol 53:
83-101.
44. Klowden MJ, Chambers GM (1991) Male accessory gland substances activate egg
development in nutritionally stressed *Aedes aegypti* mosquitoes. J Insect Physiol 37:
721-726.
45. Villarreal C, Fuentes-Maldonado G, Rodriguez MH, Yuval B (1994) Low rates of
multiple fertilization in parous *Anopheles albimanus*. J Am Mosq Control Assoc 10:
67-69.

46. Yuval B, Fritz GN (1994) Multiple mating in female mosquitoes — evidence from a field
population of *Anopheles freeborni* (Diptera: *Culicidae*). Bull Entomol Res 84: 137-
140.
47. Arnqvist G (1998) Comparative evidence for the evolution of genitalia by sexual
selection. Nature 393: 784-786.
48. Tripet F, Toure YT, Dolo G, Lanzaro GC (2003) Frequency of multiple inseminations in
field-collected *Anopheles gambiae* females revealed by DNA analysis of transferred
sperm. Am J Trop Med Hyg 68: 1-5.
49. Swanson WJ, Vacquier VD (2002) The rapid evolution of reproductive proteins. Nat Rev
Genet 3: 137-144.
50. Clifton ME, Noriega FG (2012) The fate of follicles after a blood meal is dependent on
previtellogenic nutrition and juvenile hormone in Aedes aegypti. J Insect Physiol 58:
1007-1019.
51. Chapman T, Davies SJ (2004) Functions and analysis of the seminal fluid proteins of male
Drosophila melanogaster fruit flies. Peptides 25: 1477-1490.
52. Park YI, Shu S, Ramaswamy SB, Srinivasan A (1998) Mating in *Heliothis virescens*:
transfer of juvenile hormone during copulation by male to female and stimulation of
biosynthesis of endogenous juvenile hormone. Arch Insect Biochem Physiol 38: 100-
107.
53. Kamali M, Marek PE, Peery A, Antonio-Nkondjio C, Ndo C, et al. (2014) Multigene
phylogenetics reveals temporal diversification of major African malaria vectors. PLoS
One 9: e93580.
54. Kamali M, Xia A, Tu Z, Sharakhov IV (2012) A new chromosomal phylogeny supports
the repeated origin of vectorial capacity in malaria mosquitoes of the *Anopheles*
gambiae complex. PLoS Pathog 8: e1002960.
55. Marshall JC, Powell JR, Caccone A (2005) Short report: Phylogenetic relationships of the
anthropophilic *Plasmodium falciparum* malaria vectors in Africa. Am J Trop Med
Hyg 73: 749-752.
56. Konig A, Yatsenko AS, Weiss M, Shcherbata HR (2011) Ecdysteroids affect *Drosophila*
ovarian stem cell niche formation and early germline differentiation. EMBO J 30:
1549-1562.
57. Domanitskaya E, Anllo L, Schupbach T (2014) Phantom, a cytochrome P450 enzyme
essential for ecdysone biosynthesis, plays a critical role in the control of border cell
migration in *Drosophila*. Dev Biol 386: 408-418.
58. Bai J, Uehara Y, Montell DJ (2000) Regulation of invasive cell behavior by taiman, a
Drosophila protein related to AIB1, a steroid receptor coactivator amplified in breast
cancer. Cell 103: 1047-1058.
59. Terashima J, Bownes M (2006) E75A and E75B have opposite effects on the
apoptosis/development choice of the *Drosophila* egg chamber. Cell Death Differ 13:
454-464.
60. Raikhel AS, Lea AO (1991) Control of follicular epithelium development and vitelline
envelope formation in the mosquito; role of juvenile hormone and 20-
hydroxyecdysone. Tissue Cell 23: 577-591.
61. Audit-Lamour C, Busson D (1981) Oogenesis defects in the *ecd-1* mutant of *Drosophila*
melanogaster, deficient in ecdysteroid at high temperatures. J Insect Physiol 27: 829-
837.
62. Mitchell SN, Catteruccia F (2017) Anopheline Reproductive Biology: Impacts on
Vectorial Capacity and Potential Avenues for Malaria Control. Cold Spring Harb
Perspect Med 7.

63. Reyment RA, Dingle RV (1987) Palaeogeography of Africa during the Cretaceous period.
Palaeogeogr Palaeoclimatol Palaeoecol 59: 93-116.
64. Schulte P, Alegret L, Arenillas I, Arz JA, Barton PJ, et al. (2010) The Chicxulub asteroid
impact and mass extinction at the Cretaceous-Paleogene boundary. Science 327: 1214-
1218.
65. Novacek MJ (1999) 100 Million year of land vertebrate evolution: the early Cretaceous-
early Tertiary transition. Ann Mo Bot Gard 86: 230-258.
66. Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics
Analysis (MEGA) software version 4.0. Mol Biol Evol 24: 1596-1599.
67. Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. Syst Biol 52: 696-704.
68. Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with
BEAUti and the BEAST 1.7. Mol Biol Evol 29: 1969-1973.
69. Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new
heuristics and parallel computing. Nat Methods 9: 772.
70. Posada D (2008) jModelTest: phylogenetic model averaging. Mol Biol Evol 25: 1253-
1256.
71. Ballard JW (2000) Comparative genomics of mitochondrial DNA in members of the
Drosophila melanogaster subgroup. J Mol Evol 51: 48-63.
72. Cockerell TDA (1919) The oldest mosquitoes. Nature 103: 44.
73. Zavortink TJ, Poinar JR GO (2000) *Anopheles* (*Nyssorhynchus*) *dominicanus* sp. n.
(Diptera: Culicidae) from Dominican Amber. Ann Entomol Soc Am 93: 1230–1235.
74. Borkent A, Grimaldi DA (2004) The Earliest Fossil Mosquito (Diptera: Culicidae), in
Mid-Cretaceous Burmese Amber. Ann Entomol Soc Am 97: 882–888.
75. RCoreTeam (2016) R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing.
76. Bivand S, Pebesma E, Gomez-Rubio V (2013) Applied Spatial Data Analysis with R. NY:
Springer. 405 p.
77. Wang BC, Park J, Watabe HA, Gao JJ, Xiangyu JG, et al. (2006) Molecular phylogeny of
the *Drosophila virilis* section (Diptera: Drosophilidae) based on mitochondrial and
nuclear sequences. Mol Phylogenet Evol 40: 484-500.
78. Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PD (2006) DNA barcodes
distinguish species of tropical Lepidoptera. Proc Natl Acad Sci U S A 103: 968-971.
79. Liu H, Beckenbach AT (1992) Evolution of the mitochondrial cytochrome oxidase II gene
among 10 orders of insects. Mol Phylogenet Evol 1: 41-52.
80. Krzywinski J, Wilkerson RC, Besansky NJ (2001) Evolution of mitochondrial and
ribosomal gene sequences in Anophelinae (Diptera: Culicidae): implications for
phylogeny reconstruction. Mol Phylogenet Evol 18: 479-487.
81. Lyman DF, Monteiro FA, Escalante AA, Cordon-Rosales C, Wesson DM, et al. (1999)
Mitochondrial DNA sequence variation among triatomine vectors of Chagas' disease.
Am J Trop Med Hyg 60: 377-386.
82. Zwiebel LJ, Saccone G, Zacharopoulou A, Besansky NJ, Favia G, et al. (1995) The *white*
gene of *Ceratitis capitata*: a phenotypic marker for germline transformation. Science
270: 2005-2008.
83. Bonacum J, DeSalle R, O'Grady P, Olivera D, Wintermute J, et al. (2001) New nuclear
and mitochondrial primers for systematics and comparative genomics in
Drosophilidae. Dros Inf Serv 84: 201-204.
617
618

Acknowledgments

We are grateful to L. Lambrechts, S. Blandin and J-P. Parvy for comments on the manuscript. We acknowledge D. Streicker, E. Hutchinson and M. Palmarini for discussions. We acknowledge M-T. Lecoq, S. Touron and C. Thouvenot for mosquito rearing, A-B. Failloux for providing *Ae. aegypti* and *Ae. albopictus* eggs, M. Weill (ISEM, Montpellier, France) for providing *Cx. pipiens* eggs, T. Bukhari for providing *An. arabiensis* samples, D. Valle and L. Moreira (FIOCRUZ, Brazil) for providing *An. aquasalis* and *An. albitarsis* samples, as well as F. Lardeux (IRD, Centre de Bolivie, La Paz, Bolivia) for providing *An. pseudopunctipenni*s samples. We thank E. Perret (Dynamic Imaging Platform, Institut Pasteur) for her assistance with confocal microscopy. The following reagents were obtained through BEI resources, NIAID, NIH: eggs of *An. albimanus* STECLA (MRA-126), *An. arabiensis* DONGOLA (MRA-856), *An. quadrimaculatus* ORLANDO (MRA-139), *An. quadriannulatus* SANGWE (MRA-1155), *An. merus* OPHANSI (MRA-803), *An. minimus* MINIMUS1 (MRA-729), *An. dirus* WRAIR2 (MRA-700), *An. farauti s.s.* FAR1 (MRA-489), *An. atroparvus* EBRO (MRA-493) and *An. freeborni* F1 (MRA-130).

Fundings

Support to E.P. was from an ANR-07-MIME-O25-01 award to C.B. and by the UK Medical Research Council to E.P. (MC_UU_12014/8), to C.B. from award no. ANR-10-LABX-62-IBEID. M.L. was hosted by V. Courtier-Orgogozo, financed by a FRM postdoctoral fellowship SPF20121226328 and by the CNRS.

Data availability

The sequence data reported in this paper are tabulated in the Supplementary information and archived at Genbank.

Contributions

E.P. and C.B. conceptualised and supervised the project as well as the experimental design; E.P., N.P., M.L., F.C., F.S., C.D-V. and E.B. performed the experiments; E.P., N.P., M.L., E.B. and C.B. contributed to data interpretation; E.P., N.P., M.L., F.S. and E.B. contribute to figure preparation; E.P. and C.B. wrote the manuscript; M.L., F.S., E.B. commented on the final manuscript.

Ethical compliance

This study complied with all relevant ethical regulations. Project (n° 2013-0132) approved by the Ministère de l'Enseignement Supérieur et de la Recherche – Direction Générale pour la Recherche et l'Innovation – Secrétariat « Autorisation de projet » - 1, rue Descartes, 75231 PARIS cedex 5.

657
658
659

Figures and Table

Figure 1. Distribution, phylogenetic relationships of mosquitoes and steroid production in mosquito adult males. (A) Present geographical distribution of the 16 mosquito species (*Anopheles* genus) analysed in this study represented on a zoogeographical map (modified from The geographical distribution of animals, Eckert IV projections department of geosciences, Texas tech university). Species matching numbers are shown on panel B. Pink: *Cellia* subgenus, blue : *Nyssorhynchus* subgenus, green : *Anopheles* subgenus. (B) Bayesian phylogeny of 20 *Culicidae* species, 19 species tested for ecdysteroid male production plus

Chagasia bathana (subfamily *Anophelinae,* genus *Chagasia*) used as outgroup for

phylogenetic analyses. Dominant malaria human vectors are indicated by a red star. Time is

represented in millions of years (Ma). Approximated node ages are detailed in Supplementary

Table 1. Bayesian node support values are presented on the right side of each node.

Ecdysteroid titers in whole 5-day-old virgin males are indicated on the right side of the tree.

Results are expressed as mean +/- SEM in pg E equivalents per male. Results were subjected

to statistical analysis using Kruskall-Wallis test for nonparametric data followed by Dunn's

post-test (control group: Extraction blank). The indicated p values are those obtained with

Dunn's test (***, p value < 0.001; ****, p value < 0.0001). NT: not tested. Predicted lineages

with significant male 20E production are shaded pink on the tree. The pink horizontal bar

represents the minimum/maximum 95% confidence interval (CI) estimated time in Ma for

origin of male 20E production. The geological time scale is adapted from the Geological

Society of America (http://www.geosociety.org/science/timescale/). The white coloured cases

represent the quaternary period. PAL., Paleocene; OLI., Oligocene; MIO., Miocene; P.,

Pliocene.

688
689
Figure 2. Present geographic distribution of species belonging to *Cellia* and *Anopheles*

subgenera (*Anopheles* genus). Total numbers of mosquito species belonging to the *Cellia*

(A, red) and *Anopheles* (B, green) subgenera per country (sourced from the Walter Reed

Biosystematics Unit, http://www.wrbu.org/) are represented on world maps created with R.

Numbers (nb) of mosquito species per country are represented by a coloured gradient as

depicted under each map. Grey colour means no data are available for the country.

696
697
Figure 3. Secondary follicle detachment from the germarium in virgin and mated non

blood-fed females from 12 *Anopheles species*. (A) Cartoon representing the detachment of

the secondary follicle from the germarium. Follicular cells of somatic origin are coloured in

blue. Germ cells (germline stem cells, developing cysts and nurse cells) are coloured in grey

and the future oocyte of the secondary follicle in dark grey. g: germarium, I: primary follicle,

II: secondary follicle, nc: nurse cells. (B) Secondary follicle detachment from the germarium

in ovarioles of virgin (V) and mated (M) females. Secondary follicles are either detached

from the germarium (detached, green), in the progress of detachment (constricted, yellow) or

non-detached yet (non-detached, red). Representative pictures of these three states are shown

in Supplementary Fig. 3. The secondary follicle are significantly more detached in mated

females compared to virgin females for *An. atroparvus* (p=0.0226), *An. freeborni* (p<0.0001),

An. stephensi (p<0.0001), *An. minimus* (p=0.0228) and *An. merus* (p=0.0072).

Figure 4. Egg development in virgin and mated blood-fed females from 12 species of
Anopheles **mosquitoes.** Total number of eggs in virgin (V, light colours) and mated (M, dark
colours) females 48 hours after blood feeding. Green: *Anopheles* subgenus, blue:
Nyssorhynchus subgenus*,* pink: *Cellia* subgenus. Females from *An. atroparvus* (Mann-
Whitney U=295.5, p= 0.0214), *An. albimanus* (Mann-Whitney U= 941.5, p<0.0001), *An.*
stephensi (Mann-Whitney U= 232.5, p= 0.0235), *An. dirus* (Mann-Whitney U= 298, p=
0.0240) and *An. quadriannulatus* (Mann-Whitney U= 309.5, p= 0.0373) develop significantly
more eggs when they are mated.

Subgenus	Species	DVS	Male steroid production	Follicle detachment	Egg development
Anopheles	An. quadrimaculatus	-	-	-	-
	An. atroparvus	+	-	+	+
	An. freeborni	+	-	+	-
Nyssorhynchus	An. albimanus	+	-	-	+
Cellia	An. stephensi	+	++	+	+
	An. minimus	+	+	+	-
	An. dirus	+	+	-	+
	An. farauti	+	++	-	-
	An. merus	+	+	+	-
	An. gambiae	+	+++	-	-
	An. arabiensis	+	+++	-	-
	An. quadriannulatus	+	++	-	+

Table 1. Summary of mating effect on follicle detachment and egg development in regard to malaria vector status and male steroid production in *Anopheles* species. Dominant vector species (DVS) of human malaria are signalled by a +. For male steroid production, relative low titers (mean range 500pg E equivalent per male) are indicated by +, medium (range [1000-2000] pg E equivalent per male) titers by ++, and high titers (above 3000pg E equivalent per male) by +++. For follicle detachment and increase of egg development, - indicates no effect of mating and + indicates an effect of mating on either reproductive traits in females.

732 **Supplementary Figures and Tables**

Supplementary Figure 1. Phylogenetic relationships of the genus *Anopheles* and evolution of male 20E production. Maximum likelihood phylogeny (PhyML) based on a concatenated dataset. Bootstrap supports (100 replicates) are presented on the right side of each node. Bars on the right side indicate species that belong to the same subgenus. The lineages of the subgenus *Cellia* are highlighted in blue. The lineages with male 20E production are shaded in pink.

Supplementary Figure 2. *Anopheles stephensi* males transfer ecdysteroids to females
during mating. Ecdysteroid titers were measured in virgin males (VM), in mated males
(MM) just after copulation, in virgin females (VF), and in mated females (MF) just after
copulation. Ecdysteroids were extracted from each individual mosquito and quantified by
EIA. Results are expressed as mean +/- SEM in in pg E equivalents per animal. Results were
subjected to statistical analysis using Mann-Whitney test (****, P< 0.0001).

741
Supplementary Figure 3. Secondary follicle detachment from the germarium in
***Anopheles stephensi* non blood-fed females according to the insemination status.**
Confocal pictures of ovarioles showing the germarium and the secondary follicle from 10 non
blood-fed females either virgin (A to J) or mated (K to T) stained with DAPI. g: germarium,
II: secondary follicle. Yellow arrowheads show secondary follicles detached from the
germarium and yellow stars show secondary follicles in progress of detachment. Follicles not
marked are not detached. Scale bar is 10.25 µm.
749

Anopheles albimanus

Anopheles gambiae

750
Supplementary Figure 4. Egg development in *Anopheles albimanus* (*Nyssorhynchus*
subgenus) and *Anopheles gambiae* (*Cellia* subgenus) virgin and mated females fed on
mouse or human blood. Total number of eggs in virgin (light colours) and mated (dark
colours) females was counted 48 hours after feeding on either human blood (HB) or mouse
blood (MB). *An. albimanus* is coloured in blue and *An. gambiae* in pink. Data were subjected
to Mann-Whitney non-parametric test. Mated females from *An. albimanus* develop
significantly more eggs then virgin females when fed on human blood (Mann-Whitney U=90,
p<0.0001) or mouse blood (Mann-Whitney U=122, p<0.0001). Virgin and mated females of
both species develop significantly more eggs when fed on mouse blood compared to human
blood (*An. albimanus* virgin: Mann-Whitney U=165, p<0.0001; *An. albimanus* mated: Mann-
Whitney U=52.50, p<0.0001; *An. gambiae* virgin: Mann-Whitney U=74, p<0.0001; *An.*
gambiae mated: Mann-Whitney U=90, p<0.0001).

763
764
765
766

Node	Taxon Set	Divergence Time*	Node	Taxon Set	Divergence Time*
1	all	124.2 Ma (83.7-165.8)	11	*An. atroparvus – An. quadrimaculatus*	25.6 Ma (16.7-34.8)
2	*Aedes – Culex* (genus)	78.9 Ma (51.6-106.5)	12	*An. atroparvus – An. freeborni*	18.1 Ma (11.5-24.7)
3	*Aedes* (genus)	35.6 Ma (23.1-48.3)	13	*Cellia* (subgenus)	69.2 Ma (45.6-93.0)
4	*Anopheles – Chagasia* (genus)	113.7 Ma (75.2-152.4)	14	*An. farauti – An. stephensi*	62.5 Ma (41.1-84.3)
5	*Anopheles* (genus)	84.1 Ma (55.8-112.7)	15	*An. stephensi – An. minimus*	46.6 Ma (30.0-62.8)
6	*Nyssorhynchus – Anopheles* (subgenus)	76.4 Ma (50.4-102.3)	16	*An. farauti – An. dirus*	31.6 Ma (20.5-42.9)
7	*Nyssorhynchus* (subgenus)	20.7 Ma (13.5-28.3)	17	*An. gambiae – An. quadriannulatus*	17.2 Ma (11.0-23.7)
8	*An. aquasalis – An. albitarsis*	17.4 Ma (11.1-24.1)	18	*An. gambiae – An. merus*	3.2 Ma (1.9-4.5)
9	*Anopheles* (subgenus)	71.9 Ma (47.7-97.0)	19	*An. gambiae – An. arabiensis*	1.2 Ma (0.6-1.9)
10	*An. atroparvus - An. pseudopunctipennis*	61.9 Ma (40.3-82.9)			

767   *Estimates are the posterior means with 95% highest posterior density intervals.

768   **Supplementary Table 1. Divergence time estimates of the *Anopheles* species and**

769   **outgroups**. Estimates were obtained from the Bayesian phylogenetic analysis and are based

770   on fossil data for temporal calibration.

Anopheles species	Animal species	Blood source
An. quadrimaculatus	mouse	
An. atroparvus	rabbit	
An. freeborni	mouse	
An. albimanus	mouse	human
An. stephensi	mouse	
An. minimus	mouse	
An. dirus	rabbit	
An. farauti	mouse	
An. merus	mouse	
An. gambiae	mouse	human
An. arabiensis	rabbit	
An. quadriannulatus	mouse	

773 **Supplementary Table 2. Animal species or blood source on which mosquitoes were fed in**
774 **this study.**

Name	Sequence	Locus	Reference
COI-2	TCCATTGCACTAATCTGCCA	*COI*	[77]
LepF1	ATTCAACCAATCATAAAGATATTGG		[78]
tLEU-2	ATGGCAGATTAGTGCAATGA	*COII*	modified from [79]
tLys-2	TGATTTAAGAGATCATTACTTG		modified from [79]
T7-ND5	TAATACGACTCACTATAGGGATTAACTGTATGTTATTCITTYC	*ND5*	modified from [80]
tRNAPhe-2	CCTAACATCTTCAGTGTCATGCT		modified from [80]
cytb-R2	TACTGGTCGAGCTCCAATTCA	*CYTB*	this study
T7-cytbF	TAATACGACTCACTATAGGGACAAATATCATTTTGAGGAGCIACAG		modified from [81]
18S-F2	CAGCTCCACTAGCGTATATTAA	*18S*	this study
18S-R2	TTAACCAGACAAATCIATCCAC		this study
D2F2	AGTCGTGTTGCTTGATAGTG	*28S*	[28]
D2R2	CTTGGTCCGTGTTTCAAGAC		[28]
T7-G6pdF1	TAATACGACTCACTATAGGGTGGACACGGARGGNACICAYTTYGA	*g6pd*	this study
T7-G6pdF2	TAATACGACTCACTATAGGGTCGGGAGATTTGGCTAAIAARAARATHTA		this study
G6pdR1	TGTTCCAGGTAGGGCTRAANATYTGRTT		this study
G6pdR2	CATCAGGTTYTGNACCATYTC		this study
T7-wF1	TAATACGACTCACTATAGGGCGGCTCMGIAAYTGYTGYAC	*white*	this study
T7-wz2E-2	TAATACGACTCACTATAGGGTACAACCCGGCNGAYTTYTA		modified from [82]
wR1	CCTCGACGCGGAARTTRAANGTYTC		this study
wR2	GGAACCAGGACAGGTACGANAIRTAYTTRAA		this study

Supplementary Table 3. Sequences of primers used to sequence DNA for the
phylogenetic analyses. Partial genomic regions of four nuclear genes (*g6pd*, *white*, *18S* and
28S) and four mitochondrial genes (*COI*, *COII*, *ND5* and *CYTB*) were amplified by PCR with
gene-specific or degenerate primers. Degenerate oligonucleotides were designed based on
previous studies and optionally modified or newly designed. Some oligonucleotide sequences
contained T7 or SP6 universal primer sequences at their 5' end, following Bonacum *et al*.
[83].
804

| | Locus | | | | | | | |
Species	COI	COII	ND5	CYTB	18S	28S	g6pd	white
Aedes aegypti	NC010241	NC010241	NC010241	NC010241	AAEL017915	AAEL017581	XM001660122	U73826
Aees albopictus	AY072044	AY072044	AY072044	AY072044	X57172	MQSRNAGN	---	U73828
Anopheles albimanus	AF417695	AF417731	AF311270	AF311251	MSQINSP	MSQINSP	AF317824	MSQWHITE
Anopheles albitarsis	HQ335344	HQ335344	HQ335344	HQ335344	AF417768	AF417803	AF317823	AF318198
Anopheles aquasalis	AF417697	**MG560168**	**MG560169**	**MG560174**	AF417769	AF417804	**MG560181**	***MG560185***
Anopheles arabiensis	AF417705	AF417741	SRS008420 contigs: 7466, 23167, 23406, 36079, 47611	SRS008420 contigs: 1529, 8012, 23026, 33951, 34964, 44097	AF417777	AF417812	AY118019	AARA010493-RA
Anopheles atroparvus	**MG560162**	**MG560165**	**MG560170**	**MG560175**	AM072973	**MG560178**	**MG560184**	***MG560187***
Anopheles dirus	JX219731	JX219731	JX219731	JX219731	AF417779	AF417814	ADIR005618-RA	ADIR001487-RA
Anopheles farauti	AF417708	AF417744	JX219741	HQ840893	AF121054	AF417815	AFAF008422-RA	AXCN01000101.1 pos. 115279-116093
Anopheles freeborni	AF417717	AF417753	SRS008481 contigs: 12850, 20811, 21808, 23967	SR008481 contigs: 18, 9086, 23902, 24406, 26852, 29845, 41851	AF417788	AF417824	---	AFU73830
Anopheles gambiae	MSQMTCG	MSQMTCG	MSQMTCG	MSQMTCG	AM157179	KC177663	AGAP012678	AGAP000553
Anopheles merus	**MG560163**	**MG560167**	**MG560171**	**MG560176**	**MG560177**	**MG560180**	**MG560182**	***MG560186***
Anopheles minimus	AF417710	AF417746	---	KF431913	AF417781	DQ523567	AMIN009223-RA	AMIN005306-RA
Anopheles plumbeus	**MG560164**	**MG560166**	**MG560172**	**MG560173**	AM072974	**MG560179**	**MG560183**	---
Anopheles pseudopunctipennis	AF417721	AF417757	AF311272	---	AF417792	AF417828	AF317810	AF318197
Anopheles quadriannulatus	DQ792581	DQ792581	SRS008482 contigs: 9810, 9922, 12564	SRS008482 contigs: 2436, 3560, 5140, 6641, 12944, 25648, 35018	---	KB667953 genomic scaffold supercont1.758 pos 16103-16634	AY118023	AQUA010818-RA
Anopheles quadrimaculatus	NC000875	NC000875	NC000875	NC000875	AY988423	AY569555	AF317809	AF318207
Anopheles stephensi	AF417713	AF417749	AF311273	AF311254	AF417784	AF417820	AF317808	AF318208
Chagasia bathana	AF417726	AF417762	AF311281	AF311253	AF417797	AF417831	AF317819	AF318194
Culex pipiens	NC015079	NC015079	NC015079	NC015079	AY988445	---	CPU09034	---

805
806
807
Footnote: Genbank Accession Numbers, VectorBase identifiers (shaded grey), or BlastN search hits at VectorBase (shaded dark grey). Newly generated sequences are highlighted in bold.

808
809 **Supplementary Table 4. GenBank Accession Numbers of the dataset used for the**
810 **phylogenetic analyses.**
811

Partition, site model, gamma categories, nucleotide frequencies

COI-COII-ND5-CYTB	18S	28S	g6pd	white
GTR+I+G	TN93+I+G	TN93+I+G	TN93+I+G	GTR+I+G
5	5	5	5	5
estimated	estimated	estimated	estimated	estimated

Supplementary Table 5. Partition specific site models and parameters.

826
827
828