2   Viromic analysis of wastewater input

3   to a river catchment reveals a diverse

4   assemblage of RNA viruses

Evelien M. Adriaenssens[1,*], Kata Farkas[2], Christian Harrison[1], David Jones[2],

Heather E. Allison[1], Alan J. McCarthy[1]

[1] Microbiology Research Group, Institute of Integrative Biology, University of

Liverpool, UK

[2] School of Environment, Natural Resources and Geography, Bangor University,

Bangor, LL57 2UW, UK

* Corresponding author: evelien.adriaenssens@liv.ac.uk

Abstract

Detection of viruses in the environment is heavily dependent on PCR-based approaches that require reference sequences for primer design. While this strategy can accurately detect known viruses, it will not find novel genotypes, nor emerging and invasive viral species. In this study, we investigated the use of viromics, i.e. high-throughput sequencing of the biosphere viral fraction, to detect human/animal pathogenic RNA viruses in the Conwy river catchment area in Wales, UK. Using a combination of filtering and nuclease treatment, we extracted the viral fraction from wastewater, estuarine river water and sediment, followed by RNASeq analysis on the Illumina HiSeq platform for the discovery of RNA virus genomes. We found a higher richness of RNA viruses in wastewater samples than in river water and sediment, and assembled a complete norovirus GI.2 genome from wastewater effluent, which was not contemporaneously detected by conventional qRT-PCR. To our knowledge, this is the first environmentally-derived norovirus genome sequence to be available from a public database. The simultaneous presence of diverse rotavirus signatures in wastewater indicated the potential for zoonotic infections in the area and suggested run-off from pig farms as the origin of these viruses. Our results show that viromics can be an important tool in the discovery of pathogenic viruses in the environment and can be used to inform and optimize reference-based detection methods provided appropriate and rigorous controls are included.

Importance

Enteric viruses cause gastro-intestinal illness and are commonly transmitted through the faecal-oral route. When wastewater is released into river systems, these viruses can contaminate the environment. Our results show that we can use viromics to find

37   the range of potentially pathogenic viruses that are present in the environment and

38   identify prevalent genotypes. The ultimate goal is to trace the fate of these

39   pathogenic viruses from origin to the point where they are a threat to human health,

40   informing reference-based detection methods and water quality management.

Introduction

Pathogenic viruses in water sources are likely to originate primarily from

contamination with sewage. Classic marker bacteria used for faecal contamination

monitoring, such as *Escherichia coli* and *Enterococcus* spp., are not, however, good

indicators for the presence of human enteric viruses (1). The virus component is

often monitored using qPCR approaches, which can give information on relative

abundance of specific viruses and their genotype, but only those that are both known

and characterised (2). Viruses commonly targeted in sewage contamination assays

include noroviruses (3), hepatitis viruses (4), enteroviruses (5), and various

adenoviruses (6, 7). Viral monitoring in sewage has previously yielded positive

results for norovirus, sapovirus, astrovirus, and adenovirus, indicating that people

are shedding viruses that are not necessarily detected in a clinical setting (8). This

same study found a spike in norovirus genogroup GII sequence signatures in

sewage two to three weeks before the outbreak of associated disease was reported

in hospitals and nursing homes. The suggestion, therefore, is that environmental

viromics can provide an early warning of disease outbreaks, in addition to the

monitoring of virus dissemination in watercourses.

Recent reviews have proposed the use of viral metagenomics or viromic approaches

as an alternative method to test for the presence of pathogenic viruses in the

environment (2, 9, 10). Provided the entire viral community is sampled and

sequenced, novel genotypes or even entirely novel viruses can be detected.

Potential new viral markers for faecal contamination have already been revealed,

such as pepper mild mottle virus and crAssphage (11, 12), among the huge diversity

of human viruses found in sludge samples (13).

In this pilot study, we have used viromics to investigate the presence of human

pathogenic RNA viruses in wastewater, estuarine surface water and sediment in a

single catchment. The water and sediment samples were collected at, and

downstream of, the wastewater treatment plant (Lanrwst, Wales, UK), at the estuary

of the river Conwy (Wales, UK) near Morfa beach (Figure 1). To our knowledge, this

is the first study to use unamplified environmental viral RNA for sequencing library

construction, sequence dataset production and subsequent analysis. Because we

used a directional library sequencing protocol on RNA, rather than amplifying to

cDNA, we were able to distinguish single-stranded from double-stranded RNA

genome fragments.

Results

Sample overview

Wastewater influent and effluent samples were collected from the Llanrwst

wastewater treatment plant (53°08'24.4"N 3°48'12.8"W; Figure 1) in September and

October 2016, resulting in four different samples, LI_13-9 (Llanrwst influent Sep

2016), LE_13-9 (Llanrwst effluent Sep 2016), LI_11-10 (Llanrwst influent Oct 2016),

LE_11-10 (Llanrwst effluent Oct 2016). Estuarine surface water (SW) was collected

from Morfa beach (53°17'37.7"N 3°50'22.2"W; Conwy, Wales, Figure 1) in November

2016 and sediment from the same site in October and November 2016 (Sed1, Sed2,

respectively).

Figure 1: Map of the sampling locations, indicated with blue arrows. Data in the
left panel was taken from Google Maps.

As an initial assessment, samples were tested for the presence of a subset of locally

occurring enteric RNA viruses using qRT-PCR (Table 1). Only norovirus (NoV)

genogroup GII signatures were detected in the wastewater samples. In the samples

collected in September 2016, 10^3 genome copies (gc)/l of norovirus GII were

observed in both the influent (LI_13-9) and in the effluent (LE_13-9). In the samples

collected in October 2016, approx. 10^2 gc/l (below the limit of quantification which

was approx. 200 gc/l) were observed in the influent (LI_11-10) and a considerably

higher concentration of 5×10^4 gc/l was noted in the effluent (LE_11-10). All qRT-

PCRs were negative for the presence of sapoviruses (SaV) and hepatitis A/E viruses

(HAV/HEV). None of the target enteric viruses were found in the surface water and

sediment samples.

Table 1: Summary of viromic and qRT-PCR detection of the presence of

specific RNA viruses across the samples (sewage, estuarine water and

sediment).

Sample name[a]	Sample volume/ mass	Location	# contigs (curated)	Target RNA viruses detected in contigs[b]	qRT-PCR results (gc/l)[c]
LI_13-9	1 l	Llanrwst WWTP	5721	RVA, RVC, PBV, SaV	NoVGII 1,457
LE_13-9	1 l	Llanrwst WWTP	2201	RVA, RVC, PBV	NoVGII 1,251
LI_11-10	1 l	Llanrwst WWTP	859	PBV	NoVGII detected
LE_11-10	1 l	Llanrwst WWTP	5433	NoVGI, RVA, RVC, PBV, AsV	NoVGII 50,180
SW	50 l	Morfa beach	243	-	-
Sed1	60 g	Morfa beach	550[d]	-	-
Sed2	60 g	Morfa beach	550[d]	-	-

[a] LI: sewage influent; LE: sewage effluent; SW: estuarine surface water; Sed: estuarine sediment

[b] RVA: rotavirus A; RVB: rotavirus B; PBV: picobirnavirus; SaV: sapovirus; NoVGI: norovirus genogroup I; AsV:

astrovirus

[c] Samples were tested with qRT-PCR for the following targets: NoVGI, NoVGII, SaV, HAV, HEV. Results

reported in genome copies per liter (gc/l), NoVGII was detected below limit of quantification (approx. 200 gc/l) in

sample LI_11-10. Nov GII was the only target virus detected by qRT-PCR.

[d] Samples Sed1 and Sed2 were assembled together into the contig dataset Sed.

Summary of viral diversity

The virus taxonomic diversity present in each sample was assessed by comparison

of curated read and contig datasets with both the RefSeq Viral protein database and

the non-redundant protein database of NCBI, using Diamond blastx (14) and lowest

common ancestor taxon assignment with Megan 6 (15). For wastewater samples

LI_13-9, LE_13-9 and LE_11-10, two libraries were processed (indicated with _1 and

2 in the dataset names) and one each for the wastewater influent sample LI_11-10,

the surface water sample (SW) and two sediment samples (Sed1 & Sed2). This

section focuses on those reads and contigs that have been assigned to the viral

fraction exclusively, disregarding sequences of cellular or unknown origin.

The wastewater samples showed a greater richness of known viruses and had a

larger number of curated contigs than the surface water and sediment samples

(Figure 2). At the viral family level, between 14 and 34 groups were observed for

wastewater influent and effluent samples, including the unclassified levels, 12 for the

surface estuarine water sample, and 11 and 5 for the sediment samples Sed1 and

Sed2, respectively. The unclassified viruses and unassigned bins are indicated in

red in Figure 2 and made up the majority of known reads in the estuarine sediment

samples. In most of the viromes, dsDNA and ssDNA virus families were present,

despite having performed a DNase treatment after viral nucleic acid extraction (Table

S1). These families represented only a minor (<5%) proportion of the total assigned

reads with a few exceptions. In wastewater influent sample LI_11-10, reads assigned

to the dsDNA family *Papillomaviridae* accounted for 61% of the total, while in the

surface water sample reads assigned to the ssDNA families *Circoviridae* and

Microviridae represented 50% and 12% of the total, respectively. This is most likely

to be due to incomplete digestion of the viral DNA with the DNase Max kit than to

corresponding mRNA transcripts actually being present in the viromes.

Figure 2: Taxonomic distribution of curated read data at the virus family level.
Reads were assigned to a family or equivalent group by Megan6 using a lowest
common ancestor algorithm, based on blastx-based homology using the program
Diamond with the RefSeq Viral protein database (version January 2017) and the
non-redundant protein database (version May 2017). Only viral groupings are
shown. LI: sewage influent; LE: sewage effluent; SW: estuarine surface water; Sed:
estuarine sediment.

The families of dsRNA viruses present in these datasets were *Totiviridae* (fungi and

protist hosts)*, Reoviridae* (invertebrate, vertebrate & plant hosts)*, Picobirnaviridae*

(mammals)*, Partitiviridae* (fungi & protists) and *Birnaviridae* (vertebrates and

invertebrates), with a small number of reads recognized as unclassified dsRNA

viruses (Figure 2). None of these groups were present in all libraries, but totivirus

and picobirnavirus signatures were present in all wastewater samples and reoviruses

were found in three out of the four wastewater samples. *Partitiviridae* signatures

were only found in the wastewater LE_11-10 and LI_13-9 samples, while

Birnaviridae reads were only present in the wastewater LE_13-9 libraries. The

sediment and surface water samples did not have detectable levels of dsRNA virus

sequences.

Positive sense ssRNA viruses were the most diverse class of viruses present in

these datasets. The family *Tombusviridae*, which groups plant viruses with

monopartite or bipartite linear genomes (16), was present in all samples with the sole

exception of the wastewater influent sample LI_11-10 (Figure 2, Table S1). Virus

signatures belonging to the family *Virgaviridae*, representing plant viruses, were

present in all wastewater samples at comparable levels. Other highly represented

families or groupings were the families *Dicistroviridae* (invertebrate hosts),

Nodaviridae (invertebrate & vertebrate hosts) and the bacteriophage family

Leviviridae, the plant virus genus *Sobemovirus*, and the groupings of "unclassified

ssRNA positive-strand viruses" and several unclassified/unassigned/environmental

members of the order *Picornavirales*. Sediment sample Sed1 was the only sample

with signatures of the family *Alvernaviridae*, which has as its sole member the

dinoflagellate virus Heterocapsa circularisquama RNA virus 01. The wastewater

effluent sample LE_11-10 and influent sample LI_13-9_1 were the only samples with

calicivirus signatures, and sample LE_11-10_1 and LE_1-10_2 were the only

samples with *Astroviridae* reads (vertebrate host). Several families of the order

Picornavirales were detected in the wastewater samples at different levels in

different samples, and a small number of unassigned picornaviruses was detected in

the surface water sample (SW).

We did not observe any known negative sense ssRNA viruses in any of the

sequencing libraries, but it is possible that some of the unaffiliated viral contigs

belong to this class. These types of viruses are enveloped and predicted to degrade

more rapidly than the non-enveloped enteric viruses in wastewater treatment plants

and the environment (17). Given that the known families of negative sense ssRNA

viruses consist of potentially deadly pathogens, such as Influenza A virus

(orthomyxovirus), Lassa virus (arenavirus), Zaire ebolavirus (filovirus) or Rabies

virus (rhabdovirus), the lack of signatures of these viruses in the datasets most likely

suggests reassuringly that they were not present in the investigated samples.

Potential human pathogenic viruses

An important aim of this study was to investigate the presence and genomic diversity

of potential human pathogenic RNA viruses in different sample types within the river

catchment area. To minimize miss-assignments of short sequences to taxa, we used

the assembled, curated contig dataset and looked for contigs representing near-

complete viral genomes.

Presence of a norovirus GI.2 genome

We were particularly interested in finding norovirus genomes to explore the genomic

diversity of these important and potentially abundant pathogens originating from

sewage and disseminated in watercourses, with implications for shellfisheries and

recreational waters. This is of relevance due to known issues of sewage

contamination in the region (18). Members of the genus *Norovirus* (family

Caliciviridae) are non-enveloped, icosahedral (+)ssRNA viruses with a linear,

unsegmented ~7.6 kb genome encoding three ORFs (16). These viruses are divided

into different genogroups of which GI and GII are associated with human

gastroenteritis (19, 20). Noroviruses are identified routinely by qRT-PCR, providing

an opportunity to examine correlations between qRT-PCR and metaviromic results.

We only found norovirus signatures in the libraries of wastewater effluent sample

LE_11-10. These reads assembled into a single contig of 7,542 bases, representing

a near-complete norovirus genome (GenBank accession number MG599789). Read

mapping showed an uneven coverage over the genome length between 18x and

745x (13,165 reads of library 1 and 8986 reads of library 2). This confirmed that our

contig was derived from a single-stranded genome, as all forward reads in the pairs

were oriented in the same direction. Based on this mapping, we performed variant

calling and the consensus sequence was corrected in cases where the variant was

present in more than 85% of the reads. To our knowledge, this is the only

metagenome-derived, environment-associated (i.e. non-host associated) near-

complete norovirus genome sequence deposited in a public database (INSDC

nuccore database was searched for norovirus, txid142786 sequences > 5000 nt).

A BLASTN search revealed two close relatives to our wastewater-associated

norovirus genome, norovirus Hu/GI.2/Jingzhou/2013401/CHN (KF306212) which is

7740 bases in length (21), displaying a nucleotide sequence identity of 99% over

99% of the genome length, and norovirus Hu/GI.2/Leuven/2003/BEL (FJ515294) at

95% sequence identity over 99% of the genome (Figure 3). From the 5' end of our

norovirus   contig,   62   bases   were   missing   compared   with

Hu/GI.2/Jingzhou/2013401/CHN and from the 3' end 165 bases and the polyA tail

were not present. We compared the sequence of our norovirus with

Hu/GI.2/Jingzhou/2013401/CHN base by base and observed 81 SNPs and no other

forms of variation. Of the SNPs, only eight were non-synonymous resulting in five

different amino acids incorporated in the non-structural polyprotein (ORF1); one in

the major capsid protein (ORF2) and two in the minor structural protein (ORF3)

(Table S2). According to the current classification criteria, this level of similarity

places our assembled genome in genogroup GI, genotype GI.2, with only a single

amino   acid   different   between   the   major   capsid   protein   (MCP)   of

Hu/GI.2/Jingzhou/2013401/CHN and the genome assembled here.

Figure 3: Pairwise genome comparison between the virome norovirus genome

(middle) and its closest relatives, Norovirus Hu/GI.2/Jingzhou/2013401/CHN

and Norovirus Hu/GI.2/Leuven/2003/BEL. BLASTN similarity is indicated in shades

of grey. ORFs are delineated by dark blue arrows. The deviation from the average

GC content is indicated above the genomes in a green and purple graph. The qRT-

We tested the genotype grouping of our genome in a whole genome phylogeny with

all complete genome sequences of genogroup I available in GenBank. The

phylogenomic tree clearly delineated the different genotypes within genogroup GI,

placing the newly-assembled genome within genotype GI.2, with the reference

isolate for GII used as an outgroup (Figure 4).

For further validation, the full genome of the novel norovirus GI was recovered using

RT-PCR. However, the amplicon could not be ligated into a plasmid and hence was

not fully sequenced.

Figure 4: Maximum Likelihood phylogenetic tree of norovirus genomes

belonging to genogroup GI, with the norovirus GII reference genome as outlier.

The nucleotide sequences were aligned with MUSCLE and the alignment was

trimmed to the length of the virome sequence LE_11-10 contig 6, resulting in 7758

positions analyzed for tree building. The Maximum Likelihood method was used with

a Tamura Nei model for nucleic acid substitution. The percentage of trees in which

the associated taxa clustered together is shown next to the branches. The scale bar

represents the number of substitutions per site.

Presence of diverse rotavirus segments in wastewater samples

Rotaviruses are segmented dsRNA viruses belonging to the family *Reoviridae,* causing gastroenteric illness in vertebrates and are transmitted through the faecal-oral route (16). Read signatures assigned to the genus *Rotavirus* were found in three of the four wastewater samples (all but LI_11-10). Wastewater influent sample LI_13-9 contained the most signatures with approximately 75,000 reads, assembled into 120 contigs, representing genome fragments of 10 out of the 11 rotavirus segments. At the species level, these genome fragments were assigned to either the species *Rotavirus A* or *Rotavirus C*. Comparing the amino acid sequences of the predicted proteins, some contigs showed high levels of identity (>88%) with either the segments of rotavirus A (RVA) or rotavirus C (RVC) reference genomes as available in the RefSeq database (23, 24), while others showed a lower identity with a variety of RVC isolates only. The segmented genome nature and the possibility of segment exchange make it difficult to confidently identify the number of rotavirus types present in this sample. Given the amino acid similarities with both RVA and RVC types (Supplementary Table 1), we suggest there are at least two, and possibly three types present here.

Using the RotaC 2.0 typing tool for RVA, and blast-based similarity to known genotypes, we have typed the rotavirus genome segments found here (Table 2). The combined genomic make-up of the RV community in sample LI_13-9 was G8/G10/Gx-P[1]/P[14]/P[41]/P[x]-I2/Ix-R2/Rx-C2/Cx-M2/Mx-A3/A11/Ax-Nx-T6/Tx-E2/Ex (25, 26). The potential hosts for each segment were derived from the hosts of the closest relatives. This analysis showed that the RVA viruses were possibly infecting humans or cattle, while the RVC viruses were most likely porcine (Table 2).

Table 2: Rotavirus A and C genome information and its detection in the LI_13-9 sample dataset.

Genome segment	Length (nt)	Protein	Predicted function	# contigs	Putative genotypes	Potential hosts[a]
Rotavirus A						
Segment 1	3302	VP1	RNA-dependent RNA polymerase	7	R2	Human, cow
Segment 2	2693	VP2	core capsid protein	1	C2	Human
Segment 3	2591	VP3	RNA capping protein	1	M2	Human, sheep
Segment 4	2363	VP4	outer capsid spike protein	3	P[1], P[41], P[14]	Human, pig, alpaca, monkey
Segment 5	1614	NSP1	interferon antagonist protein	6	A3, A11	Human, cow, pig, deer
Segment 6	1356	VP6	inner capsid protein	1	I2	Human
Segment 7	1105	NSP3	translation effector protein	4	T6	Human, dog, cow
Segment 8	1059	NSP2	viroplasm RNA binding protein	0	-	-
Segment 9	1062	VP7	outer capsid glycoprotein	2	G10, G8	Cow, Human
Segment 10	751	NSP4	enterotoxin	1	E2	Human, cow
Segment 11	667	NSP5;6	phosphoprotein; non-structural protein	0	-	-
Rotavirus C				(contigs RVCX)		
Segment 1	3309	VP1	RNA-dependent RNA polymerase	7 (0)	Rx	Pig, cow
Segment 2	2736	VP2	core capsid protein	4(2)	Cx	Pig, dog
Segment 3	2283	VP4	outer capsid protein	2 (4)	P[x]	Pig
Segment 4	2166	VP3	guanylyltransferase	6 (0)	Mx	Pig
Segment 5	1353	VP6	inner capsid protein	1 (0)	Ix	Pig
Segment 6	1350	NSP3		0 (1)	Tx	Human
Segment 7	1270	NSP1		0 (2)	Ax	Pig, dog
Segment 8	1063	VP7	outer capsid glycoprotein	0 (2)	Gx	Pig
Segment 9	1037	NSP2		2 (0)	Nx	Pig
Segment 10	730	NSP5		0 (0)	-	-
Segment 11	613	NSP4	enterotoxin	0 (4)	Ex	Pig

[a] Potential hosts are defined as the hosts of the reference rotavirus sequence with the highest similarity to the contigs found in the virome sample LI_13-9.

Picobirnaviruses showed a high prevalence in wastewater

All the wastewater virome libraries contained signatures assigned to the dsRNA family *Picobirnaviridae*, genus *Picobirnavirus* (Figure 2) and these reads assembled into between 42 (LE_13-9) and 510 (LI_13-9) contigs. Both picobirnavirus genome segments, segment 1 containing two hypothetical proteins and segment 2 on which the RNA-dependent RNA polymerase (RdRP) is encoded, were observed in the samples. The contigs showed little sequence similarity with the reference genome *Human picobirnavirus* (RefSeq segment accession numbers NC_007026.1 and NC_007027.1). Phylogenetic analysis of a partial region of the predicted RdRPs in the virome contigs was not able to resolve any cluster or evolutionary origin (Figure 5). Picobirnavirus RdRPs from human, animal and environmental isolates, as well as the majority of the virome sequences were grouped in one large, unsupported cluster that showed relatively little genomic diversity. While many picobirnaviruses have been isolated from humans with gastroenteritis, a review of the known cases suggested that picobirnaviruses are probably not the main cause of acute diarrhea and are secondary pathogens with potential synergistic effects (27). A qRT-PCR-based investigation into the suitability of human picobirnaviruses as indicators of human faecal contamination, showed that they were not present in a sufficient proportions of tested samples to be good water quality indicators (28), but their high diversity in our sample set warrants further investigation using metaviromic methods.

Figure 5: Maximum likelihood phylogenetic tree of RdRP amino acid sequences of isolated and virome picobirnaviruses. Sequences from isolates are indicated with open dots, virome-derived sequences with closed dots, sample LI_11-10 in purple, sample LE_11-10 in blue, and sample LI_13-9 in red. Sequences were aligned using MUSCLE providing 114 amino acid positions for tree generation. The Maximum Likelihood was used with a JTT matrix-based model [1]. The scale bar represents the number of substitutions per site. Bootstrap values of all branches were low.

Partial genomes of other potentially pathogenic RNA viruses

In sample LI_13-9, a small contig of 347 bases was found that was 94% identical at the nucleotide level to the Sapovirus Mc2 ORF1 (AY237419), in the family *Caliciviridae*. We have also identified four contigs of approximately 500 bases in sample LE_11-10 that resembled most closely the Astrovirus MLB2 isolates MLB2/human/Geneva/2014 (KT224358) and MLB2-LIHT (KX022687) at 99% nucleotide identity. In addition, we identified several reads and contigs assigned to the family *Picornaviridae* which comprises a diverse set of enteric viruses, but the closest relatives in the databases were metagenomically assembled or unidentified picornaviruses.

Discussion

We set out to explore the possibility of using viromics to find human pathogenic RNA viruses in the environment. We have been successful in identifying several potentially human pathogenic or zoonotic viral genomes from the wastewater samples, but did not find any in the surface estuarine water and sediment samples. The absence of signatures does not necessarily mean that there are no pathogenic viruses present in water or sediment, but possibly that their levels are below our limit of quantification (approximately 200 gc/l).

It is important to note here that during the RNA extraction process, many biases could have been introduced leading to a lower recovery of input viruses. Samples were first concentrated from volumes of 1 l (wastewater) or 50 l (surface water) down to 50 ml using tangential flow filtration (TFF) at a molecular weight cut-off of 100 kDa, followed by PEG 6000 precipitation. These samples were diluted in fresh buffer,

filtered through syringe filters of 0.22 µm pore size and then treated with nuclease to

remove free DNA and RNA. Previous research has shown that while any enrichment

method aimed at fractionating the viral and cellular components will decrease the

total quantity of viruses, a combination of centrifugation, filtration and nuclease

treatment increases the proportion of viral reads in sequencing datasets (29). After

implementing these steps, we used the MO BIO PowerViral® Environmental

DNA/RNA extraction kit for viral RNA extraction, which has previously been shown to

perform best overall in spiking experiments with murine norovirus, in terms of

extraction efficiency and removal of inhibitors (30). The kit has, however, given low

recoveries of viruses from sediment (31).

We did not perform an amplification step before library construction with the

NEBNext Ultra Directional RNA Library Prep Kit for Illumina, to retain the genome

sense and strand information. Instead, we increased the number of cycles of random

PCR during library preparation from 12 to 15 to counteract the low input quantity of

RNA (< 1 ng). The random amplification during library construction led to a trade-off

in which genome strand information was gained for a loss of quantitative power,

making it difficult to compare abundances of viral types within and across libraries.

This random PCR-based bias has been highlighted before, but the proposed solution

of using library preparation protocols which limit the use of PCR are only feasible

with high amounts of input nucleic acid (32), which we have found to be impossible

when processing environmental/wastewater samples to generate RNA metaviromes.

A critical issue to highlight here, is the inclusion of controls in our sequencing

libraries in order to identify potential contaminants and their origins, as has been

suggested previously (33, 34). There have been multiple reports of false positive

genome discoveries, in particular the reported discovery of a novel parvovirus-like

hybrid in hepatitis patients that was later revealed to originate from the silica-based

nucleic acid extraction columns (35–37). In this study, we included a positive control

that comprised bacterial cells (*Salmonella* Typhimurium isolate D23580 RefSeq

accession number NC_016854) and mengovirus (36), an RNA virus that serves as a

process control, as well as two negative controls, an extraction control and a library

preparation control. Analysis of the control libraries showed that while the *Salmonella*

cells and DNA were successfully removed from the positive control sample by the

enrichment protocol, the mengovirus was not recovered. Subsequent qRT-PCR

analysis revealed that the mengovirus remained detectable in the pre-processing

stages of the extraction, but was lost after RNase treatment (data not shown).

Inclusion of an inactivation step of the DNase at 75°C potentially exacerbated the

effect of the RNase step. Consequently, it is likely that we have missed several viral

types during the extraction process despite having still managed to recover an RNA

metavirome harbouring substantial diversity.

Further examination of the HiSeq and MiSeq control datasets revealed a wide range

of contaminant signatures of prokaryotic, eukaryotic and viral origin, making up 45M

read pairs per control on the HiSeq platform and 1M read pairs for the MiSeq, even

though the 16S and 18S rRNA PCR and RT-PCR reactions showed no visible bands

on an agarose gel. Most bacterial contaminant reads belonged to the phyla

Proteobacteria, Actinobacteria and *Firmicutes*. The most abundant genera included

Corynebacterium, Propionibacterium, Sphingomonas, Ralstonia, Pseudomonas,

Streptomyces, Staphylococcus and *Streptococcus* which have in the past been

identified as common lab contaminants (38). Within the eukaryotic signatures,

human-derived reads, *Beta vulgaris* and *Anopheles* reads were the most prevalent,

pointing towards potential cross-contamination of the sequencing libraries. A small

number of virus signatures were also identified, with the most prominent being *Feline*

calicivirus and *Dengue virus*. The presence of the calicivirus was traced back to the

library preparation kits after the libraries were reconstructed and resequenced. The

dengue virus signature was a <100 nt sequence which was co-extracted in all the

samples and potentially originated in one of the reagents or spin extraction column.

All sequences present in the controls were carefully removed from the sample

datasets during the quality control stage of the bioinformatics processing before

further analysis. For future experiments, we will omit the RNase treatment step

during extraction and filter out any contaminating ribosomal RNA or cellular-derived

mRNA sequences as part of the bioinformatic quality control workflow.

Our results show that while contamination is an issue when dealing with low biomass

samples, the combination of increased random PCR cycles during library

preparation, deep sequencing (i.e. HiSeq rather than MiSeq) and computational

subtraction of control sequences provides data of sufficient quantity and quality to

assemble near-complete RNA virus genomes *de novo*.

407   Norovirus

Noroviruses are one of the most common causes of gastrointestinal disease in the

developed world, with an incidence in the UK estimated as approaching 4 million

cases per annum (39). The genotype most commonly associated with disease is

GII.4 (40–42) which was not detected in the metaviromes generated here.

We retrieved one norovirus GI genome, assembled from 22,151 reads, in

wastewater effluent sample LE_11-10. This finding was in direct conflict with the

qRT-PCR analysis of this sample which did not detect any NoV GI signatures (Table

1). In contrast, NoV GII signatures were detected by qRT-PCR, but no NoV GII

genomes or genome fragments were observed in the virome libraries. One

hypothesis to explain the discrepancy between PCR and viromics approaches lies in

the differences in extraction protocol. For qRT-PCR, no viral enrichment step was

performed and RNA was not extracted with the PowerViral kit. Therefore, NoV GII

could have been lost before virome sequencing, as was the process control virus. An

alternative hypothesis is that the NoV GII signatures detected during qRT-PCR were

derived from fragmented RNA or from particles with a compromised capsid. In both

these cases, the RNA will not be detected in the virome data because of the RNase

preprocessing steps implemented in the enrichment/extraction protocol. This calls

into question the reliance of qRT-PCR for NoV detection and whether the detected

viruses are infectious or merely remnants of previous infections. Further research

using, for example, capsid integrity assays combined with infectious particle counts

will need to be conducted to assess the validity of qRT-PCR protocols for norovirus

detection.

The inability to identify NoV GI with qRT-PCR might be related to the mismatched

base present in the forward primer sequence used for detection. We subsequently

conducted a normal, long-range PCR to validate the detection of this genotype, and

this yielded a fragment of the correct size, but we were unable to clone and

sequence this fragment. While the known NoV GI.2 genotypes do not have a

mismatch in the qRT-PCR probe sequence, it is possible that the genome recovered

in this study fell below the limit of detection using the ISO standard primer/probe

combination (ISO/TS 15216-2:2013). In a recent study, researchers designed an

improved probe and observed lower Ct values and a lower limit of detection for GI.2

strains from waterborne samples (43). In general, viromics as a means of

investigating water samples for the presence of norovirus, does have the advantage

of unequivocally demonstrating the presence of intact genomes, provided the sample

processing requirements do not lead to excessive loss of viruses resulting in false

negatives. Certainly, time and money permitting, viromics is a useful adjunct to

qPCR for samples that are deemed particularly important or critical for determination

of intact viral genome presence.

Due to the difficulty of culturing noroviruses in the lab, many studies have used male-

specific coliphages such as MS2 and GA, which are ssRNA phages belonging to the

family *Leviviridae*, as alternative model systems (44, 45). Interestingly, while some

levivirus signatures were present in all wastewater samples (< 500 reads), we

observed a striking co-occurrence of these viruses with norovirus signatures in both

libraries of sample LE_11-10 (> 2500 reads). The most commonly observed viruses

in this sample were Pseudomonas phage PRR1, an unclassified levivirus, and

Escherichia phages FI and M11 in the genus *Allolevivirus*. Further studies with more

samples and replicates will indicate whether there is a significant correlation between

the presence of leviviruses and noroviruses in water samples. Furthermore, the

higher abundance of alloleviviruses compared with MS2-like viruses could indicate

that the former might be more relevant as model systems for noroviruses.

Rotavirus

Rotaviruses are, like noroviruses, agents of gastroenteritis, but the disease is

commonly associated with children under the age of 5 where severe diarrhea and

vomiting can lead to over 10,000 hospitalizations per year in England and Wales

(46). Since the introduction of the live-attenuated vaccine Rotarix, the incidence of

gastroenteritis in England has declined, specifically for children aged <2 and during

peak rotavirus seasons (47–49). Therefore, the discovery of a diverse assemblage of

rotavirus genome segments in the wastewater samples here was less expected than

the norovirus discovery. While we were unable to recover the genome of the vaccine

strain, our genomic evidence suggests that at least one RVA and one RVC

population were circulating in the Llanrwst region in September 2016.

The genome constellation for the RVA segments in sample LI_13-9, G8/G10-

P[1]/P[14]/P[41]-I2-R2-C2-M2-A3/A11-(N)-T6-E2-(H), is distinctly bovine in origin

(25) (N and H segments not recovered in this study). The closest genome segment

relatives based on nucleic acid similarity, however, have been isolated from humans

(Table 2), likely pointing towards a bovine-human zoonotic transmission of this virus

(50). The same genomic constellation has been found recently when unusual

G8P[14] RVA isolates were recovered from human strain collections in Hungary (51)

and Guatemala (52), and isolated from children in Slovenia (53) and Italy (54). Cook

and colleagues calculated that there would be approximately 5000 zoonotic human

infections per year in the UK from livestock transmission, but many would be

asymptomatic (55).

The origins of the RVC genome segments are more difficult to trace, because of

lower similarity scores with known RVC isolates. The majority of the segments were

similar to porcine RVC genomes, while others showed no nucleotide similarity at all,

only amino acid similarity. An explanation for the presence of pig-derived rotavirus

signatures can be farm run-off. While farm waste is not supposed to end up in the

sewage treatment plant, it is likely that the RVC segments originate directly from

pigs, not through zoonotic transfer. Run-off from fields onto public roads, broken

farm sewer pipes or polluted small streams might lead to porcine viruses entering the

human sewerage network, but we cannot provide formal proof from the data

available. Based on the evidence, we hypothesize that there is one, possibly two,

divergent strains of RVC circulating in the pig farms in the Llanrwst area.

Conclusion

In this study, we investigated the use of metagenomics for the discovery of RNA

viruses circulating in watercourses. We have found RNA viruses in all samples

tested, but potential human pathogenic viruses were only identified in wastewater.

The recovery of plant viruses in most samples points towards potential applications

in crop protection, for example the use of metaviromics in phytopathogen

diagnostics. However, technical limitations, including the amount of input material

necessary and contamination of essential laboratory consumables and reagents, are

currently the main bottleneck for the adoption of fine scale metagenomics in routine

monitoring and diagnostics. The discovery of a norovirus GI and a diverse set of

rotavirus segments in the corresponding metaviromes indicates that qPCR-based

approaches can miss a significant portion of relevant pathogenic RNA viruses

present in water samples. Therefore, metagenomics can, at this time, best be used

for exploration, to design new diagnostic markers/primers targeting novel genotypes

and to inform diagnostic surveys on the inclusion of specific additional target viruses.

Materials & Methods

Sample collection and processing

Wastewater samples were collected as part of a viral surveillance study described elsewhere (Farkas et al, in submission). Wastewater influent and effluent, 1 L each, was collected at the Llanrwst wastewater treatment plant by Welsh Water (Wales, UK, Figure 1) on 12[th] September (processed on 13-9, sample designations LI_13-9 and LE_13-9) and 10[th] October 2016 (processed on 11-10, sample designations LI_11-10 and LE_11-10). The wastewater treatment plant uses filter beds for secondary treatment and serves approx. 4000 inhabitants. The estuarine surface water (50 L) sample (SW) was collected at Morfa Beach (Conwy, Wales, Figure 1) approx. 22 km downstream of the Llanrwst wastewater treatment plant on 19[th] October and 2[nd] of November 2016 at low tide (only the sample from November was used for sequencing as the October sample extract failed quality control). Together with the surface water sample, 90 g of the top 1-2 cm layer of the sediment was also collected (sample designations Sed1 for the October sample and Sed2 for the November sample).

The wastewater and surface water samples were processed using a two-step concentration method as described elsewhere (Farkas et al, in submission). In brief, the 1l (wastewater) and 50l (surface water) samples were first concentrated down to 50 ml using a KrosFlo® Research IIi Tangential Flow Filtration System (Spectrumlabs, USA) with a 100 PEWS membrane. Particulate matter was then eluted from solid matter in the concentrates using beef extract buffer and then viruses were precipitated using polyethylene glycol (PEG) 6000. The viruses from the sediment samples were eluted and concentrated using beef extract elution and

532   PEG precipitation as described elsewhere (31). The precipitates were eluted in 2-10

533   mL phosphate saline buffer, (PBS, pH 7.4) and stored at -80°C.

Detection and quantification of enteric viruses with qRT-PCR

Total nucleic acids were extracted from a 0.5 mL aliquot of the concentrates using

the MiniMag NucliSENS® MiniMag® Nucleic Acid Purification System (bioMérieux

SA, France). The final volume of the nucleic acid solution was 0.05 mL (surface

water and sediment) and 0.1 mL (wastewater samples). Norovirus GI and GII,

sapovirus GI, and hepatitis A and E viruses were targeted in qRT-PCR assays as

described elsewhere (56).

Viral RNA extraction for metaviromic sequencing

Viral particles were extracted from the concentrated samples by filtration. In a first

step, the samples were diluted in 10 ml of sterile 0.5 M NaCl buffer and incubated at

room temperature with gentle shaking for 30 min to disaggregate particles. The

suspension was then filtered through a sterile, 0.22 µm pore size syringe filter

(Millex, PES membrane). The sample was desalted by centrifugation (3200 x g,

between 1 and 6h for different samples) in a sterilized spin filter (Vivaspin 20, 100

kDa molecular weight cut-off) and replacement of the buffer solution with 5 ml of a

Tris-based buffer (10 mM TrisHCl, 10 mM $MgSO_4$, 150 mM NaCl, pH 7.5). The buffer

exchange was performed twice and the volume retained after the final spin was <

500 µl. The samples were then treated with Turbo DNase (20 Units; Ambion) and

incubated for 30 minutes at 37°C, followed by inactivation at 75°C for 10 minutes. In

a next step, all samples were treated with 80 µg RNase A (Thermo Fisher Scientific)

and incubated at 37°C for 30 minutes. The RNase was inactivated with RiboLock

RNase Inhibitor (Thermo Fisher Scientific) and the inactivated complex was removed

by spin filtration (Vivaspin 500, 100 kDa molecular weight cut-off) and the samples

centrifuged until the volume was approximately 200 μl. Viral DNA and RNA were co-

extracted using the PowerViral Environmental DNA/RNA kit (MOBIO Laboratories)

according to the manufacturer's instructions. In this protocol, buffer PV1 was

supplemented with 20 μl/ml betamercaptoethanol to further reduce RNase activity.

The nucleic acid was eluted in 100 μl RNase-free water. The extracted viral DNA

was degraded using the DNase Max kit (Mobio) according to the manufacturer's

instructions. The remaining viral RNA was further purified and concentrated by

ethanol precipitation using 2.5 x sample volume of 100% ethanol and 1/10 volume of

DEPC-treated Na-acetate (3 M). The quantity and quality of RNA was determined

with Bioanalyzer Pico RNA 6000 capillary electrophoresis (Agilent Technologies). A

positive and negative extraction control sample were processed alongside the main

samples. The positive control samples contained *Salmonella enterica* strain D23580

which is not found in the UK (57) and a process control virus mengovirus (56, 58).

The viral RNA extracts were tested for bacterial and eukaryotic cellular

contamination using 16S and 18S rRNA gene PCR and RT-PCR, with primers e9F

(59) and 519R (60), and primers 1389F and 1510R (61), for the 16S and 18S rRNA

gene, respectively. Complimentary DNA was created using the SuperScript III

Reverse Transcriptase (Invitrogen) with random hexamer primers according to the

manufacturer's instructions. (RT)-PCR was performed with the MyTaq Red Mix

(Bioline) for 35 cycles (95°C for 45 sec, 50°C for 30 sec, 72°C 1 min 40 sec) and

visualized on a 1% agarose gel. Samples were considered suitable for sequencing if

no DNA bands were visible on the gel.

Library preparation and sequencing

The library preparation and sequencing were performed at the University of Liverpool Centre for Genomics Research (CGR). Twelve dual indexed, strand-specific libraries were created using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina, according to the manufacturer's instructions. These libraries were pooled and sequenced at 2 x 150 bp read lengths on the Illumina HiSeq 4000 platform. This generated between 10 and 110 million paired reads per sample.

To confirm our results, a second set of libraries was constructed from new kits and a milliQ water samples was included as a library prep control. The thirteen resulting libraries were sequenced on the Illumina MiSeq platform at CGR, at 2 x 150 bp read lengths. These data were used for verification and control purposes only as sequencing depth was insufficient for the bioinformatics analyses described in the rest of the study.

Bioinformatics

All command line programs for data analysis were run on the bioinformatics cluster of CGR (University of Liverpool) in a Debian 5 or 7 environment.

Raw fastq files were trimmed to remove Illumina adapters using Cutadapt version 1.2.1 using option -O 3 (62) and Sickle version 1.200 with a minimum quality score of 20 (63). Further quality control was performed with Prinseq-lite (64) with the following parameters: minimum read length 35, GC percentage between 5-95%, minimum mean quality 25, dereplication (removal of identical reads, leaving 1 copy), removal of tails of minimum 5 polyN sequences from 3' and 5' ends of reads.

The positive and negative control libraries described earlier were used for contaminant removal. The reads of the control samples were analyzed using

Diamond blastx (14) against the non-redundant protein database of NCBI (nr version

November 2015). The blast results were visualized using Megan6 Community

Edition (15). An extra contaminant file was created with complete genomes of

species present at over 1000 reads in the positive and negative control samples.

Then, bowtie2 (65) was used for each sample to subtract the reads that mapped to

the positive control, negative control or contaminant file. The unmapped reads were

used for assembly with SPAdes version 3.9.0 with kmer values 21, 31, 41, 51, 61,

71, and the options --careful and a minimum coverage of 5 reads per contig (66).

The contig files of each sample were compared with the contigs of the controls

(assembled using the same parameters) using blastn of the BLAST+ suite (67).

Contigs that showed significant similarity with control contigs were manually

removed, creating a curated contig dataset. The unmapped read datasets were then

mapped against this curated contig dataset with bowtie2 and only the reads that

mapped were retained, resulting in a curated read dataset.

The curated contig and read datasets were compared to the Viral RefSeq (release

January 2017) and non-redundant protein (nr, release May 2017) reference

databases using Diamond blastx at an e value of 1e-5 for significant hits (14, 68, 69).

Taxon assignments were made with Megan6 Community Edition according to the

lowest common ancestor algorithm at default settings (15). The taxon abundance

data were extracted from Megan6 and imported into RStudio for visualization (70).

Genes were predicted on the assembled contigs with Prokka (71) using the settings -

-kingdom Viruses and an e value of 1e-5. Multiple alignments of genes and genomes

were made in MEGA7 using the MUSCLE algorithm at default settings (72, 73). The

alignments were manually trimmed and phylogenetic trees were built using the

Maximum Likelihood method in MEGA7 at the default settings.

Accession numbers

Read and contig datasets are available from NCBI under the following BioProject accession numbers, PRNJA421889 (wastewater data), PRNJA421892 (sediment data) and PRJNA421894 (estuarine water data). The NoV GI genome isolate was deposited in GenBank under accession number MG599789.

Author contributions

EMA, KF, DJ, HA and AJM designed the experiments, EMA, KF, CH, performed the experiments, EMA analysed the data, EMA and KF wrote the manuscript and EMA prepared the manuscript for submission. All authors critically reviewed and edited the manuscript.

Acknowledgements

This study was funded by the Natural Environment Research Council (NERC) and the Food Standards Agency (FSA) under the Environmental Microbiology and Human Health (EMHH) Programme (NE/M010996/1). The authors gratefully acknowledge Dr James Lowther (Centre for Environment, Fisheries and Aquaculture Science; CEFAS) for providing the mengovirus sample. We also thank Gordon Steffen and Dr Nick Barcock (Dŵr Cymru Cyf - Welsh Water Ltd, UK) for facilitating sample collection at the wastewater treatment plants and Dr Julie Webb (Bangor University, UK) for assistance in sampling.

References

1.　Lin J, Ganesh A. 2013. Water quality indicators: bacteria, coliphages, enteric
viruses. Int J Environ Health Res 23:484–506.

2.　Girones R, Ferrús MA, Alonso JL, Rodriguez-Manzano J, Calgua B, de Abreu
Corrêa A, Hundesa A, Carratala A, Bofill-Mas S. 2010. Molecular detection of
pathogens in water - The pros and cons of molecular techniques. Water Res
44:4325–4339.

3.　Laverick MA, Wyn-Jones AP, Carter MJ. 2004. Quantitative RT-PCR for the
enumeration of noroviruses (Norwalk-like viruses) in water and sewage. Lett
Appl Microbiol 39:127–136.

4.　Rodriguez-Manzano J, Miagostovich M, Hundesa A, Clemente-Casares P,
Carratala A, Buti M, Jardi R, Girones R. 2010. Analysis of the evolution in the
circulation of HAV and HEV in Eastern Spain by testing urban sewage
samples. J Water Health 8:346–354.

5.　Schvoerer E, Ventura M, Dubos O, Cazaux G, Serceau R, Gournier N, Dubois
V, Caminade P, Fleury HJA, Lafon ME. 2001. Qualitative and quantitative
molecular detection of enteroviruses in water from bathing areas and from a
sewage treatment plant. Res Microbiol 152:179–186.

6.　Fong TT, Phanikumar MS, Xagoraraki I, Rose JB. 2010. Quantitative detection
of human adenoviruses in wastewater and combined sewer overflows
influencing a Michigan river. Appl Environ Microbiol 76:715–723.

7.　Bofill-Mas S, Albinana-Gimenez N, Clemente-Casares P, Hundesa A,
Rodriguez-Manzano J, Allard A, Calvo M, Girones R. 2006. Quantification and
stability of human adenoviruses and polyomavirus JCPyV in wastewater
matrices. Appl Environ Microbiol 72:7894–7896.

8.　Hellmér M, Paxéus N, Magnius L, Enache L, Arnholm B, Johansson A,
Bergström T, Norder H. 2014. Detection of pathogenic viruses in sewage
provided early warnings of hepatitis A virus and norovirus outbreaks. Appl
Environ Microbiol 80:6771–6781.

9.　Nieuwenhuijse DF, Koopmans MPG. 2017. Metagenomic Sequencing for
Surveillance of Food- and Waterborne Viral Diseases. Front Microbiol 8:1–11.

10.　Symonds EM, Breitbart M. 2015. Affordable enteric virus detection techniques
are needed to support changing paradigms in water quality management.
Clean - Soil, Air, Water 43:8–12.

11.　Rosario K, Symonds EM, Sinigalliano C, Stewart J, Breitbart M. 2009. Pepper
mild mottle virus as an indicator of fecal pollution. Appl Environ Microbiol
75:7261–7267.

12.　Stachler E, Bibby K. 2014. Metagenomic evaluation of the highly abundant
human gut bacteriophage CrAssphage for source tracking of human fecal
pollution. Environ Sci Technol Lett 1:405–409.

13.   Bibby K. 2013. Metagenomic identification of viral pathogens. Trends
689   Biotechnol 31:275–9.

690   14.   Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment
691   using DIAMOND. Nat Methods 12:59–60.

692   15.   Huson DH, Weber N. 2013. Microbial community analysis using MEGAN.
693   Methods Enzymol 531:465–85.

694   16.   2012. Virus taxonomy, 9th ed. Elsevier Inc., London, UK.

695   17.   Ye Y, Ellenberg RM, Graham KE, Wigginton KR. 2016. Survivability,
696   partitioning, and recovery of enveloped viruses in untreated municipal w
697   astewater. Environ Sci Technol 50:5077–5085.

698   18.   Winterbourn JB, Clements K, Lowther JA, Malham SK, McDonald JE, Jones
699   DL. 2016. Use of Mytilus edulis biosentinels to investigate spatial patterns of
700   norovirus and faecal indicator organism contamination around coastal sewage
701   discharges. Water Res 105:241–250.

702   19.   Patel MM, Hall AJ, Vinjé J, Parashar UD. 2009. Noroviruses: A comprehensive
703   review. J Clin Virol 44:1–8.

704   20.   Zheng DP, Ando T, Fankhauser RL, Beard RS, Glass RI, Monroe SS. 2006.
705   Norovirus classification and proposed strain nomenclature. Virology 346:312–
706   323.

707   21.   Huo Y, Cai A, Yang H, Zhou M, Yan J, Liu D, Shen S. 2014. Complete
708   nucleotide sequence of a norovirus GII.4 genotype: Evidence for the spread of
709   the newly emerged pandemic Sydney 2012 strain to China. Virus Genes
710   48:356–360.

711   22.   Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison
712   visualizer. Bioinformatics 27:1009–1010.

713   23.   Small C, Barro M, Brown TL, Patton JT. 2007. Genome heterogeneity of SA11
714   rotavirus due to reassortment with " O " agent. Virology 359:415–424.

715   24.   Chen Z, Lambden PR, Lau J, Caul EO, Clarke IN. 2002. Human group C
716   rotavirus : completion of the genome sequence and gene coding assignments
717   of a non-cultivatable rotavirus. Virus Res 83:179–187.

718   25.   Matthijnssens J, Ciarlet M, Heiman E, Arijs I, Delbeke T, McDonald SM,
719   Palombo EA, Iturriza-Gomara M, Maes P, Patton JT, Rahman M, Van Ranst
720   M. 2008. Full genome-based classification of rotaviruses reveals a common
721   origin between human Wa-Like and porcine rotavirus strains and human DS-1-
722   like and bovine rotavirus strains. J Virol 82:3204–3219.

723   26.   Matthijnssens J, Ciarlet M, Rahman M, Attoui H, Bányai K, Estes MK, Gentsch
724   JR, Iturriza-Gómara M, Kirkwood CD, Martella V, Mertens PPC, Nakagomi O,
725   Patton JT, Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K,
726   Desselberger U, Van Ranst M. 2008. Recommendations for the classification
727   of group a rotaviruses using all 11 genomic RNA segments. Arch Virol
728   153:1621–1629.

27.　Ganesh B, Bányai K, Martella V, Jakab F, Masachessi G, Kobayashi N. 2012.
Picobirnavirus infections: viral persistence and zoonotic potential. Rev Med
Virol 22:245–256.

28.　Hamza IA, Jurzik L, Überla K, Wilhelm M. 2011. Evaluation of pepper mild
mottle virus, human picobirnavirus and Torque teno virus as indicators of fecal
contamination in river water. Water Res 45:1358–1368.

29.　Hall RJ, Wang J, Todd AK, Bissielo AB, Yen S, Strydom H, Moore NE, Ren X,
Huang QS, Carter PE, Peacey M. 2014. Evaluation of rapid and simple
techniques for the enrichment of viruses prior to metagenomic virus discovery.
J Virol Methods 195:194–204.

30.　Iker BC, Bright KR, Pepper IL, Gerba CP, Kitajima M. 2013. Evaluation of
commercial kits for the extraction and purification of viral nucleic acids from
environmental and fecal samples. J Virol Methods 191:24–30.

31.　Farkas K, Hassard F, McDonald JE, Malham SK, Jones DL. 2017. Evaluation
of molecular methods for the detection and quantification of pathogen-derived
nucleic acids in sediment. Front Microbiol 8:53.

32.　Van Dijk EL, Jaszczyszyn Y, Thermes C. 2014. Library preparation methods
for next-generation sequencing: Tone down the bias. Exp Cell Res 322:12–20.

33.　Weiss S, Amir A, Hyde ER, Metcalf JL, Song SJ, Knight R. 2014. Tracking
down the sources of experimental contamination in microbiome studies.
Genome Biol 15:564.

34.　Strong MJ, Xu G, Morici L, Splinter Bon-Durant S, Baddoo M, Lin Z, Fewell C,
Taylor CM, Flemington EK. 2014. Microbial contamination in next generation
sequencing: Implications for sequence-based analysis of clinical samples.
PLoS Pathog 10:e1004437.

35.　Zhi N, Hu G, Wan Z, Zheng X, Liu X, Wong S, Kajigaya S, Zhao K, Young NS,
Africa S. 2014. Correction for Xu et al., Hybrid DNA virus in Chinese patients
with seronegative hepatitis discovered by deep sequencing. Proc Natl Acad
Sci 111:4344–4345.

36.　Zhi N, Hu G, Wong S, Zhao K, Mao Q, Young NS. 2014. Reply to Naccache et
al: Viral sequences of NIH-CQV virus, a contamination of DNA extraction
method. Proc Natl Acad Sci 111:E977–E977.

37.　Naccache SN, Greninger AL, Lee D, Coffey LL, Phan T, Rein-Weston A,
Aronsohn A, Hackett J, Delwart EL, Chiu CY. 2013. The perils of pathogen
discovery: Origin of a novel Parvovirus-like hybrid genome traced to nucleic
acid extraction spin columns. J Virol 87:11966–11977.

38.　Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P,
Parkhill J, Loman NJ, Walker AW. 2014. Reagent and laboratory
contamination can critically impact sequence-based microbiome analyses.
BMC Biol 12:87.

39.　Harris JP, Iturriza-Gomara M, O'Brien SJ. 2017. Re-assessing the total burden

of norovirus circulating in the United Kingdom population. Vaccine 35:853–
855.

40. Siebenga JJ, Vennema H, Zheng D, Vinjé J, Lee BE, Pang X, Ho ECM, Lim
W, Choudekar A, Broor S, Halperin T, Rasool NBG, Hewitt J, Greening GE, Jin
M, Duan Z, Lucero Y, O'Ryan M, Hoehne M, Schreier E, Ratcliff RM, White
PA, Iritani N, Reuter G, Koopmans M. 2009. Norovirus Illness Is a Global
Problem: Emergence and Spread of Norovirus GII.4 Variants, 2001–2007. J
Infect Dis 200:802–812.

41. Eden J-S, Tanaka MM, Boni MF, Rawlinson WD, White PA. 2013.
Recombination within the Pandemic Norovirus GII.4 Lineage. J Virol 87:6270–
6282.

42. Cannon JL, Barclay L, Collins NR, Wikswo ME, Castro CJ, Magaña LC,
Gregoricus N, Marine RL, Chhabra P, Vinjé J. 2017. Genetic and
Epidemiologic Trends of Norovirus Outbreaks in the United States from 2013
to 2016 Demonstrated Emergence of Novel GII.4 Recombinant Viruses. J Clin
Microbiol 55:2208–2221.

43. Cho H-G, Lee S-G, Mun S-K, Lee M-J, Park P-H, Jheong W-H, Yoon M-H,
Paik S-Y. 2017. Detection of waterborne norovirus genogroup I strains using
an improved real time RT-PCR assay. Arch Virol 162:3389–3396.

44. Dunkin N, Weng S, Coulter CG, Jacangelo JG, Schwab KJ. 2017. Reduction
of Human Norovirus GI, GII, and Surrogates by Peracetic Acid and
Monochloramine in Municipal Secondary Wastewater Effluent. Environ Sci
Technol 51:11918–11927.

45. Arredondo-Hernandez LJR, Diaz-Avalos C, Lopez-Vidal Y, Castillo-Rojas G,
Mazari-Hiriart M. 2017. FRNA Bacteriophages as Viral Indicators of Faecal
Contamination in Mexican Tropical Aquatic Systems. PLoS One 12:e0170399.

46. Harris JP, Jit M, Cooper D, Edmunds WJ. 2007. Evaluating rotavirus
vaccination in England and Wales. Part I. Estimating the burden of disease.
Vaccine 25:3962–3970.

47. Bawa Z, Elliot AJ, Morbey RA, Ladhani S, Cunliffe NA, O'Brien SJ, Regan M,
Smith GE, Weinstein RA. 2015. Assessing the likely impact of a rotavirus
vaccination program in England: The contribution of syndromic surveillance.
Clin Infect Dis 61:77–85.

48. Thomas SL, Walker JL, Fenty J, Atkins KE, Elliot AJ, Hughes HE, Stowe J,
Ladhani S, Andrews NJ. 2017. Impact of the national rotavirus vaccination
programme on acute gastroenteritis in England and associated costs averted.
Vaccine 35:680–686.

49. Hungerford D, Read JM, Cooke RPD, Vivancos R, Iturriza-G??mara M, Allen
DJ, French N, Cunliffe N. 2016. Early impact of rotavirus vaccination in a large
paediatric hospital in the UK. J Hosp Infect 93:117–120.

50. Wilhelm B, Waddell L, Greig J, Rajić A, Houde A, McEwen SA. 2015. A
scoping review of the evidence for public health risks of three emerging

potentially zoonotic viruses: Hepatitis E virus, norovirus, and rotavirus. Prev
Vet Med 119:61–79.

51.   Marton S, Doro R, Feher E, Forro B, Ihasz K, Varga-Kugler R, Farkas SL,
815   Banyai K. 2017. Whole genome sequencing of a rare rotavirus from archived
816   stool sample demonstrates independent zoonotic origin of human G8P[14]
817   strains in Hungary. Virus Res 227:96–103.

818   52.   Gautam R, Mijatovic-Rustempasic S, Roy S, Esona MD, Lopez B, Mencos Y,
819   Rey-Benito G, Bowen MD. 2015. Full genomic characterization and
820   phylogenetic analysis of a zoonotic human G8P[14] rotavirus strain detected in
821   a sample from Guatemala. Infect Genet Evol 33:206–211.

822   53.   Steyer A, Naglič T, Jamnikar-Ciglenečki U, Kuhar U. 2017. Detection and
823   Whole-Genome Analysis of a Zoonotic G8P[14] Rotavirus Strain Isolated from
824   a Child with Diarrhea. Genome Announc 5:e01053-17.

825   54.   Medici MC, Tummolo F, Bonica MB, Heylen E, Zeller M, Calderaro A,
826   Matthijnssens J. 2015. Genetic diversity in three bovine-like human G8P[14]
827   and G10P[14] rotaviruses suggests independent interspecies transmission
828   events. J Gen Virol 96:1161–1168.

829   55.   Cook N, Bridger J, Kendall K, Gomara MI, El-Attar L, Gray J. 2004. The
830   zoonotic potential of rotavirus. J Infect 48:289–302.

831   56.   Farkas K, Peters DE, McDonald JE, de Rougemont A, Malham SK, Jones DL.
832   2017. Evaluation of Two Triplex One-Step qRT-PCR Assays for the
833   Quantification of Human Enteric Viruses in Environmental Samples. Food
834   Environ Virol 9:342–349.

835   57.   Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, Harris
836   D, Clarke L, Whitehead S, Sangal V, Marsh K, Achtman M, Molyneux ME,
837   Cormican M, Parkhill J, MacLennan CA, Heyderman RS, Dougan G. 2009.
838   Epidemic multiple drug resistant Salmonella Typhimurium causing invasive
839   disease in sub-Saharan Africa have a distinct genotype. Genome Res
840   19:2279–2287.

841   58.   Hennechart-Collette C, Martin-Latil S, Guillier L, Perelle S. 2015.
842   Determination of which virus to use as a process control when testing for the
843   presence of hepatitis A virus and norovirus in food and water. Int J Food
844   Microbiol 202:57–65.

845   59.   Reysenbach A, Pace N. 1995. Reliable amplification of hyperthermophilic
846   archaeal 16S rRNA genes by the polymerase chain reaction, p. 101–107. *In*
847   Robb, F, Place, A (eds.), Archaea: a laboratory manual. Cold Spring Harbor
848   Laboratory Press, New York, NY, USA.

849   60.   Turner S, Pryer KM, Miao VP, Palmer JD. 1999. Investigating deep
850   phylogenetic relationships among cyanobacteria and plastids by small subunit
851   rRNA sequence analysis. J Eukaryot Microbiol 46:327–338.

852   61.   Amaral-Zettler LA, McCliment EA, Ducklow HW, Huse SM. 2009. A method for
853   studying protistan diversity using massively parallel sequencing of V9

hypervariable regions of small-subunit ribosomal RNA Genes. PLoS One 4:1–
9.

62.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal 17:10–12.

63.   Joshi N, Fass J. 2011. Sickle: A sliding-window, adaptive, quality-based
trimming tool for FastQ files (Version 1.33) [Software].

64.   Schmieder R, Edwards R. 2011. Quality control and preprocessing of
metagenomic datasets. Bioinformatics 27:863–864.

65.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2.
Nat Methods 9:357–359.

66.   Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, Lapidus A,
Prjibelsky A, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, McLean J,
Lasken R, Clingenpeel SR, Woyke T, Tesler G, Alekseyev MA, Pevzner PA.
2013. Assembling genomes and mini-metagenomes from highly chimeric
reads, p. 158–170. *In* Deng, M, Jiang, R, Sun, F, Zhang, X (eds.), Research in
Computational Molecular Biology. RECOMB 2013. Lecture Notes in Computer
Science. Springer, Berlin, Heidelberg.

67.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K,
Madden TL. 2009. BLAST+: architecture and applications. BMC Bioinformatics
10:421.

68.   Brister JR, Ako-adjei D, Bao Y, Blinkova O. 2015. NCBI Viral Genomes
Resource. Nucleic Acids Res 43:D571–D577.

69.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B,
Robbertse B, Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y,
Blinkova O, Brover V, Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM,
Goldfarb T, Gupta T, Haft D, Hatcher E, Hlavina W, Joardar VS, Kodali VK, Li
W, Maglott D, Masterson P, McGarvey KM, Murphy MR, O'Neill K, Pujar S,
Rangwala SH, Rausch D, Riddick LD, Schoch C, Shkeda A, Storz SS, Sun H,
Thibaud-Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C, Webb D, Wu W,
Landrum MJ, Kimchi A, Tatusova T, DiCuccio M, Kitts P, Murphy TD, Pruitt
KD. 2016. Reference sequence (RefSeq) database at NCBI: current status,
taxonomic expansion, and functional annotation. Nucleic Acids Res 44:D733–
D745.

70.   Racine JS. 2012. RStudio: A platform-independent IDE for R and Sweave. J
Appl Econom 27:167–172.

71.   Seemann T. 2014. Prokka: Rapid prokaryotic genome annotation.
Bioinformatics 30:2068–2069.

72.   Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy
and high throughput. Nucleic Acids Res 32:1792–1797.

73.   Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary
Genetics Analysis version 7.0 for bigger datasets. Mol Biol Evol 33:msw054.