Metatranscriptome analysis of the vaginal microbiota reveals potential mechanisms for protection against metronidazole in bacterial vaginosis

Zhi-Luo Deng[a#], Cornelia Gottschick[a*], Sabin Bhuju[b], Clarissa Masur[c], Christoph Abels[c],

and Irene Wagner-Döbler[a]

[a] Research Group Microbial Communication, Department of Medical Microbiology, Helmholtz Centre for Infection Research, Inhoffenstr. 7, 38124 Braunschweig, Germany

[b] Genome Analytics, Helmholtz Centre for Infection Research, Inhoffenstr. 7, 38124 Braunschweig, Germany

[c] Dr. August Wolff GmbH & Co. KG Arzneimittel, Sudbrackstrasse 56, 33611, Bielefeld, Germany

Running title: Functional activity profiling of the vaginal microbiota during metronidazole treatment of bacterial vaginosis

Address correspondence to Zhi-Luo Deng, Zhiluo.Deng@helmholtz-hzi.de

Z.-L.D. and C.G. contributed equally to this work.

* Present address: Institute for Medical Epidemiology, Biometrics and Computer Science, Martin-Luther-University Halle-Wittenberg, Halle (Saale), Germany

Abstract

Bacterial vaginosis (BV) is a prevalent multifactorial disease of women in their reproductive

years characterized by a shift from the healthy *Lactobacillus sp.* dominated microbial

community towards a highly diverse anaerobic community. BV can initially be cured by

antibiotic therapy in most women, but the high recurrence frequency represents a serious

challenge. Moreover, for unknown reasons, some women do not respond to therapy. In a

clinical study, out of 37 women diagnosed with BV, 31 were successfully treated with a single

peroral dose of metronidazole, while 6 still had BV after treatment. Here, we performed a

metatranscriptome analysis of the vaginal microbiota of the subgroup of those non-responding

patients (n = 6), comparing them to patients who responded (n = 8) over three months (5 time-

points). In health, *Lactobacillus iners* and *L. crispatus* contributed more than 90% of all

bacterial transcripts, and the communities were dominated by either one or the other. Their

functional activity profiles *in vivo* showed that the inerolysin was highly expressed in *L. iners*

and hydrogen peroxide producing genes were among the most highly expressed genes in *L.

crispatus*. In BV, *Gardnerella vaginalis* contributed on average 40% of all transcripts. In non-

responding patients, 7 out of 8 Cas genes and one DNA methyltransferase gene of *G.

vaginalis* were highly up-regulated, suggesting that Cas genes of *G. vaginalis*, in addition to

protecting against phages, might be involved in DNA repair thus mitigating the bactericidal

effect of DNA damaging agents like metronidazole.

Importance:

The vaginal microbiota is relatively uniform in health but shifts towards diverse during

bacterial vaginosis (BV). BV is a prevalent multifactorial disease of women in their

reproductive years and usually treated with antibiotics like metronidazole. The functional

alterations of the vaginal microbiota during the treatment have never been discovered, despite

many studies have identified the key vaginal microbial members in health and BV. Our

previous study using 16S rRNA gene sequencing approach failed to detect the difference

between patients responded to the treatment and non-responders. By using metatranscriptome,

we found out of 8 Cas genes in *G. vaginalis*, 7 were highly up-regulated in non-responding

patients. This finding suggests Cas genes of *G. vaginalis*, might play a role in defense against

DNA damaging agents like metronidazole. The *in vivo* functional activity profiles of

Lactobacillus iners and *L. crispatus* confirmed the pathogenic role of *L. iners* and protective

role of the latter.

Introduction

The healthy vaginal microbiome (VMB) is characterized by low pH and low diversity and can be categorized into community state types (CSTs) that are dominated by different *Lactobacillus sp.* such as *L. crispatus*, *L. iners*, *L. gasseri* and less frequently *L. jensenii* or a more diverse community (47). The shift of VMB towards dysbiosis has been shown associated with bacterial vaginosis (BV) (54), preterm birth (7,24) and neonatal sepsis (7). BV is a frequent multifactorial disease of women in their reproductive years that is characterized by a shift of this *Lactobacillus sp.* dominated bacterial community to a community of various mostly anaerobic bacteria (54). BV is related to a higher risk of acquiring sexually transmitted infections such as HIV (42). The most common bacteria found in BV, identified by 16S rRNA gene sequencing, are *Gardnerella*, *Atopobium*, *Prevotella*, *Bacteroides*, *Peptostreptococcus*, *Mobiluncus*, *Sneathia*, *Leptotrichia*, *Mycoplasma* and BV associated bacterium 1 (BVAB1) to BVAB3 of the order Clostridiales. Recently, three CSTs dominated by *Gardnerella vaginalis*, Lachnospiraceae and *Sneathia sanguinegens*, respectively, have been described (11) and our recent study identified *S. amnii* as the best biomarker for BV (16).

Studies have shown treatment with antibiotic metronidazole, tinidazole or/plus probiotics increased the relative abundance of *Lactobacillus* species in VMB (27,31,36). Metronidazole is a widely applied chemotherapeutic agent used to treat infectious diseases caused by anaerobic bacteria, and it is the first-line antibiotic for treating BV (6,28). It is a prodrug which requires enzymatic reduction within the cell to transform it into an active form (37). Metronidazole acts by damaging DNA, and resistance mechanisms have been studied in various pathogens (37). Previous studies have shown that failure of BV treatment by metronidazole is relatively rare (16,20), and it is unclear if it is caused by resistance of the BV pathogens to metronidazole, and which mechanisms are acting *in vivo*. A recent study has

demonstrated that failure of treatment of BV with metronidazole is not associated with higher

loads of *G. vaginalis* and *A. vaginae* (11). Therefore, the lack of response may result from the

activity of the community instead of the abundance of the pathogens and their taxonomic

composition. The high rate of recurrence is another crucial problem for BV treatment. It has

been reported that the one-year recurrence rate of BV ranges from 40% to 80% after treatment

with metronidazole (5). The recurrent rate is similar for other therapies such as clindamycin

cream (50). It has been hypothesized that patches of surviving bacterial biofilm may be one of

the reasons for recurrence (5). The predominant species of this multi-species biofilm is *G.*

vaginalis (30). *G. vaginalis* is frequently found in biofilms during BV but it is also a member

of the healthy VMB (53). *L. iners* is characteristic for a community state type in health but

may be disadvantageous when the community shifts to a dysbiotic state (32,43,53). Until now,

the majority of studies regarding the VMB have focused on 16S rRNA gene sequencing,

answering only questions on the taxonomic composition of bacterial communities but not on

their functions (54). A metatranscriptome analysis comparing vaginal swabs from two women

with BV with two healthy subjects showed that *L. iners* upregulates transcription of the

cholesterol-dependent cytolysin (CDC) and of genes belonging to the clustered regularly

interspaced short palindromic repeats (CRISPR) system in BV (32). No study has investigated

the functional shifts of the VMB during antibiotic treatment of BV.

We had previously analyzed the VMB in the context of a clinical trial using 16S rRNA gene

sequencing (16). Of 37 patients diagnosed with BV and included in this study, 31 were

initially cured by a single oral dose of metronidazole. Six patients did not respond, i.e. they

were diagnosed with BV according to Nugent score at visit 2. Moreover, some patients

developed recurrent BV during the three months of the study. Since we did not find

differences in the taxonomic composition of the microbiota of non-responding and recurrent

patients, we hypothesized that shifts in microbiota activity might be responsible for those

different responses. We therefore chose the non-responding patients (n = 6) from the clinical

study and compared their metatranscriptomes to those of 8 patients that were initially cured.

For the analysis of recurrence, we chose two recurrent patients from the lactic acid arm of the

study (see below) and compared their metatranscriptomes to those of two cured patients that

were also provided with lactic acid pessaries, and followed their transcriptomes over the three

months of the study (5 time-points). Our analysis is thus comprised of 14 patients, of which

six were cured, six did not respond to treatment, and two developed recurrence. Including the

time series samples, we analyzed 40 metatranscriptome samples in total.

We could identify genes potentially leading to lack of response to metronidazole treatment

and suggest a novel mechanism of *G. vaginalis* to mitigate the DNA damaging effect of

metronidazole. Moreover, we demonstrate the importance of *G. vaginalis* for BV, which can

be massively underestimated using 16S rRNA gene sequencing, confirm the beneficial role of

L. crispatus in comparison to *L. iners*.

Results

Study population and overview of sequencing results

We studied the vaginal microbiome of 14 patients during and after metronidazole treatment of

BV using metatranscriptome sequencing (Fig. 1). Patients were part of a clinical trial

described elsewhere (16). For ten patients, samples were derived from two time points (before

(visit 1) and after metronidazole treatment (visit 2). Six of those patients did not respond to

treatment with the antibiotic, so were still BV positive according to Nugent score and four of

them responded to the treatment and were cured. For the other four patients, 5 time points

were analyzed, covering a period of 3 months after treatment. Two of those patients

experienced recurrence, while the other two were cured without recurrence (details in Table

S1 sheet 1). In total, we analyzed 40 vaginal fluid samples, 22 with BV status and 18 without.

Metatranscriptome sequencing resulted in a total of 1,879,945,342 reads. Of these,

1,377,516,082 reads (73%) were left after quality filtering and removal of ribosomal RNA

(Table 1). On average, 34 million reads were analyzed per sample.

Construction of the reference genome and gene databases for taxonomic and functional

activity profiling

Based on the standard Kraken (58) database, ~11% of the total putative mRNA reads could be

assigned to human for BV and ~56% for health (Table S1 sheet 2). This suggests that the

bacterial load is much lower in health than in BV since the human contamination read is much

higher in health. Using this standard database, only 41% of total putative microbial (non-

human) mRNA reads could be assigned taxonomically (Table 2). Therefore, a refined Kraken

database constructed based on the genomes from HMP (38) urogenital tract database plus 16

additional genomes (see methods for details) was finally used for taxonomy classification of

putative microbial mRNA reads. Using this database, the rate of classified putative microbial

mRNA reads could be improved to 86% on average (Table 2). Interestingly, transcripts from

the most abundant 11 species contributed more than 90% of the mapped reads in BV while

the two most abundant species accounted for >90% of the classified sequences (Fig. 2), which

is much less diverse than the periodontal metatranscriptome in which more than 100 species

were required to cover 90% of mappable reads (9).

For functional assignment, we used a customized reference gene database (Table S1 sheet 4)

containing the genes from the HMP (38) urogenital tract database, and added the genes of

additional 9 genomes (see method section for details). To investigate the functional activity

shifts of the communities, we mapped the cleaned metatranscriptomic reads to this

customized reference gene database using BWA (26). In total, around 78% of total putative

microbial mRNA reads could be mapped to this microbial reference gene database with a

MAPQ >10 on average. The sequencing provided on average 8.9 million mappable microbial

mRNA reads per sample, which allowed the functional activity profiling for highly abundant

species in the community. The mapping results per sample can be seen in Table S1 sheet 2.

Shifts in the taxonomic composition of the active community during 3 months following

metronidazole treatment

The taxonomic composition of transcripts was determined using Kraken. Fig. 1 shows that all

communities with BV status (before metronidazole treatment, at recurrence, and in patients

with no response to treatment) were dominated by *G. vaginalis*, *A. vaginae*, *S. amnii* and

Prevotella timonensis. In patients that responded to treatment and were cured, the

metatranscriptomes were dominated by *L. crispatus*, *L. iners* and *L. jensenii*. Two patients

who developed recurrence were all dominated by *L. iners* after the treatment while only 3 out

of 6 patients without recurrence were dominated by *L. iners* after treatment.

The bladder was thought to be sterile in healthy women, but it has now been shown that

specific microorganisms are residing in the healthy bladder (35). Therefore, the bladder has to

be considered as a possible reservoir for pathogens. The high pH of urine may be a favorable

environment for several pathogens, including *Gardnerella sp.*. Here we observed that

Gardnerella sp. isolated from the bladder (*Gardnerells sp.* 26-12 and *Gardnerella sp.* 30-4)

(35) contributed 6% of all classified reads in BV. In several samples of BV patients,

transcripts from bladder isolates were extremely abundant (between 10-19% of all transcripts)

(samples 07_001_a/b, 13_022_a, 15_006_a/b, 17_004_a) . Some samples ((04_001_a,

06_004_c (recurrence) and 06_001_a/b, 06_006_a/b (without response to treatment)) were

dominated by *S. amnii* which has been discovered in the urine microbiome previously (17,57).

Comparison of the taxonomic composition of vaginal fluid samples between

metatranscriptome and 16S rRNA sequencing

To discover transcriptionally active bacteria which may contribute to the functional activity

shifts in BV, we compared the taxonomic composition determined using 16S rRNA

sequencing (16) and the taxonomic composition of the metatranscriptome determined by

Kraken with the refined reference database. In health, we did not observe any considerable

difference (Table S1 sheet 6), while in BV large differences between the two datasets were

found. Fig. 3A-B shows that most of the abundant species identified in the mRNA sequencing

data set were also identified using 16S rRNA gene amplicon sequencing, although usually at

different abundances. For example, *A. vaginae* comprised 15% of all reads based on 16S

rRNA gene sequencing, but only 10% in the metatranscriptome dataset. The most pronounced

difference was observed for *G. vaginalis* which comprised on average 40% of relative

abundance in the metatranscriptome and on average only 6% in the 16S rRNA sequencing

data. By contrast, the two bladder isolates of *Gardnerella sp.* were only identified in the

metatranscriptome, probably due to the low resolution of the 16S rRNA gene and the lack of

their 16S rRNA genes in the reference database. This finding suggests that *G. vaginalis* and

other *Gardnerella* species are extremely transcriptionally active in the BV community. In

addition, the commonly used 27F primer has been shown to underrepresent *G. vaginalis* (14).

***L. iners* and *L. crispatus* were dominant in health and *G. vaginalis* was the most active**

species in BV

Based on the taxonomic composition of transcripts, the samples were clearly clustered into

three groups, namely health communities dominated by *L. crispatus*, health communities

dominated by *L. iners* and BV communities (Fig. 3C). The most abundant active species after

metronidazole treatment were *Lactobacillus* species of which *L. iners* and *L. crispatus* were

the most abundant ones contributing >90% of the metatranscriptome on average. They

showed a negative correlation in occurrence (Spearman correlation coefficient =-0.851, p

=7.422e-06). High abundant *L. iners* was never accompanied by high level *L. crispatus* and

high abundant *L. crispatus* was only twice (08_003_b, 08_006_d) accompanied by low

abundant (lower than one fifth of the abundance of *L. crispatus*) *L. iners*. *L. gasseri* and *L.*

jensenii had much lower abundances and were found together with both *L. crispatus* and *L.*

iners. In one patient, the active healthy community was dominated by *L. jensenii*.

In BV, *G. vaginalis* was the most abundant active species. Only two communities of patients

with BV were dominated by other species, namely *A. vaginae* and *S. amnii*. Additional BV

associated species such as *S. sanguinegens* or different *Prevotella sp.* were frequently found

in BV samples, and M*obiluncus mulieris* was occasionally found. The only species of

Lactobacillus found to be active in BV was *L. iners*.

Functional activity profiles in health and disease

In order to profile the function of the communities, all cleaned putative mRNA reads (Table

S1 sheet 2, column C) were mapped using BWA onto a customized reference database

annotated with KEGG ortholog (KO) genes. We compared the functional activity profiles of

samples from patients in BV (22 samples) with samples from health (after successful

metronidazole treatment, 18 samples). The PCA based on the expression of KO genes clearly

showed that there was a large functional difference between BV and health; the functional

activity profiles formed clearly separated clusters (Fig. 4A). Moreover, the functional

composition in health was more variable than in BV. This is a striking finding, because 16S

rRNA gene sequencing showed a massive increase in diversity in BV (16). The reason might

be that the different species of *Lactobacillus* which dominate in different individuals in health

have different functional activity profiles, while the different species that dominate in BV all

have similar functional activity profiles.

Time course of functional activity profiles during recurrence

To identify functional difference between women with and without recurrence, the functional

activity profiles of two women with recurrence (patients 04_001 and 06_004) were compared

to those of two women without (patients 08_006 and 13_019) over a period of three months.

Five time points were analyzed of which the first represented acute BV and the others were

after metronidazole treatment. The principal components analysis of the functional activity

profiles is shown in Fig. 4B. In acute BV, samples from all four patients clustered together

(red circle). After the treatment, samples from patient 08_006 who was cured without

recurrence moved into a very dense and distinct cluster (illustrated by the arrow 1 and

enclosed by a green circle). Samples from patient 13_019 who was also cured, moved to a

different cluster after treatment, shown by arrow 2 and encircled blue.

The functional activity shifts in patient 06_004, who experienced recurrence at visit 3, were

especially noteworthy: After treatment, the community moved towards a functional activity

profile distinct from all others (arrow 3). The recurrence of BV caused the community to shift

back to the BV cluster (red circle, arrow 4). At visit 4, the community moved to the heathy

cluster (blue circle) and the patient became healthy according to the Nugent score. We

speculate that there was an unknown intervention after visit 3 which changed the microbiome

but this was not recorded. Interestingly, the other case of recurrence (patient 04_001) had a

different progression. From visit 2 to 4, patient 04_001 seemed to be cured, because these

samples clustered together in the healthy cluster indicated by the blue circle. At visit 5,

however, patient 04_001 had recurrent BV and the community shifted back again to the BV

cluster.

Interestingly, the microbial community of patient 08_006 was the only one dominated by *L.*

crispatus after metronidazole treatment, while all others, including those in the second healthy

cluster, were dominated by *L. iners*. This suggests that communities dominated by *L.*

crispatus are more protected from recurrence.

The inerolysin gene was highly expressed in *L. iners* and genes involved in the

production of H_2O_2 were highly active in *L. crispatus*

As shown above, *L. iners* and *L. crispatus* dominated different patients after treatment, and *L.*

crispatus seemed to have a protective role against recurrence. Therefore, we compared their

genomes and functional activity profiles with each other. *L. iners* has the smallest genome

encoding around 1500 proteins compared to other *Lactobacillus* species like *L. crispatus*

which encodes much more proteins (12). It is known that *L. crispatus* encodes proteins

producing hydrogen peroxide (H_2O_2) (1), while *L. iners* encodes a pore-forming toxin protein

termed inerolysin (INY) (43). Hydrogen peroxide provides protection against pathogens for

the symbiotic vaginal community (8). INY, a cholesterol-dependent cytolysin (CDC), is

activated by reducing agents and inhibited by excess cholesterol (43). CDCs were formerly

annotated as "thiol-activated cytolysins" (33). However, it is unknown that whether those

genes are transcriptionally active in the VMB *in vivo*. To gain more understanding of the

different roles of these two bacteria in the healthy vaginal microbiota, we performed a

differential expression (DE) analysis based on their KO genes comparing the expression of

KO genes of *L. crispatus* from *L. crispatus* dominated samples (n = 4) with the expression of

KO genes of *L. iners* from *L. iners* dominated samples (n = 11). The Venn diagram in Fig. 4C

shows that the two species share 569 KO genes, while 58 are unique to *L. iners* and 244 are

unique for *L. crispatus*. The DE analysis identified 584 significantly differentially expressed

KO genes, of which 355 were up-regulated in *L. crispatus* (Table S1 sheet 8). Among the top

100 most differentially expressed KO genes in terms of FDR value, 89 were up-regulated in

L. crispatus, indicating *L. crispatus* possesses far more diverse functions than *L. iners*.

Remarkably, genes assumed to be involved in the production of H_2O_2 (pyruvate oxidase,

NADH oxidase, glycolate oxidase) (19,41) were highly expressed in *L. crispatus* (Fig. 4C)

suggesting a high level of hydrogen peroxide in vaginal milieu when the VMB is dominated

by *L. crispatus.* An iron transport system was found to be highly expressed in *L. crispatus* and

it is absent in the genome of *L. iners*. On the other hand, we found that inerolysin (INY) was

highly expressed (log2 counts per million mapped reads =10.18) in *L. iners*, but absent in the

genome of *L. crispatus* (Fig. 4C, Table S1 sheet 8). Interestingly, we also found the

orthologous gene (K11031) of inerolysin known as vaginolysin (15) highly expressed in *G.*

vaginalis (details in Table S1 sheet 7). These findings show that *L. iners* exerts a similar

functional activity as the BV-associated pathogen *G. vaginalis*.

In vivo* expression of putative metronidazole resistance associated genes in *G. vaginalis

Metronidazole is a prodrug which requires anaerobic conditions for enzymatic reduction

within the cell to transform it into an active form (37). Several enzymes are involved in the

reduction of metronidazole such as ferredoxin, ferredoxin/ferredoxin-NADP reductase (FNR),

ferric uptake regulator (Fur) and nitroreductase (21,24,37,52,55). The reduced metronidazole

can introduce DNA strand breaks thus leading to the death of the pathogens. Accordingly,

genes responsible for DNA repair like recA and recA-mediated autopeptidase (Rma) might

contribute to resistance against metronidazole (51). Moreover, a gene named nitroimidazole

resistance gene (NIM) was shown to be related to the resistance against metronidazole in

Bacteroides fragilis (29). To clarify the possible contribution of these genes to metronidazole

resistance of the vaginal microbiota, we examined the expression (Table S1 sheet 10) of these

genes in *G. vaginalis* by comparing their expression in the communities with later response to

metronidazole and without at visit 1 (acute BV). For this analysis, BV communities from 10

patients were analyzed in which the level of *G. vaginalis* transcripts was >20%. Six of these

patients did not respond to treatment and four were initially cured. Although *A. vaginae* and *S.*

amnii are also key players in BV we did not analyze them here since there were two few

samples dominated by them. As shown in Fig. S1, ferredoxin, nitroreductase, Fur, FNR were

shown to be less active in *G. vaginalis* in the vaginal microbiota of no-responding patients

(Fig. S1). FNR was 4.2 times less active (Wilcoxon test P-value = 0.038, based on relative

abundance = read count of given genes of *G. vaginalis* / read count of *G. vaginalis* %) in *G.*

vaginalis in those communities. FNR is involved in activating metronidazole (21) thus its

weak expression indicates the lack of the active antibiotic, in accordance with the lack of

response in those patients. Unexpectedly, the NIM gene which was annotated with

nitroimidazole resistance gene was also less active in no-responders. RecA and RecA-

mediated autopeptidase (Rma) of *G. vaginalis* were more active in some but not all

communities from patients that did not respond to metronidazole (Fig. S1). RecA and RecA-

mediated autopeptidase are essential genes in SOS response (39) responsible for DNA repair

in bacteria. They may play a crucial role in survival of bacteria after treatment with a DNA

damaging agent like metronidazole.

CRISPR associated protein and adenine methyltransferase genes of *G. vaginalis* were
strongly up-regulated in vaginal fluids of patients not responding to treatment

We then performed a global analysis of differential expression of KO genes of *G. vaginalis* in

these same communities (visit 1, 10 BV samples with >20% transcripts from *G. vaginalis* (6

patients that did not respond to treatment and 4 that were initially cured). We observed that

there were 24 KO genes highly up-regulated with FDR <= 0.05 (log2 fold change up to 12.4)

in communities without response. Strikingly, among the top ten most strongly up-regulated

KO genes, seven were *cas* genes (44) (*cas*1-3, *cas*A-D) (Fig. 5). In total there were 8 different

G. vaginalis CRISPR-associated (Cas) genes found in its genome, namely *cas*1-3, *cas*A-E.

Cas proteins have a dual function; they can perform homology-directed repair as well as

degrade foreign DNA (3).

yhdJ and *licD* were the most strongly up-regulated genes with log2 fold change above 10, of

which *yhdJ* encodes an adenine-specific DNA-methyltransferase and *licD* encodes a

lipopolysaccharide cholinephosphotransferase. The DNA adenine methyltransferase is an

essential part of the restriction modification system that protects bacteria from digesting their

own genome via their restriction enzymes (45).

There were 32 KO genes down-regulated but the fold change values were not as high as for

the up-regulated genes. *galK* (galactokinase) was identified as the most strongly down-

regulated gene with a log2 fold change of -5.6.

These findings suggest that CRISPR-Cas, a phage defense mechanism with a dual function,

might mitigate the DNA damaging effect of metronidazole and thus may be one of the reasons

for lack of response of the vaginal microbiota to treatment. In addition, lack of expression of

enzymes (FNR, Fur, ferredoxin, nitroreductase) that help transforming the metronidazole

prodrug into the active form might also contribute to lack of response to treatment.

Discussion

The aim of this study was to identify functional activity patterns in the vaginal fluid

microbiota in health and BV. In particular, we compared metatranscriptomes of recurrent and

non-recurrent patients and of patients who did and did not respond to metronidazole

treatment. This is the first study to investigate the compositional as well as functional activity

alterations of the vaginal microbiota from patients with BV during treatment with the

antibiotic metronidazole *in vivo* using the metatranscriptomic approach. We found clear

patterns which allowed us to formulate novel hypotheses. Since our analysis is based on a

relatively small number of 14 patients, our findings should be followed up in larger patient

groups with more cases of recurrence and more cases of lack of response to antibiotic

treatment.

We observed that *G. vaginalis* was much more dominant based on the abundance of its

transcripts than based on the 16S rRNA amplicon sequencing analysis. This suggests that *G.*

vaginalis is transcriptionally more active than other vaginal bacteria; moreover, the

commonly used 27F primer was previously shown to underrepresent *G. vaginalis* (14). We

found that the functional activity profiles in health and BV were massively different.

Interestingly, *L. crispatus* and *L. iners* were the dominant species in health and communities

were dominated by either one or the other. The patients with recurrence were dominated by *L.*

iners after metronidazole treatment. There were many factors observed by laboratory or

genomic studies (13,41,46) which show that more protection is provided by *L. crispatus*

rather than by *L. iners*. *L. crispatus* has the ability to produce hydrogen peroxide to prevent

pathogen colonization, whereas *L. iners* does not (41). A recent longitudinal study also shows

that the dominance of *L. iners* in the vaginal microbiota is a risk factor for preterm birth,

whereas *L. crispatus* dominance is protective against preterm birth (22). Furthermore, an iron

transport system was found in the genome of *L. crispatus* and not in *L. iners* suggesting that

L. crispatus can capture iron released by the host during menstruation away from potential

pathogens (13). On the other hand, *L. iners* can produce inerolysin, a pore-forming cytolysin

that enables nutrient acquisition from host cells providing it with an advantage when

conditions are scarce (46), and this gene was highly expressed in *L. iners*. Strikingly, an

ortholog of inerolysin named vaginolysin was found highly expressed in *G. vaginalis* in BV.

These findings suggest that *L. iners* might initiate the dysbiotic shift of the vaginal flora and

contribute to the pathogenesis of BV as well as to recurrence.

Moreover, *Gardnerella* species isolated from the bladder were highly active in several

patients with BV, including two patients with recurrent BV. Many potential reasons are

discussed for recurrence and the occasional lack of response to metronidazole treatment. For

example, it has been reported that sexual activity can introduce pathogens thus may also

contribute to the recurrence of BV (5), dosage of metronidazole may have been too low for

the patient, or highly virulent strains of *G. vaginalis* may be residing in this particular non-

responding patient.

In our metatranscriptome analysis we found evidence for mechanisms that hinder the

activation of the metronidazole prodrug, or mitigate the damage that metronidazole inflicts on

DNA, and thus could be important reasons for the lack of response in some women.

We show for the first time *in vivo* that the FNR, Fur, ferredoxin, nitroreductase genes of *G.*

vaginalis were less active in those patients that did not respond to metronidazole. These genes

are required for the activation of metronidazole in anaerobes (21).

Adenine methyltransferase was the most strongly up-regulated gene of *G. vaginalis* in

communities from patients without response to metronidazole suggesting that it may play a

role in protecting DNA from damage. Studies have provided compelling evidence for DNA

methyltransferases 1 (DNMT1) acting as a regulator of genome integrity and as an early

responder to DNA double strand breaks in human cells (18), but the explicit role of DNMTs

in DNA strand break repair for bacteria remains to be elucidated.

Remarkably, *cas* genes of *G. vaginalis* were highly up-regulated in *G. vaginalis* from vaginal

microbiota of patients that did not respond to metronidazole treatment. The CRISPR-Cas

genes are present in about half of all Bacteria and most Archaea (34); they represent a

mechanism of adaptive immunity which protects the prokaryotic cell against foreign DNA

and has been developed into a universal tool for genome editing (10). The *cas* genes of *G.*

vaginalis belong to the *E. coli* subtype and were found in about half of the clinical isolates

(44). Their up-regulation might reflect increased phage attacks in BV. Phages have been

hypothesized to be crucial for the etiology of BV by causing the collapse of *Lactobacillus*

populations (4); accordingly, *L. iners* upregulates its CRISPR-Cas system in BV (32). More

than 400 annotated prophage sequences were found in 39 *Gardenerella* isolates from the

bladder (35). They might be induced to enter the lytic cycle by the change in pH

accompanying the shift to BV.

The upregulation of CRISPR-Cas system genes in *G. vaginalis* from those patients that did

not respond to treatment by metronidazole suggests that the CRISPR-Cas system might have a

role in not only defending the host against phages, but also in mitigating the DNA damaging

effect of metronidazole. In addition to providing adaptive immunity, CRISPR-Cas systems

can have various additional functions (56) and it was shown that they can protect the cell

against DNA damaging agents (3). The Cas1 enzyme of *E. coli* (YgbT) physically and

genetically interacts with the DNA repair system (RecBC, RuvB) and is recruited to DNA

double strand breaks; moreover, YgbT is necessary for resistance of *E. coli* to DNA damage

caused by the genotoxic antibiotic mitomycin C or UV light (3). Our findings suggest that the

CRISPR-Cas system may protect the vaginal microbiota against the DNA damaging effect of

metronidazole. If experimentally confirmed, this finding might open a new path to fight

bacterial resistance against DNA damaging agents.

Conclusions

We hypothesize that colonization by *L. iners* contribute to recurrence of BV. CRISPR-Cas, a phage defense mechanism with a dual function, might mitigate the DNA damaging effect of metronidazole and thus may be one of the reasons for lack of response of the vaginal microbiota to antibiotic treatment. This first *in vivo* study of the metatranscriptome landscape of vaginal fluid microbiota during metronidazole treatment provides a wealth of new understanding towards the possible mechanisms of lack of response to metronidazole and recurrence in BV. These findings may help to develop more efficient novel therapies to improve the treatment of BV and prevent recurrence. A completely new role for Cas proteins in microbial communities is hypothesized which warrants closer inspection.

Material and Methods

Study design

Vaginal fluid samples of women analyzed here were a subset of the samples obtained during a randomized controlled clinical trial described previously (16). The trial protocol was approved by the local ethics committee (Ärztekammer Nordrhein - Medical Association North Rhine) and written consent was obtained from all participants. The clinical trial was conducted in accordance with the Declaration of Helsinki on Ethical Principles for Medical Research Involving Human Subjects. Principles and guidelines for good clinical practice were followed. The study was registered on ClinicalTrials.gov with the identifier NCT02687789. Briefly, women were included into the clinical trial if they were BV positive according to Amsel criteria and Nugent score and were biofilm positive on vaginal epithelial cells and positive for extracellular polysaccharides (EPS) in urine. For treatment of acute BV, they received 2 g of metronidazole orally and were afterwards treated with an intravaginal pessary

for three weeks, twice a week. Samples were taken during acute BV (visit 1), after receiving

metronidazole 7 to 28 days after visit 1 (visit 2), after pessary application one week after visit

2 (visit 3), after continued pessary application two weeks after visit 3 (visit 4) and during

follow up three months after visit 4 (visit 5).

The aim of the clinical trial had been to compare the effectiveness of two different types of

pessary. The results and the taxonomic composition of the vaginal microbial communities

have been reported (16). For the metatranscriptome analysis reported here, we chose a subset

of 14 patients from the clinical trial. These 14 patients consisted of two groups named "with

response to treatment" (n = 8) and "no response to treatment" (n = 6) (Fig. 1). Among the

eight patients who responded to treatment, six were cured without recurrence and two

experienced recurrence during the three months follow-up. For the analysis of recurrence,

samples from all 5 visits were analyzed for 4 patients (two cured and two with recurrence) (20

samples in total). These four patients all received the commercially available lactic acid

pessary after metronidazole therapy at visit 3 and 4. For the analysis of lack of response to

metronidazole, samples from all 14 patients were analyzed at two time points, acute BV (visit

1) and 7 to 28 days after antibiotic treatment (visit 2) (28 samples in total). Nugent score > 6

was used to determine BV status, since it is considered the gold standard for BV diagnosis

(40) (Table S1 sheet 1). The BV status at visit 5 was determined by Amsel criteria since there

was no Nugent score available at that time point. The sample ID was obtained by

concatenating the patient ID and letters "a" to "e" indicating visit 1 to 5.

Sample collection and transport

Vaginal fluid was obtained by infusing 2 ml of saline solution into the vagina followed by

rotation against the vaginal wall with a speculum and then collecting the vaginal fluid with a

syringe. Approximately 700 µl were immediately transferred to a tube containing 2 ml

RNAprotect (Qiagen, Germany). Tubes were immediately frozen at -20°C, transported at -

20°C within a week and stored at -70°C.

RNA extraction and mRNA enrichment

RNA was extracted from 1 ml vaginal fluid suspension using the MO BIO

PowerMicrobiome™ RNA Isolation Kit (Qiagen, Germany) with pretreatment: Vaginal fluid

was centrifuged at 13,000 rpm for 1 minute. The pellet was resuspended in MoBio lysis buffer

and this suspension was added to the supplied bead tubes filled with 500 µl ice-cold phenol-

choloroformisoamylalcohol solution (Carl Roth, Germany). The bead-suspension mix was

shaken at 5 m/s for 1 minute in 3 intervals which were 2 minutes apart using the MO BIO

PowerLyzer™ (Qiagen, Germany). After centrifugation for 1 minute at 13,000 rpm and 4°C

the upper phase containing the RNA was further processed according to the manufacturer's

instructions including DNAse I treatment. RNA was eluted in 100µl nuclease free water and

vaccum concentrated to 50 µl. The Ribo-Zero Gold rRNA Removal Kit (Epidemiology) by

Illumina (USA) was then used for mRNA enrichment with ethanol precipitation according to

the manufacturer's instructions. Integrity of RNA was evaluated using a Bioanalyzer 2100

(Agilent, Germany).

Library preparation, sequencing and preprocessing of sequencing data

Paired-end mRNA Illumina sequencing libraries were constructed with the Script Seq

Illumina Kit. Strand specific paired end sequencing was performed on the HiSeq 2500

Sequencer to yield 2×110 bp paired-end reads. Primers and sequencing adaptors were

removed from raw sequencing data, followed by clipping the bases with quality score < 20

from the reads using Fastq-Mcf (2). After clipping, the remaining reads shorter than 50 were

removed. Thereafter, the ribosomal RNA reads were eliminated using SortMeRNA v2.0 (23)

with the default parameters.

Taxonomy assignment using Kraken

Kraken (58), an accurate and ultra-fast taxonomy assignment tool for metagenomes was used

to determine the taxonomic composition of the metatranscriptome data. Kraken uses the K-

mer strategy and the lowest common ancestor (LCA) algorithm to affiliate a given read to a

taxon. The standard Kraken database from the Kraken program with addition of the human

genome was used to identify human reads. This standard database consists of prokaryote

genomes (2786), virus genomes (4418) and with addition of human genome (ver. GRCh38)

was downloaded from NCBI.

The refined Kraken database contained 163 bacterial genomes from 105 species of bacteria

(Table S1 sheet 3), including 147 genomes from urogenital subset of HMP reference genome

sequence data (HMRGD) (38). To be able to identify transcripts from bladder isolates, we

added the genomes of *Gardnerella sp.* 26-12 and *Gardnerella sp.* 30-4 which were isolated

from the bladder recently (35). To make sure that the transcripts mapped to the *Gardnerella*

sp. bladder isolates did not actually originate from *G. vaginalis*, we included 5 additional

genomes (original HMP urogenital tract database contains 4 *G. vaginalis*) from distinct *G.*

vaginalis strains in the refined Kraken database. The complete list of reference genomes in the

refined Kraken database was compiled as Table S1 sheet 3. Subsequently, the cleaned

putative microbial mRNAs reads were mapped against this refined database. All results on the

taxonomic composition in this study were achieved based on this database.

Short reads alignment by BWA

Since Kraken was only used for taxonomy classification, BWA was applied to determine the

expression of functional genes. A customized reference gene database combining the genes

from the urogenital tract subset of the HMRGD (38) and 9 additional genomes (Table S1

sheet 4) was used as mapping reference for BWA to map all cleaned putative mRNA reads for

functional activity profiling. The genes of *Gardnerella sp.* 26-12, *Gardnerella sp.* 30-4 were

not included due to the poor quality of the genomes. The short reads alignment was performed

using BWA with the BWA-MEM (26) algorithm. A mapping seed length of 31 which is

much longer than the default seed length 19 was applied to achieve more reliable alignments.

To avoid a biased mapping, the reads mapped with mapping quality score (MAPQ) lower than

10 were excluded. MAPQ contains the Phred-scaled posterior probability that the mapping

position is wrong (25). A MAPQ of 10 indicates a probability of 10% that the alignment

position is wrong.

KEGG Orthologous (KO) gene annotation of customized reference database

The customized short reads alignment reference gene database which was used for BWA was

annotated using KEGG prokaryote protein sequences to calculate the expression level of KO

genes for differential KO gene expression analysis and KEGG pathway enrichment analysis.

The KEGG prokaryote protein sequence database represents a non-redundant protein dataset

of prokaryote (Bacteria and Archaea) on the species level and contains about 7 million non-

redundant peptide sequences grouped into 14,390 distinct KO genes. A KO gene contains

several genes from different species with similar function. DIAMOND (7), a much faster

alternative to BLASTX was applied to map the customized short reads alignment reference

sequences against the KEGG prokaryote protein sequence database with its "more sensitive

mode". To obtain reliable annotation, only alignments with sequence identity >=50 and E-

value <= 1e-5 and query coverage >= 70% were taken into account. By annotating the

reference genes to KO genes, we were able to determine the expression profile of KO genes,

and investigate the functional activity shifts in BV based on differential expression analysis of

KO genes and perform KEGG pathway enrichment analysis.

Differential expression (DE) analysis

All differential expression (DE) analyses were performed using the R package edgeR (48).

The Benjamini Hochberg (BH) method was used to correct the p value of DE analysis with

the false discovery rate (FDR) for multiple comparisons. Genes with FDR smaller than 0.05

were considered as significantly differentially regulated, which were then used for KEGG

pathway enrichment analysis. The sample groups defined for each comparison are listed in

Table S1 sheet 1. The groups defined in column "group for G vaginalis" is for the DE analysis

of KO genes between *G. vaginalis* in communities with no response to metronidazole and

with response. To avoid the analysis bias introduced by zero-inflated data, only the samples

with relative abundance of *G. vaginalis* higher than 20% according to the taxonomy

assignment were taken into account. In these two columns, "NR1" represents samples at visit

1 from patients with no response to metronidazole after treatment, "NR2" indicates the

samples from the same patients at visit 2. And "WR1" denotes samples at visit 1 from patients

with response to metronidazole after treatment, while "WR2" means samples from the

recurrent time point. Column "group for LC LI" is for the DE analysis for KO genes between

L. crispatus in the communities dominated by *L. crispstus* ("LC") and *L. iners* in the

communities dominated by *L. iners* ("LI").

Detection of putative metronidazole resistance related genes

To detect the expression of previously reported putative metronidazole resistant genes such as recA, recA-mediated autopeptidase (Rma), peroxiredoxin, nitroimidazole resistance protein (NIM), ferredoxin/ferredoxin-NADP reductase (FNR), nitroreductase, sialidase and ferredoxin, we examined the expression level of these genes for *G. vaginalis* in the vaginal community from patients without response to metronidazole treatment (n = 6) as well as with response (n = 4). As most of these genes do not have corresponding KO genes, we annotated the BWA reference database based on the sequences of these genes using BLASTN. These sequences were retrieved from ENA by key words of each "ferredoxin, ferric uptake regulator, NADPH flavin oxidoreductase, nitroreductase, peroxiredoxin, pyruvate ferredoxin oxidoreductase, recA, nitroimidazole resistance, sialidase" plus *G. vaginalis*. In total, 155 unique sequences of *G. vaginalis* were obtained for the annotation of BWA reference database. The identification of duplicate sequences was done by SeqKit (49).

Availability of data and material

The metatranscriptome sequencing data of this subgroup analysis have been deposited in European Nucleotide Archive and are accessible through study accession number PRJEB21446.

Authors' contributions

This study was designed by IWD and CG. CM and CA provided the clinical samples. RNA extraction and mRNA enrichment was performed by CG. SB prepared cDNA libraries, and performed Illumina sequencing. Z-LD performed all the data analyses. Data interpretation and visualization were done by IWD, Z-LD and CG. Z-LD and CG wrote the manuscript draft, and all authors reviewed the manuscript.

Acknowledgements

This work was supported by ZIM project grant KF3134201MD3 of the Bundesministerium

für Wirtschaft und Energie (BMWi), Germany. The clinical study was funded by Dr. August

Wolff GmbH & Co. KG Arzneimittel within the ZIM project KF3134201MD3.

Reference List

1. **Antonio, M. A. and S. L. Hillier**. 2003. DNA fingerprinting of Lactobacillus crispatus strain CTV-05 by repetitive element sequence-based PCR analysis in a pilot study of vaginal colonization. J.Clin.Microbiol. **41**:1881-1887.

2. **Aronesty, E.** 2011. ea-utils: Command-line tools for processing biological sequencing data. Durham, NC: Expression Analysis.

3. **Babu, M., N. Beloglazova, R. Flick, C. Graham, T. Skarina, B. Nocek, A. Gagarinova, O. Pogoutse, G. Brown, A. Binkowski, S. Phanse, A. Joachimiak, E. V. Koonin, A. Savchenko, A. Emili, J. Greenblatt, A. M. Edwards, and A. F. Yakunin**. 2011. A dual function of the CRISPR-Cas system in bacterial antivirus immunity and DNA repair. Mol.Microbiol. **79**:484-502. doi:10.1111/j.1365-2958.2010.07465.x [doi].

4. **Blackwell, A. L.** 1999. Vaginal bacterial phaginosis? Sex Transm.Infect. **75**:352-353.

5. **Bradshaw, C. S., A. N. Morton, J. Hocking, S. M. Garland, M. B. Morris, L. M. Moss, L. B. Horvath, I. Kuzevska, and C. K. Fairley**. 2006. High recurrence rates of bacterial vaginosis over the course of 12 months after oral metronidazole therapy and factors associated with recurrence. J.Infect.Dis. **193**:1478-1486. doi:JID36020 [pii];10.1086/503780 [doi].

6. **Bradshaw, C. S., M. Pirotta, G. D. De, J. S. Hocking, A. N. Morton, S. M. Garland, G. Fehler, A. Morrow, S. Walker, L. A. Vodstrcil, and C. K. Fairley**. 2012. Efficacy of oral metronidazole with vaginal clindamycin or vaginal probiotic for bacterial vaginosis: randomised placebo-controlled double-blind trial. PLoS.One. **7**:e34540. doi:10.1371/journal.pone.0034540 [doi];PONE-D-11-24250 [pii].

7. **Buchfink, B., C. Xie, and D. H. Huson**. 2015. Fast and sensitive protein alignment using DIAMOND. Nat.Methods **12**:59-60. doi:nmeth.3176 [pii];10.1038/nmeth.3176 [doi].

8. **Cardone, A., R. Zarcone, A. Borrelli, C. A. Di, A. Russo, and E. Tartaglia**. 2003.
Utilisation of hydrogen peroxide in the treatment of recurrent bacterial vaginosis.
Minerva Ginecol. **55**:483-492.

9. **Deng, Z. L., S. P. Szafranski, M. Jarek, S. Bhuju, and I. Wagner-Dobler**. 2017.
Dysbiosis in chronic periodontitis: Key microbial players and interactions with the
human host. Sci.Rep. **7**:3703. doi:10.1038/s41598-017-03804-8 [doi];10.1038/s41598-
017-03804-8 [pii].

10. **Doudna, J. A. and E. Charpentier**. 2014. Genome editing. The new frontier of
genome engineering with CRISPR-Cas9. Science. **346**:1258096. doi:346/6213/1258096
[pii];10.1126/science.1258096 [doi].

11. **Ferreira, C. S. T., G. G. Donders, C. M. G. L. Parada, A. D. R. Tristao, T.**
Fernandes, M. G. da Silva, and C. Marconi. 2017. Treatment failure of bacterial
vaginosis is not associated with higher loads of Atopobium vaginae and Gardnerella
vaginalis. J.Med.Microbiol. **66**:1217-1224. doi:10.1099/jmm.0.000561 [doi].

12. **France, M. T., H. Mendes-Soares, and L. J. Forney**. 2016. Genomic Comparisons of
Lactobacillus crispatus and Lactobacillus iners Reveal Potential Ecological Drivers of
Community Composition in the Vagina. Appl.Environ.Microbiol. **82**:7063-7073.
doi:AEM.02385-16 [pii];10.1128/AEM.02385-16 [doi].

13. **France, M. T., H. Mendes-Soares, and L. J. Forney**. 2016. Genomic Comparisons of
Lactobacillus crispatus and Lactobacillus iners Reveal Potential Ecological Drivers of
Community Composition in the Vagina. Appl.Environ.Microbiol. **82**:7063-7073.
doi:AEM.02385-16 [pii];10.1128/AEM.02385-16 [doi].

14. **Frank, J. A., C. I. Reich, S. Sharma, J. S. Weisbaum, B. A. Wilson, and G. J. Olsen**.
2008. Critical evaluation of two primers commonly used for amplification of bacterial
16S rRNA genes. Appl.Environ.Microbiol. **74**:2461-2470. doi:AEM.02272-07
[pii];10.1128/AEM.02272-07 [doi].

15. **Gelber, S. E., J. L. Aguilar, K. L. Lewis, and A. J. Ratner**. 2008. Functional and
phylogenetic characterization of Vaginolysin, the human-specific cytolysin from
Gardnerella vaginalis. J.Bacteriol. **190**:3896-3903. doi:JB.01965-07
[pii];10.1128/JB.01965-07 [doi].

16. **Gottschick, C., Z.-L. Deng, M. Vital, C. Masur, C. Abels, D. Pieper, M. Rohde, W.**
Mendling, and I. Wagner-Dobler. 2017. Treatment of biofilms in bacterial vaginosis
by an amphoteric tenside pessary - clinical study and microbiome analysis. Microbiome
in press.

17. **Gottschick, C., Z.-L. Deng, M. Vital, C. Masur, C. Abels, D. Pieper, and I. Wagner-**
Dobler. 2017. The urinary microbiota of men and women and its changes in women
during bacterial vaginosis and antibiotic treatment. Microbiome **5**:99.
doi:10.1186/s40168-017-0305--3.

18. **Ha, K., G. E. Lee, S. S. Palii, K. D. Brown, Y. Takeda, K. Liu, K. N. Bhalla, and K.**
D. Robertson. 2011. Rapid and transient recruitment of DNMT1 to DNA double-strand

breaks is mediated by its interaction with multiple components of the DNA damage
response machinery. Hum.Mol.Genet. **20**:126-140. doi:ddq451
[pii];10.1093/hmg/ddq451 [doi].

19. **Hertzberger, R., J. Arents, H. L. Dekker, R. D. Pridmore, C. Gysler, M.**
Kleerebezem, and M. J. de Mattos. 2014. H(2)O(2) production in species of the
Lactobacillus acidophilus group: a central role for a novel NADH-dependent flavin
reductase. Appl.Environ.Microbiol. **80**:2229-2239. doi:AEM.04272-13
[pii];10.1128/AEM.04272-13 [doi].

20. **Hilbert, D. W., W. L. Smith, T. E. Paulish-Miller, S. G. Chadwick, G. Toner, E.**
Mordechai, M. E. Adelson, J. D. Sobel, and S. E. Gygax. 2016. Utilization of
molecular methods to identify prognostic markers for recurrent bacterial vaginosis.
Diagn.Microbiol.Infect.Dis. doi:S0732-8893(16)30197-3
[pii];10.1016/j.diagmicrobio.2016.07.003 [doi].

21. **Ishii, M., Y. Ueda, K. S. Yoon, Y. Igarashi, and T. Kodama**. 1996. Purification and
characterization of ferredoxin from Hydrogenobacter thermophilus strain TK-6.
Biosci.Biotechnol.Biochem. **60**:1513-1515. doi:10.1271/bbb.60.1513 [doi].

22. **Kindinger, L. M., P. R. Bennett, Y. S. Lee, J. R. Marchesi, A. Smith, S. Cacciatore,**
E. Holmes, J. K. Nicholson, T. G. Teoh, and D. A. MacIntyre. 2017. The interaction
between vaginal microbiota, cervical length, and vaginal progesterone treatment for
preterm birth risk. Microbiome. **5**:6. doi:10.1186/s40168-016-0223-9
[doi];10.1186/s40168-016-0223-9 [pii].

23. **Kopylova, E., L. Noe, and H. Touzet**. 2012. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics. **28**:3211-3217. doi:bts611
[pii];10.1093/bioinformatics/bts611 [doi].

24. **Leitsch, D., A. G. Burgess, L. A. Dunn, K. G. Krauer, K. Tan, M. Duchene, P.**
Upcroft, L. Eckmann, and J. A. Upcroft. 2011. Pyruvate:ferredoxin oxidoreductase
and thioredoxin reductase are involved in 5-nitroimidazole activation while flavin
metabolism is linked to 5-nitroimidazole resistance in Giardia lamblia.
J.Antimicrob.Chemother. **66**:1756-1765. doi:dkr192 [pii];10.1093/jac/dkr192 [doi].

25. **Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.**
Abecasis, and R. Durbin. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics. **25**:2078-2079. doi:btp352 [pii];10.1093/bioinformatics/btp352 [doi].

26. **Li, H.** 2013. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. arXiv preprint arXiv:1303.3997.

27. **Ling, Z., X. Liu, W. Chen, Y. Luo, L. Yuan, Y. Xia, K. E. Nelson, S. Huang, S.**
Zhang, Y. Wang, J. Yuan, L. Li, and C. Xiang. 2013. The restoration of the vaginal
microbiota after treatment for bacterial vaginosis with metronidazole or probiotics.
Microb.Ecol. **65**:773-780. doi:10.1007/s00248-012-0154-3 [doi].

28. **Lofmark, S., C. Edlund, and C. E. Nord**. 2010. Metronidazole is still the drug of
choice for treatment of anaerobic infections. Clin.Infect.Dis. **50 Suppl 1**:S16-S23.
doi:10.1086/647939 [doi].

29. **Lofmark, S., H. Fang, M. Hedberg, and C. Edlund**. 2005. Inducible metronidazole
resistance and nim genes in clinical Bacteroides fragilis group isolates.
Antimicrob.Agents Chemother. **49**:1253-1256. doi:49/3/1253
[pii];10.1128/AAC.49.3.1253-1256.2005 [doi].

30. **Machado, D., J. Castro, A. Palmeira-de-Oliveira, J. Martinez-de-Oliveira, and N.
Cerca**. 2015. Bacterial Vaginosis Biofilms: Challenges to Current Therapies and
Emerging Solutions. Front Microbiol. **6**:1528. doi:10.3389/fmicb.2015.01528 [doi].

31. **Macklaim, J. M., J. C. Clemente, R. Knight, G. B. Gloor, and G. Reid**. 2015.
Changes in vaginal microbiota following antimicrobial and probiotic therapy.
Microb.Ecol.Health Dis. **26**:27799. doi:27799 [pii].

32. **Macklaim, J. M., A. D. Fernandes, J. M. Di Bella, J. A. Hammond, G. Reid, and G.
B. Gloor**. 2013. Comparative meta-RNA-seq of the vaginal microbiota and differential
expression by Lactobacillus iners in health and dysbiosis. Microbiome. **1**:12. doi:2049-
2618-1-12 [pii];10.1186/2049-2618-1-12 [doi].

33. **Macklaim, J. M., G. B. Gloor, K. C. Anukam, S. Cribby, and G. Reid**. 2011. At the
crossroads of vaginal health and disease, the genome sequence of Lactobacillus iners
AB-1. Proc.Natl.Acad.Sci.U.S.A **108 Suppl 1**:4688-4695. doi:1000086107
[pii];10.1073/pnas.1000086107 [doi].

34. **Makarova, K. S., Y. I. Wolf, O. S. Alkhnbashi, F. Costa, S. A. Shah, S. J. Saunders,
R. Barrangou, S. J. Brouns, E. Charpentier, D. H. Haft, P. Horvath, S. Moineau, F.
J. Mojica, R. M. Terns, M. P. Terns, M. F. White, A. F. Yakunin, R. A. Garrett, J.
van der Oost, R. Backofen, and E. V. Koonin**. 2015. An updated evolutionary
classification of CRISPR-Cas systems. Nat.Rev.Microbiol. **13**:722-736.
doi:nrmicro3569 [pii];10.1038/nrmicro3569 [doi].

35. **Malki, K., J. W. Shapiro, T. K. Price, E. E. Hilt, K. Thomas-White, T. Sircar, A. B.
Rosenfeld, G. Kuffel, M. J. Zilliox, A. J. Wolfe, and C. Putonti**. 2016. Genomes of
Gardnerella Strains Reveal an Abundance of Prophages within the Bladder Microbiome.
PLoS.One. **11**:e0166757. doi:10.1371/journal.pone.0166757 [doi];PONE-D-16-20473
[pii].

36. **Mayer, B. T., S. Srinivasan, T. L. Fiedler, J. M. Marrazzo, D. N. Fredricks, and J.
T. Schiffer**. 2015. Rapid and Profound Shifts in the Vaginal Microbiota Following
Antibiotic Treatment for Bacterial Vaginosis. J.Infect.Dis. **212**:793-802. doi:jiv079
[pii];10.1093/infdis/jiv079 [doi].

37. **Mendz, G. L. and F. Megraud**. 2002. Is the molecular basis of metronidazole
resistance in microaerophilic organisms understood? Trends Microbiol. **10**:370-375.
doi:S0966842X02024058 [pii].

38. **Methe, B. A., K. E. Nelson, M. Pop, H. H. Creasy, M. G. Giglio, C. Huttenhower,**
D. Gevers, J. F. Petrosino, S. Abubucker, J. H. Badger, A. T. Chinwalla, A. M.
Earl, M. G. FitzGerald, R. S. Fulton, K. Hallsworth-Pepin, E. A. Lobos, R.
Madupu, V. Magrini, J. C. Martin, M. Mitreva, D. M. Muzny, E. J. Sodergren, J.
Versalovic, A. M. Wollam, K. C. Worley, J. R. Wortman, S. K. Young, Q. Zeng, K.
M. Aagaard, O. O. Abolude, E. Allen-Vercoe, E. J. Alm, L. Alvarado, G. L.
Andersen, S. Anderson, E. Appelbaum, H. M. Arachchi, G. Armitage, C. A. Arze,
T. Ayvaz, C. C. Baker, L. Begg, T. Belachew, V. Bhonagiri, M. Bihan, M. J. Blaser,
T. Bloom, J. Vivien Bonazzi, P. Brooks, G. A. Buck, C. J. Buhay, D. A. Busam, J.
L. Campbell, S. R. Canon, B. L. Cantarel, P. S. Chain, I. M. Chen, L. Chen, S.
Chhibba, K. Chu, D. M. Ciulla, J. C. Clemente, S. W. Clifton, S. Conlan, J.
Crabtree, M. A. Cutting, N. J. Davidovics, C. C. Davis, T. Z. DeSantis, C. Deal, K.
D. Delehaunty, F. E. Dewhirst, E. Deych, Y. Ding, D. J. Dooling, S. P. Dugan, W.
M. Dunne, A. S. Durkin, R. C. Edgar, R. L. Erlich, C. N. Farmer, R. M. Farrell, K.
Faust, M. Feldgarden, V. M. Felix, S. Fisher, A. A. Fodor, L. Forney, L. Foster, V.
Di Francesco, J. Friedman, D. C. Friedrich, C. C. Fronick, L. L. Fulton, H. Gao, N.
Garcia, G. Giannoukos, C. Giblin, M. Y. Giovanni, J. M. Goldberg, J. Goll, A.
Gonzalez, A. Griggs, S. Gujja, B. J. Haas, H. A. Hamilton, E. L. Harris, T. A.
Hepburn, B. Herter, D. E. Hoffmann, M. E. Holder, C. Howarth, K. H. Huang, S.
M. Huse, J. Izard, J. K. Jansson, H. Jiang, C. Jordan, V. Joshi, J. A. Katancik, W.
A. Keitel, S. T. Kelley, C. Kells, S. Kinder-Haake, N. B. King, R. Knight, D.
Knights, H. H. Kong, O. Koren, S. Koren, K. C. Kota, C. L. Kovar, N. C. Kyrpides,
P. S. La Rosa, S. L. Lee, K. P. Lemon, N. Lennon, C. M. Lewis, L. Lewis, R. E. Ley,
K. Li, K. Liolios, B. Liu, Y. Liu, C. C. Lo, C. A. Lozupone, R. D. Lunsford, T.
Madden, A. A. Mahurkar, P. J. Mannon, E. R. Mardis, V. M. Markowitz, K.
Mavrommatis, J. M. McCorrison, D. McDonald, J. McEwen, A. L. McGuire, P.
McInnes, T. Mehta, K. A. Mihindukulasuriya, J. R. Miller, P. J. Minx, I. Newsham,
C. Nusbaum, M. O'Laughlin, J. Orvis, I. Pagani, K. Palaniappan, S. M. Patel,
M. Pearson, J. Peterson, M. Podar, C. Pohl, K. S. Pollard, M. E. Priest, L. M.
Proctor, X. Qin, J. Raes, J. Ravel, J. G. Reid, M. Rho, R. Rhodes, K. P. Riehle, M.
C. Rivera, B. Rodriguez-Mueller, Y. H. Rogers, M. C. Ross, C. Russ, R. K. Sanka,
J. Pamela Sankar, F. Sathirapongsasuti, J. A. Schloss, P. D. Schloss, T. M.
Schmidt, M. Scholz, L. Schriml, A. M. Schubert, N. Segata, J. A. Segre, W. D.
Shannon, R. R. Sharp, T. J. Sharpton, N. Shenoy, N. U. Sheth, G. A. Simone, I.
Singh, C. S. Smillie, J. D. Sobel, D. D. Sommer, P. Spicer, G. G. Sutton, S. M.
Sykes, D. G. Tabbaa, M. Thiagarajan, C. M. Tomlinson, M. Torralba, T. J.
Treangen, R. M. Truty, T. A. Vishnivetskaya, J. Walker, L. Wang, Z. Wang, D. V.
Ward, W. Warren, M. A. Watson, C. Wellington, K. A. Wetterstrand, J. R. White,
K. Wilczek-Boney, Y. Q. Wu, K. M. Wylie, T. Wylie, C. Yandava, L. Ye, Y. Ye, S.
Yooseph, B. P. Youmans, L. Zhang, Y. Zhou, Y. Zhu, L. Zoloth, J. D. Zucker, B.
W. Birren, R. A. Gibbs, S. K. Highlander, G. M. Weinstock, R. K. Wilson, and O.
White. 2012. A framework for human microbiome research. Nature **486**:215-221.
doi:10.1038/nature11209.

39. **Michel, B.** 2005. After 30 years of study, the bacterial SOS response still surprises us.
PLoS.Biol. **3**:e255. doi:05-PLBI-P-0386 [pii];10.1371/journal.pbio.0030255 [doi].

40. **Mohammadzadeh, F., M. Dolatian, M. Jorjani, and M. H. Alavi**. 2014. Diagnostic
value of Amsel's clinical criteria for diagnosis of bacterial vaginosis. Glob.J.Health Sci.
7:8-14. doi:10.5539/gjhs.v7n3p8 [doi].

41. **Ojala, T., M. Kankainen, J. Castro, N. Cerca, S. Edelman, B. Westerlund-**
Wikstrom, L. Paulin, L. Holm, and P. Auvinen. 2014. Comparative genomics of
Lactobacillus crispatus suggests novel mechanisms for the competitive exclusion of
Gardnerella vaginalis. BMC.Genomics **15**:1070. doi:1471-2164-15-1070
[pii];10.1186/1471-2164-15-1070 [doi].

42. **Onderdonk, A. B., M. L. Delaney, and R. N. Fichorova**. 2016. The Human
Microbiome during Bacterial Vaginosis. Clin.Microbiol.Rev. **29**:223-238. doi:29/2/223
[pii];10.1128/CMR.00075-15 [doi].

43. **Petrova, M. I., G. Reid, M. Vaneechoutte, and S. Lebeer**. 2016. Lactobacillus iners:
Friend or Foe? Trends Microbiol. doi:S0966-842X(16)30181-0
[pii];10.1016/j.tim.2016.11.007 [doi].

44. **Pleckaityte, M., M. Zilnyte, and A. Zvirbliene**. 2012. Insights into the CRISPR/Cas
system of Gardnerella vaginalis. BMC.Microbiol. **12**:301. doi:1471-2180-12-301
[pii];10.1186/1471-2180-12-301 [doi].

45. **Pukkila, P. J., J. Peterson, G. Herman, P. Modrich, and M. Meselson**. 1983. Effects
of high levels of DNA adenine methylation on methyl-directed mismatch repair in
Escherichia coli. Genetics **104**:571-582.

46. **Rampersaud, R., P. J. Planet, T. M. Randis, R. Kulkarni, J. L. Aguilar, R. I.**
Lehrer, and A. J. Ratner. 2011. Inerolysin, a cholesterol-dependent cytolysin produced
by Lactobacillus iners. J.Bacteriol. **193**:1034-1041. doi:JB.00694-10
[pii];10.1128/JB.00694-10 [doi].

47. **Ravel, J., P. Gajer, Z. Abdo, G. M. Schneider, S. S. Koenig, S. L. McCulle, S.**
Karlebach, R. Gorle, J. Russell, C. O. Tacket, R. M. Brotman, C. C. Davis, K. Ault,
L. Peralta, and L. J. Forney. 2011. Vaginal microbiome of reproductive-age women.
Proc.Natl.Acad.Sci.U.S.A **108 Suppl 1**:4680-4687. doi:1002611107
[pii];10.1073/pnas.1002611107 [doi].

48. **Robinson, M. D., D. J. McCarthy, and G. K. Smyth**. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics. **26**:139-140. doi:btp616 [pii];10.1093/bioinformatics/btp616 [doi].

49. **Shen, W., S. Le, Y. Li, and F. Hu**. 2016. SeqKit: A Cross-Platform and Ultrafast
Toolkit for FASTA/Q File Manipulation. PLoS.One. **11**:e0163962.
doi:10.1371/journal.pone.0163962 [doi];PONE-D-16-20820 [pii].

50. **Sobel, J. D., C. Schmitt, and C. Meriwether**. 1993. Long-term follow-up of patients
with bacterial vaginosis treated with oral metronidazole and topical clindamycin.
J.Infect.Dis. **167**:783-784.

51. **Steffens, L. S., S. Nicholson, L. V. Paul, C. E. Nord, S. Patrick, and V. R. Abratt**.
2010. Bacteroides fragilis RecA protein overexpression causes resistance to
metronidazole. Res.Microbiol. **161**:346-354. doi:S0923-2508(10)00085-9
[pii];10.1016/j.resmic.2010.04.003 [doi].

52. **Tsugawa, H., H. Suzuki, K. Satoh, K. Hirata, J. Matsuzaki, Y. Saito, M. Suematsu,
and T. Hibi**. 2011. Two amino acids mutation of ferric uptake regulator determines
Helicobacter pylori resistance to metronidazole. Antioxid.Redox.Signal. **14**:15-23.
doi:10.1089/ars.2010.3146 [doi].

53. **Turovskiy, Y., N. K. Sutyak, and M. L. Chikindas**. 2011. The aetiology of bacterial
vaginosis. J.Appl.Microbiol. **110**:1105-1128. doi:10.1111/j.1365-2672.2011.04977.x
[doi].

54. **van de Wijgert, J. H., H. Borgdorff, R. Verhelst, T. Crucitti, S. Francis, H.
Verstraelen, and V. Jespers**. 2014. The vaginal microbiota: what have we learned after
a decade of molecular characterization? PLoS.One. **9**:e105998.
doi:10.1371/journal.pone.0105998 [doi];PONE-D-14-20877 [pii].

55. **Wassmann, C., A. Hellberg, E. Tannich, and I. Bruchhaus**. 1999. Metronidazole
resistance in the protozoan parasite Entamoeba histolytica is associated with increased
expression of iron-containing superoxide dismutase and peroxiredoxin and decreased
expression of ferredoxin 1 and flavin reductase. J.Biol.Chem. **274**:26051-26056.

56. **Westra, E. R., A. Buckling, and P. C. Fineran**. 2014. CRISPR-Cas systems: beyond
adaptive immunity. Nat.Rev.Microbiol. **12**:317-326. doi:nrmicro3241
[pii];10.1038/nrmicro3241 [doi].

57. **Whiteside, S. A., H. Razvi, S. Dave, G. Reid, and J. P. Burton**. 2015. The
microbiome of the urinary tract--a role beyond infection. Nat.Rev.Urol. **12**:81-90.
doi:nrurol.2014.361 [pii];10.1038/nrurol.2014.361 [doi].

58. **Wood, D. E. and S. L. Salzberg**. 2014. Kraken: ultrafast metagenomic sequence
classification using exact alignments. Genome Biol. **15**:R46. doi:gb-2014-15-3-r46
[pii];10.1186/gb-2014-15-3-r46 [doi].

Figures, tables and additional files

Fig. 1: Taxonomic composition of vaginal fluid metatranscriptomes in BV and after treatment with metronidazole. Species with average relative abundance smaller than 0.5% were grouped into "Others". The red dot on top of the samples indicates BV. The digits indicate the patient ID, while the letters a-e denote visit 1-5. After the first sampling at visit 1 the patients were treated with metronidazole. Total putative bacterial mRNA reads were mapped to the refined reference database using Kraken (see Methods for details). BV status was determined by Nugent score.

Fig. 2: : The cumulative dominance of the vaginal microbiota in BV and health based on metatranscriptome. Red denotes the dominance curve in BV while blue represents the dominance in health. Y-axis indicates the cumulative relative abundance.

Fig. 3: Average taxonomic composition of vaginal fluid samples in BV determined by 16S rRNA amplicon sequencing (A) and metatranscriptome sequencing (B) and taxonomic composition of communities on the species level determined by metatranscriptome sequencing in health and BV (C). In A, amplicon sequencing was performed as described in our previous study (16) using primers V1-V2. For B and C, taxonomy was assigned based on all cleaned reads after removal of human reads using Kraken and the refined reference database. Species with average relative abundance over 2% are shown in A and B. Relative average abundance was calculated based on all mapped reads. Mean and standard error are shown. In C, only species with relative abundance over 2% in at least 1 sample are shown. Red and pink dots on top of the heatmap indicate paired samples of patients which developed recurrence.

Fig. 4: Distinct functional activity profiles in health and BV (A), temporal dynamics of

the metatranscriptome during treatment (B), and differential KO gene expression of

Lactobacillus crispatus **and** *L. iners* **(C). A)** Principal components analysis of KO gene

expression shows highly similar community activity in BV which is clearly distinct but much

more variable in health. **B)** Time course based on PCA of functional activity profiles from

four women during treatment and recurrence over a period of three months (five sampling

time points). Two women with recurrence (pink and blue color range) and two women

without recurrence (green and orange color range) are shown. In the figure legend, BV

indicates time point with BV, R indicates the recurrence and H represents health. The green

and blue circles highlight healthy clusters, respectively, while the red circle highlights

samples from BV. The arrows denote the temporal shifts of the communities during the

treatment. **C)** *L. crispatus* and *L. iners* exhibit distinct function activities. The Venn diagram

indicates the unique KO genes of *L. crispatus* and *L. iners* as well as their shared KO genes.

The innermost ring denotes the expression of KO genes by log2CPM, the outer ring illustrates

the fold change of the expression of KO genes between *L. crispatus* dominated communities

and *L. iners* dominated communities by log2FC. The small red triangle marks the inerolysin

gene, while blue triangles mark the genes involved in the production of hydrogen peroxide.

Fig. 5: Seven *cas* **genes and a methyltransferase gene of** *G. vaginalis* **were highly up-**

regulated in communities which did not respond to the treatment. KO genes with FDR

<= 0.05 are colored in red or turquoise (significantly differentially regulated) while FDR >

0.05 are in grey.

Table 1. Overview of sequencing results

Overview of sequencing results	BV (22 samples)	Health (18 samples)
raw reads	44,503,332 ± 8,934,125	50,048,447 ± 10,989,072
cleaned reads after quality control and rRNA removal	32,506,992 ± 5,955,546	36,797,904 ± 6,934,913
putative microbial mRNA reads	27,524,573 ± 6,035,462	8,686,750 ± 5,867,867

878 **Table 2. Mapping statistics using Kraken and BWA**

Mapping statistics	% mapped putative microbial mRNA reads
standard Kraken database	40.55 ± 23.93
refined Kraken database	85.88 ± 13.99
customized reference gene database using BWA	~ 78.02 ± 16.31

Fig. S1: Differential expression of putative metronidazole resistance associated genes of

***G. vaginalis* in vaginal fluid microbiota that did not respond to treatment.** The expression

value was calculated based on relative abundance of reads mapped onto *G. vaginalis* using

BWA. "NR1" (No Response 1) indicates the samples from BV that did not respond to

metronidazole treatment later; "NR2" (No Response 2) indicates the samples after treatment

with no response to the treatment. "WR1" (With Response 1) represents the samples with

acute BV which afterwards responded to metronidazole.

Supplementary Table S1: All supplementary data

Sheet 1. Sample description

Sheet 2. Read summary

Sheet 3. Genomes in the refined Kraken reference genome database

Sheet 4. Customized reference gene database for BWA

Sheet 5. Species composition determined by Kraken based on refined database

Sheet 6. The comparison of the community composition identified by 16S and

metatranscriptome (average relative abundance >= 2%)

Sheet 7. The gene expression based on customized reference gene database with KO

annotation

Sheet 8. The differential expression of KO genes between *L. crispatus* and *L. iners*

Sheet 9. KEGG pathway enriched in up-regulated KO genes in *L. crispatus*

Sheet 10. Expression of metronidazole associated genes in *G. vaginalis*

Sheet 11. The differential expression of KO genes of *G. vaginalis* from community

without response to metronidazole treatment and with response