Metatranscriptome analysis of the vaginal microbiota reveals potential

mechanisms for protection against metronidazole in bacterial vaginosis

Zhi-Luo Deng[a#], Cornelia Gottschick[a*], Sabin Bhuju[b], Clarissa Masur[c], Christoph Abels[c],

and Irene Wagner-Döbler[a]

[a] Research Group Microbial Communication, Department of Medical Microbiology, Helmholtz Centre

for Infection Research, Inhoffenstr. 7, 38124 Braunschweig, Germany

[b] Genome Analytics, Helmholtz Centre for Infection Research, Inhoffenstr. 7, 38124 Braunschweig,

Germany

[c] Dr. August Wolff GmbH & Co. KG Arzneimittel, Sudbrackstrasse 56, 33611, Bielefeld, Germany

Running title: Activity profiling of the vaginal microbiota

Address correspondence to Zhi-Luo Deng, Zhiluo.Deng@helmholtz-hzi.de

Z.-L.D. and C.G. contributed equally to this work.

* Present address: Institute for Medical Epidemiology, Biometrics and Computer Science, Martin-

Luther-University Halle-Wittenberg, Halle (Saale), Germany

Abstract

Bacterial vaginosis (BV) is a prevalent multifactorial disease of women in their reproductive

years characterized by a shift from the *Lactobacillus* spp. dominated microbial community

towards a taxonomically diverse anaerobic community. For unknown reasons, some women

do not respond to therapy. In our recent clinical study, out of 37 women diagnosed with BV,

31 were successfully treated with metronidazole, while 6 still had BV after treatment. To

discover possible reasons for the lack of response in those patients, we performed a

metatranscriptome analysis of their vaginal microbiota, comparing them to patients who

responded. Seven out of 8 Cas genes of *Gardnerella vaginalis* were highly up-regulated in

non-responding patients. Cas genes, in addition to protecting against phages, might be

involved in DNA repair thus mitigating the bactericidal effect of DNA damaging agents like

metronidazole. In the second part of our study, we analyzed the vaginal metatranscriptomes of

four patients over three months and showed high *in vivo* expression of genes for pore-forming

toxins in *L. iners* and of genes encoding enzymes for the production of hydrogen peroxide and

D-lactate in *L. crispatus.*

Importance

Bacterial vaginosis is a serious issue for women in their reproductive years. Although it can

usually be cured by antibiotics, the recurrence rate is very high, and some women do not

respond to antibiotic therapy. The reasons for that are not known. Therefore we undertook a

study to detect the activity of the complete microbiota in the vaginal fluid of women that

responded to antibiotic therapy and compared it to the activity of the microbiota in women

that did not respond. We found that one of the most important pathogens in bacterial

vaginosis, *Gardnerella vaginalis*, has activated genes that can repair the DNA damage caused

by the antibiotic in those women that do not respond to therapy. Suppressing these genes

might be a possibility to improve the antibiotic therapy of bacterial vaginosis.

Introduction

The healthy vaginal microbiome is characterized by low pH and low diversity and can be categorized into community state types (CSTs) that are dominated by different *Lactobacillus* spp. such as *L. crispatus*, *L. iners*, *L. gasseri* and less frequently *L. jensenii* or a more diverse community (1). Bacterial vaginosis (BV) is a frequent multifactorial disease of women in their reproductive years that is characterized by a shift of this *Lactobacillus* spp. dominated bacterial community to a community of various mostly anaerobic bacteria (2). BV is associated with a higher risk of preterm birth and of acquiring sexually transmitted infections such as HIV (3). The most common bacteria found in BV, identified by 16S rRNA gene sequencing, are *Gardnerella*, *Atopobium*, *Prevotella*, *Bacteroides*, *Peptostreptococcus*, *Mobiluncus*, *Sneathia*, *Leptotrichia*, *Mycoplasma* and BV associated bacterium 1 (BVAB1) to BVAB3 of the order Clostridiales. Recently, three CSTs dominated by *Gardnerella vaginalis*, Lachnospiraceae and *Sneathia sanguinegens*, respectively, have been described (4). In our recent clinical study *S. amnii* was identified as the best biomarker for BV (5).

The most important pathogen in BV is *Gardnerella vaginalis* (6). It is currently the only described species in the genus *Gardnerella*, but genome comparisons suggest that it can be separated into four genetically isolated subspecies (7). While they cannot be resolved by 16S rRNA gene sequencing, the universal target from the chaperonin-60 gene separates the species into the same four subgroups (group A, clade 4; subgroup B, clade 2; subgroup C, clade 1; subgroup D, clade 3) (6,8,9). All four subgroups of *G. vaginalis* can be detected in the vaginal microbiota of healthy women throughout the menstrual cycle (10). Subgroup A and C define distinct CSTs in health (11). Isolates from the four subgroups of *G. vaginalis* differ in their virulence as well as in their resistance against metronidazole. The sialidase activity of *G. vaginalis* is an important virulence factor and it was detected in all isolates from

subgroup B and few isolates of subgroup C but not in subgroups A and D isolates (12). The

presence of sialidase activity is used for diagnosis of BV in a commercial kit (13). Resistance

against metronidazole was found in subgroups A and D isolates, while those from subgroups

B and C were highly susceptible (14).

Metronidazole is a widely applied chemotherapeutic agent used to treat infectious diseases

caused by anaerobic bacteria, and it is the first-line antibiotic for treating BV (15,16).

Metronidazole is a prodrug which requires enzymatic reduction within the cell, which occurs

under anaerobic conditions only, to transform it into an active form (17). Activated

metronidazole acts by covalently binding to DNA, disrupting its helical structure and causing

single and double strand breaks that lead to DNA degradation and death of the pathogens

(17). Resistance can therefore be mediated by lack of activation of the prodrug, or by repair of

DNA damage, and has been studied in various pathogens. In *Helicobacter pylori* and

Campylobacter spp. ferredoxin, ferredoxin/ferredoxin-NADP reductase (FNR) and

nitroreductase contribute to metronidazole resistance (17). In *Bacteroides fragilis*, genes

responsible for DNA repair like recA and recA-mediated autopeptidase (Rma) and a gene

named nitroimidazole resistance gene (*nim*) encoding a nitroimidazole reductase were shown

to confer resistance against metronidazole (18,19). Failure of BV treatment by metronidazole

is relatively rare (5,20).It is unclear if it is caused by resistance of the BV pathogens to

metronidazole, and which mechanisms are acting *in vivo*. A recent study has demonstrated

that failure of treatment of BV with metronidazole is not associated with higher loads of *G.*

vaginalis and *A. vaginae* (21). Isolates from *G. vaginalis* subgroups A and D are intrinsically

resistant against metronidazole, but the underlying mechanism is unknown (14).

Until now, the majority of studies regarding the vaginal microbiota have focused on 16S

rRNA gene sequencing, answering only questions on the taxonomic composition of bacterial

communities but not on their functions (2). A metatranscriptome analysis comparing vaginal

swabs from two women with BV with two healthy subjects showed that *L. iners* upregulates

transcription of the cholesterol-dependent cytolysin (CDC) and of genes belonging to the

clustered regularly interspaced short palindromic repeats (CRISPR) system in BV (22). No

study has investigated the activity shifts of the vaginal microbiota during antibiotic treatment

of BV.

We had previously analyzed the vaginal microbiota in the context of a clinical trial using 16S

rRNA gene sequencing (5). Of 37 patients diagnosed with BV and included in this study, 31

were initially cured by a single oral dose of metronidazole. Six patients did not respond, i.e.

they were still diagnosed with BV according to Nugent score after antibiotic therapy. Here we

asked if differences in the activity of the microbiota might be responsible for the lack of

response in those six patients. We therefore analyzed their metatranscriptomes at the time of

diagnosis of BV (visit 1) and after treatment with metronidazole (visit 2) and compared them

to those of 8 patients that responded to treatment according to Nugent score.

The high rate of recurrence is another crucial problem for BV treatment. The one-year

recurrence rate of BV ranges from 40% to 80% after therapy with metronidazole (23) or

clindamycin cream (24). CST dominated by *L. iners* might have an increased probability to

shift to a dysbiotic state (22,25,26). In the second part of our study, we therefore followed the

activity of the microbiota from four of the patients that initially responded to metronidazole

treatment over a period of 3 months (visit 3-5) and analyzed gene expression of *L. crispatus*

and *L. iners* in *vivo*.

We show the importance of *G. vaginalis* for BV, which can be massively underestimated

using 16S rRNA gene sequencing. The relative abundance of the four subgroups of *G.*

vaginalis could be determined in responders and non-responders. Transcripts potentially

leading to lack of response to metronidazole treatment were identified. CRISPR-Cas genes are

suggested as a novel mechanism of *G. vaginalis* to mitigate the DNA damaging effect of

metronidazole. *L. iners* highly expressed genes for pore-forming toxins *in vivo*, and in *L.*

crispatus the most highly expressed transcripts *in vivo* encoded enzymes for D-lactate and

hydrogen peroxide production.

Results

Study population and overview of sequencing results

We studied the vaginal microbiome of 14 patients during and after metronidazole treatment of

BV using metatranscriptome sequencing (Fig. 1A). Patients were part of a clinical trial

described elsewhere (5). In the first part of the study, we analyzed samples from two

timepoints (diagnosis of BV, visit 1) and after metronidazole treatment (visit 2). Eight patients

responded to treatment, and six patients did not respond to treatment with the antibiotic, so

were still BV positive according to Nugent score at visit 2. In the second part of our study,

three additional timepoints were analyzed for four of the patients that initially responded to

metronidazole therapy, covering a total period of 3 months. Those four patients belonged to

the lactic acid arm of the clinical study. Two of them experienced recurrence, while the other

two were stably non-BV after treatment according to Nugent score (details in Table S1 sheet

1). In total, we analyzed 40 vaginal fluid samples, 22 with BV status and 18 without.

Metatranscriptome sequencing resulted in a total of 1,879,945,342 reads. Of these,

1,377,516,082 reads (73%) were left after quality filtering and removal of ribosomal RNA

(Table S1 sheet 2). On average, 34 million reads were analyzed per sample.

Construction of the reference genome and gene databases for taxonomic and activity profiling

Human reads comprised ~11% (BV) versus ~56% (non-BV) of the total putative mRNA reads based on the standard Kraken (27) database (Table S1 sheet 2). This suggests that the bacterial load is much lower in non-BV than in BV since the human contamination is much higher in non-BV. Using the standard Kraken database, only 41% of total putative microbial (non-human) mRNA reads could be assigned taxonomically (Table S1 sheet 2). To improve the fraction of taxonomically assignable reads we then constructed a refined database (ref_Genome) which combined the urogenital subset of the HMP (28) database (147 genomes) and all species which are not included in the urogenital subset of the HMP database but detected by the standard Kraken database with an abundance >1% (7 genomes). We also added *S. amnii* and *S. sanguinegens* which had previously been shown to be highly abundant based on 16S rRNA gene sequencing (5) but were not contained in either HMP or the standard Kraken reference database. There are four *G. vaginalis* strains in the HMP database, of which one belongs to subgroup A and three belong to subgroup C. Given the importance and high intra-species diversity of *G. vaginalis*, we added 5 additional *G. vaginalis* genomes based on the genome tree reported in the NCBI database and the completeness of the genome assembly; these five strains cover all four subgroups. We added the genomes of *Gardnerella* sp. 26-12 and *Gardnerella* sp. 30-4 which were isolated from the bladder recently (29). They were classified into *G. vaginalis* subgroup A based on sequence homology (29). In total, this database contained 163 bacterial genomes from 105 species (Table S1 sheet 3). Using this database, the rate of taxonomically classified putative microbial mRNA reads could be improved to 86% on average (Table S1 sheet 2).

For functional assignment, we constructed a reference gene database (ref_Gene) (Table S1

sheet 4). It was based on the same genomes as the ref_Genome database, except that the seven

additional *Gardnerella* spp. genomes were not included because of the low quality of the

annotation of coding sequences. The ref_Gene database contained 301,323 genes. To

investigate the activity shifts of the communities, we mapped the cleaned metatranscriptomic

reads to the ref_Gene database using BWA. In total, 78% of total putative microbial mRNA

reads could be mapped to the ref_Gene. Per sample, on average 8.9 million microbial mRNA

reads could be mapped with MAPQ >10 (Table S1 sheet 2).

Shifts in the taxonomic composition of the active community following metronidazole

treatment

The taxonomic composition of transcripts was determined using Kraken and the ref_Genome

database. Fig. 1B shows that in all communities with BV status the most abundant species

were *G. vaginalis*, *A. vaginae*, *S. amnii* and *Prevotella timonensis* In the post treatment

communities from responders (non-BV, Nugent score <6) the metatranscriptomes were

dominated by *L. crispatus*, *L. iners* and *L. jensenii*, representing typical CSTs of the healthy

female microbiota.

On average less than 14 species contributed >90% of the mapped reads in BV and 3 species

accounted for >90% of the mapped reads in non-BV (Fig. 1C). The individual dominance

plots showed the same pattern where 10 species contributed >90% of the metatranscriptomes

for most patients in BV. In non-BV, this number was 2 for most patients and the dominance

curves were extremely steep. For comparison, in the periodontal metatranscriptome more than

100 species were required to cover 90% of mapped reads (30). These data show that the active

microbiota in BV is much less diverse than suggested by 16S rRNA gene sequencing.

G. vaginalis was the most dominant active species in BV. To estimate the relative abundance

of the four subgroups of *G. vaginalis*, we extracted all reads assigned to *G. vaginalis* from the

metatranscriptomes and assigned them to the four strains representing subgroups A-D,

respectively (409-05, 00703Bmah, HMP9231, 00703Bmash) using Kraken. For this analysis,

we used samples from visit 1 where *G. vaginalis* reads comprised at least 20% of all reads,

which included all 6 patients without response to treatment, and 5 of the 8 patients that

responded to treatment. Figure 1D shows that on average >95% of the reads could be mapped

to the four subgroups and only on average 7% were assigned ambiguously. In those patients

that did not respond to treatment, subgroups A and D comprised 68.5±17.2 % of all reads,

while they accounted for 30.5±29.3% of all reads in patients that responded to treatment

(Wilcoxon test P=0.0520). We observed that *Gardnerella* spp. previously isolated from the

bladder (*Gardnerella* sp. 26-12 and *Gardnerella* sp. 30-4) (29) contributed on average 6% of

all taxonomically assigned reads in BV (Fig. S1).

Comparison of the taxonomic composition of vaginal fluid samples between

metatranscriptome and 16S rRNA gene sequencing

We compared the taxonomic composition determined using 16S rRNA sequencing previously

(5) and the taxonomic composition of the metatranscriptome determined here by Kraken with

the ref_Genome database. In non-BV, we did not observe any considerable difference for the

four most abundant species (Table S1 sheet 6), while in BV large differences between the two

datasets were found. Fig. 2 shows the top 12 most abundant taxa identified using 16S rRNA

gene sequencing and metatranscriptomics, respectively. Most of the abundant species

identified in the mRNA sequencing data set were also identified using 16S rRNA gene

amplicon sequencing, although usually at different abundances. For example, *A. vaginae*

comprised 13% of all reads based on 16S rRNA gene sequencing, but only 11% in the

metatranscriptome dataset. The most pronounced difference was observed for *G. vaginalis*

which comprised on average 47% of relative abundance in the metatranscriptome and on

average only 5% in the 16S rRNA sequencing data. Several additional differences were

found: Higher level taxa like Veillonellaceae (family) or *Parvimonas* (genus) are not listed

among the top 12 taxa of the metatranscriptome, because mapping occurred to species level,

and the abundance of individual species of these higher order taxa was too low to be found

among the top 12 taxa (Table S1 sheet 6). BVAB2 is readily detected by PCR and is an

important indicator of BV, but it is not yet cultivated and so there is no genome available to

map the reads against.

Global community profiling in non-BV and BV

In order to profile the function of the communities, all cleaned putative mRNA reads (Table

S1 sheet 2) were mapped using BWA onto the ref_Gene database annotated with KEGG

ortholog (KO) genes. We used principal component analysis (PCA) to visualize the difference

between the microbiota in BV and non-BV on the level of taxonomy (16S rRNA gene) (Fig.

3A), taxonomic composition of expressed genes (metatranscriptome) (Fig. 3B) and functional

annotation of transcripts to KEGG orthologues (KO genes) (Fig. 3C). Figure 3A shows that

the non-BV communities form a tight cluster on the level of the 16S rRNA sequencing, while

BV communities vary, in accordance with the studies using amplicon sequencing of BV. *L.*

iners, *Prevotella* spp., *G. vaginalis*, *A. vaginae* and *S. amnii* drive the separation between non-

BV and BV On the level of the taxonomic composition of the metatranscriptomes (Fig. 3B)

this pattern was reversed; samples from non-BV were much more heterogeneous than those

from BV. The non-BV communities clustered into two groups dominated by *L. iners* and *L.*

crispatus respectively, whereas *G. vaginalis*, *A. vaginae* and *S. amnii* were abundant in BV.

This reversal is even stronger on the level of KO genes (Fig. 3C): Samples from BV form a

tight cluster, while those from non-BV vary widely. It is the opposite pattern than that found

for the phylogenetic marker gene. The KO genes that contributed most to these differences in

non-BV were phosphofructokinase isozyme *pfkA* (31) and ribosomal protein coding genes

rpsI, *rpmF* and *rplU* (32,33). In BV, *msmE*, *cycB* and *pflD* genes that encode proteins

involved in carbohydrate uptake and metabolism (34,35) were stably higher expressed.

In vivo* expression of putative metronidazole resistance associated genes in *G. vaginalis

To clarify the possible contribution of genes related to metronidazole resistance in Gram

positive pathogens to the difference in response to treatment of the vaginal microbiota, we

examined their expression (Table S1 sheet 9) in *G. vaginalis*. For this analysis, BV

communities from 11 patients at visit 1 were analyzed in which the level of *G. vaginalis*

transcripts was >20%. Six of these patients did not respond to treatment and five responded.

Although *A. vaginae* and *S. amnii* are also key players in BV we could not analyze them here

since there were too few samples dominated by them. As shown in Fig. 4A, there was no clear

expression pattern for most of these genes (detailed data in Table S1 sheet 9). The only

significantly changed gene expression was that of the gene encoding ferredoxin which was

less active in *G. vaginalis* in non-responding patients (fold change=1.67, Wilcoxon test

P=0.00866 based on relative abundance = read count of given genes of *G. vaginalis* / read

count of *G. vaginalis* %).

CRISPR associated protein coding genes of *G. vaginalis* were strongly up-regulated in

vaginal fluids of patients not responding to treatment

We then performed a global analysis of differential expression of KO genes of *G. vaginalis* in

these same communities (visit 1, 11 BV samples with >20% transcripts from *G. vaginalis*

including 6 patients that did not respond to treatment and 5 that responded). We observed that

there were 9 KO genes highly up-regulated with FDR <= 0.05 (log2 fold change up to 9.46)

in communities without response. Strikingly, among the most strongly up-regulated KO

genes, seven were *cas* genes (36) (Fig. 4B). In total there were 8 different *G. vaginalis*

CRISPR-associated (Cas) genes found in the genomes, namely *cas*1-3, *cas*A-E, of which

seven were up-regulated (*cas*1-3, *cas*A-D).

There were 9 KO genes down-regulated but the fold change values were not as high as for the

up-regulated genes. *fucP* (fucose permease) was identified as the most strongly down-

regulated gene with a log2 fold change of -4.24.

Time course of activity profiles and recurrence

In the second part of our study, we analyzed the metatranscriptome of vaginal fluid samples

from four of the patients that initially responded to therapy with metronidazole for the

complete duration of the clinical trial. Two of these patients experience recurrence of BV, and

two remained stably non-BV. Five timepoints were analyzed of which the first two were

already shown in Fig. 1A. They represented acute BV (visit 1) and after metronidazole

therapy (visit 2). Here, we also show visit 3-5, which were all non-BV, with the exception of

recurrence at visit 5 in patient 04_001 and at visit 3 in patient 06_004. Fig. 5A shows the

taxonomic composition of the communities. In one of the two patients that stably maintained

a non-BV status *L. crispatus*, and in the other *L. iners* dominated the microbiota. The

principal components analysis of the activity profiles is shown in Fig. 5B. In acute BV,

samples from all four patients clustered together (red circle). After the treatment, samples

from patient 08_006 who was stably non-BV moved into a very dense and distinct cluster

(illustrated by the arrow 1 and enclosed by a green circle). Samples from patient 13_019 who

also remained stably non-BV moved to a different cluster after treatment, shown by arrow 2

and encircled blue.

The activity shifts in patient 06_004, who experienced recurrence at visit 3, were especially

noteworthy: After treatment, the community moved towards an activity profile distinct from

all others (arrow 3). The recurrence of BV caused the community to shift back to the BV

cluster (red circle, arrow 4). At visit 4, the community moved to the non-BV cluster (blue

circle) and the patient became non-BV according to the Nugent score. We speculate that there

was an unknown intervention after visit 3 which changed the microbiome but this was not

recorded. Interestingly, the other case of recurrence (patient 04_001) had a different

progression. From visit 2 to 4, patient 04_001 was non-BV and these samples clustered

together in the "non-BV" cluster indicated by the blue circle. At visit 5, however, patient

04_001 had recurrent BV and the community shifted back again to the BV cluster.

Transcriptomics of *L. iners* and *L. crispatus in vivo*

The stable colonization of the vaginal fluid of two patients that responded to antibiotic

therapy with either *L. iners* or *L. crispatus* allowed us to profile their gene expression *in vivo*

to gain more understanding of their different roles in the vaginal microbiota. We extracted the

reads mapped on *L. crispatus* in patient 08_006 (timepoint b-e) and *L. iners* in patient 13_019

(timepoint b-e) and performed a differential expression (DE) analysis using edgeR to compare

their activity profiles based on their KO genes comparing the expression of KO genes of *L.*

crispatus in *L. crispatus* dominated samples (n = 4) with the expression of KO genes of *L.*

iners in *L. iners* dominated samples (n = 4). The Venn diagram in Fig. 5C shows that the two

species share 569 KO genes, while 58 are unique to *L. iners* and 244 are unique for *L.*

crispatus, indicating *L. crispatus* possesses far more diverse functions than *L. iners.* The DE

analysis identified 654 significantly differentially expressed KO genes, of which 393 were up-

regulated in *L. crispatus* (Table S1 sheet 8). Among the top 100 most differentially expressed

KO genes in terms of FDR value, 64 were up-regulated in *L. crispatus*, Remarkably, genes

encoding enzymes involved in the production of H_2O_2 (pyruvate oxidase, NADH oxidase,

glycolate oxidase) (37,38) were highly expressed in *L. crispatus* (Table S1 sheet 8, KO genes

colored in blue). D-lactate dehydrogenase (K03778) was the most highly expressed gene in *L.*

crispatus (log2 CPM = 13.3, Table S1 sheet 8, colored in light green) and this gene is absent

in the genome of *L. iners*. On the other hand, we found that inerolysin (INY) was highly

expressed (log2 CPM =9.8) in *L. iners*, but absent in the genome of *L. crispatus* (Table S1

sheet 8, colored in red). Interestingly, we also found the orthologous gene (K11031) of

inerolysin known as vaginolysin (39) highly expressed in *G. vaginalis* (details in Table S1

sheet 7). Hemolysin C, another pore forming toxin, was highly expressed (log2 CPM = 9.8) in

L. iners but absent in *L. crispatus*.

Discussion

The aim of this study was to identify activity patterns in the vaginal fluid microbiota in BV

and after metronidazole therapy. In particular, we compared the transcriptional profiles of *G.*

vaginalis in vaginal microbiota from patients who did and did not respond to metronidazole

treatment, respectively. This is the first study to investigate the activity alterations of the

vaginal microbiota from patients with BV during treatment with the antibiotic metronidazole

using the metatranscriptomics approach. We found several changes in gene expression in non-

responding patients that might contribute to resistance against metronidazole by either not

activating the pro-drug or repairing DNA damage.

G. vaginalis was the most dominant active species in BV. *G. vaginalis* can be divided into

four phylogenetic subgroups which may in the future be described as subspecies and which

differ in virulence and susceptibility to metronidazole (6,12). We found transcripts from all

four subgroups in all patients, as previously shown based on sequencing of the universal

target cp-60 gene (6,9,11). Interestingly, in those patients that did not respond to treatment,

Gardnerella subgroups A and D which are resistant to metronidazole (14) were slightly more

abundant.

Sequencing of phylogenetic marker genes like the 16S rRNA gene or the cpn60 universal

target is a fast and sensitive method to profile the microbiota composition, but it does not

provide functional information and is prone to PCR bias. Moreover, DNA from dead cells

might also be detected. Therefore, we compared the taxonomic composition of the transcripts

with that of the 16S rRNA genes determined previously in those samples (5). We observed

that *G. vaginalis* comprised on average 47% of all transcripts in BV, while only 5% of 16S

rRNA genes were assigned to this species. This suggests that *G. vaginalis* is transcriptionally

more active than other vaginal bacteria; moreover, the commonly used 27F primer was

previously shown to underrepresent *G. vaginalis* (40). Other differences between the two

methods are caused by the low taxonomic resolution of the 16S rRNA gene, especially of

short amplicons, where a large fraction of 16S rRNA reads is assigned to higher level taxa,

e.g. genus or family. By contrast, the metatranscriptome reads are mapped to genomes and so

have species level resolution. Finally, transcripts can only be mapped if a genome is available.

If the species in question has not yet been cultivated, as for example the BVAB strains, then

reads cannot be assigned. In the periodontal pocket microbiota, about 50% of all reads cannot

be mapped to any bacterial genome (30). By contrast, the vaginal microbiota is much less

diverse and most of its representatives have been cultivated; using the improved ref_Genome

database, we were able to map 86% of all reads, indicating that uncultivated taxa did not

contribute very significantly to the active community in BV and after metronidazole therapy.

Low diversity in health and high diversity in BV is a hallmark of BV and it is so striking that

it has even been suggested to use diversity indices based on PCR amplified 16S rRNA genes

in addition to the clinical diagnosis based on Amsel criteria and Nugent score (4,41-43). Our

comparison between communities in BV and after metronidazole therapy on the levels of (1)

16S rRNA gene, (2) taxonomic composition of total transcripts, and (3) functional profiling

based on KO genes shows a reversal of this observation: BV communities, although highly

diverse on the taxonomic level, cluster tightly together on the functional level of KO genes.

On the contrary, non-BV communities are similar on the taxonomic level, but highly diverse

among individuals on the functional level.

In our metatranscriptome analysis we found evidence for mechanisms that hinder the

activation of the metronidazole prodrug, or mitigate the damage that metronidazole inflicts on

DNA, and thus could be important reasons for the lack of response in some women.

We show that the ferredoxin gene of *G. vaginalis* was less active in those patients that did not

respond to metronidazole. As an electron carrier, ferredoxin is downregulated in *H. pylori*

bacteria grown in the presence of metronidazole (17). It is required for activation of the

prodrug in *H. pylori* (44) and might have a similar role in *G. vaginalis*. Lack of response

might result from lack of activation of the prodrug. Unexpectedly, the nitroimidazole

resistance (*nim*) gene, which has been shown to mediate resistance to metronidazole in *B.*

fragilis by transforming metronidazole to a non-toxic amino derivative (16) was not highly

expressed in non-responders. This could be due to technical problems, since the Nim protein

sequence contains only partial CDS (https://www.ebi.ac.uk/ena/data/view/AGN03877).

Moreover, *nim*-negative strains of *B. fragilis* can tolerate high levels of metronidazole

indicating the importance of other mechanisms of resistance (16).

Remarkably, *cas* genes of *G. vaginalis* were highly up-regulated in samples from patients that

did not respond to metronidazole treatment. The CRISPR-Cas genes are present in about half

of all Bacteria and most Archaea (45); they represent a mechanism of adaptive immunity

which protects the prokaryotic cell against foreign DNA and has been developed into a

universal tool for genome editing (46). The *cas* genes of *G. vaginalis* belong to the *E. coli*

subtype and were found in about half of the clinical isolates (36). Their up-regulation might

reflect increased phage attacks in BV. Phages have been hypothesized to be crucial for the

etiology of BV by causing the collapse of *Lactobacillus* populations (47); accordingly, *L.*

iners upregulates its CRISPR-Cas system in BV (22). More than 400 annotated prophage

sequences were found in 39 *Gardenerella* strains (29). They might be induced to enter the

lytic cycle by the change in pH accompanying the shift to BV. However, the viral transcripts

contributed 0.1% of the total metatranscriptome in both non-responders and responders before

treatment.

The upregulation of CRISPR-Cas system genes in *G. vaginalis* from those patients that did

not respond to treatment by metronidazole suggests that the CRISPR-Cas system might have a

role in mitigating the DNA damaging effect of metronidazole. In addition to providing

adaptive immunity, CRISPR-Cas systems can have various additional functions (48) and it

was shown that they can protect the cell against DNA damaging agents (49). The Cas1

enzyme of *E. coli* (YgbT) physically and genetically interacts with the DNA repair system

(RecBC, RuvB) and is recruited to DNA double strand breaks; moreover, YgbT is necessary

for resistance of *E. coli* to DNA damage caused by the genotoxic antibiotic mitomycin C or

UV light (49). Our findings suggest that the CRISPR-Cas system may protect the vaginal

microbiota against the DNA damaging effect of metronidazole. If experimentally confirmed,

this finding might open a new path to fight bacterial resistance against DNA damaging agents.

For example, it would be worth testing if the susceptibility to metronidazole can be modified

in *Gardnerella* isolates and possibly other vaginal pathogens according to the expression level

of *cas* genes. It is not known how up-regulation of *cas* genes is regulated in the vaginal

microbiota. It might be a response to phage attack, thus by suppressing *cas* genes the

susceptibility to phages might be increased simultaneously with the susceptibility to

metronidazole. Using CRISPR engineered phages for therapy of dysbiotic communities has

been considered as one of many options of new therapeutic strategies based on a deeper

understanding of the human microbiome (50).

L. iners and *L. crispatus* dominate their respective CST in the healthy vaginal microbiota.

There were many factors observed by laboratory or genomic studies (37,51,52) which suggest

that more protection against dysbiosis is provided by *L. crispatus* rather than by *L. iners*. Here

we analyzed which genes are actually highly expressed *in vivo*: We observed that genes

encoding proteins for the production of H_2O_2 and D-lactic acid were highly expressed in *L.*

crispatus. H_2O_2 inhibits BV associated bacteria, but it has been questioned if its level in the

vaginal milieu is high enough, and it was suggested that lactic acid is more protective (53). In

L. iners, the genes for the pore-forming toxins inerolysin and hemolysin C were highly active,

supporting the hypothesis that *L. iners* may play an ambiguous role in the vaginal econiche

and is associated with vaginal dysbiosis (26,54).

Conclusions

This first study of the *in vivo* transcriptional activity of vaginal fluid microbiota during

metronidazole treatment of BV focused on possible reasons for the lack of response to

antibiotic therapy in some patients. Genes related to activation of the prodrug and repairing

the DNA damage caused by metronidazole were shown to be differentially expressed in

responders and non-responders. A completely new role for Cas proteins is hypothesized

which warrants closer inspection and may help to develop more efficient novel therapies to

improve the treatment of BV.

Material and Methods

Study design

Vaginal fluid samples of women analyzed here were a subset of the samples obtained during a

randomized controlled clinical trial described previously (5). The trial protocol was approved

by the local ethics committee (Ärztekammer Nordrhein - Medical Association North Rhine)

and written consent was obtained from all participants. The clinical trial was conducted in

accordance with the Declaration of Helsinki on Ethical Principles for Medical Research

Involving Human Subjects. Principles and guidelines for good clinical practice were followed.

The study was registered on ClinicalTrials.gov with the identifier NCT02687789. Briefly,

women were included into the clinical trial if they were BV positive according to Amsel

criteria and Nugent score and were biofilm positive on vaginal epithelial cells and positive for

extracellular polysaccharides (EPS) in urine. For treatment of acute BV, they received 2 g of

metronidazole orally and were afterwards treated with an intravaginal pessary for three weeks,

twice a week. Samples were taken during acute BV (visit 1), after receiving metronidazole 7

to 28 days after visit 1 (visit 2), after pessary application one week after visit 2 (visit 3), after

continued pessary application two weeks after visit 3 (visit 4) and during follow up three

months after visit 4 (visit 5).

The aim of the clinical trial had been to compare the effectiveness of two different types of

pessary. The results and the taxonomic composition of the vaginal microbial communities

have been reported (5). For the metatranscriptome analysis reported here, we chose a subset

of 14 patients from the clinical trial. These 14 patients consisted of two groups named "with

response to treatment" (n = 8) and "no response to treatment" (n = 6) (Fig. 1). Among the

eight patients who responded to treatment, six had no recurrence and two experienced

recurrence during the three months follow-up. For the analysis of lack of response to

metronidazole, samples from all 14 patients were analyzed at two timepoints, acute BV (visit

1) and 7 to 28 days after antibiotic treatment (visit 2) (28 samples in total). For the analysis of

recurrence, samples from all 5 visits were analyzed for 4 patients (two without recurrence and

two with) (20 samples in total). These four patients all received the commercially available

lactic acid pessary after metronidazole therapy at visit 3 and 4. Nugent score >6 was used to

determine BV status since it is considered the gold standard for BV diagnosis (55) (Table S1

sheet 1). The BV status at visit 5 was determined by Amsel criteria as there was no Nugent

score available at that timepoint. The sample ID was obtained by concatenating the patient ID

and letters "a" to "e" indicating visit 1 to 5.

Sample collection and transport

Vaginal fluid was obtained by infusing 2 ml of saline solution into the vagina followed by

rotation against the vaginal wall with a speculum and then collecting the vaginal fluid with a

syringe. Approximately 700 µl were immediately transferred to a tube containing 2 ml

RNAprotect (Qiagen, Germany). Tubes were immediately frozen at -20°C, transported at -

20°C within a week and stored at -70°C.

RNA extraction and mRNA enrichment

RNA was extracted from 1 ml vaginal fluid suspension using the MO BIO

PowerMicrobiome™ RNA Isolation Kit (Qiagen, Germany) with pretreatment: Vaginal fluid

was centrifuged at 13,000 rpm for 1 minute. The pellet was resuspended in MoBio lysis buffer

and this suspension was added to the supplied bead tubes filled with 500 µl ice-cold

Phenol:Chloroform:Isoamyl Alcohol solution (Carl Roth, Germany). The bead-suspension

mix was shaken at 5 m/s for 1 minute in 3 intervals which were 2 minutes apart using the MO

BIO PowerLyzer™ (Qiagen, Germany). After centrifugation for 1 minute at 13,000 rpm and

4°C the upper phase containing the RNA was further processed according to the

manufacturer's instructions including DNAse I treatment. RNA was eluted in 100µl nuclease

free water and vacuum concentrated to 50 µl. The Ribo-Zero Gold rRNA Removal Kit

(Epidemiology) by Illumina (USA) was then used for mRNA enrichment with ethanol

precipitation according to the manufacturer's instructions. Integrity of RNA was evaluated

using a Bioanalyzer 2100 (Agilent, Germany).

Library preparation, sequencing and preprocessing of sequencing data

Paired-end mRNA Illumina sequencing libraries were constructed with the Script Seq

Illumina Kit. Strand specific paired end sequencing was performed on the HiSeq 2500

Sequencer to yield 2×110 bp paired-end reads. Primers and sequencing adaptors were

removed from raw sequencing data, followed by clipping the bases with quality score < 20

from the reads using Fastq-Mcf (56). After clipping, the remaining reads shorter than 50 were

removed. Thereafter, the ribosomal RNA reads were eliminated using SortMeRNA v2.0 (57)

with the default parameters.

Taxonomy assignment using Kraken

Kraken (27), an accurate and ultra-fast taxonomy assignment tool for metagenomes was used

to determine the taxonomic composition of the metatranscriptome data. Kraken uses the K-

mer strategy and the lowest common ancestor (LCA) algorithm to affiliate a given read to a

taxon. The standard Kraken database with addition of the human genome was used to identify

human reads. This standard database consists of prokaryote genomes (2786), virus genomes

(4418). The human genome (ver. GRCh38) was additionally downloaded from NCBI.

The ref_Genome database contained 163 bacterial genomes from 105 species of bacteria,

including 147 genomes from the urogenital subset of the HMP reference genome sequence

data (HMRGD). The complete list of reference genomes in the ref_Genome database can be

found in Table S1 sheet 3. All results on the taxonomic composition in this study were

achieved based on this database.

Short reads alignment by BWA

Kraken was used for taxonomy classification, while BWA was applied to determine the

expression of genes. A reference gene database named ref_Gene was constructed which

contained the genes from the urogenital tract subset of the HMRGD and the genes from 9

additional genomes (Table S1 sheet 4). The genes of *Gardnerella sp.* 26-12, *Gardnerella sp.*

30-4 could not be included because only less than half of their CDS are available (around

1000 genes). The short reads alignment was performed using BWA with the BWA-MEM (58)

algorithm. A mapping seed length of 31 which is much longer than the default seed length 19

was applied to achieve reliable alignments. Reads that mapped with mapping quality score

(MAPQ) lower than 10 were excluded. MAPQ contains the Phred-scaled posterior probability

that the mapping position is wrong (59).

KEGG Orthologous (KO) gene annotation of ref_Gene database

The ref_Gene database was annotated using KEGG prokaryote protein sequences. The KEGG

prokaryote protein sequence database represents a non-redundant protein dataset of Bacteria

and Archaea on the species level and contains about 7 million non-redundant peptide

sequences grouped into 14,390 distinct KO genes. A KO gene contains several genes from

different species with similar function. DIAMOND (60), a much faster alternative to

BLASTX was applied to map the ref_Gene sequences against the KEGG prokaryote protein

sequence database with its "more sensitive mode". To obtain reliable annotation, only

alignments with sequence identity >=50 and E-value <= 1e-5 and query coverage >= 70%

were taken into account. By annotating the genes in the ref_Gene database to KO genes, we

were able to determine the expression profile of KO genes, and investigate the activity shifts

in BV based on differential expression analysis of KO genes. The cumulative dominance

analysis and PCA were carried out using Primer 7 (61).

Differential expression (DE) analysis

All differential expression (DE) analyses were performed using the R package edgeR (62).

The Benjamini Hochberg (BH) method was used to correct the p value of DE analysis with

the false discovery rate (FDR) for multiple comparisons. Genes with FDR smaller than 0.05

were considered as significantly differentially regulated. The sample groups defined for each

comparison are listed in Table S1 sheet 1.

Detection of putative metronidazole resistance related genes

To detect the expression of previously reported putative metronidazole resistance genes such

as recA, recA-mediated autopeptidase (Rma), peroxiredoxin, nitroimidazole resistance protein

(NIM), ferredoxin/ferredoxin-NADP reductase (FNR), nitroreductase and ferredoxin, we

examined the expression level of these genes for *G. vaginalis* in the vaginal community from

patients without response to metronidazole treatment (n = 6) as well as with response (n = 4).

As most of these genes do not have corresponding KO genes, we annotated the ref_Gene

database based on the sequences of these genes using BLASTN. The sequences were

retrieved from ENA by key words of each "ferredoxin, NADPH flavin oxidoreductase,

nitroreductase, peroxiredoxin, pyruvate ferredoxin oxidoreductase, recA, nitroimidazole

resistance" plus *G. vaginalis*. In total, 155 unique sequences of *G. vaginalis* were obtained for

the annotation of ref_Gene database. The identification of duplicate sequences was done by

SeqKit (63).

Availability of data and material

542   The sequencing data have been deposited in the European Nucleotide Archive with accession

543   number PRJEB21446.

Authors' contributions

545   This study was designed by IWD and CG. CM and CA provided the clinical samples. RNA

546   extraction and mRNA enrichment was performed by CG. SB prepared cDNA libraries, and

547   performed Illumina sequencing. Z-LD performed all the data analyses. Data interpretation and

548   visualization were done by IWD, Z-LD and CG. Z-LD and CG wrote the manuscript draft,

549   and all authors reviewed the manuscript.

Acknowledgements

551   This work was supported by ZIM project grant KF3134201MD3 of the Bundesministerium

552   für Wirtschaft und Energie (BMWi), Germany. The clinical study was funded by Dr. August

553   Wolff GmbH & Co. KG Arzneimittel within the ZIM project KF3134201MD3.

555   Reference

1. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle
R, Russell J, Tacket CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011.
Vaginal microbiome of reproductive-age women. Proc Natl Acad Sci U S A 108 Suppl
1:4680-4687. doi:1002611107 [pii];10.1073/pnas.1002611107 [doi].

2. van de Wijgert JH, Borgdorff H, Verhelst R, Crucitti T, Francis S, Verstraelen H,
Jespers V. 2014. The vaginal microbiota: what have we learned after a decade of
molecular characterization? PLoS One 9:e105998. doi:10.1371/journal.pone.0105998
[doi];PONE-D-14-20877 [pii].

3. Onderdonk AB, Delaney ML, Fichorova RN. 2016. The Human Microbiome during
Bacterial Vaginosis. Clin Microbiol Rev 29:223-238. doi:29/2/223
[pii];10.1128/CMR.00075-15 [doi].

4. Dols JA, Molenaar D, van der Helm JJ, Caspers MP, de KA-B, Schuren FH,
Speksnijder AG, Westerhoff HV, Richardus JH, Boon ME, Reid G, de Vries HJ, Kort R.
2016. Molecular assessment of bacterial vaginosis by Lactobacillus abundance and
species diversity. BMC Infect Dis 16:180. doi:10.1186/s12879-016-1513-3
[doi];10.1186/s12879-016-1513-3 [pii].

5. Gottschick C, Deng Z-L, Vital M, Masur C, Abels C, Pieper D, Rohde M, Mendling W,
Wagner-Dobler I. 2017. Treatment of biofilms in bacterial vaginosis by an amphoteric
tenside pessary - clinical study and microbiome analysis. Microbiome 5:119.

6. Schellenberg JJ, Patterson MH, Hill JE. 2017. Gardnerella vaginalis diversity and
ecology in relation to vaginal symptoms. Res Microbiol 168:837-844. doi:S0923-
2508(17)30060-8 [pii];10.1016/j.resmic.2017.02.011 [doi].

7. Ahmed A, Earl J, Retchless A, Hillier SL, Rabe LK, Cherpes TL, Powell E, Janto B,
Eutsey R, Hiller NL, Boissy R, Dahlgren ME, Hall BG, Costerton JW, Post JC, Hu FZ,
Ehrlich GD. 2012. Comparative genomic analyses of 17 clinical isolates of Gardnerella
vaginalis provide evidence of multiple genetically isolated clades consistent with
subspeciation into genovars. J Bacteriol 194:3922-3937. doi:JB.00056-12
[pii];10.1128/JB.00056-12 [doi].

8. Hill JE, Goh SH, Money DM, Doyle M, Li A, Crosby WL, Links M, Leung A, Chan D,
Hemmingsen SM. 2005. Characterization of vaginal microflora of healthy, nonpregnant
women by chaperonin-60 sequence-based methods. Am J Obstet Gynecol 193:682-692.
doi:S0002-9378(05)00340-6 [pii];10.1016/j.ajog.2005.02.094 [doi].

9. Paramel JT, Schellenberg JJ, Hill JE. 2012. Resolution and characterization of distinct
cpn60-based subgroups of Gardnerella vaginalis in the vaginal microbiota. PLoS One
7:e43009. doi:10.1371/journal.pone.0043009 [doi];PONE-D-12-12121 [pii].

10. Chaban B, Links MG, Jayaprakash TP, Wagner EC, Bourque DK, Lohn Z, Albert AY,
van SJ, Reid G, Hemmingsen SM, Hill JE, Money DM. 2014. Characterization of the
vaginal microbiota of healthy Canadian women through the menstrual cycle.
Microbiome 2:23. doi:10.1186/2049-2618-2-23 [doi];2049-2618-2-23 [pii].

11. Albert AY, Chaban B, Wagner EC, Schellenberg JJ, Links MG, van SJ, Reid G,
Hemmingsen SM, Hill JE, Money D. 2015. A Study of the Vaginal Microbiome in
Healthy Canadian Women Utilizing cpn60-Based Molecular Profiling Reveals Distinct
Gardnerella Subgroup Community State Types. PLoS One 10:e0135620.
doi:10.1371/journal.pone.0135620 [doi];PONE-D-15-15097 [pii].

12. Schellenberg JJ, Paramel JT, Withana GN, Patterson MH, Vaneechoutte M, Hill JE.
2016. Gardnerella vaginalis Subgroups Defined by cpn60 Sequencing and Sialidase
Activity in Isolates from Canada, Belgium and Kenya. PLoS One 11:e0146510.
doi:10.1371/journal.pone.0146510 [doi];PONE-D-15-36020 [pii].

13. Bradshaw CS, Morton AN, Garland SM, Horvath LB, Kuzevska I, Fairley CK. 2005.
Evaluation of a point-of-care test, BVBlue, and clinical and laboratory criteria for
diagnosis of bacterial vaginosis. J Clin Microbiol 43:1304-1308. doi:43/3/1304
[pii];10.1128/JCM.43.3.1304-1308.2005 [doi].

14. Schuyler JA, Mordechai E, Adelson ME, Sobel JD, Gygax SE, Hilbert DW. 2016.
Identification of intrinsically metronidazole-resistant clades of Gardnerella vaginalis.
Diagn Microbiol Infect Dis 84:1-3. doi:S0732-8893(15)00360-0
[pii];10.1016/j.diagmicrobio.2015.10.006 [doi].

15. Bradshaw CS, Pirotta M, De GD, Hocking JS, Morton AN, Garland SM, Fehler G,
Morrow A, Walker S, Vodstrcil LA, Fairley CK. 2012. Efficacy of oral metronidazole
with vaginal clindamycin or vaginal probiotic for bacterial vaginosis: randomised
placebo-controlled double-blind trial. PLoS One 7:e34540.
doi:10.1371/journal.pone.0034540 [doi];PONE-D-11-24250 [pii].

16. Lofmark S, Edlund C, Nord CE. 2010. Metronidazole is still the drug of choice for
treatment of anaerobic infections. Clin Infect Dis 50 Suppl 1:S16-S23.
doi:10.1086/647939 [doi].

17. Mendz GL, Megraud F. 2002. Is the molecular basis of metronidazole resistance in
microaerophilic organisms understood? Trends Microbiol 10:370-375.
doi:S0966842X02024058 [pii].

18. Lofmark S, Fang H, Hedberg M, Edlund C. 2005. Inducible metronidazole resistance
and nim genes in clinical Bacteroides fragilis group isolates. Antimicrob Agents
Chemother 49:1253-1256. doi:49/3/1253 [pii];10.1128/AAC.49.3.1253-1256.2005
[doi].

19. Steffens LS, Nicholson S, Paul LV, Nord CE, Patrick S, Abratt VR. 2010. Bacteroides
fragilis RecA protein overexpression causes resistance to metronidazole. Res Microbiol
161:346-354. doi:S0923-2508(10)00085-9 [pii];10.1016/j.resmic.2010.04.003 [doi].

20. Hilbert DW, Smith WL, Paulish-Miller TE, Chadwick SG, Toner G, Mordechai E,
Adelson ME, Sobel JD, Gygax SE. 2016. Utilization of molecular methods to identify
prognostic markers for recurrent bacterial vaginosis. Diagn Microbiol Infect Dis.
doi:S0732-8893(16)30197-3 [pii];10.1016/j.diagmicrobio.2016.07.003 [doi].

21. Ferreira CST, Donders GG, Parada CMGL, Tristao ADR, Fernandes T, da Silva MG,
Marconi C. 2017. Treatment failure of bacterial vaginosis is not associated with higher
loads of Atopobium vaginae and Gardnerella vaginalis. J Med Microbiol 66:1217-1224.
doi:10.1099/jmm.0.000561 [doi].

22. Macklaim JM, Fernandes AD, Di Bella JM, Hammond JA, Reid G, Gloor GB. 2013.
Comparative meta-RNA-seq of the vaginal microbiota and differential expression by

Lactobacillus iners in health and dysbiosis. Microbiome 1:12. doi:2049-2618-1-12
[pii];10.1186/2049-2618-1-12 [doi].

23. Bradshaw CS, Morton AN, Hocking J, Garland SM, Morris MB, Moss LM, Horvath
LB, Kuzevska I, Fairley CK. 2006. High recurrence rates of bacterial vaginosis over the
course of 12 months after oral metronidazole therapy and factors associated with
recurrence. J Infect Dis 193:1478-1486. doi:JID36020 [pii];10.1086/503780 [doi].

24. Sobel JD, Schmitt C, Meriwether C. 1993. Long-term follow-up of patients with
bacterial vaginosis treated with oral metronidazole and topical clindamycin. J Infect Dis
167:783-784.

25. Turovskiy Y, Sutyak NK, Chikindas ML. 2011. The aetiology of bacterial vaginosis. J
Appl Microbiol 110:1105-1128. doi:10.1111/j.1365-2672.2011.04977.x [doi].

26. Petrova MI, Reid G, Vaneechoutte M, Lebeer S. 2016. Lactobacillus iners: Friend or
Foe? Trends Microbiol. doi:S0966-842X(16)30181-0 [pii];10.1016/j.tim.2016.11.007
[doi].

27. Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification
using exact alignments. Genome Biol 15:R46. doi:gb-2014-15-3-r46 [pii];10.1186/gb-
2014-15-3-r46 [doi].

28. The Human Microbiome Project Consortium. 2012. A framework for human
microbiome research. Nature 486:215-221. doi:10.1038/nature11209.

29. Malki K, Shapiro JW, Price TK, Hilt EE, Thomas-White K, Sircar T, Rosenfeld AB,
Kuffel G, Zilliox MJ, Wolfe AJ, Putonti C. 2016. Genomes of Gardnerella Strains
Reveal an Abundance of Prophages within the Bladder Microbiome. PLoS One
11:e0166757. doi:10.1371/journal.pone.0166757 [doi];PONE-D-16-20473 [pii].

30. Deng ZL, Szafranski SP, Jarek M, Bhuju S, Wagner-Dobler I. 2017. Dysbiosis in
chronic periodontitis: Key microbial players and interactions with the human host. Sci
Rep 7:3703. doi:10.1038/s41598-017-03804-8 [doi];10.1038/s41598-017-03804-8 [pii].

31. Hellinga HW, Evans PR. 1985. Nucleotide sequence and high-level expression of the
major Escherichia coli phosphofructokinase. Eur J Biochem 149:363-373.

32. Arnold RJ, Reilly JP. 1999. Observation of Escherichia coli ribosomal proteins and their
posttranslational modifications by mass spectrometry. Anal Biochem 269:105-112.
doi:S0003-2697(98)93077-9 [pii];10.1006/abio.1998.3077 [doi].

33. Bubunenko M, Baker T, Court DL. 2007. Essentiality of ribosomal and transcription
antitermination proteins analyzed by systematic gene replacement in Escherichia coli. J
Bacteriol 189:2844-2853. doi:JB.01713-06 [pii];10.1128/JB.01713-06 [doi].

34. Webb AJ, Homer KA, Hosie AH. 2008. Two closely related ABC transporters in
Streptococcus mutans are involved in disaccharide and/or oligosaccharide uptake. J
Bacteriol 190:168-178. doi:JB.01509-07 [pii];10.1128/JB.01509-07 [doi].

35.   Shipkowski S, Brenchley JE. 2006. Bioinformatic, genetic, and biochemical evidence
679   that some glycoside hydrolase family 42 beta-galactosidases are arabinogalactan type I
680   oligomer hydrolases. Appl Environ Microbiol 72:7730-7738. doi:AEM.01306-06
681   [pii];10.1128/AEM.01306-06 [doi].

682   36.   Pleckaityte M, Zilnyte M, Zvirbliene A. 2012. Insights into the CRISPR/Cas system of
683   Gardnerella vaginalis. BMC Microbiol 12:301. doi:1471-2180-12-301
684   [pii];10.1186/1471-2180-12-301 [doi].

685   37.   Ojala T, Kankainen M, Castro J, Cerca N, Edelman S, Westerlund-Wikstrom B, Paulin
686   L, Holm L, Auvinen P. 2014. Comparative genomics of Lactobacillus crispatus suggests
687   novel mechanisms for the competitive exclusion of Gardnerella vaginalis. BMC
688   Genomics 15:1070. doi:1471-2164-15-1070 [pii];10.1186/1471-2164-15-1070 [doi].

689   38.   Hertzberger R, Arents J, Dekker HL, Pridmore RD, Gysler C, Kleerebezem M, de
690   Mattos MJ. 2014. H(2)O(2) production in species of the Lactobacillus acidophilus
691   group: a central role for a novel NADH-dependent flavin reductase. Appl Environ
692   Microbiol 80:2229-2239. doi:AEM.04272-13 [pii];10.1128/AEM.04272-13 [doi].

693   39.   Gelber SE, Aguilar JL, Lewis KL, Ratner AJ. 2008. Functional and phylogenetic
694   characterization of Vaginolysin, the human-specific cytolysin from Gardnerella
695   vaginalis. J Bacteriol 190:3896-3903. doi:JB.01965-07 [pii];10.1128/JB.01965-07 [doi].

696   40.   Frank JA, Reich CI, Sharma S, Weisbaum JS, Wilson BA, Olsen GJ. 2008. Critical
697   evaluation of two primers commonly used for amplification of bacterial 16S rRNA
698   genes. Appl Environ Microbiol 74:2461-2470. doi:AEM.02272-07
699   [pii];10.1128/AEM.02272-07 [doi].

700   41.   Srinivasan S, Hoffman NG, Morgan MT, Matsen FA, Fiedler TL, Hall RW, Ross FJ,
701   McCoy CO, Bumgarner R, Marrazzo JM, Fredricks DN. 2012. Bacterial communities in
702   women with bacterial vaginosis: high resolution phylogenetic analyses reveal
703   relationships of microbiota to clinical criteria. PLoS One 7:e37818.
704   doi:10.1371/journal.pone.0037818 [doi];PONE-D-12-04403 [pii].

705   42.   Ling Z, Kong J, Liu F, Zhu H, Chen X, Wang Y, Li L, Nelson KE, Xia Y, Xiang C.
706   2010. Molecular analysis of the diversity of vaginal microbiota associated with bacterial
707   vaginosis. BMC Genomics 11:488. doi:1471-2164-11-488 [pii];10.1186/1471-2164-11-
708   488 [doi].

709   43.   Oakley BB, Fiedler TL, Marrazzo JM, Fredricks DN. 2008. Diversity of human vaginal
710   bacterial communities and associations with clinically defined bacterial vaginosis. Appl
711   Environ Microbiol 74:4898-4909. doi:AEM.02884-07 [pii];10.1128/AEM.02884-07
712   [doi].

713   44.   Gerrits MM, van der Wouden EJ, Bax DA, van Zwet AA, van Vliet AH, de JA, Kusters
714   JG, Thijs JC, Kuipers EJ. 2004. Role of the rdxA and frxA genes in oxygen-dependent
715   metronidazole resistance of Helicobacter pylori. J Med Microbiol 53:1123-1128.
716   doi:10.1099/jmm.0.45701-0 [doi].

45.   Makarova KS, Wolf YI, Alkhnbashi OS, Costa F, Shah SA, Saunders SJ, Barrangou R,
Brouns SJ, Charpentier E, Haft DH, Horvath P, Moineau S, Mojica FJ, Terns RM, Terns
MP, White MF, Yakunin AF, Garrett RA, van der Oost J, Backofen R, Koonin EV.
2015. An updated evolutionary classification of CRISPR-Cas systems. Nat Rev
Microbiol 13:722-736. doi:nrmicro3569 [pii];10.1038/nrmicro3569 [doi].

46.   Doudna JA, Charpentier E. 2014. Genome editing. The new frontier of genome
engineering with CRISPR-Cas9. Science 346:1258096. doi:346/6213/1258096
[pii];10.1126/science.1258096 [doi].

47.   Blackwell AL. 1999. Vaginal bacterial phaginosis? Sex Transm Infect 75:352-353.

48.   Westra ER, Buckling A, Fineran PC. 2014. CRISPR-Cas systems: beyond adaptive
immunity. Nat Rev Microbiol 12:317-326. doi:nrmicro3241 [pii];10.1038/nrmicro3241
[doi].

49.   Babu M, Beloglazova N, Flick R, Graham C, Skarina T, Nocek B, Gagarinova A,
Pogoutse O, Brown G, Binkowski A, Phanse S, Joachimiak A, Koonin EV, Savchenko
A, Emili A, Greenblatt J, Edwards AM, Yakunin AF. 2011. A dual function of the
CRISPR-Cas system in bacterial antivirus immunity and DNA repair. Mol Microbiol
79:484-502. doi:10.1111/j.1365-2958.2010.07465.x [doi].

50.   Belizario JE, Napolitano M. 2015. Human microbiomes and their roles in dysbiosis,
common diseases, and novel therapeutic approaches. Front Microbiol 6:1050.
doi:10.3389/fmicb.2015.01050 [doi].

51.   France MT, Mendes-Soares H, Forney LJ. 2016. Genomic Comparisons of
Lactobacillus crispatus and Lactobacillus iners Reveal Potential Ecological Drivers of
Community Composition in the Vagina. Appl Environ Microbiol 82:7063-7073.
doi:AEM.02385-16 [pii];10.1128/AEM.02385-16 [doi].

52.   Rampersaud R, Planet PJ, Randis TM, Kulkarni R, Aguilar JL, Lehrer RI, Ratner AJ.
2011. Inerolysin, a cholesterol-dependent cytolysin produced by Lactobacillus iners. J
Bacteriol 193:1034-1041. doi:JB.00694-10 [pii];10.1128/JB.00694-10 [doi].

53.   O'Hanlon DE, Moench TR, Cone RA. 2011. In vaginal fluid, bacteria associated with
bacterial vaginosis can be suppressed with lactic acid but not hydrogen peroxide. BMC
Infect Dis 11:200. doi:1471-2334-11-200 [pii];10.1186/1471-2334-11-200 [doi].

54.   Vaneechoutte M. 2017. Lactobacillus iners, the unusual suspect. Res Microbiol
168:826-836. doi:S0923-2508(17)30151-1 [pii];10.1016/j.resmic.2017.09.003 [doi].

55.   Mohammadzadeh F, Dolatian M, Jorjani M, Alavi MH. 2014. Diagnostic value of
Amsel's clinical criteria for diagnosis of bacterial vaginosis. Glob J Health Sci 7:8-14.
doi:10.5539/gjhs.v7n3p8 [doi].

56.   Aronesty E. 2011. ea-utils: Command-line tools for processing biological sequencing
data. Durham, NC: Expression Analysis.

57. Kopylova E, Noe L, Touzet H. 2012. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics 28:3211-3217. doi:bts611
[pii];10.1093/bioinformatics/bts611 [doi].

58. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv preprint arXiv:1303 3997.

59. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics
25:2078-2079. doi:btp352 [pii];10.1093/bioinformatics/btp352 [doi].

60. Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using
DIAMOND. Nat Methods 12:59-60. doi:nmeth.3176 [pii];10.1038/nmeth.3176 [doi].

61. Clarke KR. 1993. Non-parametric multivariate analyses of changes in community
structure. Austral ecology 18:117-143.

62. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26:139-
140. doi:btp616 [pii];10.1093/bioinformatics/btp616 [doi].

63. Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. PLoS One 11:e0163962.
doi:10.1371/journal.pone.0163962 [doi];PONE-D-16-20820 [pii].

Figures, tables and additional files

Fig. 1: Study design and taxonomic composition of vaginal fluid metatranscriptomes in BV and after treatment with metronidazole. (A). Time course of the clinical study. (B) Taxonomic composition of the metatranscriptome at visit 1 (diagnosis) and visit 2 (after metronidazole therapy). (C) Cumulative dominance of the vaginal microbiota in BV and non-BV. (D) The subspecies composition of the *G. vaginalis* sub-community. Species with average relative abundance smaller than 0.5% were grouped into "Others". The red dot on top of the samples indicates BV. The digits indicate the patient ID, while the letters a-b denote visit 1-2. After the first sampling at visit 1 the patients were treated with metronidazole. Total putative bacterial mRNA reads were mapped to the ref_Genome database using Kraken (see

Methods for details). BV status was determined by Nugent score. In C, the "BV" and "NBV"

in the parenthesis indicate BV and non-BV, respectively.

Fig. 2: Average taxonomic composition of vaginal fluid samples in BV determined by

16S rRNA amplicon sequencing (A) and metatranscriptome sequencing (B). (A)

Amplicon sequencing was performed as described in our previous study (5) using primers V1-

V2. (B) The taxonomy was assigned based on all cleaned reads after removal of human reads

using Kraken and the ref_Genome database. The top 12 most abundant taxa for each approach

are shown in A and B. Relative average abundance was calculated based on all mapped reads.

Mean and standard error are shown.

Fig. 3: Principal components analysis (PCA) based on taxonomic profiles and activity

profiles in BV and after metronidazole therapy (non-BV). (A) The PCA plot based on

taxonomic profile using 16S rRNA gene sequencing. (B) The PCA based on taxonomic

composition determined by metatranscriptome. (C) The PCA plot based on KO gene

expression profile. The communities at visit 1 (BV) and visit 2 (after metronidazole therapy,

non-BV) from 14 patients are shown. In the PCA biplots of taxonomy composition (Fig. 3A-

B), the taxa with multivariate (multiple) correlation higher than 0.3 are illustrated, while for

PCA (Fig. 3C) of KO profiles the KO genes with correlation >0.2 are shown.

Fig. 4: Changes in gene expression of *G. vaginalis* in patients responding to antibiotic

treatment compared to non-responders. (A) Expression of putative metronidazole

resistance associated genes of *G. vaginalis* in vaginal fluid microbiota. (B) Differential

expression of KO genes: Seven *cas* genes of *G. vaginalis* were highly up-regulated in

communities from patients who did not respond to the treatment. (A) The expression

value was calculated based on relative abundance of reads mapped onto *G. vaginalis* using

BWA. "NR1" (No Response 1) indicates the BV samples from six patients that did not

respond to metronidazole treatment; "WR1" (With Response 1) represents the BV samples

from four patients which afterwards responded to metronidazole. The dotplot illustrates the

log2FC of the corresponding activity between *G. vaginalis* from non-responders and

responders. The values in the heatmap were scaled using Z-sore. In the figure legend, "exp."

indicates the relative expression level. (B) "NR1" samples were compared with "WR1". KO

genes with FDR <= 0.05 are colored in red or turquoise (significantly differentially regulated)

while FDR > 0.05 are in grey.

Fig. 5: Shifts in the vaginal microbiome over 3 months. (A) Taxonomic composition of

the metatranscriptome in two patients that were stably non-BV (without recurrence)

and two patients that experienced recurrence. (B) PCA of activity profiles based on KO

genes from the same patients. (C) Gene expression *in vivo* of *L. crispatus* and *L. iners*. (A)

Acute BV and recurrence according to Nugent score are indicated as red dot. (B) Two women

with recurrence (pink and blue color range) and two women without recurrence (green and

orange color range) are shown. In the figure legend, BV indicates timepoint with BV, "R"

indicates the recurrence and "H" represents health. The green and blue circles highlight

healthy clusters, respectively, while the red circle highlights samples from BV. The arrows

denote the temporal shifts of the communities during the treatment. (C) The Venn diagram

indicates the unique KO genes of *L. crispatus* and *L. iners* as well as their shared KO genes.

The innermost ring denotes the expression of KO genes by log2CPM, the outer ring illustrates

the fold change of the expression of KO genes between *L. crispatus* dominated communities

and *L. iners* dominated communities by log2FC. The KO genes are in descending order based

on log2CPM. The small red triangles mark the inerolysin and hemolysin C genes, while blue

and green triangles mark the genes encoding proteins involved in the production of D-lactic

acid and hydrogen peroxide, respectively.

Supplementary Information

Fig. S1: Taxonomic composition of communities on the species level determined by

metatranscriptome sequencing in non-BV and BV. The species present in at least 2

samples with relative abundance >1% are shown. The *Gardnerella* bladder isolates are

illustrated separately from *G. vaginalis*. The sample name in red indicates the first BV

incidence of patients with recurrence and purple depicts the second incidence.

Supplementary Table S1: All supplementary data

Sheet 1: Sample description

Sheet 2: Read summary

Sheet 3: Genomes in the ref_Genome database for taxonomic assignment

Sheet 4: Genomes in the ref_Gene database with a total of 301,323 genes for functional

assignment with BWA

Sheet 5: Species composition determined by Kraken based on the ref_Genome database

for taxonomic assignment

Sheet 6: Comparison of the community composition determined by 16S rRNA gene

amplicon sequencing (V1-V2) and metatranscriptome sequencing

Sheet 7: Gene expression based on the ref_Gene database for functional assignment with

KO annotation

Sheet 8: The differential expression of KO genes between *L. crispatus* and *L. iners*

Sheet 9: Expression of metronidazole activation and resistance associated genes in *G.*

vaginalis

Sheet 10: Differential expression of KO genes of *G. vaginalis* from communities without

response to metronidazole treatment compared to those with response

A

Metronidazole

visit1 → visit2 → Pessary → visit3 → visit4 → visit5

a — 1-2 weeks — b — 1 week — c — 2 weeks — d — To 3 months — e

B

with response to treatment | no response to treatment

% abundance

Gardnerella vaginalis
Sneathia amnii
Sneathia sanguinegens
Atopobium vaginae
Lactobacillus crispatus
Lactobacillus iners
Lactobacillus gasseri
Lactobacillus jensenii
Peptostreptococcus anaerobius
Peptoniphilus lacrimalis
Mobiluncus mulieris
Prevotella amnii
Prevotella bivia
Prevotella timonensis
Others

x-axis labels: 08_006_a, 06_006_b, 13_019_a, 13_019_b, 05_012_a, 05_012_b, 06_010_a, 06_010_b, 08_003_a, 08_003_b, 13_022_a, 13_022_b, 04_001_a, 04_001_b, 06_004_a, 06_004_b, 05_001_a, 05_001_b, 06_001_a, 06_001_b, 06_006_a, 06_006_b, 07_001_a, 07_001_b, 15_006_a, 15_006_b, 17_004_a, 17_004_b

C

% Cumulative dominance vs Species rank

04_001_a (BV) · 04_001_b (NBV)
05_012_a (BV) · 05_012_b (NBV)
06_004_a (BV) · 06_004_b (NBV)
06_010_a (BV) · 06_010_b (NBV)
08_003_a (BV) · 08_003_b (NBV)
08_006_a (BV) · 08_006_b (NBV)
13_019_a (BV) · 13_019_b (NBV)
13_022_a (BV) · 13_022_b (NBV)
average (BV) · average (NBV)

D

response | no response

% abundance

G. vaginalis
Subgroup A
Subgroup B
Subgroup C
Subgroup D
Ambiguous
unclassified

x-axis labels: 04_001_a, 06_010_a, 08_006_a, 13_019_a, 13_022_a, 05_001_a, 06_001_a, 06_006_a, 07_001_a, 15_006_a, 17_004_a