Recapitulation of human germline coding variation in an ultra-mutated infant leukemia

Alexander M Gout[1], Rishi S Kotecha[2,3,4], Parwinder Kaur[5,6], Ana Abad[1], Bree Foley[7], Kim W
Carter[8], Catherine H Cole[3], Charles Bond[9], Ursula R Kees[2], Jason Waithman[7,*], Mark N
Cruickshank[1]*

[1]Cancer Genomics and Epigenetics Laboratory, Telethon Kids Cancer Centre, Telethon Kids
Institute, University of Western Australia, Perth, Australia.

[2]Leukaemia and Cancer Research Laboratory, Telethon Kids Cancer Centre, Telethon Kids
Institute, Perth, Australia.

[3]Department of Haematology and Oncology, Princess Margaret Hospital for Children, Perth,
Australia.

[4]School of Medicine, University of Western Australia, Perth, Australia.

[5]Personalised Medicine Centre for Children, Telethon Kids Institute, Australia.

[6]Centre for Plant Genetics & Breeding, UWA School of Agriculture & Environment.

[7]Cancer Immunology Unit, Telethon Kids Institute, Perth, Australia.

[8]McCusker Charitable Foundation Bioinformatics Centre, Telethon Kids Institute, Perth,
Australia.

[9]School of Molecular Sciences, The University of Western Australia, Perth, Australia.

18
19 * These authors contributed equally to this work. Correspondence and requests for
20 materials should be addressed to J.W. (Jason.Waithman@telethonkids.org.au) or to M.N.C.
21 (Mark.Cruickshank@telethonkids.org.au).

22
Keywords: infant acute lymphoblastic leukemia, hypermutation, mismatch repair defect,
MSH2

Running title: Ultra-mutation targets germline alleles in an infant leukemia

Word Count: (4608)

Total number of Figures/Tables: 5 figures and 2 tables

Financial support: Children's Leukaemia and Cancer Research Foundation (Inc.) Fellowship
(MNC); NHMRC Career Development Fellowship 1066869 (JW); Raine Medical Research
Foundation Clinician Research Fellowship 2016-2018 (RSK); Children's Leukaemia and
Cancer Research Foundation (Inc.) project grants (URK, MNC, RSK); Ian Potter Foundation
(AG); Healy Foundation (AG). Telethon Kids Institute Blue Sky Research Grant (MNC,
URK, RSK, JW). Telethon Kids Institute Research Focus Area Precision Medicine Working
Group Funding (JW, MNC).

Conflicts of interest: The authors declare no conflicts of interest.

Abstract: *Mixed lineage leukemia/Histone-lysine N-methyltransferase 2A* gene rearrangements occur in 80% of infant acute lymphoblastic leukemia, but the role of cooperating events is unknown. Infant leukemias typically carry few somatic lesions, however we identified a case with over 100 somatic point mutations per megabase. The patient presented at 82 days of age, among the earliest onset of cancer hypermutation recorded. The transcriptional profile showed global similarities to canonical cases. Coding lesions were predominantly clonal and almost entirely targeting alleles reported in human genetic variation databases with a notable exception in the mismatch repair gene, *MSH2*. The patient's diagnostic leukemia transcriptome was depleted of rare and low-frequency germline alleles due to loss-of-heterozygosity, while somatic point mutations targeted low-frequency and common human alleles in proportions that offset this discrepancy. Somatic signatures of ultra-mutations were highly correlated with germline single nucleotide polymorphic sites indicating a common role for 5-methylcytosine deamination, DNA mismatch repair and DNA adducts. These data suggest similar molecular processes shaping population-scale human genome variation also underlie evolution of an infant ultra-mutated leukemia case.

Introduction:

Histone-lysine N-methyltransferase 2A (*KMT2A*) (also known as *Mixed lineage*

leukemia) gene rearrangements (*KMT2A*-R) at 11q23 occur in infant, pediatric, adult and

therapy-induced acute leukemias and are associated with a poor prognosis. *KMT2A*-R occurs

in around 80% of infant acute lymphoblastic leukemia (ALL) and 35-50% of infant acute

myeloid leukemia (AML) involving more than 60 distinct partner genes [1]. *KMT2A*-R infant

ALL (iALL) genomes have an exceedingly low somatic mutation burden, with recurring

mutations in *RAS-PI3K* complex genes. Compared to wild-type KMT2A, the fusion

oncoprotein has altered histone-methyltransferase activity, causing epigenetic and

transcriptional deregulation. While KMT2A-oncoproteins are strong drivers of

leukemogenesis, several lines of evidence suggest that *KMT2A*-R alone is insufficient for

tumor initiation. For example, there is significant variability in latency of disease onset in

experimental models of *KMT2A*-fusions [2, 3] and in patients with *KMT2A*-R detected in

neonatal blood spots [4]. Furthermore, *KMT2A*-R acute leukemias [5] in children are enriched for

mutations affecting epigenetic regulatory genes and on average harbor more somatic lesions

than infants. These observations have prompted speculation that iALL is a distinct

developmental and genetic entity and pre-leukemic transformation occurs *in utero*, targeting a

cell of origin with a genetic/epigenetic permissive predisposition [6].

As a step towards defining cooperating events we performed next-generation

sequencing on a *KMT2A*-R iALL patient cohort and integrated the data with a larger set of

acute leukemias. We investigated the combined germline and somatic mutation profiles,

identifying a highly mutated specimen from a patient diagnosed at 82 days of age. Here we

report sequence analysis of this unique genetic entity in relation to its canonical *KMT2A*-R

iALL counterparts, revealing common mechanisms driving leukemia ultra-mutation and

population-scale human genome variation.

Methods:

Patient specimens, external sequencing data and controls

The study cohort consisted of ten infants diagnosed at Princess Margaret Hospital for Children (Perth), of which we have previously reported RNA-seq data for five patients [7]. Additional RNA-seq data was downloaded for 39 *KMT2A*-R iALL cases at diagnosis, six cases at relapse, six cases lacking the *KMT2A*-R (*KMT2A*-germline) and for pediatric *KMT2A*-R patients at diagnosis, comprising 10 AML cases, seven ALL cases and one undifferentiated case as reported previously by Andersson *et al.* [5] (accession EGAS00001000246). RNA-seq data comprising 28 *KMT2A*-R and 52 *KMT2A*-germline adult diagnosistic AML cases downloaded from the Short Read Archive (BioProject Umbrella Accession PRJNA278766[8]; Leucegene: AML sequencing). Normal human bone marrow transcriptomes were downloaded from the Short Read Archive (BioProject PRJNA284950) comprising 20 pooled sorted human hematopoietic progenitor populations [9]. The full list of samples analysed in this study including those downloaded from external sources is summarised in Supplementary Table S1.

Sample preparation and next-generation sequencing

RNA was isolated with TRIzol™ Reagent (Invitrogen Life Science Technologies, Carlsbad, CA, USA) and purified with RNeasy Purification Kits (Qiagen, Hilden, Germany). Genomic DNA was isolated with QIAamp DNA Mini Kit (Qiagen). Nucleic acid integrity and purity was assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). RNA-seq was performed at the Australian Genome Research Facility, Melbourne with 1 μg RNA as previously reported [7]. Whole exome sequencing (WES) was performed by the Beijing Genome Institute using SureSelect XT exon enrichment (Agilent Technologies) with 200 ng of remission DNA (v4 low input kit) or 1 μg of diagnostic DNA (v5). Diagnostic RNA and remission genomic DNA sequencing was performed on a HiSeq 2000 (Illumina, Inc., San Diego, CA, USA) instrument with 100bp paired end reads; diagnostic genomic

DNA sequencing was performed on a HiSeq 2500 with 150bp paired end reads. Sequencing

libraries were constructed using Illumina TruSeq kits (Illumina, Inc.).

Bioinformatics

For gene-expression analysis, RNA-seq reads were mapped to hg19 with HISAT and

gene-counts generated with HTseq [10]. Data was normalized with RUV-seq [11] implementing

factor analysis of negative control genes identified by fitting a linear model with cell type

(hematopoietic sub-set or leukemia sub-type) as a covariate. voom! [12] was used to adjust for

library size and generate counts per million (cpm) values. *KMT2A*-fusion split-reads were

identified using RNA-seq data with *FusionFinder* [13] using default settings.

Single nucleotide variants (SNVs) were identified from RNA-seq data using the

intersection of two methodologies: (1) the "Genome Analysis Toolkit (GATK) Best Practices

workflow for SNP and indel calling on RNAseq data" and (2) *SNPiR* [14]. WES data was

processed based on the GATK Best Practices workflow. Variants were characterized using

custom scripts that extract the coverage of sequenced bases from VCF and bam files,

determining the overlap among samples, and calculating variant allele fractions (VAFs).

High confidence somatic mutations were identified using *VarScan2* (v2.3.9) [15] setting

cut-offs of *P*-value<0.005; coverage > 50X; VAF > 0.1 or VAF > 0.3 to enrich for clonal

mutations. Variants were annotated with ANNOVAR [16] and with metadata from additional

sources including seven functional/conservation algorithms (PolyPhen2 [17], Sift [18],

MutationTaster [19], likelihood ratio test [20], GERP [21], PhyloP [22], and CADD [23]) and databases of

human genetic variation (ExAC, 1000 genomes, HapMap, dbSNP, COSMIC and ClinVar).

Predicted functional alleles were assigned for variants called as damaged/deleterious by at

least 4/7 algorithms.

Somatic mutational spectra were investigated using the R package, *DeconstructSigs*

(v1.8.0) [24] calculating the fraction of somatic substitutions within the possible 96-trinucleotide

contexts relative to the exome background. Copy number analysis was performed on WES

data using *VarScan2* to calculate log2 depth ratio and GC-content and *Sequenza* (v2.1.2) [25] to

identify chromosomal segments with copy loss. Gene ontology enrichment was performed

using GREAT (v3.0.0) [26] bioinformatics tools. Mutations were inspected by mapping the

genomic position to crystal structures in the Protein DataBank[27] and analyzed with Pymol

(v1.8.6.2).

Data and code availability

Scripts used to execute variant detection software and analysis are available at

https://github.com/cruicks/ultramutated-case under an MIT license and are being developed

on Github (https://github.com/cruicks/vcfbamCompare). The sequencing data has been

deposited in the Short Read Archive with study ID PRJNA296746.

Results:

Characteristics of KMT2A-R iALL cohort and RNA-seq variant detection pipeline

We reported previously *KMT2A*-R fusion transcripts for 5/10 cases [7], and of the

remaining five *KMT2A*-R iALL we detected *KMT2A-MLLT3* (P401, P438), *KMT2A-AFF1*

(P848), *KMT2A-MLLT1* (P706) and *KMT2A-EPS15* (P809). *KMT2A*-fusion transcripts were

therefore detected in all samples and reciprocal fusion sequences detected in 5/10 samples

(Table 1). To determine the validity of our RNA-seq based pipeline for detecting rare and

common alleles, we examined the concordance of single nucleotide variants (SNVs) called

from RNA-seq and WES data. Across seven samples with matched WES and RNA-seq data,

we identified a total of 54,792 point mutations from RNA-seq reads of which 166 were absent

from matched WES (coverage>20 reads), representing a discordance rate of 0.3% (Figure 1a;

Supplementary Table S2). In one sample, P706 there were two events called with RNA-seq

data that were not identified in matched WES but with evidence extracted using the Integrated

Genome Browser at previously reported somatic mutation sites in *FLT3* (encoding FLT3

p.V491L [28]; 200/1,719 alternate RNA-seq reads; 15/214 alternate WES reads) and *KRAS*

(encoding KRAS p.A146T [29]; 24/159 alternate RNA-seq reads; 6/51 alternate WES reads).

Furthermore, analysis of 117 discordant events with coverage > 50 reads, revealed 76

recurrences in two or more samples (64%) at 26 unique sites, however none of these sites

have been validated as somatic mutations in cancer to date in the Catalog of Somatic

Mutations in Cancer (COSMIC)[30] or The Pediatric Cancer Genome Project (PCGP)[31]

databases. These results indicate a high rate of reproducibility of the RNA-seq variant

detection pipeline and suggest that RNA-DNA differences mostly comprise artifacts and/or

novel RNA-editing events [32] and a minority of putative somatic alterations exclusively

detected by RNA-seq.

Identification of a highly mutated infant leukemia

Expressed somatic mutations were defined as variants detected in both diagnostic

leukemia RNA-seq and WES, but absent in remission WES at sites with >20 reads (Figure

1b-d). Most patients (5/6) showed the expected "silent genomic landscape" (Supplementary

Table S3) except for patient P337 (Figure 1b-d; Supplementary Table S4). Three patients

expressed RAS-family hot-spot somatic mutations (P848 and P706: KRAS p.A146T and

P438: NRAS.p.G12S). We detected no expressed somatic missense mutations in patients

P399 and P401. The pair of monozygotic twins (P809 and P810) lacked private expressed

mutations but we detected two shared expressed missense mutations clustered in the *PER3*

gene (encoding PER3.p.M1006R and PER3.p.K1007E). These results are consistent with

recurring RAS-pathway mutations and low mutation rate reported previously [5, 33]. However,

in patient P337 RNA-seq reads we detected 1,420 expressed somatic mutations

(Supplementary Table S4). The overlap of both rare and common alleles across the matched

datasets verifies the common origin of RNA-seq and diagnostic/remission WES data ruling

out sample contamination or mislabeling (Figure 1a-c). We previously reported a complex

KMT2A translocation involving 2q37, 19p13.3 and 11q23[34] in the highly mutated leukemia

sample and established two cell lines from the diagnostic sample [7].

Characterisation and clonal analysis of somatic lesions

Somatic lesions were further evaluated in six cases by comparing diagnostic and remission

WES. Two different variant allele fraction (VAF)-cut-offs, either VAF>0.1 or VAF>0.3, were

chosen to enrich for sub-clonal and clonal events respectively. There were on average ~82.2

(range 69-95) sub-clonal and ~3.4 clonal (range 3-4) somatic point mutations in canonical

KMT2A-R iALL exomes, confirming previous studies of a low somatic mutation rate in the

dominant clone (Table 1). In contrast, there were 198 sub-clonal and 5,054 clonal somatic

point mutations in the highly-mutated sample, equating to a rate of 139 mut/Mb, classifying

this sample as ultra-mutated. There were also ~40-fold increased number of indels in the

dominant clone of the ultra-mutated sample (n=260) compared to typical cases (average =

6.6; range: 3-9) including 13 predicted to cause a frame-shift protein-coding truncation (Table

1; Supplementary Table S5). There were 2,420 exonic substitutions encoding 1,109 missense

alleles and 12 stop-gain alleles (Supplementary Table S5). We also noted that the ultra-

mutated case carried an excess of loss-of-heterozygosity (LOH) positions, with 6,555 high

confidence sites; vastly more than typical cases which had a mean of 57.2 LOH positions

(range 28-100) (Figure 2a-b; Table 1).

WES was performed on two independent cell lines derived from the ultra-mutated diagnostic

specimen, PER-784A and PER-826A, to characterize clonal somatic events. (Figure 2c-d).

Most somatic and LOH-events detected in the primary ultra-mutated sample were also called

in both, PER-784A and PER-826A cell lines using P337 remission WES for all comparisons

(Figure 2d). These data suggest that the somatic lesions are largely clonal in agreement with

the VAF-distributions (Figure 2b, d). In total, 4,754/5,054 (94%) somatic events

(Supplementary Table S6) and 6,150/6,575 (93%) LOH-events (Supplementary Table S7)

were called in the derived cell lines. Moreover, there were relatively few discordant sites

when directly comparing cell lines with the primary diagnostic sample (PER-784A: 34

discordant sites; PER-826A: 44 discordant sites; Supplementary Table S6)). We further

investigated evidence of clonal somatic lesions in cell lines using an alternative approach, by

directly extracting quality filtered WES reads mapping to coding somatic point mutations

(n=2,411; Q>50 in diagnosis WES; Figure 2e) and LOH-sites (n=3,117; Q>50 in remission

WES; Figure 2f; Supplementary Table S8).   All coding clonal somatic mutations were

supported by reads in both cell lines of which 121 (5%) were homozygous (VAF>0.95 in cell

line and VAF>0.9 in primary sample). Most LOH-sites were evidenced by reference alleles

(VAF<10%; 2,179/3,117) in cell lines, with ~30% of LOH-sites evidenced by alternate alleles

(VAF> 90%; 926/3,117) and only 12 sites (0.3%) with both reference and alternate alleles

(VAF range 10-90%).

We performed copy number analysis of the ultra-mutated sample to map LOH-sites to

chromosomal segments with evidence of deletions. Most genomic regions were called as

diploid with limited chromosomal copy loss detected (Supplementary Figure S1a-b). There

was a total of 382/6,555 (5.8%) LOH-sites mapping to homozygous deletions, suggesting

only a minority of the identified LOH-sites likely arose through chromosome segment

deletion. Infant *KMT2A*-R ALL typically carry a small number of copy-number alterations

which may coincide with LOH-sites. We determined the distance of LOH-sites, somatic

mutations and germline variants (called by *VarScan2*) to the nearest homozygous SNV in

each leukemia diagnostic sample finding that in canonical cases LOH-sites tend to be located

closer to homozygous sites than either germline or somatic variants (Supplementary Figure

S1c; left panel) and show a unique bimodal distribution (Supplementary Figure S1c; right

panel). In contrast the average distance of LOH-sites to homozygous SNVs in the ultra-

mutated sample was similar to that of germline variants and somatic mutations with each

displaying a similar distribution (Supplementary Figure S1c). Thus, most LOH-sites in the

ultra-mutated specimen do not show evidence of localising within deleted or copy-neutral

loss-of-heterozygosity chromosomal segments.

In summary, these data demonstrate that somatic lesions in the primary sample were

predominantly clonal and LOH-sites appear to comprise a large proportion of somatic

substitutions rather than chromosomal alterations. We also note relatively few coding

sequence changes in derived cell lines compared to the dominant diagnostic leukemic clone,

both which are capable of long-term propagation. These results could reflect loss of the

hypermutator phenotype during *in vitro* culture or in diagnostic leukemia cells.

Signatures of somatic mutations and LOH-sites correlate with profiles of germline alleles

240   We next investigated the sequence features of somatic and LOH substitutions in the

ultra-mutated sample (Figure 3). We hypothesised that LOH-sites and somatic mutations may

show common feature biases depending on the context of alteration. Somatic events were

evaluated separately that induce a heterozygous mutation (n=4,773) or homozygous mutation

(n=281), and LOH causing reference→alternative changes (n=2,057) were evaluated

separately to alternative→reference changes (n=4,453). Since we noted an overlap in the

somatic mutations of the ultra-mutated sample with variants in the ExAC database (Figure 1a-

c), we first examined the distribution of these sites with respect to the derived allele frequency

in ExAC. We observe that each of these categories of sequence alterations in the ultra-

mutated sample were distributed across the range of population allele-frequencies (Figure 3a-

b). While the distributions of LOH and somatic sites were similar, somatic heterozygous

changes were relatively more frequent at higher frequency alleles as were LOH

reference→alternate alleles.

253   Somatic signatures of ultra-mutations were investigated in comparison to

heterozygous and homozygous single nucleotide polymorphisms (SNPs) in patient remission

samples. Heterozygous positions were defined by VAF<0.9 yielding 51,347 heterozygous and

31,237 homozygous SNPs in canonical remission samples and 10,681 heterozygous and 6

617 homozygous SNPs in the ultra-mutated remission sample. Analysis of WES remission

variants from canonical patients and the ultra-mutated sample revealed similar ExAC allele

frequency distributions with distinctions between heterozygous and homozygous alleles

(Figure 3c-d). The overall enrichment of each category of somatic substitution and LOH

across the 96 possible trinucleotide contexts resembled patient remission SNPs with elevated

proportions of C>T transitions at NpCpG motifs (Figure 3e).

Pre-defined signatures were used to infer mutational profiles revealing Signature 1A

(associated with spontaneous deamination of 5-methylcytosine), Signature 12 (associated

with unknown process with strand-bias for T→C substitutions) and Signature 20 (associated

with defective DNA mismatch repair, MMR) as enriched in each of the mutation/variant sets,

however with distinctive proportions (Figure 3f). Heterozygous somatic mutations and LOH

alternative→reference sites were additionally enriched for Signature 1B (variant of Signature

1A) and showed highly correlated profiles (Pearson's correlation 0.919). LOH

reference→alternative sites were additionally enriched for Signature 5 (associated with

unknown process with strand-bias for T→C substitutions) and were highly correlated with

homozygous somatic mutation profiles (0.912) and to profiles of patient heterozygous

positions (P337: 0.935; canonical: 0.926). Homozygous somatic mutations also displayed

similarities to heterozygous patient SNPs (P337: 0.981; canonical: 0.985). Overall the somatic

mutational profiles indicate highly related patterns of both somatic ultra-mutations and LOH-

sites. Notably LOH alternative→reference sites showed unique and overlapping features to

somatic heterozygous mutations. We also observe underlying similarities in the profiles of

somatic substitutions and germline alleles.

Ultra-mutation distribution recapitulates common and rare germline variation

We noted a similar spectrum of rare and common alleles in the ultra-mutated

leukemia transcriptome compared to typical *KMT2A*-R iALL cases; but there were

comparatively fewer rare/low-frequency alleles shared with the remission sample (Figure 1a-

c; ExAC<0.5%). We further explored this trend to disentangle the contribution of somatic

acquired alleles and LOH-sites. Analysis of the remission and diagnostic samples from this

patient in isolation, revealed a similar proportion of rare/low-frequency coding alleles

287 detected by WES and both were proportionally within the same range as canonical

288 counterparts (Figure 4a). However, when directly comparing the remission and diagnostic

289 WES, we noted a modest depletion of germline rare/low-frequency coding SNV alleles

290 defined by *VarScan2* (Figure 4a; 2.9% rare alleles in P337 compared to 3.5±0.2% in

291 canonical cases). This apparent depletion of germline alleles in the ultra-mutated leukemic

292 sample was coincident with an increased proportion of somatic acquired rare/low-frequency

293 coding SNV alleles compared to LOH-sites (Figure 4a). Therefore, when considering the total

294 pool of SNVs in the leukemic coding genome, the somatic gains and losses of rare/low-

295 frequency coding SNV alleles were in proportions that resulted in a similar distribution as

296 canonical cases (Figure 4a). We extended this observation by investigating the rare and

297 common allele abundance of expressed germline-encoded variants and somatic mutations in

298 the ultra-mutated leukemic sample compared with a larger cohort of *KMT2A*-R iALL

299 transcriptomes (Figure 4b; n=32 St. Jude patients and n=8 Perth patients). Most typical

300 *KMT2A*-R iALL diagnostic specimens expressed similar numbers of rare and low-frequency

301 expressed SNVs, except for 4/32 cases sequenced by St Jude investigators, which had an

302 elevated number of expressed rare and low-frequency ExAC alleles (Figure 4b). Excluding

303 these samples, typical *KMT2A*-R iALL patients expressed on average 42.3 ±10.4 (standard

304 deviation) variants that were absent from ExAC and 212 ± 32.5 variants with an allele

305 frequency of <0.5% in ExAC. The total number of expressed variants in the ultra-mutated

306 sample was slightly lower than typical cases (38 absent from ExAC and 174 AF <0.5% in

307 ExAC), and were approximately comprised of half germline (17/38 and 88/174 respectively)

308 and half somatic acquired alleles. Overall, the ultra-mutated transcriptome had the lowest

309 prevalence of germline-encoded rare and low-frequency alleles among the larger sample size

310 (Table 2). Furthermore, the somatic mutation abundance within each allele-frequency bin was

311 proportional to their abundance in prototypical silent cases. Consequently, the low numbers of

312 rare germline alleles in the ultra-mutated transcriptome were off-set by somatic mutations

313 such that the total numbers were within the same range as typical cases (Figure 4b-c). We

314 conclude that despite carrying thousands of somatic mutations, the coding-genome of ultra-

mutated patient P337 shows similarities to canonical *KMT2A*-R iALL cases, vis-à-vis rare

allele prevalence and representation of common human genetic variation.

RAS-pathway and DNA-replication and repair gene mutations

Gene ontology analysis of the expressed non-synonymous somatic mutations (n=535)

revealed over-representation of RAS-pathway genes (7/23 genes; 7-fold enriched; *P*=0.049;

RALGDS, *RAC1*, *CHUK*, *PIK3R1*, *NFKB1*, *PLD1* and *PIK3CA*) and "*MLL* signature 1 genes"

(33/369 genes; 2-fold enriched; *P*=0.046). Of the 535 non-synonymous expressed somatic

mutations, 45 were predicted deleterious/damaging, including in the MMR gene *MSH2*

(p.E489K). The *MSH2* somatic mutation localizes to a hot-spot, within the MMR ATPase

domain in a region at the periphery of the DNA binding site, approximately 20 Å from the

DNA molecule in the complex structure of full-length MSH2, a fragment of MSH6, ADP and

DNA containing a G:T mispair [35]. The mutation encodes a substitution of a negatively

charged glutamate with a positively charged lysine (Figure 4d). There was a non-synonymous

somatic mutation in another MMR gene, *MUTYH*; however, this site is a common human

allele with a derived AF of 0.297 and was not called as damaged/deleterious by our prediction

pipeline. There were no somatic mutations, LOH-sites nor rare/deleterious coding germline

alleles detected in *POLE* or *POLD1* which are frequently co-mutated with *MSH2* in ultra-

mutated tumours [36, 37] and in sporadic early-onset colorectal cancer [38]. Among the 45

expressed and predicted damaging somatic mutations, there were two rare substitutions in

DNA polymerase encoding genes: *POLG2* (p.D308V; ExAC AF 0.0000165), which encodes

a mitochondrial DNA polymerase and *POLK* (p.R298H; ExAC AF 0.0003), which encodes

an error prone replicative polymerase that catalyzes translesion DNA synthesis. The *POLK*

mutation maps to a conserved impB/mucB/samB family domain (Pfam: PF00817) involved in

UV-protection. The encoded substitution of arginine to histidine is located approximately 20

Å from the DNA molecule in the structure of a 526 amino acid N-terminal Pol K fragment

complexed with DNA containing a bulky deoxyguanosine adduct induced by the carcinogen

benzo[a]pyrene (Figure 4e) [39]. Both wild-type and mutant residues are positively charged,

however arginine is strongly basic (pI 11) while histidine is weakly basic (pI 7) and titratable

at physiological pH [40]. We additionally observed an expressed and predicted functional

germline rare allele in *ATR*, a master regulator of the DNA damage response (p.L2076V;

ExAC AF 0.0002).

Global transcriptional similarities of the ultra-mutated and canonical infant leukemias

We next examined the transcriptional profile of the highly mutated *KMT2A*-R iALL

specimen in comparison to a collection of patient leukemia samples and purified blood

progenitor populations (Figure 5). As expected cell type was the major source of variation

distinguishing normal hematopoietic progenitor, lymphoblastic leukemias and myeloid

leukemias. We also observe clustering according to *KMT2A* translocation status and partner

genes as reported previously [41]. Replicate samples of the highly mutated *KMT2A*-R iALL

specimen cluster with canonical counterparts demonstrating global transcriptional similarities

(Figure 5).

Discussion:

We report sequence analysis of an ultra-mutated acute lymphoblastic leukemia

diagnosed in an infant at 82 days of age which, to our knowledge, represents the earliest onset

of a hypermutant tumor recorded. The mutation rate observed was at the upper limit of cases

reported previously in childhood cancers [36, 37]. By integrating analysis of somatic mutations

and germline alleles with a larger cohort of *KMT2A*-R iALL patients we found that the high

somatic mutation rate does not generate large numbers of rare alleles compared to typical

cases. Rather, we observe that the ultra-mutated *KMT2A*-R leukemic coding genome closely

resembles the rare allele landscape of canonical cases. This apparent paradox is explained by

the net effect of somatic gains and losses, both of which were targeted to rare and common

alleles in proportions that off-set the depleted pool of germline alleles observed in the

leukemic coding sequences. Our data indicate both the rare allele burden and transcriptional

profile of the ultra-mutated leukemia were comparable to its canonical counterparts.

Therefore, despite its radically different genetic route to leukemogenesis, the molecular

profile of the ultra-mutated specimen seems to have converged towards that of canonical

cases.

Most ultra-mutated childhood cancers reported previously have a combined MMR

and proof reading polymerase deficiency [36, 37] - we detect evidence for the former (*MSH2*

point mutation), but not the latter. However, we identify somatic mutations in two genes

encoding DNA polymerases (*POLK* and *POLG2*) and an additional MMR gene (*MUTYH*).

Notably, the *POLK* mutation is a rare allele, predicted functional and encodes pol κ, a

replicative error-prone translesion DNA polymerase that plays a role in bypassing unrepaired

bases that otherwise block DNA replication [42, 43]. For example, pol κ preferentially

incorporates dAMP at positions complementary to 7,8-dihydro-8-oxo-2′-deoxyguanosine

adducts [44]. As a corollary to these observations, *POLK* mutations have been found in prostate

cancer with elevated C→T transition [45]. Further studies are required to determine if any of

these mutations cooperate to specify the unique mutational pattern observed in the ultra-

mutated *KMT2A*-R iALL genome.

383   The largest fraction of ultra-mutations was attributed to Signature 1, associated with

5-methylcytosine deamination inducing C→T transitions, which accumulates in normal

somatic cells with age over the human lifespan [46]. This signature is found ubiquitously across

tumour sub-types [47], and prominently in hypermutated early onset AML patients with

germline *MBD4* mutations [48]. LOH-sites (reference→alternative) were enriched for Signature

5 which is also correlated with human aging. We observe enrichment of Signature 20 in

LOH-sites and somatic alterations, consistent with an association with a MMR-defect [47]. This

signature has also been reported as enriched in tumours with germline-MMR and secondary

mutation in the *POLD1* proof-reading gene [47]. We also find enrichment of Signature 12 which

arises from a poorly characterized underlying molecular process; however, the dominant base

transitions, A→G and T→C have been observed in association with DNA damage (e.g. by

adducts) in normal and cancer tissues [49, 50]. These observations suggest common mutagenic

and repair deficits driving age-related mutation in humans were operative in the cell of origin

during pre-natal evolution of the ultra-mutated infant leukemia genome. Collectively, these

results implicate the *MSH2* point mutation as the primary MMR-hit, followed by an

additional event effecting a proof reading polymerase such as *POLK*, however, our data imply

that *POLE* and *POLD1* mutations are not cooperative in this process.

400   The preponderance of ultra-mutated sites to overlap germline alleles in proportion to

their population allele frequencies demonstrates that hypermutation can recapitulate aspects

of population scale human genetic variation in a cancer genome. Aproximately half of the

somatic substitutions appear to target heterozygous SNPs resulting in loss-of-heterozygosity.

We cannot rule out a role for focal chromosomal alterations undetectable by our copy-number

and zygosity analyses. However collectively our data suggest that a major fraction of the

LOH-sites arose by point mutagenesis of heterozygous positions. They more frequently target

the alternate allele inducing reversion to the reference base, with around 30% of events

resulting in a homozygous non-reference allele in the diagnostic sample. Most ultra-mutations

are therefore likely passengers coinciding with common human alleles, consistent with

findings that hypermutated tumours tend to have relatively few drivers proportional to their

mutation burden [51]; however, as proposed previously by Valentine and colleagues[52], a

minority of rare alleles may contribute to infant leukemogenesis. While our results do not

resolve the temporal order of the key oncogenic events - namely *MSH2* point mutation,

genomic hypermutation and *KMT2A* translocation – due to the clonal nature of these events

they appear to have occurred in rapid succession. This could be explained by a sentinel DNA

repair defect inducing hypermutation and genome instability followed by positive selection of

cooperating lesions conferring a growth advantage resulting in clonal expansion and fixation

of passengers [53]. An alternative model is also possible in which the *MSH2* MMR-defect

impaired double-strand-break repair [54, 55] promoting *KMT2A*-rearrangement, but with neutral

selection of somatic point mutations. Such a scenario appears inconsistent with the observed

predominance of clonal mutations, but could be reconciled if the hypermutator phenotype was

lost rapidly after the *KMT2A*-rearrangement and prior to clonal expansion. Since pervasive

positive selection has been observed in cancer and non-cancerous tissues [51, 56], we favour the

former model as being more plausible. However, since we noted loss of hypermutator

phenotype in patient-derived cell lines, the later model is not ruled out by our data.

Collectively our results illuminate a link between mutagenesis during human aging, germline

variation and somatic ultra-rmutation in cancer. Further studies are required to determine if

combinations of somatic mutations and/or germline alleles promote *KMT2A*-R iALL

leukemogenesis/hypermutation and if somatic mutated sites significantly overlap human

germline alleles in other malignancies.

Financial support: Children's Leukaemia and Cancer Research Foundation (Inc.) Fellowship (MNC); NHMRC Career Development Fellowship 1066869 (JW); Raine Medical Research Foundation Clinician Research Fellowship 2016-2018 (RSK); Children's Leukaemia and Cancer Research Foundation (Inc.) project grants (URK, MNC, RSK); Ian Potter Foundation (AG); Healy Foundation (AG). Telethon Kids Institute Blue Sky Research Grant (MNC, URK, RSK, JW). Telethon Kids Institute Research Focus Area Precision Medicine Working Group Funding (JW, MNC).

Acknowledgments: We thank Daniel MacArthur and Fengmei Zhao for assisting with exome sequence analysis; and researchers at The Pediatric Cancer Genome Project at St. Jude Children's Research Hospital for provision of data. We thank all the patients, families and hospital staff for providing samples.

Authorship contributions: AG wrote computer code, performed bioinformatics, conceived analysis strategy, interpreted and analyzed data and edited the manuscript. AA and BF performed experiments. KWC, PK and CB performed additional data analysis and interpretation. RSK provided patient samples and clinical data, assisted with data interpretation and edited the manuscript. CHC, URK and JW acquired funding support, assisted with data interpretation and edited the manuscript. MNC performed bioinformatics, formulated research, conceived analysis strategy, acquired funding support, interpreted data, designed experiments and drafted the manuscript.

Disclosure of Conflicts-of-Interest: The authors declare no conflicts of interest.

451 **References:**

452　1.　Meyer C, *et al.* The MLL recombinome of acute leukemias in 2017. *Leukemia*,
453　　　(2017).

454
455　2.　Bergerson RJ, *et al.* An insertional mutagenesis screen identifies genes that
456　　　cooperate with Mll-AF9 in a murine leukemogenesis model. *Blood* **119**, 4512-4523
457　　　(2012).

458
459　3.　Chen W, O'Sullivan MG, Hudson W, Kersey J. Modeling human infant MLL leukemia
460　　　in mice: leukemia from fetal liver differs from that originating in postnatal marrow.
461　　　*Blood* **117**, 3474-3475 (2011).

462
463　4.　Gale KB, *et al.* Backtracking leukemia to birth: identification of clonotypic gene
464　　　fusion sequences in neonatal blood spots. *Proc Natl Acad Sci U S A* **94**, 13950-13954
465　　　(1997).

466
467　5.　Andersson AK, *et al.* The landscape of somatic mutations in infant MLL-rearranged
468　　　acute lymphoblastic leukemias. *Nat Genet* **47**, 330-337 (2015).

469
470　6.　Sanjuan-Pla A, *et al.* Revisiting the biology of infant t(4;11)/MLL-AF4+ B-cell acute
471　　　lymphoblastic leukemia. *Blood* **126**, 2676-2685 (2015).

472
473　7.　Cruickshank MN, *et al.* Systematic chemical and molecular profiling of MLL-
474　　　rearranged infant acute lymphoblastic leukemia reveals efficacy of romidepsin.
475　　　*Leukemia*, (2016).

476
477　8.　Lavallee VP, *et al.* The transcriptomic landscape and directed chemical interrogation
478　　　of MLL-rearranged acute myeloid leukemias. *Nat Genet* **47**, 1030-1037 (2015).

479
480　9.　Casero D, *et al.* Long non-coding RNA profiling of human lymphoid progenitor cells
481　　　reveals transcriptional divergence of B cell and T cell lineages. *Nat Immunol* **16**,
482　　　1282-1291 (2015).

483
484　10.　Anders L, *et al.* Genome-wide localization of small molecules. *Nat Biotechnol* **32**, 92-
485　　　96 (2014).

486
487　11.　Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor
488　　　analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).

489
490　12.　Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model
491　　　analysis tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).

493   13.   Francis RW, Thompson-Wicking K, Carter KW, Anderson D, Kees UR, Beesley AH.
494   FusionFinder: a software tool to identify expressed gene fusion candidates from
495   RNA-Seq data. *PLoS One* **7**, e39987 (2012).

497   14.   Piskol R, Ramaswami G, Li JB. Reliable identification of genomic variants from RNA-
498   seq data. *Am J Hum Genet* **93**, 641-651 (2013).

500   15.   Koboldt DC*, et al.* VarScan 2: somatic mutation and copy number alteration
501   discovery in cancer by exome sequencing. *Genome Res* **22**, 568-576 (2012).

503   16.   Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants
504   from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).

506   17.   Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense
507   mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**, Unit7 20 (2013).

509   18.   Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous
510   variants on protein function using the SIFT algorithm. *Nat Protoc* **4**, 1073-1081
511   (2009).

513   19.   Schwarz JM, Rodelsperger C, Schuelke M, Seelow D. MutationTaster evaluates
514   disease-causing potential of sequence alterations. *Nat Methods* **7**, 575-576 (2010).

516   20.   Chun S, Fay JC. Identification of deleterious mutations within three human genomes.
517   *Genome Res* **19**, 1553-1561 (2009).

519   21.   Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, Sidow A. Distribution and
520   intensity of constraint in mammalian genomic sequence. *Genome Res* **15**, 901-913
521   (2005).

523   22.   Cooper GM*, et al.* Single-nucleotide evolutionary constraint scores highlight disease-
524   causing mutations. *Nat Methods* **7**, 250-251 (2010).

526   23.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general
527   framework for estimating the relative pathogenicity of human genetic variants. *Nat
528   Genet* **46**, 310-315 (2014).

530   24.   Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C. DeconstructSigs:
531   delineating mutational processes in single tumors distinguishes DNA repair
532   deficiencies and patterns of carcinoma evolution. *Genome Biol* **17**, 31 (2016).

534 25. Favero F, *et al.* Sequenza: allele-specific copy number and mutation profiles from
535    tumor sequencing data. *Ann Oncol* **26**, 64-70 (2015).

536
537 26. McLean CY, *et al.* GREAT improves functional interpretation of cis-regulatory
538    regions. *Nat Biotechnol* **28**, 495-501 (2010).

539
540 27. Berman HM, *et al.* The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000).

541
542 28. He J, *et al.* Integrated genomic DNA/RNA profiling of hematologic malignancies in
543    the clinical setting. *Blood* **127**, 3004-3014 (2016).

544
545 29. Chang MT, *et al.* Identifying recurrent mutations in cancer reveals widespread
546    lineage diversity and mutational specificity. *Nat Biotechnol* **34**, 155-163 (2016).

547
548 30. Forbes SA, *et al.* COSMIC: somatic cancer genetics at high-resolution. *Nucleic Acids*
549    *Res* **45**, D777-D783 (2017).

550
551 31. Downing JR, *et al.* The Pediatric Cancer Genome Project. *Nat Genet* **44**, 619-622
552    (2012).

553
554 32. Pickrell JK, Gilad Y, Pritchard JK. Comment on "Widespread RNA and DNA sequence
555    differences in the human transcriptome". *Science* **335**, 1302; author reply 1302
556    (2012).

557
558 33. Chang VY, Basso G, Sakamoto KM, Nelson SF. Identification of somatic and germline
559    mutations using whole exome sequencing of congenital acute lymphoblastic
560    leukemia. *BMC Cancer* **13**, 55 (2013).

561
562 34. Henderson MJ, *et al.* A xenograft model of infant leukaemia reveals a complex MLL
563    translocation. *Br J Haematol* **140**, 716-719 (2008).

564
565 35. Warren JJ, Pohlhaus TJ, Changela A, Iyer RR, Modrich PL, Beese LS. Structure of the
566    human MutSalpha DNA lesion recognition complex. *Mol Cell* **26**, 579-592 (2007).

567
568 36. Campbell BB, *et al.* Comprehensive Analysis of Hypermutation in Human Cancer. *Cell*
569    **171**, 1042-1056 e1010 (2017).

570
571 37. Shlien A, *et al.* Combined hereditary and somatic mutations of replication error
572    repair genes result in rapid onset of ultra-hypermutated cancers. *Nat Genet* **47**, 257-
573    262 (2015).

575   38.   Palles C, *et al.* Germline mutations affecting the proofreading domains of POLE and
576   POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet* **45**, 136-144
577   (2013).

578
579   39.   Jha V, Bian C, Xing G, Ling H. Structure and mechanism of error-free replication past
580   the major benzo[a]pyrene adduct by human DNA polymerase kappa. *Nucleic Acids*
581   *Res* **44**, 4957-4967 (2016).

582
583   40.   Szpiech ZA, *et al.* Prominent features of the amino acid mutation landscape in
584   cancer. *PLoS One* **12**, e0183273 (2017).

585
586   41.   Stam RW, *et al.* Gene expression profiling-based dissection of MLL translocated and
587   MLL germline acute lymphoblastic leukemia in infants. *Blood* **115**, 2835-2844 (2010).

588
589   42.   Kanemaru Y, *et al.* DNA polymerase kappa protects human cells against MMC-
590   induced genotoxicity through error-free translesion DNA synthesis. *Genes Environ*
591   **39**, 6 (2017).

592
593   43.   Maddukuri L, Ketkar A, Eddy S, Zafar MK, Eoff RL. The Werner syndrome protein
594   limits the error-prone 8-oxo-dG lesion bypass activity of human DNA polymerase
595   kappa. *Nucleic Acids Res* **42**, 12027-12040 (2014).

596
597   44.   Irimia A, Eoff RL, Guengerich FP, Egli M. Structural and functional elucidation of the
598   mechanism promoting error-prone synthesis by human DNA polymerase kappa
599   opposite the 7,8-dihydro-8-oxo-2'-deoxyguanosine adduct. *J Biol Chem* **284**, 22467-
600   22480 (2009).

601
602   45.   Yadav S, Mukhopadhyay S, Anbalagan M, Makridakis N. Somatic Mutations in
603   Catalytic Core of POLK Reported in Prostate Cancer Alter Translesion DNA Synthesis.
604   *Hum Mutat* **36**, 873-880 (2015).

605
606   46.   Alexandrov LB, *et al.* Clock-like mutational processes in human somatic cells. *Nat*
607   *Genet* **47**, 1402-1407 (2015).

608
609   47.   Alexandrov LB, *et al.* Signatures of mutational processes in human cancer. *Nature*
610   **500**, 415-421 (2013).

611
612   48.   Mathijs A Sanders EC, Christoffer Flensburg, Annelieke Zeilemaker, Sarah E Miller,
613   Adil al Hinai, Ashish Bajel, Bram Luiken, Melissa Rijken, Tamara Mclennan, Remco M
614   Hoogenboezem, François G Kavelaars, Marnie E Blewitt, Eric M Bindels, Warren S
615   Alexander, Bob Löwenberg, Andrew W Roberts, Peter J M Valk, Ian Majewski.
616   Germline loss of MBD4 predisposes to leukaemia due to a mutagenic cascade driven
617   by 5mC. *bioRxiv*, (2018).

619   49.   Chen L, Liu P, Evans TC, Jr., Ettwiller LM. DNA damage is a pervasive cause of
620   sequencing errors, directly confounding variant identification. *Science* **355**, 752-756
621   (2017).

623   50.   Lin J, Shi T. Error-prone DNA polymerase and oxidative stress increase the incidences
624   of A to G mutations in tumors. *Oncotarget* **8**, 45154-45163 (2017).

626   51.   Martincorena I*, et al.* Universal Patterns of Selection in Cancer and Somatic Tissues.
627   *Cell* **171**, 1029-1041 e1021 (2017).

629   52.   Valentine MC*, et al.* Excess congenital non-synonymous variation in leukemia-
630   associated genes in MLL- infant leukemia: a Children's Oncology Group report.
631   *Leukemia*, (2013).

633   53.   Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer therapies.
634   *Nat Rev Clin Oncol*, (2017).

636   54.   van Oers JM*, et al.* The MutSbeta complex is a modulator of p53-driven
637   tumorigenesis through its functions in both DNA double-strand break repair and
638   mismatch repair. *Oncogene* **33**, 3939-3946 (2014).

640   55.   Burdova K, Mihaljevic B, Sturzenegger A, Chappidi N, Janscak P. The Mismatch-
641   Binding Factor MutSbeta Can Mediate ATR Activation in Response to DNA Double-
642   Strand Breaks. *Mol Cell* **59**, 603-614 (2015).

644   56.   Martincorena I*, et al.* Tumor evolution. High burden and pervasive positive selection
645   of somatic mutations in normal human skin. *Science* **348**, 880-886 (2015).

648   Table 1: Characterisation of somatic lesions in *KMT2A*-R patient samples.

Patient ID	KMT2A-fusion	Reciprocal KMT2A-fusion	Diagnostic leukemia DNA	Constitutional remission DNA	Point Mutations 0.1 > VAF[1] < 0.3	Point Mutations VAF > 0.3	Indels VAF > 0.3	LOH[2]-sites	Point mutations or hotspots with RNA-SEQ reads
P337	*KMT2A-MLLT1*	Not detected	Exome	Exome	198	5,054	260	6,555	> 1,000 expressed somatic mutations
P401	*KMT2A- MLLT 3*	Not detected	Exome	Exome	78	3	9	45	No non-silent somatic mutations
P399	*KMT2A-AFF1*	*AFF1-KMT2A*	Exome	Exome	95	3	7	67	No non-silent somatic mutations
P438	*KMT2A- MLLT3*	Not detected	Exome	Exome	74	4	9	100	NRAS.p.G12S
P810	*KMT2A-EPS15*	*EPS15-KMT2A*	Exome	Exome	95	4	5	46	PER3.p.M1006R, PER3.p.K1007E
P809	*KMT2A-EPS15*	*EPS15-KMT2A*	Exome	Exome	69	3	3	28	PER3 M1006R, PER3.p.K1007E
P848	*KMT2A-AFF1*	*AFF1-KMT2A*	no	Exome	NA	NA	NA	NA	KRAS.p.A146T
P287	*KMT2A-AFF1*	*AFF1-KMT2A*	no	no	NA	NA	NA	NA	No COSMIC[3] hotspots
P706	*KMT2A- MLLT1*	Not detected	no	Exome	NA	NA	NA	NA	KRAS.p.A146T
P272	*KMT2A-AFF1*	Not detected	no	no	NA	NA	NA	NA	No COSMIC[3] hotspots

649   [1]VAF: Variant allele fraction; [2]LOH: Loss-of-heterozygosity; [3]COSMIC: Catalogue of Somatic Mutations in Cancer

650 Table 2: Mean and standard deviation of expressed rare (Not in ExAC or <0.5% allele frequency), low-frequency (0.5-1% allele frequency) and common
651 (>1% allele frequency) alleles in 39 *KMT2A*-R patient samples compared to ultra-mutated (P337) sample and its constituent expressed germline alleles and
652 somatic mutations.

	Not in ExAC	<0.5% AF[1] in ExAC	0.5-1% AF in ExAC	>1% AF in ExAC
KMT2A-R iALL	42.3±10.4	212.1±32.6	76.8±14.7	7722.3±266.0
P337 expressed	38	174	89	7590
P337 Germline	17	88	47	6235
P337 Somatic	21	86	42	1355

[1]AF: allele frequency; [2]ExAC: Exome aggregation consortium

Figure Legends:

Figure 1: Characterization of germline and somatic point mutations identifies an ultra-

mutated infant leukemia case. (a) Overlap of variants identified from diagnostic leukemia

specimen by RNA-seq and whole exome sequencing (WES). (b) Overlap of variants called in

diagnostic RNA-seq and remission bone marrow WES. (c) Overlap of variants called in

diagnostic and remission WES. Barplots colored showing overlap binned by derived allele

frequency as follows: present with AF<0.5% in ExAC (green), present with AF 0.5-1% in

ExAC (yellow), present with AF>1% in ExAC (purple), absent from the ExAC (orange), non-

overlapping variants are colored to show sites with coverage <20 reads (grey) or coverage

>20 reads (black) in the comparator. (d) Scatterplots of diagnostic leukemia (x-axis) versus

remission (y-axis) variant allele fraction (VAF) from WES for coding variants detected in

diagnostic leukemia RNA-seq.

Figure 2:   Clonal analysis of somatic ultra-mutations. Histograms showing the VAF

density distribution of somatic mutations (left, x-axis: dark shaded bars) or LOH-sites (right;

x-axis: light shaded bars) in diagnostic leukemia WES from canonical cases (a) and the ultra-

mutated case (b). (c) Illustration of cell lines derived from ultra-mutated specimen and

histograms displaying the VAF density distribution of somatic mutations (left, x-axis: dark

shaded bars) or LOH-sites (right; x-axis: light shaded bars) in cell lines (PER-784A: red;

PER-826A: green). (d) Venn diagram displaying the overlap in somatic mutations (left) and

LOH-sites (right) identified by *VarScan2* comparing the primary diagnostic specimen and

derived cell lines (PER-784A: red; PER-826A: green) to the primary remission sample. (e)

Somatic mutation and (f) LOH coding sites showing total read coverage (left y-axis) and VAF

(right y-axis) from primary specimens (left panels) and cell lines (right panels) ordered by

decreasing VAF in the diagnostic sample (x-axis).

Figure 3: Germline-like somatic signatures of ultra-mutations. (a-d) Histograms showing

the number (a-b) or frequency (c-d) of events overlapping the spectrum of ExAC derived

allele frequency bins with 5% increments. (a) Somatic mutations called by *VarScan2*

grouping homozygous and heterozygous events separately. (b) Loss-of-heterozygosity sites

called by *VarScan2* grouping reference→alternative and alterantive→reference sites

separately. (c-d) All quality filtered coding variants from each of the canonical patients or

P337 ultra-mutated specimen called by GATK grouped by single nucleotide polymorphisms

(SNPs) that were homozygous and heterozygous events from (c) remission sample or (d)

diagnostic sample. (e) Normalized fraction of variants within unique trinucleotide context

(ordered alphabetically on the x-axis) for *VarScan2* calls (left panel; as shown for panels a-b),

or remission variants (right panel; as shown for panel c). (f) Somatic signatures identified by

deconstructSigs for categories of mutations/SNPs (displayed in panel e) showing their relative

proportions (bar plot) and pair-wise Pearson's correlation coefficients (heatmap), ordered by

unsupervised hierarchical clustering.

Figure 4: Ultra-mutations target rare and common human alleles and are associated

with somatic point mutations encoding a novel *MSH2* allele and a rare *POLK* allele.

(a) Proportion of coding single nucleotide variants in diagnostic (D) and remission (R) whole

exome sequencing (WES) data binned by derived allele frequency as follows: present with

AF<0.5% in ExAC (green), present with AF 0.5-1% in ExAC (yellow), present with AF>1%

in ExAC (purple), absent from the ExAC (orange). Grey shading denotes data from the ultra-

mutated specimen. The distribution of variants called as germline, somatic or LOH by

VarScan2 are displayed for comparison. (b) Number of variants (y-axis) called from RNA-seq

reads from St Jude and Perth cohorts (open circles) are plotted with the total variants detected

in the ultra-mutated sample (shaded circles), germline-encoded variants (shaded triangle

pointing up) and somatic point mutations (shaded triangle pointing down) binned according to

the allele count in ExAC database (red: not in ExAC; green < 0.5%; yellow: 0.5-1% and blue

> 0.5%). (c) Bar plot displaying fraction of variants within rare, low-frequency and common

allele frequency bins (as described for panel a) ordered by increasing proportion of common

alleles (i.e. derived AF>1%) and excluding outliers (SJINF018, SJINF021, SJINF010 and

SJINF008). Proportion of RNA-seq germline and somatic variants identified by overlap with

remission/diagnostic WES data are displayed for comparison. (d) Crystal structure of the

MutS lesion recognition complex (PDB CODE 2O8D[35]) MSH2:MSH6 mismatch repair

proteins (MSH2: green, MSH6: blue) bound to ADP and G:T mismatch DNA (orange)

showing the location of the amino acid change (NM_000251:exon9:c.G1465A:p.E489K). (e)

Crystal structure of Pol k (green and blue representing different molecules) in complex with a

bulky DNA adduct (orange) (PDB CODE 4U6P[39]) showing locations of the amino acid

change (NM_016218:exon7:c.G893A:p.R298H).

Figure 5: Global transcriptional concordance of ultra-mutated infant leukemia. Multi-

dimensional scaling of RNA-seq count data from acute leukemias and normal bone marrow

progenitor populations displaying translocation partner gene by color and blood/leukemia

type by symbol shape according to the legend. Labels show replicate measurements from the

ultra-mutated diagnostic specimen, P337.

Supplementary Fig. S1. Copy-number alterations and loss-of-heterozygosity in ultra-

mutated *KMT2A*-R iALL diagnostic sample. (a) Genomic copy number ratios of the ultra-

mutated sample comparing diagnostic and remission WES data. (b) Proportion of reads

mapping to different copy-number bins (up to 4). (c) Box and whisker plots (median with

upper and lower quartiles) displaying the nearest distance of a variant/mutation to a

homozygous variant in the diagnostic WES data for canonical samples (left) compared to the

ultra-mutated sample (right) for *VarScan2* germline, somatic and LOH SNV calls.

Supplementary Table S1: Summary of samples

Supplementary Table S2: Read evidence of RNA-seq variants that were not detected in

matched WES data.

Supplementary Table S3: Comparison of RNA-seq reads from diagnosis sample with WES

reads.

Supplementary Table S4: Comparison of RNA-seq reads from diagnosis sample with WES

reads from diagnosis, remission and derived cell lines.

Supplementary Table S5: Annotation of somatic mutations and indels from ultra-mutated

patient P337.

Supplementary Table S6: Summary of LOH-sites called in primary P337 and derived cell

lines.

Supplementary Table S7: Summary of somatic SNVs called in primary P337 and derived

cell lines.

FIGURE 1

a

Diagnosis RNA vs exome

b

Diagnosis RNA vs remission exome

c

Diagnosis exome vs remission exome

P810, P809, P438, P399, P401, P337

Number of variants

Not in ExAC · <0.5% AF in ExAC · 0.5-1% AF in ExAC · >1% AF in ExAC · Discordant >20 reads · Discordant <20 reads

d

"Silent genomic landscape" · Ultra-mutated

P399 · P401 · P438 · P809 · P810 · P337

Remission exome VAF

Diagnosis exome VAF

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

Blood progenitor
MLL/KMT2A-Germline
MLL/KMT2A-MLLT1
MLL/KMT2A-AFF1
MLL/KMT2A-MLLT3
MLL/KMT2A-MLLT4
MLL/KMT2A-MLLT10
MLL/KMT2A-EPS15
MLL/KMT2A-USP2
MLL/KMT2A-CASC5
MLL/KMT2A-MLLT6
MLL/KMT2A-ELL
MLL/KMT2A-ENL
MLL/KMT2A-SEPT9
MLL/KMT2A-ENAH
MLL/KMT2A-GAS7

○ Diagnostic infant ALL
□ Relapse infant ALL
△ Childhood ALL
◇ Childhood AML
▽ Leucegene AML

SUPPLEMENTARY FIGURE S1