Recapitulation of human germline coding variation in an ultra-mutated infant leukemia

Alexander M Gout[1], Rishi S Kotecha[2,3,4], Parwinder Kaur[5,6], Ana Abad[1], Bree Foley[7], Kim W
Carter[8], Catherine H Cole[3], Charles Bond[9], Ursula R Kees[2], Jason Waithman[7,]*, Mark N
Cruickshank[1]*

[1]Cancer Genomics and Epigenetics Laboratory, Telethon Kids Cancer Centre, Telethon Kids
Institute, University of Western Australia, Perth, Australia.

[2]Leukaemia and Cancer Research Laboratory, Telethon Kids Cancer Centre, Telethon Kids
Institute, Perth, Australia.

[3]Department of Haematology and Oncology, Princess Margaret Hospital for Children, Perth,
Australia.

[4]School of Medicine, University of Western Australia, Perth, Australia.

[5]Personalised Medicine Centre for Children, Telethon Kids Institute, Australia.

[6]Centre for Plant Genetics & Breeding, UWA School of Agriculture & Environment.

[7]Cancer Immunology Unit, Telethon Kids Institute, Perth, Australia.

[8]McCusker Charitable Foundation Bioinformatics Centre, Telethon Kids Institute, Perth,
Australia.

[9]School of Molecular Sciences, The University of Western Australia, Perth, Australia.

*Corresponding author

*Correspondence to: Dr Mark Nicholas Cruickshank; address: 100 Roberts Road, Subiaco,
Western Australia, 6008; email: Mark.Cruickshank@telethonkids.org.au; telephone: +61 8
9489 7777; fax: +61 8 9489 7700. Dr Jason Waithman; address: 100 Roberts Road, Subiaco,
Western Australia, 6008; email: Jason.Waithman@telethonkids.org.au; telephone: +61 8 9489
7777; fax: +61 8 9489 7700.

Keywords: infant acute lymphoblastic leukemia, hypermutation, mismatch repair defect,
MSH2

Running title: Ultra-mutation targets germline alleles in an infant leukemia

Word Count: (4240)

Total number of Figures/Tables: 4 figures and 2 tables

Financial support: Children's Leukaemia and Cancer Research Foundation (Inc.) Fellowship
(MNC); NHMRC Career Development Fellowship 1066869 (JW); Raine Medical Research
Foundation Clinician Research Fellowship 2016-2018 (RSK); Children's Leukaemia and
Cancer Research Foundation (Inc.) project grants (URK, MNC, RSK); Ian Potter Foundation
(AG); Healy Foundation (AG). Telethon Kids Institute Blue Sky Research Grant (MNC,
URK, RSK, JW). Telethon Kids Institute Research Focus Area Precision Medicine Working
Group Funding (JW, MNC).

Conflicts of interest: The authors declare no conflicts of interest.

Abstract: *Mixed lineage leukemia/Histone-lysine N-methyltransferase 2A* gene

rearrangements occur in 80% of infant acute lymphoblastic leukemia, but the role of

cooperating events is unknown. We identified a case with over 1,000 non-synonymous coding

lesions. The patient presented at 82 days of age, representing the earliest recorded

manifestation of cancer hypermutation. The transcriptional profile showed global similarities

to canonical cases. Coding lesions were predominantly clonal and almost entirely targeting

alleles reported in human genetic variation databases with a notable exception in the

mismatch repair gene, *MSH2*. While the patient's germline was relatively depleted of rare and

low-frequency alleles, point mutations targeted rare and common alleles in proportion such

that their total abundance was equivalent to canonical cases. Somatic signatures suggest a role

for 5-methylcytosine deamination, DNA mismatch repair and DNA adducts. These data

indicate that common molecular processes shaping population-scale human genome variation

were operative during evolution of an ultra-mutated infant leukemia.

Introduction:

Histone-lysine N-methyltransferase 2A (*KMT2A*) (also known as *Mixed lineage*

leukemia) gene rearrangements (*KMT2A*-R) at 11q23 occur in infant, pediatric, adult and

therapy-induced acute leukemias and are associated with a poor prognosis. *KMT2A*-R occurs

in around 80% of infant acute lymphoblastic leukemia (ALL) and 35-50% of infant acute

myeloid leukemia (AML) involving more than 60 distinct partner genes [1]. *KMT2A*-R infant

ALL (iALL) genomes have an exceedingly low somatic mutation burden, with recurring

mutations in *RAS-PI3K* complex genes. Compared to wild-type KMT2A, the fusion

oncoprotein has altered histone-methyltransferase activity, causing epigenetic and

transcriptional deregulation. While KMT2A-oncoproteins are strong drivers of

leukemogenesis, several lines of evidence suggest that *KMT2A*-R alone is insufficient for

tumor initiation. For example, there is significant variability in latency of disease onset in

experimental models of *KMT2A*-fusions [2, 3] and in patients with *KMT2A*-R detected in

neonatal blood spots [4]. Furthermore, *KMT2A*-R acute leukemias [5] in children are enriched for

mutations affecting epigenetic regulatory genes and on average harbor more somatic lesions

than infants. These observations have prompted speculation that iALL is a distinct

developmental and genetic entity and pre-leukemic transformation occurs *in utero*, targeting a

cell of origin with a genetic/epigenetic permissive predisposition [6].

As a step towards defining cooperating events we performed next-generation

sequencing on a *KMT2A*-R iALL patient cohort and integrated the data with a larger set of

acute leukemias. We investigated the combined germline and somatic mutation profiles,

identifying a highly mutated specimen from a patient diagnosed at 82 days of age. Here we

report sequence analysis of this unique genetic entity in relation to its canonical *KMT2A*-R

iALL counterparts, revealing common mechanisms driving leukemia ultra-mutation and

population-scale human genome variation.

Results:

Characteristics of KMT2A-R iALL cohort and RNA-seq variant detection pipeline

We reported previously *KMT2A*-R fusion transcripts for 5/10 cases [7], and of the

remaining five *KMT2A*-R iALL cases two express *KMT2A-MLLT3* (P401, P438), two express

KMT2A-AFF1 (P272 and P848) and one case expresses *KMT2A-MLLT1* (P706). *KMT2A*-

fusion transcripts were detected in all samples and reciprocal fusion sequences detected in

5/10 samples (Table 1). To determine the validity of our RNA-seq based pipeline for

detecting rare and common alleles, we examined the concordance of single nucleotide

variants (SNVs) called from RNA-seq and WES data. Across seven samples with matched

WES and RNA-seq data, we identified a total of 54,792 point mutations from RNA-seq reads

of which 166 were absent from matched WES (coverage>20 reads), representing a

discordance rate of 0.3% (Figure 1a; Supplementary Table S2). In one sample, P706 there

were two events called with RNA-seq data that were not identified in matched WES but with

evidence extracted using the Integrated Genome Browser at previously reported somatic

mutation sites in *FLT3* (encoding FLT3 p.V491L [8]; 200/1,719 alternate RNA-seq reads;

15/214 alternate WES reads) and *KRAS* (encoding KRAS p.A146T [9]; 24/159 alternate RNA-

seq reads; 6/51 alternate WES reads). Furthermore, analysis of 117 discordant events with

coverage > 50 reads, revealed 76 recurrences in two or more samples (64%) at 26 unique

sites, however none of these sites have been validated as somatic mutations in cancer to date

in the Catalog of Somatic Mutations in Cancer (COSMIC)[10] or The Pediatric Cancer Genome

Project (PCGP)[11] databases. These results indicate a high rate of reproducibility of the RNA-

seq variant detection pipeline and suggest that RNA-DNA differences mostly comprise

artifacts and/or novel RNA-editing events [12] and a minority of putative somatic alterations

exclusively detected by RNA-seq. The full list of samples analysed in this study including

those downloaded from external sources is summarised in Supplementary Table S2.

Identification of a highly mutated infant leukemia

Expressed somatic mutations were defined as variants detected in both diagnostic

leukemia RNA-seq and WES, but absent in remission WES at sites with >20 reads (Figure

1b-d). Most patients (5/6) showed the expected "silent genomic landscape" (Supplementary

Table S3) except for patient P337 (Figure 1b-d; Supplementary Table S4). Three patients

expressed RAS-family hot-spot somatic mutations (P848 and P706: KRAS p.A146T and

P438: NRAS.p.G12S). We detected no expressed somatic mutations in patients P399 and

P401. The pair of monozygotic twins (P809 and P810) lacked private mutations but we

detected two shared expressed missense mutations clustered in the *PER3* gene (encoding

PER3.p.M1006R and PER3.p.K1007E). These results are consistent with recurring RAS-

pathway mutations and low mutation rate reported previously [5, 13]. However, in patient P337

RNA-seq reads we detected 1,559 expressed somatic mutations (Supplementary Table S4).

The overlap of both rare and common alleles across the matched datasets verifies the common

origin of RNA-seq and diagnostic/remission WES data ruling out sample contamination or

mislabeling (Figure 1a-c). We previously reported a complex *KMT2A* translocation involving

2q37, 19p13.3 and 11q23[14] in the highly mutated leukemia sample and established two cell

lines from the diagnostic sample [7].

Characterisation and clonal analysis of somatic lesions

Somatic lesions were further evaluated in six cases by comparing diagnostic and remission

WES. Two different variant allele fraction (VAF)-cut-offs, either VAF>0.1 or VAF>0.3, were

chosen to enrich for sub-clonal and clonal events respectively. There were on average ~82.2

(range 69-95) sub-clonal and ~3.4 clonal (range 3-4) somatic point mutations in canonical

KMT2A-R iALL exomes, confirming previous studies of a low somatic mutation rate in the

dominant clone (Table 1). In contrast, there were 198 sub-clonal and 5,054 clonal somatic

point mutations in the highly-mutated sample, equating to a rate of 139 mut/Mb, classifying

this sample as ultra-mutated. There were also ~40-fold increased number of indels in the

dominant clone of the ultra-mutated sample (n=260) compared to typical cases (average =

6.6; range: 3-9) including 13 predicted to cause a frame-shift protein-coding truncation (Table

1; Supplementary Table S5). Somatic substitutions encoded a total of 1,109 exonic missense

alleles and 12 exonic stop-gain alleles (Supplementary Table S5). We also noted that the

ultra-mutated case carried an excess of loss-of-heterozygosity (LOH) positions, with 6,555

high confidence sites; vastly more than typical cases which had a mean of 57.2 LOH positions

(range 28-100) (Figure 2a-b; Table 1).

WES was performed on two independent cell lines derived from the ultra-mutated diagnostic

specimen, PER-784A and PER-826A, to characterize clonal somatic events. (Figure 2c-d).

Most somatic and LOH-events detected in the primary ultra-mutated sample were also called

in both, PER-784A and PER-826A cell lines using P337 remission WES for all comparisons

(Figure 2d). These data suggest that the somatic lesions are largely clonal in agreement with

the VAF-distributions (Figure 2b, d). In total, 4,754/5,054 (94%) somatic events

(Supplementary Table S6) and 6,150/6,575 (93%) LOH-events (Supplementary Table S7)

were called in the derived cell lines. Moreover, there were relatively few discordant sites

when directly comparing cell lines with the primary diagnostic sample (PER-784A: 34

discordant sites; PER-826A: 44 discordant sites; Supplementary Table S6)). We further

investigated evidence of clonal somatic lesions in cell lines using an alternative approach, by

directly extracting quality filtered WES reads mapping to coding somatic point mutations

(n=2,411; Q>50 in diagnosis WES; Figure 2e) and LOH-sites (n=3,117; Q>50 in remission

WES; Figure 2f; Supplementary Table S8).   All coding clonal somatic mutations were

supported by reads in both cell lines of which 121 (5%) were homozygous (VAF>0.95 in cell

line and VAF>0.9 in primary sample). Most LOH-sites were evidenced by reference alleles

(VAF<10%; 2,179/3,117) in cell lines, with ~30% of LOH-sites evidenced by alternate alleles

(VAF> 90%; 926/3,117) and only 12 sites (0.3%) with both reference and alternate alleles

(VAF range 10-90%).

We performed copy number analysis of the ultra-mutated sample to map LOH-sites to

chromosomal segments with evidence of deletions. Most genomic regions were called as

diploid with limited chromosomal copy loss detected (Supplementary Figure S1a-b). There

was a total of 382/6,555 (5.8%) LOH-sites mapping to homozygous deletions, suggesting

only a minority of the identified LOH-sites likely arose through chromosome segment

deletion. Infant *KMT2A*-R ALL typically carry a small number of copy-number alterations

which may coincide with LOH-sites. We determined the distance of LOH-sites, somatic

mutations and germline variants (called by VarScan2) to the nearest homozygous SNV in

each leukemia diagnostic sample finding that in canonical cases LOH-sites tend to be located

closer to homozygous sites than either germline or somatic variants (Supplementary Figure

S1c; left panel) and show a unique bimodal distribution (Supplementary Figure S1c; right

panel). In contrast the average distance of LOH-sites to homozygous SNVs in the ultra-

mutated sample was similar to that of germline and somatic variants with each displaying a

similar distribution (Supplementary Figure S1c). Thus, LOH-sites in the ultra-mutated do not

show evidence of localising within deleted or copy-neutral loss-of-heterozygosity

chromosomal segments.

In summary these data demonstrate that somatic lesions in the primary sample were

predominantly clonal and LOH-sites appear to comprise a large proportion of somatic

substitutions rather than chromosomal alterations. We also note relatively few coding

sequence changes in derived cell lines compared to the dominant diagnostic leukemic clone,

both which are capable of long-term propagation. These results suggest that loss of the

hypermutator phenotype was either acquired during *in vitro* culture or in the patient sample

prior to clonal expansion of the bulk diagnostic leukemia.

Signatures of somatic mutations and LOH-sites correlate with profiles of germline alleles

We next investigated the sequence features of somatic and LOH substitutions in the

ultra-mutated sample (Figure 3). We hypothesised that LOH-sites and somatic mutations may

show common feature biases depending on the context of alteration. Somatic events were

evaluated separately that induce a heterozygous mutation (n=4,773) or homozygous mutation

(n=281), while LOH causing reference→alternative changes (n=2,057) were evaluated

separately to alternative→reference changes (n=4,453). Since we noted an overlap in the

somatic mutations of the ultra-mutated sample with variants in the ExAC database (Figure 1a-

c), we first examined the distribution of these sites with respect to the derived allele frequency

in ExAC. We observe that each of these categories of sequence alterations in the ultra-

mutated sample were distributed across the range of population allele-frequencies (Figure 3a-

b). While the distributions of LOH and somatic sites were similar, somatic heterozygous

changes were relatively more frequent at higher frequency alleles as were LOH

reference→alternate alleles.

Somatic signatures of ultra-mutations were investigated in comparison to

heterozygous and homozygous single nucleotide polymorphisms (SNPs) in patient remission

samples. Heterozygous positions were defined by VAF<0.9 yielding 51,347 heterozygous and

31,237 homozygous SNPs in canonical remission samples and 10,681 heterozygous and 6

617 homozygous SNPs in the ultra-mutated remission sample. Analysis of WES remission

variants from canonical patients and the ultra-mutated sample revealed similar ExAC allele

frequency distributions with distinctions between heterozygous and homozygous alleles

(Figure 3c-d). The overall enrichment of each category of somatic substitution and LOH

across the 96 possible trinucleotide contexts resembled patient remission SNPs (Figure 3e).

Pre-defined signatures were used to infer mutational profiles revealing Signature 1A

(associated with spontaneous deamination of 5-methylcytosine), Signature 12 (associated

with unknown process with strand-bias for T→C substitutions) and Signature 20 (associated

with defective DNA mismatch repair) as enriched in each of the mutation/variant sets,

however with distinctive proportions (Figure 3f). Heterozygous somatic mutations and LOH

alternative→reference sites were additionally enriched for Signature 1B (variant of Signature

1A) and showed highly correlated profiles (Pearson's correlation 0.919). LOH

reference→alternative sites were additionally enriched for Signature 5 (associated with

unknown process with strand-bias for T→C substitutions) and were highly correlated with

homozygous somatic mutation profiles (0.912) and to profiles of patient heterozygous

positions (P337: 0.935; canonical: 0.926). Homozygous somatic mutations also displayed

similarities to heterozygous patient SNPs (P337: 0.981; canonical: 0.985). Overall the somatic

mutational profiles indicate highly related patterns of both somatic ultra-mutations and LOH-

sites. Notably LOH alternative→reference sites showed unique and overlapping features to

somatic heterozygous mutations. We also observe underlying similarities in the profiles of

somatic substitutions and germline alleles. While copy-number alterations or zygosity shifts

may underlie some of the observed LOH in the ultra-mutated genome, collectively our data

suggests most these sites arise through point mutagenesis.

Ultra-mutation distribution recapitulates common and rare germline variation

We noted a similar spectrum of rare and common alleles in the ultra-mutated

leukemia transcriptome compared to typical *KMT2A*-R iALL cases; however, the remission

sample had the lowest number of rare/low-frequency alleles (Figure 1a-c; ExAC<0.5%). We

extended this observation by investigating the rare and common allele abundance of germline

variants and somatic mutations in the ultra-mutated genome compared with a larger cohort of

KMT2A-R iALL transcriptomes (n=32 St. Jude patients and n=8 Perth patients). Most typical

KMT2A-R iALL diagnostic specimens showed similar numbers of rare and low-frequency

expressed SNVs, except for 4/32 cases sequenced by St Jude investigators, which had an

elevated number of rare and low-frequency ExAC alleles (Figure 4a). Excluding these

samples, typical *KMT2A*-R iALL patients expressed on average 42.3 ±10.4 (standard

deviation) variants that were absent from ExAC and 212 ± 32.5 variants with an allele

frequency of <0.5% in ExAC. The total number of expressed variants in the ultra-mutated

sample was slightly lower than typical cases (38 absent from ExAC and 174 AF <0.5% in

ExAC), and were approximately comprised of half germline (17/38 and 88/174 respectively)

and half somatic acquired alleles. Overall, the germline of the ultra-mutated sample had the

236 lowest prevalence of rare and low-frequency alleles among the larger sample size (Table 2).

237 Furthermore, the somatic mutation abundance within each allele-frequency bin was

238 proportional to their abundance in prototypical silent cases. Consequently, the low numbers of

239 rare germline alleles in the ultra-mutated transcriptome were off-set by somatic mutations

240 such that the total numbers were within the same range as typical cases (Figure 4a-b). A

241 similar trend was also observed in WES from the Perth cohort, however with a smaller

242 magnitude of difference in the prevalence of rare alleles (Figure 4b; 2.9% rare alleles in P337

243 compared to 3.5±0.2% in canonical cases). We conclude that despite carrying thousands of

244 somatic mutations, the coding-genome of ultra-mutated patient P337 shows similarities to

245 typical *KMT2A*-R iALL cases, vis-à-vis rare allele prevalence and representation of common

246 human genetic variation.

247 *RAS-pathway and DNA-replication and repair gene mutations*

248   Gene ontology analysis of the expressed non-synonymous somatic mutations (n=535)

249 revealed over-representation of RAS-pathway genes (7/23 genes; 7-fold enriched; *P*=0.049;

250 *RALGDS*, *RAC1*, *CHUK*, *PIK3R1*, *NFKB1*, *PLD1* and *PIK3CA*) and "*MLL* signature 1 genes"

251 (33/369 genes; 2-fold enriched; *P*=0.046). Of the 535 non-synonymous expressed somatic

252 mutations, 45 were predicted deleterious/damaging, including in the mismatch repair gene

253 *MSH2* (p.E489K). The *MSH2* somatic mutation localizes to a hot-spot, within the mismatch

254 repair ATPase domain in a region at the periphery of the DNA binding site, approximately 20

255 Å from the DNA molecule in the complex structure of full-length MSH2, a fragment of

256 MSH6, ADP and DNA containing a G:T mispair [15]. The mutation encodes a substitution of a

257 negatively charged glutamate with a positively charged lysine (Figure 4c). There was a non-

258 synonymous somatic mutation in another mismatch repair gene, *MUTYH*; however, this site is

259 a common human allele with a derived AF of 0.297 and was not called as

260 damaged/deleterious by our prediction pipeline. There were no somatic mutations nor

261 rare/deleterious germline alleles detected in *POLE* or *POLD1* which are frequently co-

262 mutated with *MSH2* in ultra-mutated tumours [16, 17] and in sporadic early-onset colorectal

cancer [18]. Among the 45 expressed and predicted damaging somatic mutations, there are two

rare substitutions in DNA polymerase encoding genes: *POLG2* (p.D308V; ExAC AF

0.0000165), which encodes a mitochondrial DNA polymerase and *POLK* (p.R298H; ExAC

AF 0.0003), which encodes an error prone replicative polymerase that catalyzes translesion

DNA synthesis. The *POLK* mutation maps to a conserved impB/mucB/samB family domain

(Pfam: PF00817) involved in UV-protection. The encoded substitution of arginine to histidine

is located approximately 20 Å from the DNA molecule in the structure of a 526 amino acid

N-terminal Pol K fragment complexed with DNA containing a bulky deoxyguanosine adduct

induced by the carcinogen benzo[a]pyrene (Figure 4d) [19]. Both wild-type and mutant residues

are positively charged, however arginine is strongly basic (pI 11) while histidine is weakly

basic (pI 7) and titratable at physiological pH [20]. Expressed and predicted functional germline

rare alleles were also observed in *ATR*, a master regulator of the DNA damage response

(p.L2076V; ExAC AF 0.0002), and RAS-pathway gene *PLD2* p.R330Q; ExAC AF 0.0032).

Global transcriptional similarities of the ultra-mutated and canonical infant leukemias

We next examined the transcriptional profile of the highly mutated *KMT2A*-R iALL

specimen in comparison to a collection of patient leukemia samples and purified blood

progenitor populations (Figure 5). As expected cell type was the major source of variation

distinguishing normal hematopoietic progenitor, lymphoblastic leukemias and myeloid

leukemias. We also observe clustering according to *KMT2A* translocation status and partner

genes as reported previously [21]. Replicate samples of the highly mutated *KMT2A*-R iALL

specimen cluster with canonical counterparts demonstrating global transcriptional similarities

(Figure 5).

Discussion:

We report sequence analysis of an ultra-mutated acute lymphoblastic leukemia

diagnosed in an infant at 82 days of age which, to our knowledge, represents the earliest onset

of a hypermutant tumor recorded. The mutation rate observed was at the upper limit of cases

reported previously in childhood cancers [16, 17]. By integrating analysis of somatic mutations

and germline alleles with a larger cohort of *KMT2A*-R iALL patients we found that the high

somatic mutation rate does not generate large numbers of rare alleles compared to typical

cases. Rather, we observe that the ultra-mutated *KMT2A*-R leukemic coding genome closely

resembles the rare allele landscape of canonical cases. This apparent paradox is explained by

two factors: (1) the germline of this patient was comparatively depleted of rare alleles, and (2)

somatic point mutations were targeted to rare and common alleles in proportions that off-set

this discrepancy. Indeed, both the rare allele burden and transcriptional profile of the ultra-

mutated leukemia were similar to its prototypic counterparts. Therefore, despite its radically

different genetic route to leukemogenesis, the molecular profile of the ultra-mutated specimen

converged towards that of canonical cases.

Most ultra-mutated childhood cancers reported previously have a combined mismatch

repair and proof reading polymerase deficiency [16, 17] - we detect evidence for the former

(*MSH2* point mutation), but not the latter. However, we identify somatic mutations in two

genes encoding DNA polymerases (*POLK* and *POLG2*) and an additional mismatch repair

gene (*MUTYH*). Notably, the *POLK* mutation is a rare allele, predicted functional and

encodes pol κ, a replicative error-prone translesion DNA polymerase that plays a role in

bypassing unrepaired bases that otherwise block DNA replication [22, 23]. For example, pol κ

preferentially incorporates dAMP at positions complementary to 7,8-dihydro-8-oxo-2′-

deoxyguanosine adducts [24]. As a corollary to these observations, *POLK* mutations have been

found in prostate cancer with elevated C→T transition [25]. Further studies are required to

determine if any of these mutations cooperate with the *MSH2*-lesion in a pre-natal cell-of-

origin to specify the unique mutational pattern observed in the ultra-mutated *KMT2A*-R iALL

genome.

The largest fraction of mutations was attributed to Signature 1, associated with 5-

methyl-cytosine deamination inducing C→T transitions, which accumulates in normal

somatic cells with age over the human lifespan [26]. This signature is found in many cancers,

prominently in AML patients with germ-line mutations in *MBD4* [27]. LOH-sites

(reference→alternative) were enriched for Signature 5 which is also correlated with human

aging. Consistent with previous reports, we observe enrichment of Signature 20 associated

with mismatch repair defective tumors [28]. We also find enrichment of Signature 12 which

arises from a poorly characterized underlying molecular process; however, the dominant base

transitions, A→G and T→C have been observed in association with DNA damage (e.g. by

adducts) in normal and cancer tissues [29, 30]. These observations suggest common mutagenic

and repair deficits driving age-related mutation were operative in the cell of origin during pre-

natal evolution of the ultra-mutated infant leukemia genome.

The preponderance of ultra-mutated sites to overlap germline alleles in proportion to

their population allele frequencies demonstrates that hypermutation can recapitulate aspects

of population scale human genetic variation in a cancer genome. Remarkably, approximately

half of the somatic substitutions appear to target heterozygous SNPs resulting in loss-of-

heterozygosity. They, more frequently target the alternate allele inducing reversion to the

reference base but around 30% of events mutate a heterozygous site resulting in a

homozygous variant allele in the diagnostic sample. Most ultra-mutations are therefore likely

passengers coinciding with common human alleles, consistent with findings that

hypermutated tumours tend to have relatively few drivers compared to their mutation burden

[31]; however, as proposed previously by Valentine and colleagues[32], a minority of rare alleles

may contribute to infant leukemogenesis. While our results do not resolve the temporal order

of the key oncogenic events - namely *MSH2* point mutation, genomic hypermutation and

KMT2A translocation – due to the clonal nature of these events they appear to have occurred

in rapid succession. This could be explained by a sentinel DNA repair defect inducing

hypermutation and genome instability followed by positive selection of cooperating lesions

conferring a growth advantage resulting in clonal expansion and fixation of passengers [33]. An

alternative model is also possible in which the DNA-repair defect induced by the *MSH2* point

mutation may impair double-strand-break repair [34, 35] promoting *KMT2A*-rearrangement, but

with neutral selection of somatic point mutations. Such a scenario appears inconsistent with

the observed predominance of clonal mutations, but could be reconciled if the hypermutator

phenotype was lost prior to clonal expansion. Since we have observed loss of hypermutator

phenotype in cell lines derived from the diagnostic leukaemia specimen the later model is not

ruled out by our data. These data illuminate a link between mutagenesis during human aging,

germline variation and somatic hypermutation in cancer. Further studies are required to

determine if combinations of somatic mutations and/or germline alleles promote *KMT2A*-R

iALL leukemogenesis and if somatic mutated sites significantly overlap human germline

alleles in other malignancies.

Methods:

Patient specimens, external sequencing data and controls

The study cohort consisted of ten infants diagnosed at Princess Margaret Hospital for

Children (Perth), of which we have previously reported RNA-seq data for five patients [7].

Additional RNA-seq data was downloaded for 39 *KMT2A*-R iALL cases at diagnosis, six

cases at relapse, six cases lacking the *KMT2A*-R (*KMT2A*-germline) and for pediatric

KMT2A-R patients at diagnosis, comprising 10 AML cases, seven ALL cases and one

undifferentiated case as reported previously by Andersson *et al.* [5] (accession

EGAS00001000246). RNA-seq data comprising 28 *KMT2A*-R and 52 *KMT2A*-germline adult

diagnosistic AML cases downloaded from the Short Read Archive (BioProject Umbrella

Accession PRJNA278766[36]; Leucegene: AML sequencing). Normal human bone marrow

transcriptomes were downloaded from the Short Read Archive (BioProject PRJNA284950)

comprising 20 pooled sorted human hematopoietic progenitor populations [37].

Sample preparation and next-generation sequencing

RNA was isolated with TRIzol™ Reagent (Invitrogen Life Science Technologies,

Carlsbad, CA, USA) and purified with RNeasy Purification Kits (Qiagen, Hilden, Germany).

Genomic DNA was isolated with QIAamp DNA Mini Kit (Qiagen). Nucleic acid integrity

and purity was assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies, Santa

Clara, CA). RNA-seq was performed at the Australian Genome Research Facility, Melbourne

with 1 μg RNA as previously reported [7]. Whole exome sequencing (WES) was performed by

the Beijing Genome Institute using SureSelect XT exon enrichment (Agilent Technologies)

with 200 ng of remission DNA (v4 low input kit) or 1 μg of diagnostic DNA (v5). Diagnostic

RNA and remission genomic DNA sequencing was performed on a HiSeq 2000 (Illumina,

Inc., San Diego, CA, USA) instrument with 100bp paired end reads; diagnostic genomic

DNA sequencing was performed on a HiSeq 2500 with 150bp paired end reads. Sequencing

libraries were constructed using Illumina TruSeq kits (Illumina, Inc.).

Bioinformatics

For gene-expression analysis, RNA-seq reads were mapped to hg19 with HISAT and

gene-counts generated with HTseq [38]. Data was normalized with RUV-seq [39] implementing

factor analysis of negative control genes identified by fitting a linear model with cell type

(hematopoietic sub-set or leukemia sub-type) as a covariate. voom! [40] was used to adjust for

library size and generate counts per million (cpm) values. *KMT2A*-fusion split-reads were

identified using RNA-seq data with *FusionFinder* [41] using default settings.

Single nucleotide variants (SNVs) were identified from RNA-seq data using the

intersection of two methodologies: (1) the "Genome Analysis Toolkit (GATK) Best Practices

workflow for SNP and indel calling on RNAseq data" and (2) *SNPiR* [42]. WES data was

processed based on the GATK Best Practices workflow. Variants were characterized using

custom scripts that extract the coverage of sequenced bases from VCF and bam files,

determining the overlap among samples, and calculating variant allele fractions (VAFs).

High confidence somatic mutations were identified using *VarScan2* (v2.3.9) [43] setting

cut-offs of P-value<0.005; coverage > 50X; VAF > 0.1 or VAF > 0.3 to enrich for clonal

mutations. Variants were annotated with ANNOVAR [44] and with metadata from additional

sources including seven functional/conservation algorithms (PolyPhen2 [45], Sift [46],

MutationTaster [47], likelihood ratio test [48], GERP [49], PhyloP [50], and CADD [51]) and databases of

human genetic variation (ExAC, 1000 genomes, HapMap, dbSNP, COSMIC and ClinVar).

Predicted functional alleles were assigned for variants called as damaged/deleterious by at

least 4/7 algorithms.

Somatic mutational spectra were investigated using the R package, *DeconstructSigs*

(v1.8.0) [52] calculating the fraction of somatic substitutions within the possible 96-trinucleotide

contexts relative to the exome background. Copy number analysis was performed on WES

data using *VarScan2* to calculate log2 depth ratio and GC-content and *Sequenza* (v2.1.2) [53] to

identify chromosomal segments with copy loss. Gene ontology enrichment was performed

using GREAT (v3.0.0) [54] bioinformatics tools. Mutations were inspected by mapping the

genomic position to crystal structures in the Protein DataBank[55] and analyzed with Pymol

(v1.8.6.2).

Data and code availability

Scripts used to execute variant detection software and analysis are available at

https://github.com/cruicks/ultramutated-case under an MIT license and are being developed

on Github (https://github.com/cruicks/vcfbamCompare). The sequencing data has been

deposited in the Short Read Archive with study ID PRJNA296746

Financial support: Children's Leukaemia and Cancer Research Foundation (Inc.) Fellowship

(MNC); NHMRC Career Development Fellowship 1066869 (JW); Raine Medical Research

Foundation Clinician Research Fellowship 2016-2018 (RSK); Children's Leukaemia and

Cancer Research Foundation (Inc.) project grants (URK, MNC, RSK); Ian Potter Foundation

(AG); Healy Foundation (AG). Telethon Kids Institute Blue Sky Research Grant (MNC,

URK, RSK, JW). Telethon Kids Institute Research Focus Area Precision Medicine Working

Group Funding (JW, MNC).

Acknowledgments: We thank Daniel MacArthur and Fengmei Zhao for assisting with exome

sequence analysis; and researchers at The Pediatric Cancer Genome Project at St. Jude

Children's Research Hospital for provision of data. We thank all the patients, families and

hospital staff for providing samples.

Authorship contributions: AG wrote computer code, performed bioinformatics, conceived

analysis strategy, interpreted and analyzed data and edited the manuscript. AA and BF

performed experiments. KWC, PK and CB performed additional data analysis and

interpretation. RSK provided patient samples and clinical data, assisted with data

interpretation and edited the manuscript. CHC, URK and JW acquired funding support,

assisted with data interpretation and edited the manuscript. MNC performed bioinformatics,

formulated research, conceived analysis strategy, acquired funding support, interpreted data,

designed experiments and drafted the manuscript.

Disclosure of Conflicts-of-Interest: The authors declare no conflicts of interest.

References:

1.   Meyer C, *et al.* The MLL recombinome of acute leukemias in 2017. *Leukemia*,
(2017).

435
436   2.   Bergerson RJ, *et al.* An insertional mutagenesis screen identifies genes that
437   cooperate with Mll-AF9 in a murine leukemogenesis model. *Blood* **119**, 4512-4523
438   (2012).

439
440   3.   Chen W, O'Sullivan MG, Hudson W, Kersey J. Modeling human infant MLL leukemia
441   in mice: leukemia from fetal liver differs from that originating in postnatal marrow.
442   *Blood* **117**, 3474-3475 (2011).

443
444   4.   Gale KB, *et al.* Backtracking leukemia to birth: identification of clonotypic gene
445   fusion sequences in neonatal blood spots. *Proc Natl Acad Sci U S A* **94**, 13950-13954
446   (1997).

447
448   5.   Andersson AK, *et al.* The landscape of somatic mutations in infant MLL-rearranged
449   acute lymphoblastic leukemias. *Nat Genet* **47**, 330-337 (2015).

450
451   6.   Sanjuan-Pla A, *et al.* Revisiting the biology of infant t(4;11)/MLL-AF4+ B-cell acute
452   lymphoblastic leukemia. *Blood* **126**, 2676-2685 (2015).

453
454   7.   Cruickshank MN, *et al.* Systematic chemical and molecular profiling of MLL-
455   rearranged infant acute lymphoblastic leukemia reveals efficacy of romidepsin.
456   *Leukemia*, (2016).

457
458   8.   He J, *et al.* Integrated genomic DNA/RNA profiling of hematologic malignancies in
459   the clinical setting. *Blood* **127**, 3004-3014 (2016).

460
461   9.   Chang MT, *et al.* Identifying recurrent mutations in cancer reveals widespread
462   lineage diversity and mutational specificity. *Nat Biotechnol* **34**, 155-163 (2016).

463
464   10.   Forbes SA, *et al.* COSMIC: somatic cancer genetics at high-resolution. *Nucleic Acids
465   Res* **45**, D777-D783 (2017).

466
467   11.   Downing JR, *et al.* The Pediatric Cancer Genome Project. *Nat Genet* **44**, 619-622
468   (2012).

469
470   12.   Pickrell JK, Gilad Y, Pritchard JK. Comment on "Widespread RNA and DNA sequence
471   differences in the human transcriptome". *Science* **335**, 1302; author reply 1302
472   (2012).

474   13.   Chang VY, Basso G, Sakamoto KM, Nelson SF. Identification of somatic and germline
475   mutations using whole exome sequencing of congenital acute lymphoblastic
476   leukemia. *BMC Cancer* **13**, 55 (2013).

478   14.   Henderson MJ, *et al.* A xenograft model of infant leukaemia reveals a complex MLL
479   translocation. *Br J Haematol* **140**, 716-719 (2008).

481   15.   Warren JJ, Pohlhaus TJ, Changela A, Iyer RR, Modrich PL, Beese LS. Structure of the
482   human MutSalpha DNA lesion recognition complex. *Mol Cell* **26**, 579-592 (2007).

484   16.   Campbell BB, *et al.* Comprehensive Analysis of Hypermutation in Human Cancer. *Cell*
485   **171**, 1042-1056 e1010 (2017).

487   17.   Shlien A, *et al.* Combined hereditary and somatic mutations of replication error
488   repair genes result in rapid onset of ultra-hypermutated cancers. *Nat Genet* **47**, 257-
489   262 (2015).

491   18.   Palles C, *et al.* Germline mutations affecting the proofreading domains of POLE and
492   POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet* **45**, 136-144
493   (2013).

495   19.   Jha V, Bian C, Xing G, Ling H. Structure and mechanism of error-free replication past
496   the major benzo[a]pyrene adduct by human DNA polymerase kappa. *Nucleic Acids*
497   *Res* **44**, 4957-4967 (2016).

499   20.   Szpiech ZA, *et al.* Prominent features of the amino acid mutation landscape in
500   cancer. *PLoS One* **12**, e0183273 (2017).

502   21.   Stam RW, *et al.* Gene expression profiling-based dissection of MLL translocated and
503   MLL germline acute lymphoblastic leukemia in infants. *Blood* **115**, 2835-2844 (2010).

505   22.   Kanemaru Y, *et al.* DNA polymerase kappa protects human cells against MMC-
506   induced genotoxicity through error-free translesion DNA synthesis. *Genes Environ*
507   **39**, 6 (2017).

509   23.   Maddukuri L, Ketkar A, Eddy S, Zafar MK, Eoff RL. The Werner syndrome protein
510   limits the error-prone 8-oxo-dG lesion bypass activity of human DNA polymerase
511   kappa. *Nucleic Acids Res* **42**, 12027-12040 (2014).

513   24.   Irimia A, Eoff RL, Guengerich FP, Egli M. Structural and functional elucidation of the
514   mechanism promoting error-prone synthesis by human DNA polymerase kappa

515   opposite the 7,8-dihydro-8-oxo-2'-deoxyguanosine adduct. *J Biol Chem* **284**, 22467-
516   22480 (2009).

517
518   25.   Yadav S, Mukhopadhyay S, Anbalagan M, Makridakis N. Somatic Mutations in
519   Catalytic Core of POLK Reported in Prostate Cancer Alter Translesion DNA Synthesis.
520   *Hum Mutat* **36**, 873-880 (2015).

521
522   26.   Alexandrov LB*, et al.* Clock-like mutational processes in human somatic cells. *Nat
523   Genet* **47**, 1402-1407 (2015).

524
525   27.   Mathijs A Sanders EC, Christoffer Flensburg, Annelieke Zeilemaker, Sarah E Miller,
526   Adil al Hinai, Ashish Bajel, Bram Luiken, Melissa Rijken, Tamara Mclennan, Remco M
527   Hoogenboezem, François G Kavelaars, Marnie E Blewitt, Eric M Bindels, Warren S
528   Alexander, Bob Löwenberg, Andrew W Roberts, Peter J M Valk, Ian Majewski.
529   Germline loss of MBD4 predisposes to leukaemia due to a mutagenic cascade driven
530   by 5mC. *bioRxiv*, (2018).

531
532   28.   Alexandrov LB*, et al.* Signatures of mutational processes in human cancer. *Nature
533   **500**, 415-421 (2013).

534
535   29.   Chen L, Liu P, Evans TC, Jr., Ettwiller LM. DNA damage is a pervasive cause of
536   sequencing errors, directly confounding variant identification. *Science* **355**, 752-756
537   (2017).

538
539   30.   Lin J, Shi T. Error-prone DNA polymerase and oxidative stress increase the incidences
540   of A to G mutations in tumors. *Oncotarget* **8**, 45154-45163 (2017).

541
542   31.   Martincorena I*, et al.* Universal Patterns of Selection in Cancer and Somatic Tissues.
543   *Cell* **171**, 1029-1041 e1021 (2017).

544
545   32.   Valentine MC*, et al.* Excess congenital non-synonymous variation in leukemia-
546   associated genes in MLL- infant leukemia: a Children's Oncology Group report.
547   *Leukemia*, (2013).

548
549   33.   Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer therapies.
550   *Nat Rev Clin Oncol*, (2017).

551
552   34.   van Oers JM*, et al.* The MutSbeta complex is a modulator of p53-driven
553   tumorigenesis through its functions in both DNA double-strand break repair and
554   mismatch repair. *Oncogene* **33**, 3939-3946 (2014).

556 35. Burdova K, Mihaljevic B, Sturzenegger A, Chappidi N, Janscak P. The Mismatch-
557   Binding Factor MutSbeta Can Mediate ATR Activation in Response to DNA Double-
558   Strand Breaks. *Mol Cell* **59**, 603-614 (2015).

559
560 36. Lavallee VP, *et al.* The transcriptomic landscape and directed chemical interrogation
561   of MLL-rearranged acute myeloid leukemias. *Nat Genet* **47**, 1030-1037 (2015).

562
563 37. Casero D, *et al.* Long non-coding RNA profiling of human lymphoid progenitor cells
564   reveals transcriptional divergence of B cell and T cell lineages. *Nat Immunol* **16**,
565   1282-1291 (2015).

566
567 38. Anders L, *et al.* Genome-wide localization of small molecules. *Nat Biotechnol* **32**, 92-
568   96 (2014).

569
570 39. Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor
571   analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).

572
573 40. Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model
574   analysis tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).

575
576 41. Francis RW, Thompson-Wicking K, Carter KW, Anderson D, Kees UR, Beesley AH.
577   FusionFinder: a software tool to identify expressed gene fusion candidates from
578   RNA-Seq data. *PLoS One* **7**, e39987 (2012).

579
580 42. Piskol R, Ramaswami G, Li JB. Reliable identification of genomic variants from RNA-
581   seq data. *Am J Hum Genet* **93**, 641-651 (2013).

582
583 43. Koboldt DC, *et al.* VarScan 2: somatic mutation and copy number alteration
584   discovery in cancer by exome sequencing. *Genome Res* **22**, 568-576 (2012).

585
586 44. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants
587   from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).

588
589 45. Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense
590   mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**, Unit7 20 (2013).

591
592 46. Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous
593   variants on protein function using the SIFT algorithm. *Nat Protoc* **4**, 1073-1081
594   (2009).

595
596 47. Schwarz JM, Rodelsperger C, Schuelke M, Seelow D. MutationTaster evaluates
597   disease-causing potential of sequence alterations. *Nat Methods* **7**, 575-576 (2010).

599   48.   Chun S, Fay JC. Identification of deleterious mutations within three human genomes.
600   *Genome Res* **19**, 1553-1561 (2009).

602   49.   Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, Sidow A. Distribution and
603   intensity of constraint in mammalian genomic sequence. *Genome Res* **15**, 901-913
604   (2005).

606   50.   Cooper GM, *et al.* Single-nucleotide evolutionary constraint scores highlight disease-
607   causing mutations. *Nat Methods* **7**, 250-251 (2010).

609   51.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general
610   framework for estimating the relative pathogenicity of human genetic variants. *Nat*
611   *Genet* **46**, 310-315 (2014).

613   52.   Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C. DeconstructSigs:
614   delineating mutational processes in single tumors distinguishes DNA repair
615   deficiencies and patterns of carcinoma evolution. *Genome Biol* **17**, 31 (2016).

617   53.   Favero F, *et al.* Sequenza: allele-specific copy number and mutation profiles from
618   tumor sequencing data. *Ann Oncol* **26**, 64-70 (2015).

620   54.   McLean CY, *et al.* GREAT improves functional interpretation of cis-regulatory
621   regions. *Nat Biotechnol* **28**, 495-501 (2010).

623   55.   Berman HM, *et al.* The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000).

626   Table 1: Characterisation of somatic lesions in *KMT2A*-R patient samples.

Patient ID	KMT2A-fusion	Reciprocal KMT2A-fusion	Diagnostic leukemia DNA	Constitutional remission DNA	Point Mutations 0.1 > VAF[1] < 0.3	Point Mutations VAF > 0.3	Indels VAF > 0.3	LOH[2]-sites	Point mutations or hotspots with RNA-SEQ reads
P337	KMT2A-KMT2AT1	Not detected	Exome	Exome	198	5,054	260	6,555	> 1,000 expressed somatic mutations
P401	KMT2A-KMT2AT3	Not detected	Exome	Exome	78	3	9	45	No non-silent somatic mutations
P399	KMT2A-AFF1	AFF1-KMT2A	Exome	Exome	95	3	7	67	No non-silent somatic mutations
P438	KMT2A-KMT2AT3	Not detected	Exome	Exome	74	4	9	100	NRAS.p.G12S
P810	KMT2A-EPS15	EPS15-KMT2A	Exome	Exome	95	4	5	46	PER3.p.M1006R, PER3.p.K1007E
P809	KMT2A-EPS15	EPS15-KMT2A	Exome	Exome	69	3	3	28	PER3 M1006R, PER3.p.K1007E
P848	KMT2A-AFF1	AFF1-KMT2A	no	Exome	NA	NA	NA	NA	KRAS.p.A146T
P287	KMT2A-AFF1	AFF1-KMT2A	no	no	NA	NA	NA	NA	No COSMIC[3] hotspots
P706	KMT2A-KMT2AT1	Not detected	no	Exome	NA	NA	NA	NA	KRAS.p.A146T
P272	KMT2A-AFF1	Not detected	no	no	NA	NA	NA	NA	No COSMIC[3] hotspots

627   [1]VAF: Variant allele fraction; [2]LOH: Loss-of-heterozygosity; [3]COSMIC: Catalogue of Somatic Mutations in Cancer

628   Table 2: Mean and standard deviation of expressed rare (Not in ExAC or <0.5% allele frequency), low-frequency (0.5-1% allele frequency) and common
629   (>1% allele frequency) alleles in 39 *KMT2A*-R patient samples compared to ultra-mutated (P337) sample and its constituent expressed germline alleles and
630   somatic mutations.

	Not in ExAC	<0.5% AF[1] in ExAC	0.5-1% AF in ExAC	>1% AF in ExAC
KMT2A-R iALL	42.3±10.4	212.1±32.6	76.8±14.7	7722.3±266.0
P337 expressed	38	174	89	7590
P337 Germline	17	88	47	6235
P337 Somatic	21	86	42	1355

[1]AF: allele frequency; [2]ExAC: Exome aggregation consortium

Figure Legends:

Figure 1: Characterization of germline and somatic point mutations identifies an ultra-

mutated infant leukemia case. (a) Overlap of variants identified from diagnostic leukemia

specimen by RNA-seq and whole exome sequencing (WES). (b) Overlap of variants called in

diagnostic RNA-seq and remission bone marrow WES. (c) Overlap of variants called in

diagnostic and remission WES. Barplots colored showing overlap binned by derived allele

frequency as follows: present with AF<0.5% in ExAC (green), present with AF 0.5-1% in

ExAC (yellow), present with AF>1% in ExAC (purple), absent from the ExAC (orange), non-

overlapping variants are colored to show sites with coverage <20 reads (grey) or coverage

>20 reads (black) in the comparator. (d) Scatterplots of diagnostic leukemia (x-axis) versus

remission (y-axis) variant allele fraction (VAF) from WES for coding variants detected in

diagnostic leukemia RNA-seq.

Figure 2: Clonal analysis reveals somatic signatures operative during infant leukemia

ultra-mutation. Histograms showing the VAF density distribution of somatic mutations (left,

x-axis: dark shaded bars) or LOH-sites (right; x-axis: light shaded bars) in diagnostic

leukemia WES from canonical cases (a) and the ultra-mutated case (b). (c) Illustration of cell

lines derived from ultra-mutated specimen and histograms displaying the VAF density

distribution of somatic mutations (left, x-axis: dark shaded bars) or LOH-sites (right; x-axis:

light shaded bars) in cell lines (PER-784A: red; PER-826A: green). (d) Venn diagram

displaying the overlap in somatic mutations (left) and LOH-sites (right) identified by

VarScan2 comparing the primary diagnostic specimen and derived cell lines (PER-784A: red;

PER-826A: green) to the primary remission sample. (e) Somatic mutation and (f) LOH

coding sites showing total read coverage (left y-axis) and VAF (right y-axis) from primary

specimens (left panels) and cell lines (right panels) ordered by decreasing VAF in the

diagnostic sample (x-axis).

Figure 3: Germline-like somatic signatures of ultra-mutations. (a-d) Histograms showing

the number (a-b) or frequency (c-d) of events overlapping the spectrum of ExAC derived

allele frequency bins with 5% increments. (a) Somatic mutations called by VarScan2 were

plotted grouping homozygous and heterozygous events separately. (b) Loss-of-heterozygosity

sites   called   by   VarScan2   were   plotted   grouping   reference→alternative   and

alterantive→reference sites separately. All quality filtered coding variants from each of the

canonical patients or P337 ultra-mutated specimen called by GATK grouped by homozygous

and heterozygous events from (c) remission sample or (d) diagnostic sample. Normalised

fraction of variants within unique trinucleotide context (ordered alphabetically on the x-axis)

for VarScan calls (left panel) as shown for (a-b), or remission variants (right panel) as shown

for   (c).   (f)   Somatic   signatures   identified   by   *deconstrucSigs*   for   categories   of

mutations/variants shown in (e) showing their relative proportions (bar plot) and pair-wise

Pearson's correlation coefficients (heatmap), with both plots ordered by unsupervised

hierarchical clustering.

Figure 4: Ultra-mutations target rare and common human alleles. (a) Number of variants

(y-axis) called from RNA-seq reads from St Jude and Perth cohorts (open circles) are plotted

with the total variants detected in the ultra-mutated sample (shaded circles), germline variants

(shaded triangle pointing up) and somatic mutations (shaded triangle pointing down) binned

according to the allele count in ExAC database (red: not in ExAC; green $< 0.5\%$; yellow: 0.5-

1% and blue $> 0.5\%$). (b) Bar plot displaying fraction of variants within rare, low-frequency

and common allele frequency bins (as described for b) ordered by increasing proportion of

common alleles (i.e. derived AF>1%) and excluding outliers (SJINF018, SJINF021,

SJINF010 and SJINF008). Variants extracted from RNA-seq data are shown in the left panel.

Variants extracted from diagnostic WES samples and for ultra-mutation germline, somatic

mutations and LOH-sites defined by VarScan2 are shown in the right panel. (c) Crystal

structure of the MutS lesion recognition complex (PDB CODE 2O8D[15]) MSH2:MSH6

mismatch repair proteins (MSH2: green, MSH6: blue) bound to ADP and G:T mismatch

DNA (orange) showing the location of the p.E423K amino acid change. (d) Crystal structure

of Pol k (green and blue representing different molecules) in complex with a bulky DNA

adduct (orange) (PDB CODE 4U6P[19]) showing locations of the p.R298H amino acid change.

Figure 5: Global transcriptional concordance of ultra-mutated infant leukemia. Multi-

dimensional scaling of RNA-seq count data from acute leukemias and normal bone marrow

progenitor populations displaying translocation partner gene by color and blood/leukemia

type by symbol shape according to the legend. Labels show replicate measurements from the

ultra-mutated diagnostic specimen, P337.

Supplementary Fig. S1. Copy-number alterations and loss-of-heterozygosity overlap in

typical and ultra-mutated *KMT2A*-R iALL. (a) Genomic copy number ratios of the ultra-

mutated sample comparing diagnostic and remission WES data.   (b) Proportion of reads

mapping to different copy-number bins (up to 4). (c) Box and whisker plots (median with

upper and lower quartiles) displaying the nearest distance of a variant/mutation to a

homozygous variant in the diagnostic WES data for canonical samples (left) compared to the

ultra-mutated sample (right) for VarScan2 germline, somatic and LOH SNV calls.

Supplementary Table S1: Read evidence of RNA-seq variants that were not detected in

matched WES data.

Supplementary Table S2: Summary of samples

Supplementary Table S3: Comparison of RNA-seq reads from diagnosis sample with WES

reads.

Supplementary Table S4: Comparison of RNA-seq reads from diagnosis sample with WES

reads from diagnosis, remission and derived cell lines.

Supplementary Table S5: Annotation of somatic mutations and indels from ultra-mutated

patient P337.

Supplementary Table S6: Summary of LOH-sites called in primary P337 and derived cell

lines.

Supplementary Table S7: Summary of somatic SNVs called in primary P337 and derived

cell lines.

FIGURE 1

FIGURE 2

FIGURE 4

FIGURE 5

Blood progenitor
MLL/KMT2A-Germline
MLL/KMT2A-MLLT1
MLL/KMT2A-AFF1
MLL/KMT2A-MLLT3
MLL/KMT2A-MLLT4
MLL/KMT2A-MLLT10
MLL/KMT2A-EPS15
MLL/KMT2A-USP2
MLL/KMT2A-CASC5
MLL/KMT2A-MLLT6
MLL/KMT2A-ELL
MLL/KMT2A-ENL
MLL/KMT2A-SEPT9
MLL/KMT2A-ENAH
MLL/KMT2A-GAS7

○ Diagnostic infant ALL
□ Relapse infant ALL
△ Childhood ALL
◇ Childhood AML
▽ Leucegene AML

SUPPLEMENTARY FIGURE S1