Adaptation decorrelates shape representations

Marcelo G. Mattar[1*], Maria Olkkonen[2,3*], Russell A. Epstein[1], & Geoffrey K. Aguirre[4]

1. Department of Psychology, University of Pennsylvania, Philadelphia, PA 19104, USA

2. Department of Psychology, Durham University, Durham, DH1 3LE, United Kingdom

3. Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, 00100, Finland

4. Department of Neurology, University of Pennsylvania, Philadelphia, PA 19104, USA

* These authors contributed equally to this work

Abstract

Perception and neural responses are modulated by sensory history. Visual adaptation, an example of such an effect, has been hypothesized to improve stimulus discrimination by decorrelating responses across a set of neural units. While a central theoretical model, behavioral and neural evidence for this theory is limited and inconclusive. Here, we use a parametric 3D shape-space to test whether adaptation decorrelates shape representations in humans. In a behavioral experiment with 20 participants, we find that adaptation to a shape class improves discrimination of subsequently presented stimuli with similar features. In a BOLD fMRI experiment with 10 participants we observe that adaptation to a shape class decorrelates the multivariate representations of subsequently presented stimuli with similar features in object-selective cortex. These results support the long-standing proposal that adaptation improves perceptual discrimination and decorrelates neural representations, offering insights into potential underlying mechanisms.

Keywords: adaptation, decorrelation, repetition suppression, shape perception, psychophysics, multi-voxel pattern analysis

Introduction

Neural responses to visual stimuli are modulated by the preceding temporal context, a phenomenon know as "adaptation" (Enroth-Cugell and Shapley, 1973; Ohzawa et al., 1982; Dragoi et al., 2000; Kohn and Movshon, 2004; Engel, 2005; Krekelberg et al., 2006; Kusunoki et al., 2006; Schwartz et al., 2007; Wark et al., 2009). Adaptation has been proposed to serve as a mechanism of efficient sensory coding, tuning the response properties of neural populations to the current sensory environment (Barlow and Földiák, 1989; Clifford et al., 2007; Kohn, 2007). In particular, adaptation may reduce the correlation between neural activity patterns corresponding to frequently encountered stimuli (Barlow and Földiák, 1989; Barlow, 1990), either by shifting neuronal tuning curves away from one another or by narrowing neuronal selectivity (Clifford et al., 2000; Kohn, 2007; Grill-Spector et al., 2006; Seriès et al., 2009; Cortes et al., 2012). Empirical support for this hypothesis has been found in some animal studies. Neurophysiological recordings from monkey primary visual cortex show that adaptation to stimulus orientation decorrelates neural responses (Mueller et al., 1999; Gutnisky and Dragoi, 2008), and recordings from cat primary visual cortex show that adaptation promotes population homeostasis (Benucci et al., 2013).

Adaptation also alters perception (Webster, 2015). Psychophysical data in humans collected with low-level stimuli (e.g., color, motion direction, speed, orientation) demonstrate an improvement in perceptual discrimination of stimulus features following adaptation to stimuli with similar features (Regan and Beverley 1985; Krauskopf et al. 1992; Phinney et al. 1997; Krekelberg et al. 2006; Giesel et al. 2009; although see Barlow et al. 1976). For more complex stimuli, however, results are mixed. Adaptation to faces, for example, does not reliably improve discrimination of face-related attributes (Oruç et al., 2011; Keefe et al., 2013; Rhodes et al., 2007; Ng et al., 2008).

Theoretical work suggest possible links between the neural decorrelation effects of adaptation and the improvements in perceptual discrimination that follow (Wainwright, 1999; Seriès et al., 2009), although direct experimental evidence of simultaneous neural response and perceptual effects is lacking. Here, we tested the predictions of the efficient coding model in both the psychophysical performance of humans and in the corresponding multivariate patterns of BOLD fMRI response. Specifically, we tested whether brief exposure to a set of synthetic visual shapes drawn from a common prototype would enhance the discriminability of subsequently presented stimuli with similar features. The use of complex 3D shapes allowed us to measure the distributed pattern of voxel responses within higher-level visual cortex. To anticipate our results, we observed that perceptual discrimination of 3D shapes is enhanced after adaptation, and that this effect is accompanied by a decorrelation of the multi-voxel fMRI patterns evoked by these stimuli.

Figure 1: Experimental design. *a)* We sampled 3D shapes from a 24-dimensional, radial frequency stimulus space in which the dimensions were defined by the frequencies, amplitudes, and phases of 8 sinusoids. Two visually distinct shapes from this space were chosen as prototypes "A" and "B". For each prototype, 50 adaptor stimuli were generated in the multidimensional space by jittering the orientation and amplitude of the 8 sinusoidal components, producing two distinct shape classes. *b)* In the beginning of each experimental run (for both psychophysics and fMRI experiments), participants were presented with a series of shapes from one of the two shape classes for 60 seconds at a rate of 5 Hz (*adapt* phase). Trial presentations began immediately following the adaptation period. All trials began with a *top-up* adaptation period of 4 seconds at 5 Hz, followed by trial events that differed for the psychophysics and fMRI experiments (panels *c,d*). In all experiments, adaptation condition (A or B) was blocked in counterbalanced order across participants. *c)* We investigated the effect of adaptation on perceptual discrimination performance with a delayed match-to-sample (DMTS) task. "Morph" stimuli were generated along the vector in the shape space that connected the two prototype shapes. In the DMTS task, participants first saw a morph and, after a brief delay, the same morph and a prototype. The participant was asked to report which of the two shapes in the second interval matched the first shape (i.e., which of the two shapes was a morph). To avoid pixel-based matches, each morph and prototype shape was displayed as a static image in one of two possible rotations. Trials using A and B as prototypes were interleaved in a block, and separate 1-up, 3-down staircases controlled the morph value for each prototype. *d)* We used fMRI to investigate the effect of adaptation on neural pattern similarity. Two distinct probe stimuli were created for each shape class (A1, A2 and B1, B2) along axes passing through the prototype stimuli. After a top-up adaptation phase and a variable interstimulus interval, participants saw one of the four probe stimuli and were asked to report whether the probe was larger or smaller than the adaptors in the top-up phase. We measured the neural response to each of the four probe stimuli and calculated the multivariate pattern similarity between the two probe shapes (e.g. A1 and A2) after adaptation to the same shape class (A) compared to adaptation to the different shape class (B).

Results

Adaptation improves perceptual shape discriminability

To test the effect of adaptation on discrimination of high-level visual stimuli, we created two sets of computer-generated 3-dimensional shapes (Figure 1a). Each set of shapes was based upon a different "prototype" shape, and all items in a given stimulus set were more similar to each other than they were to items in the other stimulus set. We then asked if perceptual adaptation to stimuli from one of the sets would improve subsequent shape discrimination thresholds for items from that same set.

Twenty participants performed multiple runs of a delayed match-to-sample task. Each of two experimental runs started with an *adaptation* phase during which the participants viewed a rapid, serial presentation of shapes from one of the sets for 60 seconds (Figure 1b). After the adaptation period, participants performed a series of discrimination trials. Each trial started with a 4 s of *top-up* adaptation, followed by the presentation a single sample stimulus and, after a brief delay, the same sample stimulus shown simultaneously with one of the two prototype stimuli. Participants indicated with a button press which of two shapes displayed in the second interval matched the shape in the first interval (Figure 1c). A match-to-sample task was convenient for measuring discrimination thresholds because it does not require verbalizing the dimension on which participants are to make the match – they merely have to pick the more similar shape, instead of having to pick the (e.g.) spikier one (see Macmillan and Creelman, 2004). To generate a series of samples for each prototype on an arbitrary axis in shape space, we morphed the two prototypes with fine spacing. The similarity between the sample and the prototype stimulus shown on each trial was adjusted based on a 1-up, 3-down staircase procedure. We estimated the discrimination thresholds for each prototype from the last five reversals of the staircase and averaged the discrimination thresholds within subject across the two experimental runs. Trials in which the sample and prototype stimuli were drawn from the same stimulus set as the adaptors were interleaved with trials in which the sample and prototype stimuli were drawn from the other stimulus set.

The efficient coding hypothesis predicts a lower discrimination threshold (i.e. better performance) for stimuli from the adapted stimulus set as compared to the unadapted stimulus set. This result was found for all twenty subjects (Figure 2a). We tested this effect in a three-way repeated measures ANOVA with adaptation condition (adapted/unadapted), prototype (A/B), and run (1/2) as factors. Adaptation improved discrimination thresholds, evidenced by a main effect of adaptation condition ($F(1, 77) = 58.5$, $p < 0.001$; Figure 2a). In post-hoc tests, we also observed a main effect of prototype stimulus on thresholds ($F(1, 77) = 14.4$, $p < 0.001$), suggesting that participants found it easier to discriminate stimuli within shape space "A" as compared to "B" (Figure 2b). The adaptation effect was present for both shape sets, however: mean decrease in threshold was 4.6 morph units for prototype A (one-sided t-test $t(19) = 4.8, p < 0.001$) and 6.4 units for prototype

B $(t(19) = 8.8, p < 0.001)$. Finally, there was no significant difference in the adaptation effect for the two runs $(F(1, 77) = 0.33, p = 0.57$; Figure 2c).

Figure 2: Perceptual shape discriminability. *a)* Shape discrimination thresholds are plotted for each participant after adapting and testing for the same shape class (x-axis) versus adapting and testing to different shape classes (y-axis). Each data point is the average over two repetitions and over both prototype shapes. All data points lie above the diagonal, indicating that shape discrimination thresholds were lower after adapting to the same shape class for every subject. *b)* Thresholds shown separately for the two prototype shapes, averaged over runs. The two distributions are partially non-overlapping, suggesting that shape space "A" was easier to discriminate. *c)* Thresholds shown separately for the two runs, averaged over shape.

Adaptation improves neural pattern discriminability

We next asked whether adaptation enhances the discriminability of neural patterns evoked in object-selective regions of the brain. We tested this hypothesis in 10 new participants using BOLD fMRI to measure the pattern of neural response to probe stimuli from each shape class (Figure 1d). The same procedure as in the psychophysics study was followed for initial adaptation on each experimental run and the top-up adaptation at the beginning of each trial. Following top-up adaptation, participants were presented with single probe stimuli derived either from the same shape class as the adaptors or the other shape class. The ordering of the four probe stimuli across trials was counterbalanced, and a 10% size modulation was applied to each probe stimulus. Participants performed a cover task in which they indicated with a button press whether the probe was smaller or larger than the

stimuli in the adapting phase. Four fMRI acqusitions were conducted using each of the shape classes as the adapting stimulus.

We focus our initial analysis upon the left lateral occipital cortex (left LO) (Grill-Spector et al., 2001), as this lateralized region has been found to exhibit larger adaptation effects for repeated presentations of different exemplars of the same shape category (Koutstaal et al., 2001; Simons et al., 2003). We used the data from an independent functional localizer scan collected for each participant to identify the 100-vertices on the cortical surface within a left LO parcel that had the greatest differential response to shapes vs. scrambled shapes. In each participant, we measured the average amplitude of evoked response for each of the four probe stimuli at each of the 100 vertices, and then obtained the pairwise, Pearson correlation between the patterns of response to each of the four probes (Figure 3a).

We tested this effect in a three-way repeated measures ANOVA with adaptation condition (adapted/unadapted), prototype (A/B), and run (1/2) as factors.

The efficient coding hypothesis predicts that neural representations of adapted stimuli become less correlated with one another, in keeping with an increased ability to discriminate their identity. In our experiment, this effect would be manifest as a decrease in the correlation between the neural patterns evoked by a pair of probe stimuli (e.g. A1 and A2) when those stimuli are preceded by a matching adapting phase. We find evidence for this effect in our data (Figure 3b). Using a repeated measures ANOVA, we examine the influence of adaptation condition (adapted/unadapted) and prototype (A/B) on the correlation between the two probe shapes. We observed that the main effect of adaptation condition was significant ($F(1,9) = 14.6895, p = 0.004$), indicating that pairwise similarity after adapting to the same space was lower than after adapting to the other space. Neither the main effect of prototype ($F(1,9) = 0.9211, p = 0.36$) nor the interaction term were significant ($F(1,9) = 1.6976, p = 0.22$), suggesting that the decorrelation effect was not significantly different between the two prototypes. In post-hoc tests we examined this effect in the two stimulus spaces separately. We find that the correlation between the "B" probe stimuli (B1 and B2) was lower when participants were adapted to stimulus set "B" than when they were adapted to stimulus set "A" ($t(9) = 3.83$, $p = 0.004$), although the complementary test with the "A" shape class did not show a significant difference ($t(9) = 1.59$, $p = 0.15$).

We examined the consistency of the effect across subjects. We measured for each subject the Pearson correlation of the neural pattern of response for a pair of probe stimuli when those stimuli were from the same class as the adaptors, and when they were from the unadapted shape class (Figure 3c). All ten participants had a lower correlation between probe stimuli in the adapted condition (paired t-test on Fisher-Z transformed correlation coefficients: $t(9) = 5.81$, $p < 0.001$).

Next, we examined the cortical extent of these effects. We performed a searchlight analysis (Kriegeskorte et al., 2006), calculating within each volumetric searchlight (5 mm radius) the magnitude of the pattern decorrelation induced by adaptation (Figure 3d). A broad swath of posterior occipital-temporal cortex evidenced pattern decorrelation, with the largest effect sizes observed within the occipital pole, lateral occipital cortex, and the

posterior region of fusiform gyrus.

Taken together, the behavioral and fMRI results support the efficient coding hypothesis: adaptation to a given shape class produces both a behavioral improvement in the discriminability of shapes from the adapted class, as well as an increased separability of their multi-voxel patterns in object-selective cortical regions and other visual areas.

Mechanisms of decorrelation

As adaptation is known to reduce the amplitude of neural response, one possible mechanism for our findings is a reduction in stimulus-evoked responses. If this reduction in response occurs in the setting of independent, unchanged measurement noise, the reduced correlation we observe in the patterns evoked by adapted stimuli may be the product only of a lower signal-to-noise ratio. Consistent with this mechanism, we find that the evoked BOLD fMRI signal amplitude is smaller in the adapted as compared to the unadapted condition within left LO (percent signal change in adapted condition: $1.30\% \pm 0.16$ vs. unadapted: $1.45\% \pm 0.16$; paired t-test: $t(9) = 2.78, p = 0.0213$). We find, however, that this reduction in amplitude varies markedly across vertices. We calculated the magnitude of suppression for each vertex as a scaling factor between adapted and unadapted responses, and then identified sets of vertices that had different degrees of suppression. The 50 vertices with lowest values for this index had a mean suppression value of 0.25 ± 0.32 (i.e., the BOLD fMRI signal evoked by the stimuli was reduced by 75% in the adapted condition). In contrast, the 50 vertices with the largest values for this index actually demonstrated response *enhancement* in the adaptation condition (1.36 ± 0.14, or an increase by 36% of response amplitude in the adaptation condition). If a reduction in response amplitude alone accounts for the decorrelation of patterns that we observe in the adaptation condition, then a decorrelation effect should not be present in the sub-set of vertices with response enhancement. In contrast to this account, a significant decorrelation effect was still found (two-sample t-test on Fisher-Z transformed correlation coefficients: $t(9) = 2.85, p = 0.0095$). This result indicates that response suppression alone does not account for the observed pattern decorrelation effects.

Finally, we examined if the effect of adaptation upon neural responses in our data are better described as *sharpening* or *fatigue*(Grill-Spector et al., 2006), as the former could account for a decorrelation of neural patterns following adaptation. In the sharpening model, suppression is largest in neurons that that are weakly tuned to the adaptor, as opposed to the fatigue model which describes either a uniform scaling of responses in all neurons, or a larger suppression in neurons that are strongly tuned to the adaptor. These models can therefore be distinguished by examining the degree of response suppression that occurs within neurons that are either strongly or weakly responsive to the stimulus.

We measured in our data the suppressive effect of adaptation upon responses in those vertices that had either weak or strong responses to the stimuli. For the 50% of vertices that responded most weakly to unadapted stimuli, we observed that adapted responses were

Figure 3: Neural shape discriminability *a)* Schematic of the fMRI decorrelation analysis. We measured responses in left LO to the a pair of probe shapes after adaptation to either stimulus space "A" or "B". We then computed the similarity of neural pattern response to the pairs of probe shapes, and compared the resulting Pearson's correlation for the probe shapes that were either adapted or unadapted. A decorrelation effect was defined as the difference between these two correlation values. The logic of the analysis is illustrated for the test in which shape class "B" served as the probe. *b)* Pattern dissimilarity for for probe stimuli from the "A" (left) and "B" (right) shape spaces. Patterns were more distinct (less correlated) for probe shape pairs after undergoing adaptation to the same shape space. Error bars are standard error of the mean across participants. *c)* Correlation between multi-voxel patterns for each subject, comparing the case in which stimuli were from the *same* space as the adaptors (x-axis), to when they were from the *different* space (y-axis). For every subject, patterns from adapted pairs of stimuli were less correlated than patterns from unadapted stimuli. Each data point represents one participant and error bars represent variability (SEM) across scan acquisitions within subject. *d)* A whole-brain searchlight analysis presents the mean, across-subject magnitude of the decorrelation effect across the cortex.

scaled by approximately 0.74 ± 0.073 in relation to the unadapted responses. In the 50% of vertices that responded most strongly to unadapted stimuli, we observed a scaling factor of 0.91 ± 0.073 (one-sided, paired t-test: $t(9) = 2.10, p = 0.032$). This suggests that weakly tuned vertices undergo a larger response suppression than strongly tuned vertices, in line with the sharpening model.

Discussion

We used behavioral and neural data to test the hypothesis that adaptation decorrelates stimulus representations. In a psychophysical experiment, we found that adaptation improved the perceptual discrimination of similar shapes. In a BOLD fMRI experiment, we found that adaptation enhanced the distinctiveness of neural responses to perceptually similar stimuli. While the fMRI discrimination enhancement effect was concentrated in object-selective cortex, a searchlight analysis demonstrated similar results in nearby regions of the occipital and ventral temporal lobes. Our results offer three novel contributions: (i) we provide evidence for enhanced perceptual discrimination of high-level stimuli (3D shapes) following adaptation, thus clarifying earlier findings whose results were mixed (Rhodes et al., 2007; Ng et al., 2008; Oruç et al., 2011; Keefe et al., 2013); (ii) we demonstrate a decorrelation of neural representations in human observers undergoing visual adaptation; (iii) we offer joint behavioral and neural evidence using a similar experimental paradigm, thus offering a link between the neural effect and its behavioral consequences.

Our analyses provide some insight into the mechanism of the neural decorrelation effect. We find that a uniform reduction in neural response (in the face of an unchanged level of measurement noise) cannot completely account for our results. Decorrelation could instead be driven by a non-homogenous scaling of responses, such as a larger response reduction in weakly responsive units. Such an effect would serve to maintain population homeostasis in sensory cortices (Benucci et al., 2013), and has found empirical support as a "sharpening" effect in fMRI studies of adaptation (Grill-Spector et al., 2006).

Computationally, a sharpening of neural tuning curves have been demonstrated to produce both perceptual biases and a decrease in perceptual discrimination thresholds (Seriès et al., 2009; Cortes et al., 2012). While our study demonstrates perceptual improvements in threshold discrimination and neural changes in representation using the same adaptation procedure, we note that our findings do not directly relate these phenomena. An ideal model would provide a quantitative mapping between neural and perceptual effects on a trial-by-trial basis. A challenge to such an effort is that measurement of the neural pattern of response is complicated in the face of a perceptual task that requires the subject to explicitly process the similarity of presented stimuli. A complete model would also account for the cortical extent over which this decorrelation effect is observed. We find that visual cortex broadly demonstrates the decorrelation effect, although it is of greater strength in object-responsive areas that have been previously shown to exhibit coarse spatial coding

for object shape (Drucker and Aguirre, 2009).

Our paradigm bears some resemblance to category learning, especially as discussed in prototype theory (e.g. Ashby and Maddox, 2005). However, instead of learning to discriminate A from B, our subjects had to discriminate *within* category A or B, which is a question not typically addressed in category learning studies. Our study also shares similarities to perceptual learning paradigms, in which participants learn to discriminate between two initially indistinguishable stimuli after several hours of practice (Goldstone and Gibson, 1962). While the phenomena of category and perceptual learning manifest over much longer time scales than the seconds-to-minutes of adaptation studied here, we note that adaptation and long time-scale perceptual representations may share underlying neural mechanisms (Mattar et al., 2016).

In summary, we find evidence in support of adaptation as a mechanism for representational efficiency. In line with theoretical predictions, decorrelated neural representations accompany an improvement in perceptual discrimination.

Methods

Experiment 1: Psychophysics

Participants

20 participants (15 women, ages 18-51) with normal or corrected-to-normal vision, recruited through the University of Pennsylvania participant pool took part in the study. Participants were compensated $10/hour. The experimental protocol was approved by the University of Pennsylvania Institutional Review Board.

Stimulus design and experimental protocol

To measure shape discrimination thresholds, we constructed a parametric shape space with ShapeToolbox (URL: http//github.com/saarela/ShapeToolbox). ShapeToolbox generates three-dimensional, radial frequency patterns by modulating basis shapes such as spheres with an arbitrary combination of sinusoidal modulations in different frequencies, phases, amplitudes, and orientations. Specifically, we generated shapes by modulating a sphere with different combinations of six different sinusoids. Each sinusoid had four parameters: component frequency, amplitude, phase, and orientation, resulting in a 24-dimensional shape space. We then chose two shapes as prototype shapes, which were sufficiently different that they were intuitively perceived as members of two different shape categories (see Fig 1a). The same two prototype shapes (prototype shape A and prototype shape B) were used for all participants.

Participants viewed the stimuli in a dim room from a distance of approximately 57 cm on a Macbook pro laptop (15" retina display, resolution 2889 × 1800). Discrimination thresholds for the two prototype shapes were measured in interleaved trials embedded in

three different types of block. In the baseline blocks, discrimination thresholds were measured without adaptation to shape. In the "adapt A" blocks, a rapid stream of adaptors that were similar to prototype shape A were shown in the beginning of the block for 60 seconds, and correspondingly for "adapt B" blocks. In the adapting phase, each adaptor was presented for 150 ms with a 50 ms ISI, or at a rate of 5 Hz (Figure 1b). The adaptors for each prototype shape were created by jittering the orientation and amplitude of the sinusoidal modulation of the particular prototype while maintaining the Euclidean distances in the multidimensional shape space between the prototype and the adaptors, thus creating a region in shape space around the prototype shape (Figure 1a). The variability in parameter space of these adaptors (50 total) was held constant and smaller than the variability between the two stimulus categories. To ensure focus on the stimuli during the adaptation periods, a fixation cross in the middle of the display randomly changed color from black to gray, and participants were asked to indicate the color changes with a key press. The end of the adaptation period was indicated to the participant with the text "start experiment". Participants first practiced the baseline discrimination task, after which they ran one baseline run. They then practiced the adaptation task, and ran each adaptation block twice in ABAB or BABA order (counterbalanced across participants).

Shape discrimination was measured with a delayed match-to-sample procedure. On each trial, participants saw a test stimulus in the center of the screen for 200 ms, and then after a delay of 250 ms the same test stimulus was paired with either prototype A or B and displayed until response. Participants were asked to indicate which of the two shapes in the second interval best matched the first shape. To avoid pixel matches, the tests and the prototype were shown in one of two possible vertical rotations. This ensured that the participant had to extract the 3D shape of the stimuli to make the match.

Test stimuli were generated by morphing the two prototype shapes at equal intervals along an 80-step morph continuum (Figure 1c). The test shape on each trial was selected according to an adaptive staircase procedure that was separately run for the two prototypes, with stimuli in each case drawn from the 40 steps closest to the prototype. Each of the two staircases converged on the 79% discrimination threshold (1-up ,3-down rule). The staircases ended after eight reversals. The discrimination threshold was taken as the average of the last five reversals. In adaptation blocks, a top-up adaptation period at 5 Hz for 4 seconds preceded each trial. To characterize the effect of adaptation on discrimination, we compared thresholds for each prototype shape after adapting to spaces A and B.

Experiment 2: fMRI

Participants

10 healthy human subjects (6 females, ages 21-37) with normal or corrected-to-normal vision were recruited through the University of Pennsylvania participant pool. Participants were compensated $20/hour. The experimental protocol was approved by the University

of Pennsylvania Institutional Review Board. None of the 10 subjects participated in the psychophysics experiment.

Stimulus design and experimental protocol

The same prototype shapes and pool of adaptors from the behavioral experiment were used in the fMRI experiment. We additionally generated two "probe" stimuli for each shape class as follows. For each prototype, we defined a random direction in our parametric shape space and selected two shapes along this dimension, equally distant from the prototype in the positive and negative direction (Figure 1d). While the random direction was not constrained to be orthogonal to the shape space vector that connects the two prototype shapes, the high-dimensionality of the shape space essentially guaranteed this property. The same probe stimuli were used for all subjects and across all runs.

Each experimental block (corresponding to one scan acquisition run) began with a 60 second pre-adaptation phase during which a stream of adaptors from one of the shape class was presented at a rate of 5Hz in random order. Participants then completed 16 trials, each starting with a top-up adaptation phase for 4 seconds at the same 5 Hz rate, followed by a jittered ISI (1-4 seconds) and a test stimulus selected from the four possible probe (A_1, A_2, B_1, B_2), presented for 1 second. We used a "cover" task on each trial to ensure attention to the stimuli. A size modulation of 10% (larger of smaller) was applied to the probe stimuli, and participants were instructed to select with a left/right button press whether the stimulus was smaller/larger than the stimuli in the top-up phase. Trial onsets were separated by 14 seconds, so the interval between the offset of the probe stimulus and the onset of the top-up phase of the next trial was 5-8 seconds. The size modulation of each shape and the order of the test stimuli within each run were counterbalanced (4 instances of each test stimulus per run, each presented twice in smaller/larger versions). Participants completed a total of eight scans, four with each shape class used in the adaptation phases, in counterbalanced order. Stimuli occupied 10 degrees of visual angle ($\pm$ 10% for the tests) and were presented on a gray background.

In addition to the main experimental scans, participants completed one or two functional localizer scans consisting of 16-second blocks of faces, objects, scenes, and scrambled objects. Images were presented for 800 ms with a 200 ms ISI, and participants performed a one-back task on image repetition.

Data acquisition and preprocessing

Magnetic resonance images were obtained at the Hospital of the University of Pennsylvania using a 3.0 T Siemens Trio MRI scanner equipped with a 32-channel head coil. Stimuli were displayed on a Sanyo-PLC-XT35 LCD projector via a mirror mounted on the head coil. The viewing area of the display was 50.5×38 cm or 23×17 degrees of visual angle. Stimulus size was approximately 10×10 degrees of visual angle. T1-weighted structural images of the

whole brain were acquired on the first scan session using a three-dimensional magnetization-prepared rapid acquisition gradient echo pulse sequence (repetition time (TR) 1620 ms; echo time (TE) 3.09 ms; inversion time 950 ms; voxel size 1 mm × 1 mm × 1 mm; matrix size 190 × 263 × 165). A field map was also acquired at each scan session (TR 1200 ms; TE1 4.06 ms; TE2 6.52 ms; flip angle 60°voxel size 3.4 mm × 3.4 mm × 4.0 mm; field of view 220 mm; matrix size 64 × 64 × 52) to correct geometric distortion caused by magnetic field inhomogeneity. Functional data were acquired with T2*-weighted images sensitive to blood oxygenation level-dependent contrasts using a slice accelerated multiband echo planar pulse sequence (TR 2,000 ms; TE 25 ms; flip angle 60°; voxel size 1.5 mm × 1.5 mm × 1.5 mm; field of view 192 mm; matrix size 128 × 128 × 80, acceleration factor 4). 105 volumes were acquired for the functional localizer(s) scans, and 142 volumes were acquired for each experimental scan.

Cortical reconstruction and volumetric segmentation of the structural data was performed with the Freesurfer image analysis suite (Dale et al., 1999). Boundary-Based Registration between structural and mean functional image was performed with Freesurfer bbregister (Greve and Fischl, 2009). Preprocessing of the fMRI data was carried out using FEAT (FMRI Expert Analysis Tool) Version 6.00, part of FSL (FMRIB's Software Library, www.fmrib.ox.ac.uk/fsl). The following pre-statistics processing was applied: EPI distortion correction using FUGUE (M. Jenkinson, 2004); motion correction using MCFLIRT (Jenkinson et al., 2002); slice-timing correction using Fourier-space time series phase-shifting; non-brain removal using BET (Smith, 2002); grand-mean intensity normalization of the entire 4D dataset by a single multiplicative factor; highpass temporal filtering (Gaussian-weighted least-squares straight line fitting, with sigma=50.0s). Spatial smoothing was performed with a 5mm FWHM kernel using FSL's `fslmaths`.

General Linear Model

Statistical analyses were performed upon the time-series data from each functional run using a General Linear Model in FSL. Analysis of the main experimental scans included one binary covariate for the pre-adaptation phase, one binary covariate for the 16 top-up phases, and one binary covariate for each of the four test stimuli. The resulting statistical maps were projected to surface space using Freesurfer's `mri_vol2surf`.

Analysis of the functional localizer scans included one binary covariate for each stimulus type (faces, objects, scenes, and scrambled objects) and one binary covariate for button-press. Parameter estimates were averaged across scans for those participants that completed two functional localizers. The resulting statistical maps were projected to surface space using Freesurfer's `mri_vol2surf`.

ROI definition

We focused our analyses on a well-characterized object-selective area of the human brain, the Lateral Occipital Complex (LOC; Grill-Spector et al., 2001). The human LOC is divided into a ventral part in the posterior fusiform gyrus (pFus) and a region in the lateral occipital cortex (LO). Previous studies suggest the existence of a coarse spatial coding of shape features in LO and a more focused coding of the entire shape space within pFus (Drucker and Aguirre, 2009). Studies focusing on laterality differences in LOC have reported larger adaptation effects for repeated presentations of different exemplars of the same object category in the left hemisphere (Koutstaal et al., 2001; Simons et al., 2003), and greater sensitivity to changes in viewpoint in the right hemisphere (Vuilleumier et al., 2002). We therefore focused on the left lateral occipital cortex (left LO) as our region of interest due to its greater sensitivity to shape similarity at the multi-voxel scale, and greater invariance to category and size information.

We defined left LO for each participant as the 100 surface vertices on the cortical surface that responded most strongly in the functional localizer scan to the contrast of objects vs. scrambled objects within a larger group-defined left LO parcel warped to the participant's own surface space (Figure 3a). This method ensures sensitivity to the between-subject variability of the spatial location of this ROI (Julian et al., 2012).

Multi-voxel pattern analyses and searchlight

ROI pattern analyses were performed on the parameter estimates within each ROI. Whole-brain pattern analyses were performed using a volumetric searchlight procedure (Kriegeskorte et al., 2006), with analyses performed on the parameter estimates within every 5mm sphere in the brain. We considered separately the multi-voxel patterns evoked by adapted shapes (when the test stimulus was from the same space as the adaptors) and the multi-voxel patterns evoked by non-adapted shapes (when the test stimulus was from the other, unadapted space). For each ROI or searchlight, the correlation between the two patterns evoked by the adapted shapes and the correlation between the two patterns evoked by the non-adapted shapes were calculated for each experimental run, their Fisher z-transformations averaged across scans, and the averaged values compared (Figure 2a,b). With the goal of visualizing the magnitude of the effect across the cortical surface, we used a threshold based on effect size (difference in correlation equal to 0.05) rather than statistical significance.

Mechanisms of decorrelation

We calculated the average amplitude of evoked response for each vertex in both adapted and unadapted states, averaging the patterns evoked by each stimulus across runs. To examine the magnitude of a scaling effect across vertices in our region of interest, we calculated the best linear fit by least squares estimation between unadapted and adapted responses,

where a slope of less than one indicates a divisive effect. We then assessed the existence of a differential suppressive effect across vertices by performing the linear fit separately for the 50% vertices that responded on average most weakly to unadapted stimuli (weakly tuned units), and for the 50% vertices that responded on average most strongly to unadapted stimuli (strongly tuned units).

Acknowledgements

This project was supported by NIH R21-EY-022751-02, Core Grant for Vision Research P30 EY001583, and the Neuroscience Neuroimaging Center Core Grant P30 NS045839. MO received support from the Academy of Finland Academy Fellow program. We thank Toni Saarela for valuable help with stimulus generation and Toni Saarela and Stacey Aston for helpful comments on the manuscript.

References

Ashby, F. G. and Maddox, W. T. (2005). Human category learning. *Annual review of psychology*, 56(1):149–78.

Barlow, H. B. (1990). A theory about the functional role and synaptic mechanism of visual after-effects. In Blakemore, C. B., editor, *Vision: coding and efficiency*, pages 363–375. Cambridge University Press, Cambridge.

Barlow, H. B. and Földiák, P. (1989). Adaptation and decorrelation in the cortex. In Durbin, R., Miall, C., and Mitchison, G., editors, *The Computing Neuron*, number 1988, pages 54–72. T. J. Press, Cornwall.

Barlow, H. B., MacLeod, D. I. A., and van Meeteren, A. (1976). Adaptation to gratings: No compensatory advantages found. *Vision Research*, 16(10):1043–1045.

Benucci, A., Saleem, A. B., and Carandini, M. (2013). Adaptation maintains population homeostasis in primary visual cortex. *Nat Neurosci*, 16(6):724–729.

Clifford, C. W. G., Webster, M. A., Stanley, G. B., Stocker, A. A., Kohn, A., Sharpee, T. O., and Schwartz, O. (2007). Visual adaptation: Neural, psychological and computational aspects. *Vision Research*, 47(25):3125–3131.

Clifford, C. W. G., Wenderoth, P., and Spehar, B. (2000). A functional angle on some after-effects in cortical vision. *Proceedings. Biological sciences / The Royal Society*, 267(1454):1705–1710.

Cortes, J. M., Marinazzo, D., Series, P., Oram, M. W., Sejnowski, T. J., and Van Rossum, M. C. (2012). The effect of neural adaptation on population coding accuracy. *Journal of computational neuroscience*, 32(3):387–402.

Dale, A. M., Fischl, B., and Sereno, M. I. (1999). Cortical Surface-Based Analysis. *NeuroImage*, 9(2):179–194.

Dragoi, V., Sharma, J., and Sur, M. (2000). Adaptation-induced plasticity of orientation tuning in adult visual cortex. *Neuron*, 28(1):287–298.

Drucker, D. M. and Aguirre, G. K. (2009). Different spatial scales of shape similarity representation in lateral and ventral LOC. *Cerebral Cortex*, 19(10):2269–2280.

Engel, S. A. (2005). Adaptation of oriented and unoriented color-selective neurons in human visual areas. *Neuron*, 45(4):613–623.

Enroth-Cugell, C. and Shapley, R. (1973). Adaptation and dynamics of cat retinal ganglion cells. *Journal of Physiology*, 233:271–309.

Giesel, M., Hansen, T., and Gegenfurtner, K. R. (2009). The discrimination of chromatic textures. *Journal of Vision*, 9(9):1–28,11.

Goldstone, R. L. and Gibson, E. J. (1962). Perceptual Learning. *Learning*, 49(1):585–612.

Greve, D. N. and Fischl, B. (2009). Accurate and robust brain image alignment using boundary-based registration. *NeuroImage*, 48(1):63–72.

Grill-Spector, K., Henson, R., and Martin, A. (2006). Repetition and the brain: neural models of stimulus-specific effects. *Trends in cognitive sciences*, 10(1):14–23.

Grill-Spector, K., Kourtzi, Z., and Kanwisher, N. G. (2001). The lateral occipital complex and its role in object recognition. *Vision Research*, 41(10-11):1409–1422.

Gutnisky, D. A. and Dragoi, V. (2008). Adaptive coding of visual information in neural populations. *Nature*, 452(7184):220–4.

Jenkinson, M., Bannister, P., Brady, M., and Smith, S. (2002). Improved optimisation for the robust and accurate linear registration and motion correction of brain images. *NeuroImage*, 17(2):825–841.

Julian, J. B., Fedorenko, E., Webster, J., and Kanwisher, N. (2012). An algorithmic method for functionally defining regions of interest in the ventral visual pathway. *NeuroImage*, 60(4):2357–2364.

Keefe, B. D., Dzhelyova, M., Perrett, D. I., and Barraclough, N. E. (2013). Adaptation improves face trustworthiness discrimination. *Frontiers in psychology*, 4(June):358.

Kohn, A. (2007). Visual Adaptation: Physiology, Mechanisms, and Functional Benefits. *Journal of Neurophysiology*, 10461:3155–3164.

Kohn, A. and Movshon, J. A. (2004). Adaptation changes the direction tuning of macaque MT neurons. *Nature neuroscience*, 7(7):764–72.

Koutstaal, W., Wagner, A. D., Rotte, M., Maril, A., Buckner, R. L., and Schacter, D. L. (2001). Perceptual speci city in visual object priming: functional magnetic resonance imaging evidence for a laterality difference in fusiform cortex. *English*, 39(2):184 – 199.

Krauskopf, J., Gegenfurtner, K., and Karl, G. (1992). Color discrimination and adaptation. *Vision Research*, 32(11):2165–2175.

Krekelberg, B., van Wezel, R. J. A., and Albright, T. D. (2006). Adaptation in macaque MT reduces perceived speed and improves speed discrimination. *Journal of neurophysiology*, 95(1):255–70.

Kriegeskorte, N., Goebel, R., and Bandettini, P. a. (2006). Information-based functional brain mapping. *Proceedings of the National Academy of Sciences of the United States of America*, 103(10):3863–8.

Kusunoki, M., Moutoussis, K., and Zeki, S. (2006). Effect of background colors on the tuning of color-selective cells in monkey area V4. *Journal of Neurophysiology*, 95:3047–3059.

M. Jenkinson (2004). Improving the registration of B0-disorted EPI images using calculated cost function weights. In *Tenth Int. Conf. on Functional Mapping of the Human Brain 2004*.

Macmillan, N. A. and Creelman, C. D. (2004). *Detection theory: A user's guide*. Psychology press.

Mattar, M. G., Kahn, D. A., Thompson-Schill, S. L., and Aguirre, G. K. (2016). Varying timescales of stimulus integration unite neural adaptation and prototype formation. *Current Biology*, 26(13):1669–1676.

Mueller, J. R., Metha, A. B., Krauskopf, J., and Lennie, P. (1999). Rapid Adaptation in Visual Cortex to the Structure of Images. *Science*, 285(5432):1405–1408.

Ng, M., Boynton, G. M., and Fine, I. (2008). Face adaptation does not improve performance on search or discrimination tasks. *Journal of Vision*, 8(1):1.1–20.

Ohzawa, I., Sclar, G., and Freeman, R. D. (1982). Contrast gain control in the cat visual cortex. *Nature*, 298(5871):266–268.

Oruç, I., Barton, J. J. S., Oruc, I., and Barton, J. J. S. (2011). Adaptation improves discrimination of face identity. *Proceedings. Biological sciences / The Royal Society*, 278(1718):2591–2597.

Phinney, R. E., Bowd, C., and Patterson, R. (1997). Direction-selective coding of stereoscopic (cyclopean) motion. *Vision Research*, 37(7):865–869.

Regan, D. and Beverley, K. I. (1985). Postadaptation orientation discrimination. *Journal of the Optical Society of America. A*, 2(2):147–155.

Rhodes, G., Maloney, L. T., Turner, J., and Ewing, L. (2007). Adaptive face coding and discrimination around the average face. *Vision Research*, 47(7):974–989.

Schwartz, O., Hsu, A., and Dayan, P. (2007). Space and time in visual context. *Nature reviews. Neuroscience*, 8(7):522–535.

Seriès, P., Stocker, A. A., and Simoncelli, E. P. (2009). Is the homunculus "aware" of sensory adaptation? *Neural Computation*, 21(12):3271–3304.

Simons, J. S., Koutstaal, W., Prince, S., Wagner, A. D., and Schacter, D. L. (2003). Neural mechanisms of visual object priming: Evidence for perceptual and semantic distinctions in fusiform cortex. *NeuroImage*, 19(3):613–626.

Smith, S. M. (2002). Fast, Robust Automated Brain Extraction. *Human Brain Mapping*, 17(3):143–155.

Vuilleumier, P., Henson, R. N., Driver, J., and Dolan, R. J. (2002). Multiple levels of visual object constancy revealed by event-related fMRI of repetition priming. *Nature Neuroscience*, 5(5):491–499.

Wainwright, M. J. (1999). Visual adaptation as optimal information transmission. *Vision Research*, 39(23):3960–3974.

Wark, B., Fairhall, A., and Rieke, F. (2009). Timescales of Inference in Visual Adaptation. *Neuron*, 61(5):750–761.

Webster, M. A. (2015). Visual adaptation. *Annual review of vision science*, 1:547–567.