A hypermorphic *Nfkbid* allele represents an *Idd7* locus gene contributing to impaired

thymic deletion of autoreactive diabetogenic CD8[+] T-cells in NOD mice.[1,2]

Maximiliano Presa[*], Jeremy J. Racine[*], Deanna J Lamont[*], Jennifer Dwyer[*], Jeremy Ratiu[*],

Vishal Kumar Sarsani[*], Yi-Guang Chen[†], Aron Geurts[†], Ingo Schmitz[‡§], Timothy Stearns[*],

Jennifer Allocco[*], Harold D. Chapman[*], and David V. Serreze[*]

[*] The Jackson Laboratory, Bar Harbor, Maine USA.

[†] Medical College of Wisconsin, Milwaukee, Wisconsin USA

[‡] Systems-Oriented Immunology and Inflammation Research Group, Helmholtz

Centre for Infection Research, Inhoffenstraße 7, 38124 Braunschweig

[§] Institute of Molecular and Clinical Immunology, Otto-von-Guericke University
12

Running Title: *Nfkbid* contributes to *Idd7*-control of negative selection.

Correspondence:

Dr. David Serreze, PhD

The Jackson Laboratory

600 Main St.

Bar Harbor, Maine 04609

ORCID: 0000-0001-7614-5925

dave.serreze@jax.org

(207) 288-6403

Abstract: 234/250 words

Text: 5786 words

Abstract:

In both NOD mice and humans the development of type 1 diabetes (T1D) is dependent at least in part on autoreactive CD8$^+$ T-cells recognizing pancreatic ß-cell peptides presented by often quite common MHC class I variants. Studies in NOD mice previously revealed the common K^d and/or D^b class I molecules expressed by this strain acquire an aberrant ability to mediate pathogenic CD8 T-cell responses through interactions with T1D susceptibility (*Idd*) genes outside the MHC.   A gene(s) mapping to the *Idd7* locus on proximal Chromosome 7 was previously shown to be an important contributor to the failure of the common class I molecules expressed by NOD mice to mediate the normal thymic negative selection of diabetogenic CD8$^+$ T-cells. Using an inducible model of thymic negative selection and mRNA transcript analyses we initially identified an elevated *Nfkbid* expression variant as a strong candidate for an NOD *Idd7* region gene contributing to impaired thymic deletion of diabetogenic CD8$^+$ T-cells. CRISPR/Cas9-mediated genetic attenuation of *Nfkbid* expression in NOD mice resulted in improved negative selection of autoreactive diabetogenic AI4 and NY8.3 CD8$^+$ T-cells. These results indicated allelic variants of *Nfkbid* represent an *Idd7* gene contributing to the efficiency of intrathymic diabetogenic CD8$^+$ T-cell deletion. However, while enhancing thymic deletion of pathogenic CD8$^+$ T-cells, ablation of *NFkbid* expression surprisingly accelerated T1D onset in NOD mice likely by eliciting decreases in both regulatory T- and B-lymphocytes (Tregs/Bregs) mediating peripheral immunosuppressive effects.

Introduction:

In both the NOD mouse model and in humans, type 1 diabetes (T1D) results from the autoimmune destruction of insulin-producing pancreatic β-cells mediated by the combined activity of $CD4^+$ and $CD8^+$ T-cells as well as B-lymphocytes (1). T1D is highly polygenic in nature, as more than 50 loci have been associated with disease susceptibility or resistance in both humans and NOD mice (2, 3). However, while its complete pathogenic etiology remains unsolved there is a wide acceptance that disease develops as a consequence of interactions between T1D susceptibility (*Idd*) genes resulting in breakdowns in mechanisms controlling induction of immunological tolerance to self-proteins (4, 5). Some *Idd* genes appear to contribute to defects in central tolerance induction mechanisms that normally represent a first checkpoint engendering the thymic deletion of autoreactive $CD4^+$ and $CD8^+$ T-cells during early stages of their development (6-8). However, even in healthy individuals, some autoreactive T-cells escape central tolerance mechanisms (9). Such autoreactive effectors are normally prevented from mediating pathogenic effects by a second checkpoint, provided by multiple mechanisms of peripheral tolerance induction, with significant evidence for some of these processes also being disrupted by various *Idd* genes (5).

In the thymus, double positive (DP, $CD4^+CD8^+$) immature thymocytes are able to recognize self-peptide-MHC complexes displayed on antigen presenting cells (APC, thymic epithelial cells, dendritic cells (DC) or macrophages) through interactions with clonally distributed T-cell receptor (TCR) molecules (10). The strength of TCR signaling resulting from the trimolecular complex interaction, dictates the fate of immature DP thymocytes (9, 10). DP thymocytes undergoing high avidity interactions with self-peptide-MHC complexes are normally deleted by apoptosis or diverted to a regulatory T-cell lineage (Tregs, $CD4^+Foxp3^+$) (9).

Deficiencies in this process strongly contribute to the survival and presence in the periphery of

potentially autoreactive T-cells or inadequacies in self-antigen specific Tregs (5). Both in

humans and NOD mice, the defects in central tolerance can be exerted at three levels: reduced

presentation of self-antigen in thymus; defects in post-TCR engagement signaling events; and

variability in the basal or induced expression level of molecules controlling the final pro-/anti-

apoptotic balance (4, 5). Several *Idd* genes and loci have been identified as contributing to such

central tolerance induction defects (4, 5). Among these are MHC haplotypes long known to

provide the primary T1D genetic risk factor (11). Within the MHC, certain unusual class II

molecules play an important role in the pathogenesis of T1D by allowing the development and

functional activation of autoreactive CD4[+] T-cells (12). However, MHC class II expression is

largely restricted to thymic epithelial cells and hematopoietically derived APC (10, 13). Thus,

while class II restricted autoreactive CD4[+] T-cells clearly play a critical role in T1D

development, they cannot do so by directly engaging insulin producing pancreatic β cells (14).

Conversely, like most other tissues, pancreatic β cells do express MHC class I molecules. Hence,

due to an ability to directly engage and destroy pancreatic β cells, MHC class I restricted

autoreactive CD8[+] T-cells are likely the ultimate mediators of T1D development in both humans

and NOD mice (14, 15). Epidemiological studies have shown that in addition to class II effects,

particular HLA class I variants provide an independent T1D risk factor in humans (16, 17).

Interestingly, this includes some quite common MHC class I variants that in both humans and

NOD mice only acquire an aberrant ability to contribute to T1D when expressed in the context of

other *Idd* genes.

Previous studies found interactive contributions from some polymorphic *Idd* genes

outside of the MHC determine the extent to which particular class I molecules can allow the

development and functional activation of diabetogenic CD8$^+$ T-cells. This was initially

demonstrated by congenically transferring a transgenic TCR (Vα8/Vβ2) from the H2^{g7} class I

restricted diabetogenic CD8$^+$ T-cell clone AI4 from a NOD genetic background strain (NOD-

AI4) to a T1D resistant C57BL/6-stock congenic for the *H2^{g7}* haplotype (B6.*H2^{g7}*). Autoreactive

AI4 TCR transgenic T-cells were deleted to a significantly greater extent at the DP stage of

thymic development in B6.*H2^{g7}* than NOD mice (7). This result indicated non-MHC genes

allelically differing in NOD and B6 background mice regulate the extent to which autoreactive

diabetogenic CD8$^+$ T-cells undergo thymic deletion. Such non-MHC genes controlling differing

thymic negative selection efficiency in NOD and B6 background mice were also found to

function in a T-cell intrinsic manner (7). Subsequent linkage analyses and congenic strain

evaluation identified a region on proximal Chromosome (Chr.) 7 to which the *Idd7* locus had

been previously mapped *(18, 19)*, as containing a gene strongly contributing to the differential

thymic deletion efficiency of diabetogenic AI4 CD8$^+$ T-cells in NOD and B6.*H2^{g7}* mice (8).

In this study we identify *Nfkbid* as a likely *Idd7* region gene regulating the efficiency of

diabetogenic CD8 T-cells thymic negative selection. *Nfkbid* (aka IκBNS) is a family member of

the non-conventional IkB modulators of the transcription factor NF-κB (20). Depending on cell

type and physiological context *Nfkbid* can either positively or negatively modulate NFkB activity

(20). *Nfkbid* was first discovered as a gene expressed in thymocytes undergoing negative

selection upon antigen specific TCR stimulation (21). Curiously, genetic ablation of *Nfkbid* in B6

background mice did not elicit any alteration in thymocyte subset distribution or any overt signs

of autoimmunity (22). Using TCR transgenic mouse models to analyze antigen specific thymic

negative selection, we show in this report that a higher expression variant of *Nfkbid* in NOD than

B6 mice contributes to diminished negative selection in the former strain of both the AI4 and

NY8.3 MHC class I restricted autoreactive diabetogenic CD8[+] T-cell clonotypes. While genetic

attenuation of *Nfkbid* expression to levels similar to that in B6 background mice results in

improved thymic deletion of pathogenic CD8[+] T-cells, total ablation of this gene also diminishes

levels of peripheral immunoregulatory T- and B-lymphocytes (Tregs, Bregs) resulting in

accelerated T1D development in NOD mice.

Materials and Methods

Mouse strains

NOD/ShiLtDvs (NOD), B6.NOD-(D17Mit21-D17Mit10)/LtJ (B6.$H2^{g7}$) and NOD.Cg-

Prkdc^{scid} Emv30b/Dvs (NOD.*scid*) mice are maintained in a specific pathogen free research

colony at The Jackson Laboratory. NOD.*Foxp3-eGFP* knockin mice, carrying the reporter

construct *Foxp3$^{tm2(eGFP)-Tch}$* were previously described (23, 24). NOD mice carrying a transgenic

TCR (Vα8.3, Vβ2) derived from the AI4 diabetogenic CD8^{+} T-cell clone (NOD-AI4) have also

been previously described (25). Previously generated NOD.LCMV mice transgenically express a

TCR (Vα2/Vβ8) from a CD8^{+} T-cell clone recognizing the H2-D^{b} restricted gp33 peptide

(KAVYNFATM) derived from Lymphocytic Choriomeningitis Virus (LCMV) (26, 27). An

NOD stock transgenically expressing the TCR derived from the H2-K^{d} restricted diabetogenic

NY8.3 CD8^{+} T-cell clone (28) (here designated NOD.NY8.3) were acquired from the type 1

diabetes resource (http://type1diabetes.jax.org/) and maintained in an heterozygous state. An

NOD stock congenic for a segment of B6-derived Chr. 7 delineated by the flanking markers *Gpi^{b}*

and *D7Mit346* (29), designated as NOD.*Ch7^{B6}*FL was used to introduce the interval into NOD-

AI4 mice (8).

Congenic truncation analysis

The NOD.*Ch7^{B6}*FL-AI4 stock was intercrossed with NOD-AI4. Resultant F1 progeny

were intercrossed and all F2 offspring were analyzed for recombination events in the Chr. 7

congenic region delineated by the flanking markers *Gpi^{b}* (34.2 Mb) and *D7Mit346* (58.7 Mb).

Genomic tail DNA samples were screened for the microsatellite markers *D7MIT267, D7MIT117,*

D7MIT155, Gpi1^{b}, D7MIT79, D7MIT225, D7MIT78, D7MIT247, D7MIT270, D7MIT230, and

D7mit346 by PCR. Identified recombinant mice were backcrossed with NOD-AI4 and then

intercrossed to generate progeny homozygous for the sub-congenic regions. The homozygous

state for each congenic line generated was verified by PCR analyses of the markers listed above.

Generation of NOD Nfkbid deficient mice

CRISPR/Cas9 technology was utilized to directly ablate the *Nfkbid* gene in NOD mice

(NOD-*Nfkbid*$^{-/-}$). NOD/ShiLtDvs embryos were intra-cytoplasmic microinjected with 3 pl of a

solution containing Cas9 mRNA and single guide RNA (sgRNA) at respective concentrations of

100 ng/μl and 50 ng/μl. The sgRNA sequence (5'-AGGCCCATTTCCCCTGGTGA-3') was

designed to delete the transcriptional start site of *Nfkbid* in exon 3. Genomic tail DNA was

screened by targeted Sanger sequencing: the genomic region around exon 3 was amplified by

PCR with primers Nfkbid-KO-F1 (5'-TGCTTGAGATCCAGTAG-3') and Nfkbid-KO-R1 (5'-

CCCTGACATCTCAGAATA-3'), the resulting PCR product was purified and sequenced using

an ABI 3730 DNA analyzer (Applied Biosystems). Analyses of Sanger sequencing data was

done using the software Poly Peak Parser (30) which allows identification of the different alleles

present in heterozygous mutant mice. During the screening of N1 mutants we identified an allele

characterized by an 8 nt deletion and an insertion of 154 nt originated by copy and inversion of a

DNA segment from *Nfkbid* exon 2. This produced a major disruption of the *Nfkbid* gene and

deletion of the transcription start site (Supplementary Fig. 1). N1F1 NOD-*Nfkbid*$^{+/-}$ mutant mice

were intercrossed and the disrupted allele fixed to homozygosity. The new stock, formally

designated NOD/ShiLtDvs-*Nfkbid*$^{<em3Dvs>/}$Dvs (abbreviated as NOD-*Nfkbid*$^{-/-}$), was maintained

by brother-sister matting.

Previous reports indicated that *Nfkbid* deficient mice have an altered Foxp3$^+$ Treg

compartment (31). Thus, we introduced the *Foxp3$^{tm2(eGFP)-Tch}$* reporter construct previously

congenically transferred to the NOD strain (24) into the NOD-*Nfkbid$^{-/-}$* stock and subsequently

fixed to homozygosity.

Gene expression analysis

Total RNA from whole thymus or sorted cell lysates was extracted using the RNeasy

mini kit (Qiagen, Germantown, MD, USA) following the vendor instructions. For whole thymus

tissue RNA, the complete organ was put in 2 ml of RNAlater stabilization solution (Thermo

Fisher Scientific, Waltham, MA, USA), incubated 24 h at room temperature and then stored at -

20˚C until processing. For RNA extraction from sorted cells, 2×10^5 cells were sorted in 100%

FBS, washed 2 times in cold PBS and resuspended in RLT lysis buffer (Qiagen, Germantown,

MD, USA) and stored at -20˚C until processing. Extracted total RNA was quantified by

NanoDrop (Thermo Fisher Scientific, Waltham, MA, USA) and quality assessed in Bioanalyzer

(Agilent Technologies, Santa Clara, CA, USA). Samples with a RNA integrity index (RIN

index) of 7 or higher were used for gene expression analyses. For cDNA synthesis 500 ng of

total RNA was diluted to 5 μl in DEPC treated water and mixed with 5 μl of SuperScript IV

VILO Master Mix (Thermo Fisher Scientific, Waltham, MA, USA) following vendor

instructions.

Expression of *Nfkbid* was assessed by a predesigned and validated TaqMan qPCR assay

using Mm.PT.58.12759232 (IDT Integrated Technologies, Coralville, Iowa, USA) targeting the

exon 3-4 region and able to identify all four possible transcripts. The assay consists of a FAM-

labeled   and   double-quenched   probe   (5'-/56-FAM-CCTGGTGAT-/ZEN/-

GGAGGACTCTCTGGAT-/3IABkFQ/-3') and the primers p1 (5'-

GACAGGGAAGGCTCAGGATA-3') and p2 (5'-GCTTCCTGACTCCTGATTTCTAC-3').

The assay was reconstituted in 500µl of TE buffer (10 mM Tris-HCl pH 8.0, 0.1 mM EDTA)

generating a 20x solution that yield a final 1x solution of 500 nM primers and 250 nM probe. As

an endogenous control we used a predesigned and validated TaqMan assay for *Gapdh* (ID:

Mm99999915_g1) (VIC-labeled) (Applied Biosystems -Thermo Fisher Scientific, Waltham,

MA, USA). The qPCR analysis was done in a multiplexing setting combining both *Nfkbid* and

Gapdh assays using IDT prime Gene expression 2x master mix (IDT Integrated Technologies,

Coralville, Iowa, USA) and 6 ng of cDNA template in a final volume of 10µl. The experiment

was run in a ViiA-7 real time PCR system (Thermo Fisher Scientific, Waltham, MA, USA) with

four technical and four biological replicates, using the following program: 2 min at 50 ˚C, 3 min

at 95˚C, and 40 cycles of 3 min at 95˚C, 1 min 60˚C. An internal calibration curve was included

for both *Nfkbid* and *Gapdh*. The efficiency of each TaqMan assay was calculated from their

respective standard curves using the application SC, while relative gene expression was

determined by the ddCt method using the application RQ both in Thermo Fisher Cloud Software,

Version 1.0 (Thermo Fisher Scientific, Waltham, MA, USA).

Microarray analysis

Three to five 5-week-old, NOD.LCMV and NOD.Ln82-LCMV female mice were

injected i.v. with 0.5 µg of gp33 or D^b binding control peptide. Two hours post-injection whole

thymus tissue was processed as described above for total RNA extraction. Total RNA samples

were hybridized to Affymetrix Mouse Gene 1.0 ST Arrays. Average signal intensities for each

probe set within arrays were calculated by and exported from Affymetrix's Expression Console

(Version 1.1) software using the RMA method which incorporates convolution background

correction, summarization based on a multi-array model fit robustly using the median polish

algorithm, and sketch-quantile normalization. For this experiment, two pairwise comparisons

were used to statistically resolve gene expression differences between experimental groups using

the R/maanova analysis package (32). Specifically, differentially expressed genes were detected

by using F1, the classical F-statistic that only uses data from individual genes. Statistical

significance levels of the pairwise comparisons were calculated by permutation analysis (1000

permutations) and adjusted for multiple testing using the false discovery rate (FDR), q-value,

method (33). Differentially expressed genes are declared at an FDR q-value threshold of 0.05.

Two contrast groups were established: control D^b peptide (NOD.LCMV vs NOD.Ln82-LCMV),

and gp33 peptide (NOD.LCMV vs NOD.Ln82-LCMV). Data sets for contrast group Db and

gp33 were analyzed in IPA analysis software QIAGEN Bioinformatics (Redwood City, CA,

USA). Data were filtered for a expression fold change (EFC) > 2 and an expression FDR (q-

value) < 0.05.

Flow cytometry analysis

Single cell suspensions of thymus (Thy), spleen (SPL) and pancreatic lymph nodes

(PLN) from 5-6 week old female mice were prepared. Red blood cells in splenocytes samples

were lysed with Gey's buffer. $2x10^6$ cells were stained with the following monoclonal

antibodies: CD4-BV650 (GK1.5), CD8-PE-Cy7 (53-6.72), TCRVα8.3-FITC or PE (B21.14),

TCRVβ2-A647 (B20.6), TCRVβ8.1,2,3-FITC (F23.1) CD19-A700 (ID3), CD11b-PB (M1/70),

CD11c-PB (N418), CD21-FITC (7G6), CD23-PE (B3B4), CD5-PE (53-7.313) acquired from

BD Bioscience (San Jose, CA, USA) or BioLegend (San Diego, CA, USA), MHC class I

tetramers Tet-AI4-PE (H-2D^b/MimA2, sequence YAIENYLEL) (34) and Tet-NRPv7-PE (H-

2K^d/NRPv7) were acquired from the National Institutes of Health tetramer core facility (Atlanta,

GA, USA). Dead cells were excluded by DAPI staining. Stained cells were acquired using a

FACS LSRII instrument (BD Bioscience). All flow cytometric data were analyzed with FlowJo

(FlowJo LLC, Ashland, OR, USA).

T-cell Proliferation assay

Total splenic T-cells were purified from 5-week-old NOD and NOD-*Nfkbid$^{-/-}$* female

mice by negative depletion of CD11c$^+$, CD11b$^+$, B220$^+$, Ter19$^+$, CD49b$^+$ cells with biotin-

conjugated antibodies and streptavidin magnetic beads over MACS columns (Miltenyi Biotech)

following manufacturer's protocols. Purified T-cells were washed with cold PBS, counted and

adjusted to a concentration of $2x10^7$/ml for labeling with 10 μM cell tracer eFluor670

(eBioscience). Labeled cells were adjusted to $4x10^6$/ml. Whole collagenase digested splenocytes

from NOD.*scid* mice were used as APCs. Labeled T-cells were seeded into a 96 well plate at a

density of $2x10^5$/well with $4x10^5$ APC/well in triplicate. Anti-CD3 monoclonal antibody (145-

2C11) was added at final concentrations of 4, 2, 1, 0.5, 0.25 and 0.125 μg/ml. The cells were

incubated at 37˚C, 5% CO$_2$ for 48h. T-cell proliferation was assessed by flow cytometric analysis

of cell tracer eFluor670 dilution. Proliferation parameters were obtained using the proliferation

platform analysis in FlowJo (FlowJo LLC, Ashland, OR, USA).

Cytokine secretion analysis

IL-2 and IFN-γ secretion was assessed in supernatants of T-cell proliferation cultures by

ELISA using BD OptEIA kits (BD Bioscience, San Diego, CA, USA).

Western blot

Whole thymus lysates were prepared in RIPA buffer (Cell signaling Technologies) containing Halt protease inhibitor cocktail (Thermo Fisher Scientific). Total protein content was quantified by Bradford assay (Thermo Fisher Scientific). 20µg of total protein were separated by SDS-PAGE using Miniprotean TGX precast gradient 4-15% gels (Bio-rad Laboratories). Proteins were electrotransferred to a PVDF membrane and incubated in blocking buffer (5% Milk, TBST) overnight at 4˚C. The membrane was probed with a polyclonal rabbit anti-NFkbid (35), , diluted at 1/1000 in 0.5% Milk/TBST, and incubated 1 hr at room temperature. Anti-Rabbit-HRP diluted at 1/40000 in 0.5% Milk/TBST, was used as secondary Ab. Supersignal Wes-fempto (Thermo Fisher Scientific) was used for chemiluminescent detection. β-Actin was detected as loading control, over the same membrane after stripping, using mouse mAb anti-β-Actin, and anti-mouse IgG-HRP.

Diabetes incidence

Diabetes was monitored once a week using urine glucose strips (Diastix, Bayer). Mice with two consecutive readings >250mg/dl (corresponding to a blood glucose of >300mg/dl) were considered diabetic.

Statistical analysis

Data analysis and graphs were made using Prism 6 software (GraphPad, San Diego, CA, USA). Statistical analysis is detailed in the correspondent Figure Legends.

Results

Mapping an Idd7 region gene(s) regulating thymic numbers of diabetogenic AI4 CD8$^+$ T-

cells to a 5.4 Mb region on Chr. 7.

We previously developed a mouse stock, designated NOD.*Chr7^{B6}*FL-AI4, expressing the

transgenic AI4 TCR and congenic for a C57BL/6-derived segment of Chr.7 overlapping the

previously identified *Idd7* locus delineated by the flanking markers *Gpi1^b* (34.2 Mb) and

D7Mit346 (58.7Mb) (29). AI4 T-cells underwent significantly higher levels of thymic deletion

at the DP stage of development in the NOD.*Chr7^{B6}*FL-AI4 congenic stock than in full NOD

background mice (8).   This indicated a gene(s) allelically varying between NOD and B6

background mice mapping within the *Idd7* locus overlapping congenic interval played a strong

role in the extent diabetogenic AI4 TCR transgenic CD8$^+$ T-cells underwent thymic deletion (8).

We were subsequently able to derive from the original NOD.*Chr7^{B6}*FL-AI4 stock two sub-

congenic lines designated NOD.Ln82-AI4 (markers *D7Mit117–D7Mit247*) and NOD.Ln16-AI4

(markers *D7Mit79–D7Mit247*) (Fig. 1A). Analyses by flow cytometry found that compared to

NOD-AI4 controls, the yield of DP AI4 thymocytes (identified by staining with the TCR Vα8.3

specific monoclonal antibody B21.14) (Fig. 1B) was significantly decreased in the NOD.Ln82-

AI4 sub-strain (Fig. 1C). In contrast, the yield of AI4 DP thymocytes in the NOD.Ln16-AI4

stock were similar to that in NOD-AI4 controls (Fig. 1B, C). These differences indicated a 5.4

Mb region on Chr.7 delineated by the markers *D7Mit117-D7Mit225* contains a polymorphic

gene(s) influencing the extent to which DP AI4 thymocytes undergo negative selection.

Expression levels of the clonotypic Vα8.3 TCR element was higher in the NOD.Ln82-AI4 stock

compared to NOD-AI4 controls (Fig. 1D), a trait previously observed in NOD.*Chr7^{B6}*FL-AI4

mice (8). These results indicated a gene(s) in the 5.4 Mb region on Chr. 7 between *D7Mit117*

and *D7Mit225* may modulate how efficiently AI4 T-cell undergo thymic deletion through

regulation of TCR expression levels (8).

Nfkbid is the only differentially expressed Idd7 region gene in diabetogenic CD8$^+$ T-cells

undergoing low versus high levels of thymic deletion.

Due to differing cellular composition, we did not feel it appropriate to utilize thymii from

NOD-AI4 and NOD.Ln82-AI4 mice to carry out initial mRNA transcript analyses to identify

genes potentially contributing to the differential negative selection efficiency of diabetogenic

CD8$^+$ T-cells in these strains. Instead, we reasoned a more appropriate platform to carry out

such analyses would be provided by a previously produced NOD background stock

transgenically expressing a TCR (Vα2/Vβ8) recognizing the H2-D^b class I restricted gp33

peptide (KAVYNFATM) derived from Lymphocytic Choriomeningitis Virus (LCMV) (26, 27).

The basis for choosing this NOD.LCMV stock was that LCMV reactive CD8$^+$ T-cells

developing in the thymus would not be under any thymic negative selection pressure until such

time their cognate antigen is exogenously introduced. We generated an additional NOD.LCMV

stock homozygous for the Ln82 congenic interval. NOD.LCMV and NOD.Ln82-LCMV mice

were injected i.v. with 0.5µg of the gp33 or an H2-D^b binding control peptide (ASNENMETM).

At 24 hours post-injection, we assessed the proportion of LCMV specific DP thymocytes (TCR

Vα2$^+$) remaining present in NOD.LCMV and NOD.Ln82-LCMV mice that had received the

gp33 peptide relative to control cohorts. A significantly greater proportion of gp33 reactive DP

thymocytes were deleted in NOD.Ln82-LCMV than NOD.LCMV mice (Fig. 2A). These results

further indicated a polymorphic gene(s) in the *Idd7* region regulates the sensitivity of DP

thymocytes to undergo antigen-induced deletion.

The LCMV TCR transgenic system provided an opportunity to compare the influence of

the *Idd7* region over the transcriptional profile of thymocytes undergoing different levels of

antigen-induced negative selection. Time-course experiments indicated that at 2 and 4 hours

post-injection with the cognate antigen or peptide control there was no significant reduction in

the number of Vα2^{+} DP thymocytes (Supplemental Fig. 2). However, we hypothesized that at 2-

hour post-antigen injection, a differential gene expression profile may have already been induced

in NOD.LCMV and NOD.Ln82-LCMV thymocytes contributing to their subsequent low and

high level of deletion seen at 24 hours. Using cutoff criteria of a q value <0.05, and minimum

two fold strain variation, microarray-based mRNA profiling found 16 and 212 genes were

respectively differentially expressed in thymocytes from NOD.LCMV and NOD.Ln82-LCMV

mice injected with the D^{b} binding control or gp33 peptide (Fig. 2B). This included 12 genes that

were commonly differentially expressed in thymocytes from NOD.LCMV and NOD.Ln82-

LCMV mice treated with either the control or gp33 peptide (Fig. 2B). None of the genes

differentially expressed in thymocytes from NOD.LCMV and NOD.Ln82-LCMV mice treated

with the control peptide mapped to the earlier defined 5.4 Mb *Idd7* support interval (genomic

coordinates 7:29,926,306-35,345,930). Intriguingly, *Nfkbid* was the only gene mapping to this

5.4 Mb *Idd7* support interval that was differentially expressed in thymocytes from NOD.LCMV

and NOD.Ln82-LCMV mice treated with the gp33 peptide (Fig. 3C). *Nfkbid* was expressed at a

3.6 fold higher level in thymocytes from gp33 treated NOD.LCMV than NOD.Ln82-LCMV

mice (Fig. 3C). We also subsequently found by RT-qPCR analyses that *Nfkbid* is expressed at

~2 fold higher levels in flow cytometrically sorted equalized numbers of DP thymocytes from

NOD-AI4 than NOD.Ln82-AI4 mice (Fig. 2D).

Nfkbid (aka IκBNS) encodes a non-conventional modulator of the transcription factor

NF-κB (20), and ironically was first discovered as a gene expressed in thymocytes undergoing

negative, but not positive selection (21). Depending on cell type and physiological context

NFkbid can either positively or negatively modulated NF-κB activity (20). Low to moderate

levels of NF-κB activity (primarily the p50/p65 heterodimeric form) reportedly inhibits thymic

negative selection of CD8 lineage destined T-cells (36). On the other hand, high NF-κB activity

levels can elicit thymic negative selection of CD8$^+$ T-cells (36). Thus, either negative or positive

Nfkbid-mediated modulation of NF-κB activity could potentially contribute to the extent

diabetogenic CD8$^+$ T-cells undergo thymic deletion. For this reason, the rest of the current study

focused on testing the hypothesis that the higher *Nfkbid* expression variant in NOD than B6 mice

contributes to the less efficient thymic deletion of diabetogenic CD8 T-cells in the former strain.

Generation of NOD-Nfkbid deficient mice

In order to test whether the higher relative to B6 *Nfkbid* expression variant in NOD mice

contributes to diminished thymic deletion of diabetogenic CD8$^+$ T-cells, we utilized a

CRISPR/Cas9 approach to ablate this gene. The mouse *Nfkbid* gene is located on Chr.7 (genomic

coordinates GRCm38.p5 7: 30,421,732-30,428,746 and NOD_ShiLtJ_V1 7:31068239-

31076307), and has four alternative transcripts, three of which generate the same protein (327

amino acids) with a forth representing a non-coding sequence (37). Based on the reference

sequence (NM_172142.3), *Nfkbid* consist of 11 exons, 9 of them protein coding, with the start

codon located at the end of exon 3 (Fig. 3A). We designed a single guide RNA (gRNA) (Fig.

3A) with the intention to disrupt the start codon in exon 3 and thus eliminate translation of all

potential transcripts. We identified via PCR and targeted DNA sequencing a mutant line with a

deletion of 8 nucleotides including the start codon, and an additional Indel mutation consisting of

a 154 nucleotides copied from exon 2 and inserted in an inverted orientation in the Cas9 cutting

site (Fig. 3B). The mutation was fixed to homozygosity and the new stock was designated as

NOD-*Nfkbid*[-/-] (formally NOD/ShiLtDvs-*Nfkbid*[<emDvs3>]/Dvs. Analyses of gene expression by

RT-qPCR indicated an absence of thymic *Nfkbid* transcript expression in NOD-*Nfkbid*[-/-] mice

compared to the easily detectable levels in wild type NOD controls (Fig. 3C). We performed

western blot analyses using total thymus lysates from NOD, NOD-*Nfkbid*[-/-], and B6.*Nfkbid*[-/-]

mice (22, 31). Nfkbid was absent in both NOD and B6 Nfkbid deficient mice, while it was

present in wild type NOD mice (Fig. 3D).

Diminution of Nfkbid expression enhances thymic deletion of both AI4 and NY8.3

diabetogenic CD8[+] T-cells in NOD mice.

We next evaluated whether diminution of *Nfkbid* expression enhanced the thymic

deletion of diabetogenic CD8[+] T-cells in NOD mice. To initially do this, the inactivated *Nfkbid*

gene was transferred in both a heterozygous and homozygous state to the NOD-AI4 strain. As

expected analysis of gene expression by qPCR showed respective high and absent *Nfkbid* mRNA

transcript levels in thymocytes from NOD-AI4 and NOD-AI4-Nfkbid[-/-] mice (Fig. 4A). Also as

expected, *Nfkbid* mRNA transcript levels were significantly higher in thymocytes from NOD-

AI4 than B6.*H2*[g7] control mice (Fig. 4A). A finding of potential physiological relevance was that

thymic *Nfkbid* mRNA transcript expression in NOD-AI4-Nfkbid[+/-] heterozygous mice was

reduced to a level not differing from B6.*H2*[g7] controls (Fig. 4A). Numbers of DP, but not CD8[+]

SP thymocytes positively staining with the H2-D^b/MimA2 tetramer capable of binding the AI4

TCR (Tet-AI4) (gating strategy shown in Fig. 4B) were significantly less in both heterozygous

and homozygous *Nfkbid* knockout mice, compared to gene intact controls (Fig. 4C). The same

result was obtained by staining with Vα8.3 and Vβ2 specific antibodies binding the AI4 TCR

(results not shown). As shown in Fig. 1D, TCR expression was significantly higher on DP

thymocytes from NOD.*Ch7^{B6}*Ln82-AI4 congenic mice than in controls in which the *Idd7* region

was of NOD origin. In contrast, partial or complete ablation of *Nfkbid* did not alter AI4 TCR

expression levels on either DP or CD8$^+$ SP thymocytes (Fig. 4D). Thus, some *Idd7* region

gene(s) other than *Nfkbid* regulates thymic TCR expression levels. These results also disprove

an earlier hypothesis (8) that an *Idd7* region gene(s) may control the thymic deletion efficiency

of diabetogenic CD8$^+$ T-cell by modulating TCR expression levels.

We next tested whether the apparent ability of varying *Nfkbid* expression levels to

modulate the thymic deletion efficiency of diabetogenic CD8$^+$ T-cells extended to clonotypes

beyond AI4. Thus, we introduced the *Nfkbid* knockout allele into a NOD stock with CD8$^+$ T-

cells transgenically expressing the NY8.3 TCR (NOD.NY8.3) that recognizes a peptide derived

from the ß-cell protein islet-specific glucose-6-phosphatase catalytic subunit-related protein

(IGRP) (28, 38). As observed for the AI4 clonotype, numbers of DP thymocytes staining with a

tetramer (designated NRPV7) recognized by the NY8.3 TCR (Fig. 5A) were significantly less in

6-week-old NOD.NY8.3-*Nfkbid$^{+/-}$* heterozygous mice than in controls (Fig. 5B). When examined

at 4 weeks of age, numbers of both NY8.3 DP and CD8$^+$ SP thymocytes were significantly less

in NOD.NY8.3-*Nfkbid$^{-/-}$* homozygous mice than in controls (Fig. 5C). Also, similar to the case

observed for the AI4 clonotype, expression levels of the NY8.3 TCR on DP or CD8$^+$ SP

thymocytes was not altered by the presence or absence of Nfkbid (Fig. 5D). Thus, based on

assessments of two separate clonotypes we conclude the *Nfkbid* expression variant characterizing

NOD mice represents the *Idd7* region gene contributing to impaired thymic deletion of

diabetogenic CD8$^+$ T-cells in this strain.

Nfkbid deficiency paradoxically contributes to accelerated T1D in NOD mice by

impairing development of regulatory lymphocyte populations.

We previously found that *Idd7* region gene effects were restricted to the thymus. This

was based on findings that while numerically reduced in the thymus, the AI4 T-cells that did

reach the periphery of B6.*H2^{g7}* and NOD.*Ch7^{B6}*FL background mice were still able to expand

and mediate T1D development (7). However, we were surprised to find T1D development was

actually accelerated in NOD-*Nfkbid$^{-/-}$* mice (Fig. 6A). This prompted us to assess if ablation of

the *Nfkbid* gene in NOD mice induced any numerical or functional differences in peripheral

leukocyte populations varying from those previously reported in B6 mice carrying a similar

mutation, but that exhibited no signs of autoimmunity (22, 31, 39). As previously observed for

the B6 background stock, numbers of splenic CD4$^+$ and CD8$^+$ T cells in NOD-*Nfkbid$^{-/-}$* mice did

not differ from wild type controls (data not shown). As previously observed in B6 (40), marginal

zone (MZ) B-lymphocytes in the spleen and B1a cells in the peritoneum were both significantly

reduced in *Nfkbid* deficient NOD mice (Fig. 6B, C).   The proliferative capacity of CD4$^+$ and

CD8$^+$ T-cells and their ability to secrete the cytokines IL-2 and IFNγ was reported to be

diminished in *Nfkbid* deficient B6 mice (22). Similar phenotypes characterized CD4$^+$ and CD8$^+$

T-cells from NOD-*Nfkbid$^{-/-}$* mice (Fig. 6D, E). Ablation of *Nfkbid* has also been reported to

diminish numbers of FoxP3 expressing Tregs in B6 background mice (31). A similar effect was

elicited by ablation of *Nfkbid* in the NOD strain (Fig. 6F). Previous reports indicates that Nfkbid

is a direct regulator of IL-10 (35). We found that proportions of FoxP3^{+} and FoxP3^{-} CD4^{+} T-cells

producing immunosuppressive IL-10 upon *ex vivo* stimulation were significantly lower in *Nfkbid*

deficient than intact NOD mice (Fig. 6G). We recently found that partially overlapping

populations of CD73 expressing and IL-10 producing B-lymphocytes can mediate peripheral

regulatory effects (Bregs) inhibiting autoimmune T1D developments in NOD mice (41).

Ablation of *Nfkbid* also reduced levels of such IL-10 producing and CD73 expressing B-

lymphocytes in NOD mice (Fig. 6H, I). Thus, while diminishing *Nfkbid* expression enhances

thymic negative selection of diabetogenic CD8^{+} T cells in NOD mice, the coincidental decrease

in numbers of immunoregulatory lymphocyte populations likely accounts for why any

pathogenic effectors that do reach the periphery still aggressively induce disease development.

Discussion

It has become increasingly appreciated that in addition to the strong contributions

provided by some class II region genes, that when expressed in the proper genomic context,

particular common MHC class I variants are also key contributors to T1D development.

Previous work demonstrated primarily due to interactions with some non-MHC gene(s) in the

Idd7 locus on proximal Chr. 7 the common class I variants expressed by NOD mice (K^d, D^b)

have a significantly lessened ability to mediate the thymic negative selection of diabetogenic

$CD8^+$ T-cells (7). In this study, through congenic truncation analyses we initially fine mapped

the *Idd7* region gene(s) controlling the efficiency of diabetogenic $CD8^+$ T-cell thymic deletion to

a 5.4 Mb interval on proximal Chr. 7. Identification of the responsible *Idd7* region gene was

greatly aided by the exploitation of NOD.LCMV TCR transgenic stocks that did or did not carry

a B6 derived congenic interval encompassing the relevant region of proximal Chr. 7. The

advantage of these stocks is that their thymocytes are under no negative selection pressure until

such time the LCMV cognate gp33 antigenic peptide is deliberately introduced. Upon

introduction of the gp33 peptide LCMV DP thymocytes were deleted to a significantly greater

extent in the presence of B6 rather than NOD derived genome across the relevant *Idd7* region.

Subsequent microarray based mRNA transcript analyses found following introduction of the

gp33 antigenic peptide *Nfkbid* was the only gene within the relevant 5.4 Mb region on Chr. 7 that

was differentially expressed in NOD.LCMV TCR mice carrying NOD or B6 genomic material

across the relevant *Idd7* region. The presence of NOD rather than B6 derived genome across this

Idd7 region resulted in *Nfkbid* being expressed at 3-4 times higher levels in antigen stimulated

LCMV reactive thymocytes. This was similarly true for thymocytes transgenically expressing

the TCR from the naturally occurring diabetogenic AI4 $CD8^+$ T-cell clone derived from NOD

mice. Thus, we tested through CRISPR/Cas9 technology if diminishing or eliminating *Nfkbid*

expression in NOD mice resulted in improved thymic negative selection. NOD stocks in which

Nfkbid expression was either completely eliminated or reduced to B6-like levels were found to

have an improved capacity to mediate thymic negative selection of AI4 clonotypic cells as well

as those transgenically expressing the TCR from the diabetogenic NY8.3 CD8$^+$ T-cell clone.

These collective results indicated the higher *Nfkbid* expression variant in NOD than B6 mice is

likely the *Idd7* region gene strongly contributing to diminished thymic deletion of at least

diabetogenic CD8$^+$ T-cells in the former strain.

Nfkbid is mostly a nuclear protein formed by 7 ankyrin repeated domains (21). Due to the

lack of DNA binding and transactivation domains, Nfkbid function is mediated through protein-

protein interactions with NF-κB family members. Indeed Nfkbid can function as inhibitor or

enhancer of NF-κB activities depending on the protein partner and cell type (20). This includes

previously reported interactions of *Nfkbid* with the NF-κB subunit p50 (Nfkb1) in thymocytes

undergoing negative selection (21). However, our current results indicate *Nfkbid* is not required

for at least thymocytes destined for the CD8$^+$ lineage to undergo negative selection, and instead

its particular expression level in NOD mice appears to diminish the efficiency of this process.

Three different *Nfkbid* deficient mouse models have been previously produced, all on the

B6 background (22, 39, 42). These previously produced models have been useful to identify

several functional aspects of *Nfkbid*. In the B-cell compartment, *Nfkbid* expression is necessary

for development of the MZ and B1-a subsets as well as the generation of antibody responses to

T-dependent and independent antigens (22, 40, 42-44). *Nfkbid* deficiency was also associated

with an impairment in IL-10 secreting B-lymphocytes (45). *Nfkbid* deficient T-cells have

impaired *in vitro* proliferation and cytokine secretion (IL-2, IFN-γ) (22). We found these

phenotypes also characterize our *Nfkbid* deficient NOD mouse stock.

Interaction of Nfkbid with c-Rel has also been reported to contribute to Foxp3 expression

in Treg cells (31). Thus, we found it of interest that in addition to enhancing the thymic deletion

of diabetogenic CD8$^+$ T-cells, diminution of *Nfkbid* expression also resulted in decreased

numbers of peripheral Tregs in NOD mice. We recently reported that overlapping populations of

CD73 expressing and IL-10 producing B-lymphocytes can exert T1D protective

immunoregulatory activity in NOD mice (41).   Ablation of *Nfkbid* expression also diminished

levels of both CD73 expressing and IL-10 producing B-lymphocytes in the periphery. Thus, we

conjecture it is the loss of both Tregs and these immunoregulatory B-lymphocytes that accounts

for the unexpected finding that T1D development is actually accelerated in *Nfkbid* deficient

NOD mice despite their improved, albeit not completely restored, ability to mediate the thymic

deletion of pathogenic CD8$^+$ T-cells. Hence, *Nfkbid* expression levels appears to act as a two-

edged sword, low levels in thymus can result in better negative selection of autoreactive CD8$^+$ T-

cells, but at the same time can contribute to autoimmunity in the periphery by diminishing levels

of regulatory lymphocyte populations.

In conclusion, *Nfkbid* appears to be an important contributory gene to how efficiently

autoreactive diabetogenic CD8$^+$ T-cells undergo thymic negative selection. However, this study

also revealed a caveat that would have to be considered if it might ultimately become possible to

diminish *Nfkbid* expression through pharmacological means. That is while diminishing *Nfkbid*

expression allows for increased thymic deletion of diabetogenic CD8$^+$ T-cells, it does so at the

price of decreasing levels of regulatory lymphocytes allowing any pathogenic effectors that do

reach the periphery to more rapidly induce disease onset. This explains our previous findings

that the *Idd7* region exerts two edged sword effects where the presence of B6 derived genome in

the region exerted T1D protective effects at the thymic level, but in the periphery actually

promotes disease development (8). Thus, when contemplating possible future interventions

designed to diminish numbers of diabetogenic T-cells, a consideration that should be taken into

account is whether the treatment might also impair regulatory cell numbers or activity allowing

the pathogenic effectors which do remain present to become more aggressive.

Acknowledgements

We would like to thank the Jackson Laboratory Research Animal Facility, Genetic Engineering Technology group, Flow Cytometry service, and Transgenic Genotyping core for technical support on this project. We thank the National Institutes of Health tetramer core facility for the tetramers used in this study.

Footnotes

[1] MP was supported by JDRF Fellowship 3-PDF-2014-219-A-N. For parts of this work, JJR1 was supported by either NIH Fellowship 1F32DK111078 or JDRF Fellowship 3-PDF-2017-372-A-N. He was also supported by a grant from the Diabetes Research Connection DRC 006887 JR. DVS is supported by NIH grants DK-46266, DK-95735, and OD-020351-5022. This work was also partly supported by Cancer Center Support Grant CA34196. The authors would also like to thank Carl Stiewe and his wife, Maike Rohde, for their generous donation toward T1D research at The Jackson Laboratory, which contributed to this work. IS was supported by grants of the Deutsche Forschunggemeinschaft (SCHM1586/6-1 and project A23 of SFB854)

[2] MP designed and conducted experimentation, interpreted data, and wrote the manuscript. JJR contributed to experimental design, data interpretation, and writing of the manuscript. JDL, JA and HDC conducted experimentation. JD and JR designed and conducted experiments and interpreted data. VKS and TS contributed to statistical analyses. YGC, AG and IS contributed to experimental design. DVS contributed to study conception, supervised experimental effort, and writing of the manuscript.

References

1. Atkinson, M. A., G. S. Eisenbarth, and A. W. Michels. 2014. Type 1 diabetes. *The Lancet* 383: 69-82.
2. Ram, R., M. Mehta, Q. T. Nguyen, I. Larma, B. O. Boehm, F. Pociot, P. Concannon, and G. Morahan. 2016. Systematic Evaluation of Genes and Genetic Variants Associated with Type 1 Diabetes Susceptibility. *The Journal of Immunology* 196: 3043-3053.
3. Driver, J., D. Serreze, and Y.-G. Chen. 2011. Mouse models for the study of autoimmune type 1 diabetes: a NOD to similarities and differences to human disease. *Seminars in Immunopathology* 33: 67-87.
4. Geenen, V. 2012. Thymus and type 1 diabetes: an update. *Diabetes Res Clin Pract* 98: 26-32.
5. Jeker, L. T., H. Bour-Jordan, and J. A. Bluestone. 2012. Breakdown in Peripheral Tolerance in Type 1 Diabetes in Mice and Humans. *Cold Spring Harbor Perspectives in Medicine* 2.
6. Kishimoto, H., and J. Sprent. 2001. A defect in central tolerance in NOD mice. *Nat Immunol* 2: 1025-1031.
7. Choisy-Rossi, C.-M., T. M. Holl, M. A. Pierce, H. D. Chapman, and D. V. Serreze. 2004. Enhanced Pathogenicity of Diabetogenic T Cells Escaping a Non-MHC Gene-Controlled Near Death Experience. *The Journal of Immunology* 173: 3791-3800.
8. Serreze, D. V., C. M. Choisy-Rossi, A. E. Grier, T. M. Holl, H. D. Chapman, J. R. Gahagan, M. A. Osborne, W. Zhang, B. L. King, A. Brown, D. Roopenian, and M. P. Marron. 2008. Through Regulation of TCR Expression Levels, an Idd7 Region Gene(s) Interactively Contributes to the Impaired Thymic Deletion of Autoreactive Diabetogenic CD8+ T Cells in Nonobese Diabetic Mice. *The Journal of Immunology* 180: 3250-3259.
9. von Boehmer, H., and F. Melchers. 2010. Checkpoints in lymphocyte development and autoimmune disease. *Nat Immunol* 11: 14-20.
10. Klein, L., M. Hinterberger, G. Wirnsberger, and B. Kyewski. 2009. Antigen presentation in the thymus for positive selection and central tolerance induction. *Nat Rev Immunol* 9: 833-844.
11. Todd, J. A. 2010. Etiology of Type 1 Diabetes. *Immunity* 32: 457-467.
12. Pociot, F., B. Akolkar, P. Concannon, H. A. Erlich, C. Julier, G. Morahan, C. R. Nierras, J. A. Todd, S. S. Rich, and J. Nerup. 2010. Genetics of Type 1 Diabetes: What's Next? *Diabetes* 59: 1561-1571.
13. Drozina, G., J. Kohoutek, N. Jabrane-Ferrat, and B. M. Peterlin. 2005. Expression of MHC II Genes. In *Molecular Analysis of B Lymphocyte Development and Activation*. H. Singh, and R. Grosschedl, eds. Springer Berlin Heidelberg. 147-170.
14. Varanasi, V., L. Avanesyan, D. M. Schumann, and A. V. Chervonsky. 2012. Cytotoxic Mechanisms Employed by Mouse T Cells to Destroy Pancreatic β-Cells. *Diabetes* 61: 2862-2870.
15. Knight, R. R., D. Kronenberg, M. Zhao, G. C. Huang, M. Eichmann, A. Bulek, L. Wooldridge, D. K. Cole, A. K. Sewell, M. Peakman, and A. Skowera. 2013. Human β-Cell Killing by Autoreactive Preproinsulin-Specific CD8 T Cells Is Predominantly

Granule-Mediated With the Potency Dependent Upon T-Cell Receptor Avidity.
Diabetes 62: 205-213.

16.   Nejentsev, S., J. M. Howson, N. M. Walker, J. Szeszko, S. F. Field, H. E. Stevens, P.
Reynolds, M. Hardy, E. King, J. Masters, J. Hulme, L. M. Maier, D. Smyth, R. Bailey, J. D.
Cooper, G. Ribas, R. D. Campbell, D. G. Clayton, and J. A. Todd. 2007. Localization of
type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. *Nature*
450: 887-892.

17.   Noble, J. A., A. M. Valdes, M. D. Varney, J. A. Carlson, P. Moonsamy, A. L. Fear, J. A.
Lane, E. Lavant, R. Rappner, A. Louey, P. Concannon, J. C. Mychaleckyj, H. A. Erlich,
and f. t. T. D. G. Consortium. 2010. HLA Class I and Genetic Susceptibility to Type 1
Diabetes: Results From the Type 1 Diabetes Genetics Consortium. *Diabetes* 59:
2972-2979.

18.   Ghosh, S., S. M. Palmer, N. R. Rodrigues, H. J. Cordell, C. M. Hearne, R. J. Cornall, J. B.
Prins, P. McShane, G. M. Lathrop, L. B. Peterson, and et al. 1993. Polygenic control of
autoimmune diabetes in nonobese diabetic mice. *Nat Genet* 4: 404-409.

19.   McAleer, M. A., P. Reifsnyder, S. M. Palmer, M. Prochazka, J. M. Love, J. B. Copeman, E.
E. Powell, N. R. Rodrigues, J.-B. Prins, D. V. Serreze, N. H. DeLarato, L. S. Wicker, L. B.
Peterson, N. J. Schork, J. A. Todd, and E. H. Leiter. 1995. Crosses of NOD Mice With
the Related NON Strain: A Polygenic Model for IDDM. *Diabetes* 44: 1186-1195.

20.   Schuster, M., M. Annemann, C. Plaza-Sirvent, and I. Schmitz. 2013. Atypical IkappaB
proteins - nuclear modulators of NF-kappaB signaling. *Cell Communication and*
Signaling 11: 23.

21.   Fiorini, E., I. Schmitz, W. E. Marissen, S. L. Osborn, M. Touma, T. Sasada, P. A. Reche,
E. V. Tibaldi, R. E. Hussey, A. M. Kruisbeek, E. L. Reinherz, and L. K. Clayton. 2002.
Peptide-Induced Negative Selection of Thymocytes Activates Transcription of an NF-
kB Inhibitor. *Molecular cell* 9: 637-648.

22.   Touma, M., V. Antonini, M. Kumar, S. L. Osborn, A. M. Bobenchik, D. B. Keskin, J. E.
Connolly, M. J. Grusby, E. L. Reinherz, and L. K. Clayton. 2007. Functional Role for
IκBNS in T Cell Cytokine Regulation As Revealed by Targeted Gene Disruption. *The*
Journal of Immunology 179: 1681-1692.

23.   Haribhai, D., W. Lin, L. M. Relland, N. Truong, C. B. Williams, and T. A. Chatila. 2007.
Regulatory T cells dynamically control the primary immune response to foreign
antigen. *J Immunol* 178: 2961-2972.

24.   Presa, M., Y.-G. Chen, A. E. Grier, E. H. Leiter, M. A. Brehm, D. L. Greiner, L. D. Shultz,
and D. V. Serreze. 2015. The Presence and Preferential Activation of Regulatory T
Cells Diminish Adoptive Transfer of Autoimmune Diabetes by Polyclonal Nonobese
Diabetic (NOD) T Cell Effectors into NSG versus NOD-scid Mice. *The Journal of*
Immunology.

25.   Graser, R. T., T. P. DiLorenzo, F. Wang, G. J. Christianson, H. D. Chapman, D. C.
Roopenian, S. G. Nathenson, and D. V. Serreze. 2000. Identification of a CD8 T cell
that can independently mediate autoimmune diabetes development in the complete
absence of CD4 T cell helper functions. *J Immunol* 164: 3913-3918.

26.   Pircher, H., K. Burki, R. Lang, H. Hengartner, and R. M. Zinkernagel. 1989. Tolerance
induction in double specific T-cell receptor transgenic mice varies with antigen.
Nature 342: 559-561.

27. Serreze, D. V., E. A. Johnson, H. D. Chapman, R. T. Graser, M. P. Marron, T. P. DiLorenzo, P. Silveira, Y. Yoshimura, S. G. Nathenson, and S. Joyce. 2001. Autoreactive Diabetogenic T-Cells in NOD Mice Can Efficiently Expand From a Greatly Reduced Precursor Pool. *Diabetes* 50: 1992-2000.

28. Verdaguer, J., D. Schmidt, A. Amrani, B. Anderson, N. Averill, and P. Santamaria. 1997. Spontaneous Autoimmune Diabetes in Monoclonal T Cell Nonobese Diabetic Mice. *The Journal of Experimental Medicine* 186: 1663-1676.

29. Chen, J., P. Reifsnyder, F. Scheuplein, W. Schott, M. Mileikovsky, S. Soodeen-Karamath, A. Nagy, M. Dosch, J. Ellis, F. Koch-Nolte, and E. Leiter. 2005. "Agouti NOD": identification of a CBA-derived Idd locus on Chromosome 7 and its use for chimera production with NOD embryonic stem cells. *Mammalian Genome* 16: 775-783.

30. Hill, J. T., B. L. Demarest, B. W. Bisgrove, Y.-C. Su, M. Smith, and H. J. Yost. 2014. Poly peak parser: Method and software for identification of unknown indels using sanger sequencing of polymerase chain reaction products. *Developmental Dynamics* 243: 1632-1636.

31. Schuster, M., R. Glauben, C. Plaza-Sirvent, L. Schreiber, M. Annemann, S. Floess, Anja A. Kühl, Linda K. Clayton, T. Sparwasser, K. Schulze-Osthoff, K. Pfeffer, J. Huehn, B. Siegmund, and I. Schmitz. 2012. IκBNS Protein Mediates Regulatory T Cell Development via Induction of the Foxp3 Transcription Factor. *Immunity* 37: 998-1008.

32. Wu, H., M. K. Kerr, X. Cui, and G. A. Churchill. 2003. MAANOVA: A Software Package for the Analysis of Spotted cDNA Microarray Experiments. In *The Analysis of Gene Expression Data: Methods and Software*. G. Parmigiani, E. S. Garrett, R. A. Irizarry, and S. L. Zeger, eds. Springer New York, New York, NY. 313-341.

33. Storey, J. D. 2002. A direct approach to false discovery rates. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 64: 479-498.

34. Lieberman, S. M., T. Takaki, B. Han, P. Santamaria, D. V. Serreze, and T. P. DiLorenzo. 2004. Individual Nonobese Diabetic Mice Exhibit Unique Patterns of CD8+ T Cell Reactivity to Three Islet Antigens, Including the Newly Identified Widely Expressed Dystrophia Myotonica Kinase. *The Journal of Immunology* 173: 6727-6734.

35. Annemann, M., Z. Wang, C. Plaza-Sirvent, R. Glauben, M. Schuster, F. Ewald Sander, P. Mamareli, A. A. Kühl, B. Siegmund, M. Lochner, and I. Schmitz. 2015. IκBNS Regulates Murine Th17 Differentiation during Gut Inflammation and Infection. *The Journal of Immunology* 194: 2888-2898.

36. Jimi, E., I. Strickland, R. E. Voll, M. Long, and S. Ghosh. 2008. Differential Role of the Transcription Factor NF-κB in Selection and Survival of CD4+ and CD8+ Thymocytes. *Immunity* 29: 523-537.

37. Aken, B. L., S. Ayling, D. Barrell, L. Clarke, V. Curwen, S. Fairley, J. Fernandez Banet, K. Billis, C. García Girón, T. Hourlier, K. Howe, A. Kähäri, F. Kokocinski, F. J. Martin, D. N. Murphy, R. Nag, M. Ruffier, M. Schuster, Y. A. Tang, J.-H. Vogel, S. White, A. Zadissa, P. Flicek, and S. M. J. Searle. 2016. The Ensembl gene annotation system. *Database* 2016: baw093-baw093.

38. Amrani, A., J. Verdaguer, P. Serra, S. Tafuro, R. Tan, and P. Santamaria. 2000. Progression of autoimmune diabetes driven by avidity maturation of a T-cell population. *Nature* 406: 739-742.

39.   Kuwata, H., M. Matsumoto, K. Atarashi, H. Morishita, T. Hirotani, R. Koga, and K.
Takeda. 2006. IκBNS Inhibits Induction of a Subset of Toll-like Receptor-Dependent
Genes and Limits Inflammation. *Immunity* 24: 41-51.
40.   Touma, M., D. B. Keskin, F. Shiroki, I. Saito, S. Koyasu, E. L. Reinherz, and L. K.
Clayton. 2011. Impaired B Cell Development and Function in the Absence of IκBNS.
The Journal of Immunology 187: 3942-3952.
41.   Ratiu, J. J., J. J. Racine, M. G. Hasham, Q. Wang, J. A. Branca, H. D. Chapman, J. Zhu, N.
Donghia, V. Philip, W. H. Schott, C. Wasserfall, M. A. Atkinson, K. D. Mills, C. M. Leeth,
and D. V. Serreze. 2017. Genetic and Small Molecule Disruption of the AID/RAD51
Axis Similarly Protects Nonobese Diabetic Mice from Type 1 Diabetes through
Expansion of Regulatory B Lymphocytes. *The Journal of Immunology*.
42.   Arnold, C. N., E. Pirie, P. Dosenovic, G. M. McInerney, Y. Xia, N. Wang, X. Li, O. M. Siggs,
G. B. Karlsson Hedestam, and B. Beutler. 2012. A forward genetic screen reveals
roles for Nfkbid, Zeb1, and Ruvbl2 in humoral immunity. *Proceedings of the National
Academy of Sciences* 109: 12286-12293.
43.   Pedersen, G. K., M. Àdori, S. Khoenkhoen, P. Dosenovic, B. Beutler, and G. B. Karlsson
Hedestam. 2014. B-1a transitional cells are phenotypically distinct and are lacking
in mice deficient in IκBNS. *Proceedings of the National Academy of Sciences* 111:
E4119-E4126.
44.   Pedersen, G. K., M. Ádori, j. M. Stark, S. Khoenkhoen, C. Arnold, B. Beutler, and G. B.
Karlsson Hedestam. 2016. Heterozygous mutation in IκBNS leads to reduced levels
of natural IgM antibodies and impaired responses to T-independent type 2 antigens.
Frontiers in Immunology 7.
45.   Miura, M., N. Hasegawa, M. Noguchi, K. Sugimoto, and M. Touma. 2016. The atypical
IκB protein IκBNS is important for Toll-like receptor-induced interleukin-10
production in B cells. *Immunology* 147: 453-463.
713
714

Figure 1

Figure 1: Mapping an *Idd7* region gene(s) controlling the thymic negative selection
efficiency of diabetogenic AI4 CD8[+] T-cells to a 5.4 Mb region on Chr.7. (**A**) Schematic
diagram of B6 derived Chr.7 congenic regions in NOD-AI4 mouse strains. Markers positions are
indicated in Mb based on genome assembly release GRCm38.p5 (37) (**B**) Flow cytometric gating
strategy to enumerate AI4 DP thymocytes.   Thymocytes from 5-week-old female NOD-AI4
(n=32), NOD.Ln82-AI4 (n=20) and NOD.Ln16-AI4 (n=14) were stained with CD4, CD8 and
Vα8.3 specific antibodies. (**C**) Numbers of live (PI negative) CD4[+]CD8[+] (DP) AI4 (Vα8.3[+])
thymocytes in the indicated strains. (**D**) Mean fluorescence intensity (MFI) of staining of DP
thymocytes from the indicated strains by the monoclonal antibody Vα8.3-PE. All values
represent mean±SEM of more than 3 independent experiments, statistical significance was
determined by Log-rank (Mantel-Cox) t-test (ns p > 0.05, ** p < 0.001, **** p < 0.00001).
728

729
730

Figure 2: *Nfkbid* is the only differentially expressed *Idd7* region gene in diabetogenic
CD8[+] T-cells undergoing low versus high levels of thymic deletion. (**A**) For antigen-induced
negative selection, either 0.5 μg of gp33 (cognate antigen) or D[b] binding control peptide were
injected i.v. into five-week-old female NOD.LCMV (n=18 for controls and n=6 for gp33 treated)
or NOD.Ln82-LCMV (n=22 for controls and n=6 for gp33 treated) mice. Remaining DP Vα2[+]
thymocytes were quantified at 24 hours post-injection in each group. The efficiency of deletion
was calculated as a ratio of the number DP Vα2[+] thymocytes remaining in gp33 treated mice to
the average numbers in controls ($\frac{x\ gp33\ DP\ V\alpha2+}{yDb\ DP\ V\alpha2+} \times 100$) and expressed as a percentage. Statistical
significance was determined by Log-rank (Mantel-cox) t-test (ns p > 0.05, * p < 0.05). (**B**) As
assessed by microarray analyses number of genes deferentially expressed declared at a false
discovery rate (q-value) <0.05 and a fold change ≥2 in thymocytes from NOD.LCMV and
NOD.Ln82-LCMV 2 hours after i.v. injection with Db control or gp33 peptide. (**C**) Graph
showing the average log$_2$ (Intensity of probe set) for differentially expressed probe sets (same
criteria as noted for panel B) between NOD.LCMV *vs* NOD.Ln82-LCMV mice injected i.v. two
hours earlier with gp33 peptide. The arrow indicates *Nfkbid*. (**D**) Total RNA was extracted from
2x10[5] DP Vß8[+] thymocytes sorted from 6-week-old female NOD.AI4 and NOD.Ln82-AI4 mice.
Nfkbid mRNA relative quantification was done using the ddCt method and *Gapdh* as endogenous
control and NOD.Ln82-AI4 mice as reference, mean relative fold change (RFC) ± SEM is
indicated. * p <0.05 unpaired t-test.

Figure 3: Targeting *Nfkbid* by CRISPR/Cas9 in NOD mice. (**A**) Representation of
mouse *Nfkbid* gene showing the translation start site at the end of exon 3 and underlined the
single guide RNA designed to target this region by CRISPR/Cas9. The PAM sequence is
indicated in lower case.. (**B**) The resulting targeted mutation consisting of an 8 nucleotide
deletion and an insertion of 154 nucleotides originated by copy and inversion of a segment from
Nfkbid exon 2. (**C**) Total thymic RNA was extracted from NOD.*Nfkbid*$^{+/+}$ and NOD.*Nfkbid*$^{-/-}$
mice. Relative *Nfkbid* expression was analyzed by qPCR using *Gapdh* as endogenous control and
NOD.*Nfkbid*$^{+/+}$ as a reference. Relative fold change (RFC) ± SEM is indicated. (**D**) Nfkbid
protein expression was assessed by western blot analysis. Total thymus lysates from NOD,
NOD.*Nfkbid*$^{-/-}$, and B6.*Nfkbid*$^{-/-}$ (negative control) were separated by SDS-PAGE and electro-
transferred to a PVDF membrane. Nfkbid was detected using a previously validated polyclonal
rabbit anti-Nfkbid. Nfkbid is detected as two isoforms, a 35 kDa and a high molecular weight of
54 kDa probably resulting from post-translational modifications. Lower panel indicates the
detection of the loading control β-Actin obtained using the same membrane after stripping.
768

Figure 4

Figure 4: Genetic attenuation of *Nfkbid* expression controls the numbers of AI4 DP thymocytes. (**A**) *Nfkbid* gene expression was analyzed by qPCR of total thymus RNA from 6-week-old *B6.H2^{g7}* (n=2), NOD-AI4 (n=4), NOD-AI4-Nfkbid$^{+/-}$ (n=4) and NOD-AI4-Nfkbid$^{-/-}$ (n=3) mice. *Gapdh* was used as endogenous control and B6.*H2^{g7}* as the reference group. Graph shows the mean±SEM of relative fold change of *Nfkbid* expression normalized to *Gapdh*. Statistical significance was determined by one-way ANOVA and Bonferroni's multiple comparison test. The p-value of samples compared to the reference group is indicated. Note: We did not use B6.*H2^{g7}*-AI4 mice as a reference in this study because the numbers of thymocytes is extremely low and would not provide a fair comparison for gene expression analysis. (**B**) Illustration of gating to enumerate DP and CD8 SP thymocytes staining with the AI4 TCR specific tetramer. Depicted profiles are for thymocytes from a NOD-AI4 control. (**C**) Numbers of AI4 TCR expressing DP and CD8$^+$ SP thymocytes from 6-week-old NOD-AI4, NOD-AI4-Nfkbid$^{+/-}$, and NOD-AI4-Nfkbid$^{-/-}$ female mice. Data represent the mean±SEM of DP or CD8$^+$ SP thymocytes staining positive with the AI4 specific tetramer reagent. (D) Geometric MFI of AI4 tetramer staining of the same DP and CD8$^+$ SP thymocytes evaluated in panel B. Bars represent the mean±SEM. Statistical significance was analyzed by 2-way ANOVA and Bonferroni's multiple comparison test. ns p >0.05, * p <0.05, ** p <0.01.

Figure 5: *Nfkbid* expression levels modulate thymic negative selection efficiency of
diabetogenic NY8.3 clonotypic CD8$^+$ T-cells. (A) Illustration of gating to enumerate DP and
CD8 thymocytes staining with the NY8.3 TCR specific NRPv7 tetramer. Depicted profiles are
for thymocytes from a NOD.NY8.3 control. (B) Numbers of NY8.3 TCR expressing DP and
CD8$^+$ thymocytes from 6-week-old female NOD.NY8.3 and NOD.NY8.3-*Nfkbid*$^{+/-}$ heterozygous
mice. Data represent the mean±SEM of DP or CD8$^+$ SP thymocytes staining positive with the
NY8.3 specific NRPv7 tetramer reagent. (C) Numbers of NY8.3 TCR expressing DP and CD8$^+$
SP thymocytes from 4-week-old female NOD.NY8.3 and NOD.NY8.3-*Nfkbid*$^{-/-}$ homozygous
mice. Bars represent the mean±SEM of 3 independent experiments. (D) Geometric MFI of
NY8.3 tetramer staining of DP and CD8$^+$ SP thymocytes from NOD.NY8.3, NOD.NY8.3-
Nfkbid$^{+/-}$, and NOD.NY8.3-*Nfkbid*$^{-/-}$ mice. Bars represent mean±SEM. Statistical significance
was analyzed by 2-way ANOVA and Bonferroni's multiple comparison test. ns p >0.05, ** p
<0.01.
807

Figure 6: *Nfkbid* deficiency paradoxically accelerates T1D onset in NOD mice by impairing development of regulatory lymphocyte populations.
(A) T1D development was analyzed in a cohort of NOD and NOD.*Nfkbid*$^{-/-}$ female mice. Survival curve comparisons were analyzed through Log-rank (Mantel-Cox) test. (B) Splenocytes from 7 week old female NOD (n=3) and NOD.*Nfkbid*$^{-/-}$(n=5) mice were labeled with Ab for CD19, CD21 and CD23 and analyzed by flow cytometry. Singlet, DAPI$^-$ CD19$^+$ cells were identified as total B-cells. Based on expression of CD23 and CD21, B-cells were classified as follicular (FO: CD23$^+$CD21^{lo}), marginal zone (MZ: CD23$^-$CD21^{hi}) transitional 1 (T1: CD23$^-$CD21$^-$) and transitional 2 (T2: CD23^{int} CD21^{hi}) (C) Peritoneal cavity B-cells were isolated by

lavage with cold HBSS from the same group of mice described in (B), and stained for CD19,
B220, CD5 and analyzed by flow cytometry. Total B-cells (Singlet, DAPI$^-$ CD19$^+$) were
classified based in the expression of B220 and CD5 as B1-a (CD5$^+$ B220$^{int/lo}$) and B1-b (CD5$^-$
B220^{lo}) (D) The proliferation capacity of purified total T-cells from NOD and NOD.*Nfkbid*$^{-/-}$
mice was assessed as described in methods section. Briefly, MACS purified total T-cells were
labeled with cell tracer eFluor670 and stimulated *in vitro* with increasing amounts of anti-CD3
mAb. After 48 hours of culture the cells were labeled with Ab for CD4 and CD8. Proliferation
in CD4 and CD8 T-cell subsets was analyzed by flow cytometric analyses of cell tracer
eFluor670 dilution. Proliferation index was determined using the Flow Jo cell proliferation tool.
Data represent the mean±SEM of 3 technical replicates. (E) Cell culture supernatants of purified
T-cells used in the proliferation assay were assessed for secretion of the cytokines IL-2 and IFN-
γ by ELISA. Data represent mean±SEM of 3 technical replicates. (F) Enumeration of splenic
CD4$^+$ FoxP3$^+$ (based on *Foxp3$^{tm2(eGFP)-Tch}$* reporter expression) Tregs in 6-week-old female NOD
(n=5) and NOD.*Nfkbid*$^{-/-}$ mice (n=9). (G) Analysis by intracellular staining and subsequent flow
cytometric analyses of proportion of Foxp3$^+$ and FoxP3$^-$ splenic CD4$^+$ T-cells from 6-week-old
female NOD (n=5) and NOD.*Nfkbid*$^{-/-}$ mice (n=9) producing IL-10. (H) Analysis by intracellular
staining and subsequent flow cytometric analyses of proportions of IL-10 producing splenic B-
lymphocytes in 6-week-old female NOD (n=5) and NOD.*Nfkbid*$^{-/-}$ mice (n=9). (I) Proportions of
splenic CD73$^+$ B-lymphocytes in 6-week-old female NOD (n=5) and NOD.*Nfkbid*$^{-/-}$ mice (n=9).
All bar graphs represent mean±SEM. Statistical significance was determined by 2-way ANOVA
and Bonferroni's multiple comparison test. ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.