Title

Methionine coordinates a hierarchically organized anabolic program through *GCN4*

Adhish S. Walvekar[1], Rajalakshmi Srinivasan[1], Ritu Gupta[1] and Sunil Laxman[1*]

[1] Institute for Stem Cell biology and Regenerative Medicine,

NCBS-TIFR campus, GKVK Post, Bellary Road, Bangalore 560065.

[*] Correspondence: sunil@instem.res.in

Keywords: methionine, S-adenosyl methionine, Gcn4, growth, pentose phosphate pathway,

NADPH, reductive biosynthesis

Abstract

Methionine availability during overall amino acid limitation transforms cells to a proliferative state, the underlying metabolic basis of which remains unclear. Here, we investigate the nature and mechanism of this anabolic transformation, using yeast. Combining comparative transcriptome analysis, biochemical and metabolic flux based approaches, we discover that methionine, predominantly through Gcn4p, controls specific regulatory nodes in metabolism. Methionine increases activity of the pentose phosphate pathway, reductive biosynthetic capacity, and transamination capacity, including the synthesis of glutamate/glutamine. These processes converge on key reactions in the synthesis of costly amino acids, and subsequently nucleotides, which are also induced in a methionine dependent manner. This thereby enables the arrangement of a sequential cascade driving an overall anabolic transformation. Thus, methionine switches cells to anabolism, co-opting a "starvation stress response" regulator, Gcn4p, to achieve this. These results suggest a hierarchical metabolic framework explaining how methionine mediates an anabolic transformation.

Introduction

Cell growth is expensive, and is therefore tightly co-ordinated with the intrinsic cellular metabolic state. Changes in the cellular metabolic state influence global transcription and translation, which determine the cell fate appropriate for that condition (*Broach, 2012*; *Ljungdahl and Daignan-Fornier, 2012*; *Dechant and Peter, 2008*; *Brauer et al., 2008*; *Boer et al., 2010*). In general, the enormous metabolic costs incurred during growth and proliferation come from two well-studied phenomena. First, to successfully complete division, a cell must make substantial metabolic investments, in order to replicate its genome, and synthesize proteins, lipids, nucleotides, and other macromolecules. Correspondingly, the

process of protein synthesis required for growth itself consumes large amounts of energy

(*Warner et al., 2001*; *Warner, 1999*), as the translational output of cells increases (*Jorgensen*

and Tyers, 2004). Therefore, understanding the key metabolic transformations driving growth

are areas of intense study, particularly in cancer biology, ranging from the famous "Warburg

effect" (*Warburg, 1956*; *Vander Heiden et al., 2009*), to a broader identification of the

biosynthetic and metabolic requirements for cell growth (*DeBerardinis et al., 2008*; *Tong et*

al., 2009; *Vander Heiden et al., 2009*). However, many fundamental concepts explaining how

cell growth and proliferation are coordinated with the metabolic state of a cell remain poorly

addressed.

Nonetheless, several studies particularly using unicellular models like *Saccharomyces*

cerevisiae have revealed dedicated strategies utilized by eukaryotic cells to integrate growth

with metabolic state (*Brauer et al., 2008*; *Boer et al., 2010*; *Gray et al., 2004*; *Broach, 2012*;

Ljungdahl and Daignan-Fornier, 2012; *Cai et al., 2011*; *Slavov and Botstein, 2011*). A

powerful approach enabling a systematic dissection of molecular mechanisms and metabolic

hierarchies has been to first limit preferred carbon or nitrogen sources, thereby slowing down

overall growth. By slowing down overall growth, and then reintroducing individual factors

and studying responses, several studies have untangled hierarchies of complex metabolic

phenomena, revealed specific nutrient sensing mechanisms in cells, and identified how the

build-up and utilization of specific, "sentinel" metabolites control growth outputs (*Brauer et*

al., 2008; *Boer et al., 2010, 2008*; *Cai et al., 2011*; *Tu et al., 2005, 2007*; *Slavov and Botstein,*

2013; *Xu et al., 2013*; *Klosinska et al., 2011*; *Laxman and Tu, 2010*; *Sutter et al., 2013*;

Laxman et al., 2013). Utilizing such a strategy, *S. cerevisiae* subjected to amino acid

limitations during a shift from a rich, complex medium with lactate as the carbon source, to a

defined minimal medium with the same carbon source, revealed that a single amino acid,

methionine (and its metabolite S-adenosylmethionine (SAM)), strongly promoted growth

(*Wu and Tu, 2011, Sutter et al., 2013*; *Laxman et al., 2013*; *Laxman et al., 2014*). These

observations reinforce a host of data from other eukaryotic systems including mammals

suggesting that methionine is a strong growth signal, or "growth metabolite" (*Troen et al.,*

2007; *Lee et al., 2016, 2014*; *Cavuoto and Fenech, 2012*; *Sugimura et al., 1959*; *Breillout et*

al., 1990). Therefore, this phenomenon in yeast provides a system using which two

fundamental, related questions can be addressed: First, what is the methionine mediated

anabolic transformation, and the metabolic logic therein? Second, what is the mechanism by

which methionine mediates this anabolic transformation, even in overall amino acid limiting

conditions? We address these related questions in this study.

Here, building from known models of growth regulation during amino acid limitation,

we uncover how methionine uniquely mediates an anabolic transformation. Unexpectedly,

methionine availability during challenging nutrient conditions results in cells co-opting

Gcn4p, a key mediator of a nutrient stress/survival response, for anabolism. This mediator is

necessary for the methionine mediated anabolic transformation, and therefore growth. The

methionine mediated anabolic transformation itself is due to an activation of very specific

metabolic nodes. When these nodes are coincidently activated, they consequently permit a

cascade of events leading to the overall biosynthesis of "costly" amino acids and nucleotides.

These results position methionine at an apex of an overall anabolic network, and provide a

metabolic logic to how methionine results in cells committing to growth.

Results

Methionine induced growth requires Gcn4p.

When wild-type, prototrophic yeast cells are shifted from a complex, amino acid rich medium with lactate as the sole carbon source (RM), to a synthetic minimal medium containing nitrogen base and lactate (MM), they show a significant lag phase and slower growth (Figure 1A). Supplementation with all 20 standard amino acids restores growth after this nutrient downshift (Figure S1A) (*Sutter et al., 2013*; *Laxman et al., 2014*). Importantly, methionine supplementation alone substantially increases growth (Figure 1A), comparable to or better than adding all eighteen other non-sulfur amino acids (nonSAAs) together (*Sutter et al., 2013*; *Laxman et al., 2014*). Given that methionine is itself not a good "nutrient source", and adding methionine would not transform the medium to a nutrient replete condition, we wondered how methionine might mediate an anabolic and proliferative transformation. We therefore thought that examining key growth/survival regulating nodes that are known to be altered during growth under nutrient stress (in the presence or absence of methionine) might serve as a good starting point from which to address this question. In the currently understood model for cellular responses to amino acid limitation (Figure 1B), the Gcn2p kinase is activated and phosphorylates eukaryotic initiation factor eIF2α. This results in global translational repression, but selective translation of *GCN4* (or Atf4 in mammals), where the Gcn4 protein controls general amino acid biosynthesis (Figure 1B) (*Hinnebusch, 2005*). This process is well conserved across eukaryotes (*Vattem and Wek, 2004*). By monitoring the activity status of these steps within this model, and introducing methionine as a variable, we reasoned that clues might emerge that will lead us to the core metabolic phenomenon regulated by methionine. We therefore monitored outputs of responses during amino acid limitation (illustrated in Figure 1B), to determine if any of these aspects were unusually altered in the presence of methionine.

We first monitored global translational output by standard polysome profiling (Figure

1C). Expectedly, RM grown cells showed a 3:1 polysome: monosome ratio indicating robust

translation, and cells shifted to MM showed a drastic reduction in the polysome pool, and an

increased monosome peak (polysome:monosome ratio of ~1:1) (Figure 1C, Figure S1B).

Supplementation with either methionine, or all the nonSAAs combined, during the shift to

MM also showed an increase in the monosome pool, and a partial restoration of the polysome

levels resulting in a polysome: monosome ratio of ~1.7:1 (or ~1.3:1, for nonSAAs) (Figure

1C, Figure S1B), suggesting slightly increased translation due to methionine. We also

monitored the phosphorylation status of eIF2α (Figure 1D), and expectedly observed a

substantial increase in the phosphorylation levels of eIF2α after a RM to MM shift. However,

methionine supplementation only marginally reduced the levels of phosphorylated eIF2α, and

levels of phosphorylation remained high (Figure 1D). Collectively, the phosphorylation status

of the eIF2α, and the results obtained with polysome profiling, suggest that methionine has a

modest impact on global translation during amino acid limited growth, but the cells continue

to appear "nutrient stressed", while showing increased proliferation. These observations were

largely in line with expected outcomes.

We next monitored the amounts of endogenous Gcn4p (chromosomally tagged with a

C-terminal HA epitope) after a shift to MM, with and without supplementation of different

amino acids, expecting a decrease in Gcn4p amounts upon the addition of methionine.

Surprisingly, Gcn4p accumulated substantially upon methionine supplementation, compared

to MM, or supplementing all 18 other nonSAAs (Figure 1E). This was independently

confirmed using immunofluorescence based experiments (Figure S1C). We further asked

whether Gcn4p was necessary for the increased growth upon methionine supplementation.

Notably, the *gcn4Δ* cells did not show any increased growth in methionine supplemented

medium, but instead grew comparably to WT cells growing in MM (Figure 1F). As controls,

in all other conditions (lacking methionine, or in RM) the growth of *gcn4Δ* cells was

indistinguishable from the WT cells (Figure S1D). Collectively, these data indicate that

Gcn4p is necessary uniquely for the methionine-mediated growth in these amino acid poor

conditions. Given the established role of Gcn4p as a regulator of metabolic homeostasis, this

also suggested that methionine triggered a core transcriptional response, with a key

component of this response being mediated by the Gcn4p transcription factor. Thus, we

reasoned that dissecting the methionine and Gcn4p mediated transcriptional response might

provide insight into the anabolic transformation mediated by methionine, and allow the

elucidation of a core metabolic response that drives proliferation.

Methionine induces a transcriptional remodelling program driving key anabolic nodes.

These data raise two questions: (i) what is the underlying logic of the anabolic

transformation mediated by methionine?, and (ii) what is the contribution of Gcn4p towards

this anabolic transformation? We started with a comparative RNA-seq approach in order to

address aspects of both questions. In this section we address how methionine transforms cells

into an anabolic state. We performed RNA-seq analysis on distinct sample sets of wild-type

cells- (i) RM grown, or cells shifted to (ii) MM for 2h, (iii) MM+Met for 2h (Met set), and

(iv) MM+nonSAAs for 2h. Transcript reads from the biological replicates showed

exceptional correlation across all conditions (Pearson correlation coefficient, R≥0.99) (Figure

S2). Setting a stringent cut-off, we only considered differentially expressed genes with $\geq \log_2$

1.5 fold changes, and a p-value cut-off $<10^{-4}$ for further analysis. We first compared global

transcription trends in WT cells growing in RM, MM+Met or MM+nonSAAs to MM,

focusing on a unique transcriptional signature in the presence of methionine. We examined

overall global gene expression trends in these conditions (compared to MM), looking at the

distribution of the most induced or downregulated genes (Figure 2A, Figure S3). Strikingly,

broad trends of up- or down- regulated genes exhibited by cells in MM+Met were more

similar to cells in RM, than the cells in MM+nonSAAs (which were nearly indistinguishable

from MM) (Figure 2A, Figure S3 and S4). This suggests that methionine alone is perceived

by cells as a stronger "growth cue" than all non-sulfur amino acids combined, and transforms

cells into a transcriptional state resembling that of rapidly proliferating cells in complex,

amino acid rich media.

We next more closely examined the transcriptional response unique to methionine, by

comparing transcriptomes of cells growing in MM vs MM+Met (the only variable being

methionine). This comparison identified 372 genes, of which 262 genes were upregulated in

the +Met set (Figure 2B, Figure S4, Supplementary file E1). Using gene ontology (GO),

these genes were grouped into related processes (Figure 2C, Figure S5A, Supplementary file

E2). We observed a substantial transcriptional response related to the core translational

machinery. Additionally, GO also grouped multiple induced genes into categories that would

fit generally under "nucleotide metabolism" (Figure 2C). GO broadly fits these under

"purine/pyrimidine" or "nucleobase and nucleotide metabolism", along with the biosynthesis

of secondary metabolites (Figure 2C). All of this would be entirely expected for any cell in a

proliferative state, since proliferation relies on increased translation and replication. Largely,

and unsurprisingly, the GO grouping showed a signature of a cell in a "proliferative state".

This form of GO grouping does not lend any insight into the metabolic logic of the

methionine mediated response, beyond their obvious requirements for growth. This is largely

a limitation of using GO, which builds groups by looking for enriched pathway terms (relying

on multiple genes within a pathway to be overrepresented). However, metabolic regulation

does not require entire pathways to be regulated, since most regulation happens by

controlling key nodes or "rate-limiting steps" in metabolism (*Nelson and Cox, 2017*). We

therefore rebuilt connections from metabolic first principles, in order to examine if there were

unique nodes that were affected, and to determine if any logical hierarchy of metabolic

responses could be built. These groupings will not necessarily be picked up in GO

enrichments, but may be central to understanding the metabolic state. By this reconstruction,

we compartmentalized the metabolic response regulated by methionine into a group of key

biochemical reactions, which could drive (and explain) a majority of the other metabolic

processes upregulated (based on gene expression), as described below.

Only a few genes involved in classical "carbon/carbohydrate metabolism" were

upregulated in the presence of methionine, and strikingly none of them group to glycolysis,

the TCA cycle or gluconeogenesis (Figure 2D). Instead, the genes encoding three key

enzymes of the pentose phosphate pathway (PPP) (*GND1*, *RKI1* and *TKL1*), which regulate

four steps in the PPP, were highly induced in the presence of methionine (Figure 2D).

Furthermore, two other genes (*SOL3* and *TAL1*), which control two other steps in the PPP,

were also induced by methionine (just below the $\log_2$ 1.5 fold cut-off limit we had arbitrarily

set) (Figure 2D). Gnd1p catalyzes the last step in the oxidative arm of the PPP, generating

NADPH and producing ribulose-5-phosphate. Most of the genes of the non-oxidative arm of

the PPP, which make ribose-5-phosphate and other critical intermediates, are also

upregulated (Figure 2D). Additionally, *HXK2*, encoding a hexokinase is highly upregulated

when methionine is present. While this is not classified under the PPP (say by GO grouping),

this enzyme produces glucose-6-phosphate, which is required as the substrate for the first,

rate-limited step of the PPP, so we included it under the PPP in our grouping (Figure 2D).

Thus, only the PPP arm of carbon metabolism was induced by methionine. Second, we noted

that key regulator resulting in the formation of pyridoxal-5-phosphate or PLP (encoded by

SNO1), required for all transamination reactions was induced (Figure 2D). Third, transcripts

of Gdh1p, which regulates the key nitrogen assimilation reaction resulting in the formation of

glutamate (and *GLN1*, which is further required to make glutamine), were highly induced in

methionine (Figure 2D). This reaction itself requires NADPH (coming from the PPP, Figure

2D), and also is critical for the subsequent formation of all other amino acids, and nucleotides

(Figure 2D). Thus, this grouping suggested that methionine induced the PPP-GDH-PLP

nodes in metabolism, which could be central for all the downstream anabolic outputs.

Methionine sets-up a hierarchical metabolic response leading to anabolism

We therefore inspected the other up-regulated genes that could be grouped broadly

under "amino acid biosynthesis", "nucleotide synthesis", and

"oxidoreduction/transamination" categories, particularly focusing on the substrates or co-

factors required for their function. Notably, essentially all enzymes encoded by these

upregulated genes utilized either a PPP intermediate/product, and/or NADPH, and/or

glutamate/glutamine, or combinations of all of these (Figure 3A). We next more closely

examined the steps in these respective biosynthetic pathways that these genes regulated

(coming from Figure 3A), categorizing all steps in the respective amino acid biosynthesis

pathway as either the rate-limiting/initiation step and/or final step in the production of that

amino acid, organizing them by the use of costly and complex precursors or cofactors (Figure

3B). Strikingly, we observed that the methionine induced genes in these pathways regulated

the critical, rate-limiting or costly steps in amino acid biosynthesis (p-value of $3.8e^{-09}$,

Fisher's exact test), but had little or no significant role in regulating the multiple other genes

in the pathway that encode enzymes for the other, inexpensive steps (Figure 3B). Finally,

these methionine-induced genes in the "amino acid biosynthesis" bin do not just broadly

represent all amino acid biosynthesis, but synthesize what are viewed as the costliest amino

acids to synthesize (*Barton et al., 2010*), namely the aromatic amino acids, the branched-

chain amino acids, and lysine, which is itself highly overrepresented in ribosomal and core

translational machinery proteins (*Laxman et al., 2013*). Similarly, nucleotide biosynthesis

involves very large, multi-step pathways. The methionine dependent, upregulated genes

encoded enzymes that control only a few of the critical, limiting steps in nucleotide synthesis,

and utilize glutamate/glutamine and/or NADPH, as well as pentose sugars from the PPP

(Figure 3A). Furthermore, *RNR1*, which encodes the key enzyme in converting

ribonucleotides to deoxyribonucleotides (and hence the critical hub for DNA synthesis), is

strongly upregulated upon methionine addition (Figure 3A). Separately, we expanded this

analysis and compared the methionine response to minimal medium supplemented with all

other nonSAAs, and the overall metabolic grouping or organization remained unchanged

(Figure S4), with nonSAAs resembling MM. Finally, in MM+nonSAAs, the few highly

induced genes (compared to RM) function in methionine (and sulfur-amino acid) related

biosynthesis or salvage (Figure S5B, Supplementary file E1), and not additional anabolic

reactions. This further supports our overall observations for the role of methionine as an

"anabolic signal". Collectively, this comparative transcriptome analysis suggests not just a

general anabolic remodelling due to methionine, but a hierarchical metabolic organization

induced by methionine (Figure 3C). In this hierarchical organization, methionine induces

genes regulating the PPP, key transamination reactions and the synthesis of

glutamine/glutamate. These three processes directly allow critical steps in synthesis of the

costliest amino acids and nucleotides, and these key steps are also induced by methionine,

collectively setting up a structured anabolic program (Figure 3C). These data thus uniquely

position methionine as an anabolic cue.

Gcn4p regulates the metabolic response due to methionine

We next addressed the contribution of Gcn4p towards this methionine mediated

transformation, at the level of transcription. We first examined global trends of gene

expression (similar to those in Figure 2A) in the absence of Gcn4p, compared to WT cells.

For this, a relative comparison of transcriptomes of *gcn4Δ* cells grown in RM, MM,

MM+Met and MM+nonSAAs to wild-type cells in MM was carried out. To our surprise,

global gene expression trends in the Met set even more strongly resembled the RM set than

WT cells (in methionine, from Figure 2A) (Figure 4A). The transcriptional response in all

other conditions was almost unaffected (Figure 4A). This suggested that, unexpectedly, in the

absence of *GCN4*, methionine invokes a very strong transcriptional response with global

trends seemingly resembling a strong "growth state" (Figure 4A). We more closely compared

transcriptomes of WT cells with *gcn4Δ* cells, under the same combination of conditions used

earlier, and analyzed our data with the same stringent filters used in the previous section. The

overall changes in transcriptomes of WT vs *gcn4Δ* cells are shown (Figure 4B, Figure S6,

Supplementary file E1). Notably, in all other conditions except methionine, WT and *gcn4Δ*

cells showed similar gene expression profiles (Figure 4B). To dissect the role of Gcn4p in the

methionine response, we compared relative transcript amounts, for the transcripts induced by

methionine, in cells lacking Gcn4p compared to MM. When grouped using GO, we found

that a very large number of genes (~200) induced in the presence of methionine, grouped into

the general groups of "ribosome/translation", and "nucleotide synthesis" (Figure 4C,

Supplementary file E2). Again, this representation was even more striking than that seen in

WT cells (Figure 2C), suggesting a strong "growth signature" in the presence of methionine

in cells lacking Gcn4p.

Note that (as shown earlier in Figure 1F) Gcn4p was essential for the growth

induction due to methionine. These data therefore paradoxically suggested that the overall

"growth signature" response due to methionine was being amplified in the absence of Gcn4p,

which might mean that Gcn4p only directly regulated a small number of critical steps in this

growth program. We therefore more carefully compared transcript amounts from WT and

gcn4Δ cells, growing in the presence of methionine (+met), and separated our functional

groupings (based on GO) into two broad "bins". One bin represented all genes related to

ribosome function and translation (as seen in Figure 2C), while the other bin included the

core metabolic genes (similar to those in Figures 2 and 3) (Figure 4D). Strikingly, transcripts

of every gene that mapped to ribosome/translation function (from Figure 2C) was higher in

gcn4Δ cells compared to WT cells in the presence of methionine (Figure 4D). Contrastingly,

every gene related to amino acid biosynthesis (and related metabolism) was significantly

lower in the *gcn4Δ* cells (Figure 4D). Thus, this organization revealed that in the presence of

methionine, while the induction of the translation machinery, and many "nucleotide

synthesis" genes was not Gcn4p- dependent, the entire core metabolic transformation might

hinge upon Gcn4p.

Therefore, we carefully examined the methionine- and Gcn4p- dependent

transcription of genes that functionally regulated the PPP, key transamination reactions, or

the specific nodes in amino acid biosynthesis (Figure 4E). Strikingly, the genes from the PPP,

and transamination reactions were strongly downregulated in *gcn4Δ* cells compared to WT

cells in methionine (Figure 4E). Gdh1 (required for glutamate synthesis) was induced by

methionine, but was not Gcn4p dependent (Figure 4E). Next, the transcripts of genes

encoding key, rate-limiting steps in multiple branched-chain or aromatic amino acid, and

lysine and arginine biosynthesis were induced by methionine in a Gcn4p-dependent manner

(Figure 4). Finally, while most nucleotide biosynthesis genes were not Gcn4p- dependent, the

entire RNR complex (critical for NTP to dNTP conversion) was strongly methionine- and

Gcn4p- dependent (Figure 4E). Collectively, these data suggest that the anabolic program

induced by methionine requires Gcn4p.

Methionine stimulates key transamination and reductive biosynthetic processes in amino acid biosynthesis.

We therefore wanted to more directly examine the extent of (i) methionine controlling this anabolic transformation, through the PPP, reductive biosynthesis coupled reactions, transamination reactions and glutamine/glutamate synthesis feeding subsequent amino acid, cofactor, or nucleotide synthesis; and (ii), the extent of Gcn4p dependence for these steps. We first tested these possibilities at the biochemical level, comparing the actual enzyme amounts of three key targets that represent the coupling of the PPP with other processes. These are Snz1p, Gnd2p and Gdh1p. Snz1p is required for pyridoxal phosphate (PLP) biosynthesis (*Dong et al; 2004*), and PLP is essential for all transamination reactions (*Nelson and Cox, 2017*). As illustrated earlier in Figure 2D, PLP biosynthesis itself requires the PPP intermediate erythrose-4-phosphate as a substrate. Gnd2p is the key NADPH generating enzyme in the oxidative branch of the PPP. Gdh1p consumes NADPH and makes glutamate from 2-ketoglutarate. We measured amounts of these enzymes from WT and *gcn4Δ* cells growing in MM, and MM+methionine (Figure 5A). Notably, Snz1p and Gnd2p showed a strong induction requiring both methionine and Gcn4p (Figure 5A, Figure S7). Gdh1p was strongly dependent upon methionine, but was not on Gcn4p (Figure 5A). We next measured *in vitro* Gdh1p activity (NADPH-GDH activity) in lysates from cells growing in MM or with methionine, and found that overall Gdh1p activity was higher in cells growing with methionine (Figure 5B). However, as seen in the Western blot analysis (Figure 5A), the *in vitro* Gdh1p activity was not dependent on Gcn4p (not shown). These data support the observation that methionine drives these key, coupled steps in biosynthesis, and that they are largely mediated by Gcn4p. Finally, an expected crude readout of this biochemical coupling

should be changes in steady-state nucleotide amounts, requiring Gcn4p in methionine replete

conditions. Comparing relative amounts of nucleotides in wild-type and *gcn4Δ* cells, we

noted a decrease in the nucleotide amounts in *gcn4Δ* cells in the presence of methionine

(Figure 5C). Collectively, direct, steady-state biochemical read-outs support the hypothesis

emerging from the transcriptional response seen to methionine, suggesting a coupled

regulation of the PPP, and key transamination reactions, leading to increased amino acid and

nucleotide synthesis.

Methionine stimulates *de novo* biosynthesis of key amino acids in a Gcn4p- dependent

manner.

Validating these findings using metabolite measurements is challenging, since any

steady-state metabolite measurement (as in Figure 5C) will undoubtedly show differences

between cells, but cannot directly distinguish synthesis from consumption. Therefore, to

directly resolve this, we resorted to a stable-isotope based pulse labelling and LC-MS/MS

based approach to directly measure the new synthesis of amino acids. To WT or *gcn4Δ* cells

in the respective medium with or without methionine, we pulsed ^{15}N labelled ammonium

sulfate, and measured the ^{15}N incorporation into amino acids (Figure 6A, Table S2), before

an effective "steady-state" of labelled amino acid synthesis and consumption was reached.

This will permit the detection of newly synthesized amino acids, which will have

incorporated the ^{15}N label, before a steady-state is reached. We observed that biosynthesis of

all the aromatic amino acids, lysine, histidine, proline, arginine and asparagine is strongly

dependent on methionine (Figure 6B). For technical reasons it was not possible to measure

label-incorporation into branched-chain amino acids. Notably, the label immediately

percolated in asparagine and aromatic amino acid biosynthesis, and showed a very strong

methionine dependence (Figure 6B). Asparagine, proline and phenylalanine biosynthesis

were methionine dependent even in *gcn4Δ* cells, pointing towards some *GCN4*-independent

roles of methionine. For all the other amino acids measured, their biosynthesis increased in

the presence of methionine in a *GCN4*-dependent manner (Figure 6B). These data directly

validate our earlier observations, strongly suggesting that methionine availability controls the

key nodes around PPP-PLP-GDH axis thereby generating the amino acid pool required for

proliferation.

Methionine increases nucleotide biosynthesis in a Gcn4p- dependent manner.

Given that the PPP and amino acid biosynthesis (leading to nucleotide biosynthesis)

are directly regulated by methionine and Gcn4p, we predicted that collectively, perturbing

this node should have a severe consequence on nucleotide biosynthesis, even if most

nucleotide biosynthesis genes themselves were not regulated by Gcn4p. We therefore adopted

a direct, quantitative estimation of methionine and Gcn4p dependent increases in nucleotide

synthesis (similar to the approach in Figure 6). This would directly reflect flux through the

coupled steps of the PPP, glutamate/glutamine synthesis, and the use of intermediates from

amino acid biosynthetic pathways for carbon and nitrogen assimilation into nucleotides

(Figure 7A). The carbon skeleton of nucleotides comes from the PPP, the nitrogen base is

directly derived from glutamine/glutamate and aspartate, and glutamate synthesis is itself

coupled to NADPH (from the PPP) (Figure 7A). Furthermore, nucleotide biosynthesis is also

coupled to histidine and tryptophan synthesis. Using a stable-isotope based nitrogen or

carbon pulse labelling approach, coupled to targeted LC-MS/MS based measurement of

nucleotides, we separately measured the incorporation of the nitrogen and carbon label into

nucleotides, to estimate newly synthesized nucleotides before homeostasis was reached, as

illustrated in Figure 7B and 7C. We observed a dramatic increase in ^{15}N-labelled nucleotides

upon the addition of methionine (Figure 7B, Table S2). Furthermore, this methionine-

mediated incorporation of ^{15}N-label in nucleotides was entirely Gcn4p- dependent (Figure 7B

and Figure S8). Monitoring carbon flux is extremely challenging in a non-fermentable carbon

source like lactate (as compared to glucose), given the difficulties of following the labelled

carbon molecules. Despite that, like the ^{15}N-labeling experiments described above, a similar

experimental design of 0.5X-label spiking was adopted to measure the ^{13}C-label

incorporation into AMP (Figure 7C, Table S2). We observed a significant increase in ^{13}C-

labelled AMP upon the addition of methionine, and this methionine dependent incorporation

of ^{13}C-label in AMP was not observed in cells lacking Gcn4p (Figure 7C). Collectively, this

shows a methionine and Gcn4p dependent increase in *de novo* synthesis of nucleotides,

coupling carbon and nitrogen flux that is dependent on the PPP, and glutamate synthesis.

Discussion

In this study, we show how methionine availability drives cellular proliferation

through an anabolic transformation, even under amino acid limited, apparently challenging

conditions. We uncover a regulated, hierarchical activation of metabolic processes in the

presence of methionine, which lead to an anabolic state. We also provide a mechanism of

how methionine mediates this anabolic transformation.

Starting with a framework of existing models for responses to amino acid limitation,

we find that the overall methionine mediated growth depends upon Gcn4p (Figure 1).

Surprisingly, methionine results in increased Gcn4p, and Gcn4p is necessary for the

methionine mediated anabolic transformation (Figure 1). More pertinently, methionine,

substantially through Gcn4p, mediates a transcriptional remodelling in cells controlling the

anabolic transformation, and not the translation component (Figures 2, 4). Understanding the

core metabolic logic within the transcriptional response to methionine required a

reorganization of GO based groupings based on metabolic first principles, and the logic that

emerged was striking (Figure 2, 3 and 4). Methionine positively regulates the pentose

phosphate pathway (PPP) (Figure 2). The PPP provides the pentose sugar backbones for

nucleotides, along with reducing equivalents (NADPH) allowing reductive biosynthesis for a

variety of anabolic molecules (*Nelson and Cox, 2017*). Key among these are amino acids and

nucleotides. For amino acid and nucleotide synthesis, pyridoxal phosphate, which controls all

transamination reactions, is essential (*Eliot and Kirsch, 2004*) (Figure 2). Notably, genes

controlling these three key nodes in metabolism are all strongly induced by methionine

(Figure 2). These steps feed the subsequent metabolic nodes induced by methionine.

Methionine induces the expression of genes that control rate limiting or final steps in the

synthesis of "costly" amino acids, the aromatic amino acids and the branch chain amino acids

(Figure 3). Almost every one of these (methionine regulated) steps use cofactors or

intermediates from the pentose phosphate pathway, are transamination reactions requiring

PLP, or/and require glutamate or glutamine (which depends on the PLP mediated

transamination) (Figure 2 and 3). Thus, it appears that methionine sets up this metabolic

hierarchy, as illustrated in a schematic in Figure 3C and 7D. Interestingly, the methionine-

induced activity in most of these metabolic nodes, and thus the overall anabolic

transformation, depends upon Gcn4p (Figure 4). Gcn4p is best understood as a regulator of

amino acid biosynthesis during starvation (*Natarajan et al., 2001*). However, while many of

the *GCN4* targets picked up in our study compare well with the landmark study of *GCN4*

targets (*Natarajan et al., 2001*) (see Figure S9 and Table S3), the role of Gcn4p in the

presence of methionine in controlling these key hubs of the PPP, glutamate biosynthesis, and

rate regulating steps in costly amino acid and nucleotide biosynthesis has not been previously

appreciated, particularly in contexts where growth is enhanced. A combination of rigorous

biochemical, and metabolic flux based analysis using stable-isotopes directly demonstrate

this coupling of the PPP, NADPH utilization and transamination reactions (in both nitrogen

assimilation and carbon assimilation) first towards the increased synthesis of aromatic and

branch-chain amino acids, and next towards nucleotides, in a methionine- and Gcn4p-

dependent manner (Figures 6, 7). Interestingly, Gcn4p appears to regulate only the core

metabolic program induced by methionine, and not the induction in the synthesis of the

translation machinery (as seen in Figure 4). Collectively, our data not just shows how

methionine can co-opt Gcn4 to achieve a "growth state", but suggests how methionine sits

atop a pyrimidical hierarchy of metabolic events, that set up an anabolic transformation

(model in Figure 3C).

The central role of the PPP in anabolism is now text-book knowledge (*Nelson and*

Cox, 2017). Yet, a better appreciation of the importance of the PPP in mediating anabolic and

proliferative transformations is now emerging due to the association of the PPP to cancer

metabolism (*Patra and Hay, 2014*; *Cairns et al., 2011*). Many anabolic transformations require contributions from the PPP, however, the metabolic cues regulating the PPP (and coupling to other processes) are not often obvious. More generally, our study highlights how methionine (likely through its downstream metabolite SAM), acts as an anabolic signal for cells. This is analogous to another central metabolite, acetyl-CoA, better known to determine cellular decisions towards growth (*Cai et al., 2011*; *Comerford et al., 2014*; *Shi and Tu, 2013*; *Pietrocola et al., 2015*; *Mariño et al., 2014*). Interesting correlations can be made from our observations to known roles of methionine in cancer cell metabolism, and metazoan growth. The earliest observations of methionine as important for some cancers to proliferate dates back to the 1950s (*Cavuoto and Fenech, 2012*; *Sugimura et al., 1959*; *Breillout et al., 1990)*, and several types of cancer cells are addicted to methionine (*Halpern et al., 1974*; *Stern and Hoffman, 1986*; *Guo et al., 1993*; *Lu and Epner, 2000*; *Poirson-Bichat et al., 2000*; *Kokkinakis et al., 2001*; *Cellarier et al., 2003*; *Cavuoto and Fenech, 2012*; *Clarke et al., 2016*). Other, distinct studies show that *Drosophila* fed on methionine rich diets exhibit rapid growth, high fecundity, and shorter lifespans (*Troen et al., 2007*; *Lee et al., 2016, 2014*), all hallmarks of what a "proliferative" metabolite will do. Studies from yeast show how methionine inhibits autophagy, or regulates the TORC1 to boost growth (*Sutter et al., 2013*; *Laxman et al., 2013, 2014*). One of the earliest known cell cycle entry check-points is linked to methionine (*Unger and Hartwell, 1976*). Upon sulfate (and thereby methionine) starvation, cells arrest their growth to promote survivability (*Boer et al., 2008*), and transform their proteome to preferentially express proteins containing fewer cysteine/methionine residues to save sulfur (*Fauchon et al., 2002*). There are other, less appreciated observations connecting methionine metabolism and the pentose phosphate pathway. Yeast cells lacking *ZWF1* (encoding glucose-6-phosphate dehydrogenase, the first enzyme in the PPP) exhibit methionine auxotrophy (*Thomas et al., 1991*), and methionine supplementation increases the

oxidative stress tolerance of *zwf1Δ* (*Campbell et al., 2016*). While these studies highlight this

critical role of methionine in anabolism, an overarching logic of the core anabolic

transformation mediated by methionine, and the mechanisms by which this is mediated, have

remained unclear. Our study directly elucidates this.

Note that our study emphasizes the use of prototrophic ("wild type") yeast strains to

study responses to amino acids. Typically, studies utilize laboratory strains derived from an

auxotrophic backgrounds (eg. S288C/BY4741), which require supplemented uracil, histidine,

leucine and methionine for survival (*Sherman, 1991*; *Pronk, 2002*; *Corbacho et al., 2011*;

Brachmann et al., 1998; *Gerashchenko and Gladyshev, 2014*; *Alam et al., 2016*). This

precludes experiments in amino acid limitation. Additionally, our use of a "less-preferred"

carbon source, lactate, can help reveal regulatory phenomena otherwise hidden in glucose and

amino acid rich laboratory conditions, where a surfeit of costly metabolic resources are

present. Notably, several recent reports emphasize the importance of lactate as a carbon

source in rapidly proliferating cells (*Kennedy et al., 2013*; *Faubert et al., 2017*; *Hu et al.,*

2017), and our observations provide mechanistic and molecular insight into growth under

these conditions. Finally, the so called integrated stress response (ISR), and the Gcn4p

ortholog in mammals, Atf4, play important roles in cancer cell proliferation (*Bi et al., 2005*;

Ye et al., 2010; *Palam et al., 2015*), and many cancers continue to grow in apparently poor

nutrient environments. Our study suggests how, in methionine rich (but otherwise amino acid

limiting) conditions, Gcn4/Atf4 could function to promote growth, and not just help cells

recover from nutrient stress. A separate emerging question will be to understand how Gcn4p

is itself regulated under these otherwise amino acid limited conditions, by methionine.

We close by reiterating a possible metabolic cost based hypothesis for what might

make methionine a strong growth cue. The *de novo* synthesis of methionine and its

immediate metabolites (notably SAM) is exceptionally costly in terms of NADPH molecules

invested (*Thomas and Surdin-Kerjan, 1997*; *Kaleta et al., 2013*; *Mampel et al., 2005*; *Sutter*

et al., 2013). Cells require at least 6 molecules of NADPH to reduce sulfur to synthesize a

single molecule of methionine. Since biology has tied multiple anabolic processes to

reductive biosynthesis (dependent on NADPH from the PPP), the availability of methionine

might be an ancient signal to represent a metabolic state where reductive equivalents are

sufficiently available for all other reductive biosynthetic processes as a whole.

Materials and Methods:

Yeast strains and growth media

The prototrophic CEN.PK strain (referred to as wild type, WT) was used in all experiments (*van Dijken et al., 2000*). Strains with gene deletions or chromosomally tagged proteins (at the C-terminus) were generated as described. Strains used in this study are listed in Table S1.

The growth media used in this study are RM (1% yeast extract, 2% peptone and 2% lactate) and MM (0.17% yeast nitrogen base without amino acids, 0.5% ammonium sulfate and 2% lactate). All amino acids were supplemented at 2 mM. NonSAAs refers to the mixture of all standard amino acids (2 mM each) except methionine, cysteine and tyrosine.

The indicated strains were grown in RM with repeated dilutions (~36 hours), and the culture in the log phase (absorbance at 600 nm of ~1.2) was subsequently switched to MM, with or without addition of the indicated amino acids. For growth curves, the RM acclimatized cultures were used and diluted in a fresh medium with the starting absorbance of ~0.2 and the growth was monitored at the indicated time intervals.

Polysome analysis

The cells (50 OD) were treated with cycloheximide (100 μg/ml) for 15 minutes before harvesting in ice-filled, cold centrifuge bottles. The pellet obtained after centrifugation (8000g, 5 min, 4°C) was once washed, and then re-suspended in 0.5 ml lysis buffer (10 mM Tris-HCl pH 7, 0.1 M NaCl, 30 mM $MgCl_2$, 1U/μl Ribolock RNase inhibitor and 80 μg/ml cycloheximide) with chilled, baked acid-washed glass beads. After bead-beating, 0.5 ml lysis buffer was added and the crude extract obtained was centrifuged (18407g, 10 min, 4°C) and the supernatant (200 μl) containing polysomes is fractionated on a sucrose gradient (7%-47% sucrose; containing 20 mM Tris-HCl pH 7, 140 mM KCl, 5 mM $MgCl_2$, 1 mM DTT and 50

515 μg/ml cycloheximide) by ultra-centrifugation (35000 rpm, 3 h, 4°C). The polysome profile

516 was obtained on an ISCO fractionator, wherein the absorbance at 254 nm was traced. For

517 polysome: monosome ratio calculation, area under the curve was quantified using ImageJ

518 software (*Rasband, ImageJ*, U. S. National Institutes of Health, Bethesda, Maryland, USA,

519 http://imagej.nih.gov/ij/, 1997-2016).

Western blot analysis

Approximately ten OD_{600} cells were collected from respective cultures, pelleted and flash-

frozen in liquid nitrogen until further use. The cells were re-suspended in 400 μl of 10%

trichloroacetic acid and lysed by bead-beating three times: 30 sec of beating and then 1 min

of cooling on ice. The precipitates were collected by centrifugation, re-suspended in 400 μl of

SDS-glycerol buffer (7.3% SDS, 29.1% glycerol and 83.3 mM Tris base) and heated at

100°C for 10 min. The supernatant after centrifugation was treated as the crude extract.

Protein concentrations from extracts were estimated using bicinchoninic acid assay (Thermo

Scientific). Equal amounts of samples were resolved on 4 to 12% Bis-Tris gels (Invitrogen).

Coomassie blue–stained gels were used as loading controls. Western blots were developed

using the antibodies against the respective tags. We used the following primary antibodies:

monoclonal FLAG M2 (Sigma), HA (12CA5, Roche), and phosphor-eIF2α Ser51 (9721, Cell

Signaling). Horseradish peroxidase–conjugated secondary antibodies (mouse and rabbit)

were obtained from Sigma. For Western blotting, standard enhanced chemiluminescence

reagents (GE Healthcare) were used. ImageJ was used for quantification.

Immunofluorescence measurements

Yeast cells were fixed with 3.7% formaldehyde, washed and resuspended in spheroplasting

buffer (40 mM potassium phosphate buffer, pH 6.5; 0.5 mM $MgCl_2$; 1.2 M sorbitol).

Spheroplasts were prepared by zymolyase (MP Biomedicals, 08320921) treatment and spread

on a slide pretreated with 50 μl of 1 mg/ml polylysine (Sigma-Aldrich, P6407). Gcn4-HA

was stained with the mouse monoclonal anti-HA (12CA5) primary antibody (Roche,

11583816001) and Alexa Fluor 488-conjugated Goat anti-Mouse IgG (H+L) secondary

antibody (Thermofisher, A32723). DNA was stained with 1μg/ml DAPI for 2 minutes,

washed and mounted in Fluoromount-G (Southern Biotech, 0100-01). The cells were imaged

using Olympus FV1000 confocal microscope.

RNA-seq analysis

Total RNA from yeast cells was extracted using hot acid phenol method (*Collart and*

Oliviero, 2001). The quality of RNA was checked on Bioanalyzer using an RNA 6000 Nano

kit (Agilent) and the libraries were prepared using TruSeq RNA library preparation kit V2

(Illumina). The samples were sequenced on Illumina platform HiSeq2500. The raw data is

available with NCBI-SRA under the accession number SRP101768. Genome and the

annotation files of *S. cerevisiae* S288C strain were downloaded from Saccharomyces

Genome Database (SGD; http://www.yeastgenome.org/). 100-mer, single-end reads obtained

from RNA sequencing experiments were mapped to the S288C genome using Burrows

Wheeler Aligner (BWA) (*Li and Durbin, 2009*). Mapped reads with the mapping quality of $\geq$

20 were used for the further analysis. The number of reads mapped to each gene was

calculated and the read count matrix was generated. The read count matrix was fed into

EdgeR, a Bioconductor package used for analyzing differential gene expression (*Robinson et*

al., 2010). Genes which are differentially expressed by at least 3-fold with the p-value of

<0.0001 were considered for further analysis. Normalized gene expression was calculated by

dividing the number of reads by the gene length and the total number of reads for those

samples, then dividing each of these values with the mode of its distribution (*Srinivasan et*

al., 2013). Absolute expression levels of the genes between the replicates are well correlated

with the Pearson correlation coefficient (R) values more than 0.99 (see Figure S2). Mapping

of genes to the related pathways and gene ontology analysis were carried out using public

databases such as Yeastcyc (*Caspi et al., 2014*), GeneCodis (*Tabas-Madrid et al., 2012*;

Nogales-Cadenas et al., 2009; *Carmona-Saez et al., 2007*) and SGD (*Dwight et al., 2002*).

Metabolite extractions and measurements by LC-MS/MS

Cells were grown in RM for ~36 hours and transferred to MM with and without methionine

for the indicated time. After incubation, cells were rapidly harvested and metabolite extracted

as described earlier (*Tu et al., 2007*). Metabolites were measured using LC-MS/MS method

as described earlier (*Laxman et al., 2014*). Standards were used for developing multiple

reaction monitoring (MRM) methods on Thermo Scientific TSQ Vantage triple stage

quadrupole mass spectrometer or Sciex QTRAP 6500. For positive polarity mode,

metabolites were separated using a Synergi 4μ Fusion-RP 80A column (150 × 4.6 mm,

Phenomenex) on Agilent's 1290 infinity series UHPLC system coupled to mass spectrometer.

Buffers used for separation were: buffer A: 99.9% H_2O/0.1% formic acid and buffer B:

99.9% methanol/0.1% formic acid (Flow rate, 0.4 ml/min; T = 0 min, 0% B; T = 3 min, 5%

B; T = 10 min, 60% B; T = 10.1 min, 80% B; T = 12 min, 80% B; T = 14 min, 5% B; T = 15

min, 0% B; T = 20 min, stop). For negative polarity mode, metabolites were separated using

a Luna HILIC 200A column (150 × 4.6 mm, Phenomenex). Buffers used for separation were:

buffer A: 5 mM ammonium formate in H_2O and buffer B: 100% acetonitrile (flow rate: 0.4

ml/min; T = 0 min, 95% B; T = 1 min, 40% B; T = 7 min, 10% B; T = 11 min, 1% B; T = 13

min, 95% B; T = 17 min, stop). The area under each peak was calculated using Thermo

Xcalibur software (Qual and Quan browsers).

^{15}N- and ^{13}C- based metabolite labelling experiments

For detecting ^{15}N-label incorporation in amino acids and nucleotides, ^{15}N-ammonium sulfate

with all nitrogens labelled (Sigma- Aldrich) was used. For ^{13}C-labeling experiment, ^{13}C-

lactate with all carbons labelled (Cambridge Isotope Laboratories) was used. In the labelling

experiments, 0.5X refers to 0.25% ammonium sulfate or 1% lactate. All the parent/product

masses measured are enlisted in Table S2. Amino acid measurements were done in the

positive polarity mode. For all the nucleotide measurements, release of the nitrogen base was

monitored in positive polarity mode. For the ^{13}C-label experiment, the phosphate release was

monitored in negative polarity mode. Under these conditions, the nitrogen base release cannot

be monitored here as the nitrogen base itself has carbon skeleton, which will complicate the

analysis. The HPLC and MS/MS protocol was similar to those explained above.

GDH assays

Glutamate dehydrogenase activity was measured as described in (*Doherty, 1970*), with some

modifications. Yeast cells were lysed by bead-beating in lysis buffer (100 mM potassium

phosphate buffer, pH 7; 5% glycerol; 1 mM PMSF; 0.1% Tween-20; 1 mM EDTA; 1 mM 2-

mercaptoethanol). NADP-dependent activity was measured by monitoring oxidation of

NADPH (assay buffer: 100 mM Tris-HCl, pH 7.2; 10 mM 2-ketoglutarate, pH adjusted to

7.2; 100 mM ammonium chloride; 0.1 mM NADPH) at 340 nm. Protein concentrations from

extracts were estimated using bicinchoninic acid assay (Thermo Scientific). One enzyme unit

corresponds to the amount of enzyme required to oxidize one µmol of NADPH min^{-1} at room

temperature.

Statistical analysis

In most of the experiments, Student's t-test was applied for calculating the p-values.

Wherever necessary, other tests were applied and indicated accordingly.

Acknowledgements:

We acknowledge Dhananjay Shinde, Padma Ramakrishnan and the NCBS/inStem/C-CAMP

Mass spectrometry facility for LC-MS/MS support, Avadheesh Pandit and the

NCBS/inStem/C-CAMP next-generation sequencing facility for assistance with library

preparation, and Ravi Muddashetty for the polysome fractionator. We thank Utpal Banerjee,

Mark Sharpley, Marco Foiani, Christopher Bruhn, Krishnamurthy Natarajan and PJ Bhat for

critical comments on this manuscript. This work was supported by a Wellcome Trust-DBT

IA intermediate fellowship (IA/I/14/2/501523) and inStem/DBT institutional support to SL.

ASW, RG, RS acknowledge bridging fellowships (from inStem), and ASW, RG and RS are

supported by DST SERB-national postdoctoral fellowships (PDF/2015/000225,

PDF/2016/000416 and PDF/2016/001877 respectively).

Figure legends:

Figure 1: Methionine induced growth requires Gcn4p.

A) Methionine and growth during amino acid limitation. Shown are growth profiles of WT cells grown in RM and shifted to MM with or without the indicated amino acid supplements (2 mM each; nonSAAs indicates all the non-sulfur amino acids except tyrosine). The growth profile with methionine is in blue. Also see Figure S1A.

B) The current model explaining cellular responses to amino acid starvation. Amino acid starvation activates the Gcn2p kinase, leading to decreased levels of the ternary complex. The key outputs indicating amino acid starvation (highlighted in green) are increased phosphorylation of eIF2α, global translational repression, and induced Gcn4p translation.

C) Global translational repression is moderately rescued by methionine. WT cells were grown to mid-log phase in RM and shifted to MM, MM plus methionine or MM plus nonSAAs for two hours. The cells were treated with cycloheximide for 15 min before harvesting and polysome pool was fractionated on sucrose gradients using ultracentrifugation. The polysome/monosome ratio for the corresponding samples is shown. Also see Figure S1B.

D) Methionine and eIF2α phosphorylation. Yeast cells grown in RM were shifted to the indicated media. The samples were analysed for phosphorylation status of eIF2α (Sui2p in yeast) by Western blot. Rapamycin (100 nM) treated cells served as a positive control. WT cells expressing FLAG-tagged eIF2α were used to monitor the protein abundance under identical conditions. A representative blot is shown.

E) Gcn4p amounts in the presence of methionine during amino acid limited growth. Gcn4p was detected by Western blotting of WT cells (expressing Gcn4p with an HA epitope, tagged at the endogenous locus) shifted from RM to MM, or MM supplemented with the indicated combinations of amino acids. RM grown cells treated with rapamycin (100 nM) served as a positive control for Gcn4p induction. A representative blot is shown.

F) *gcn4Δ* cells and methionine-mediated growth. The WT and *gcn4Δ* strains were shifted from RM to MM with or without methionine supplementation and growth was monitored. Also see Figure S1D.

In all panels, data shows mean±SD. $^{*}p < 0.05$.

Figure 2: Methionine induces a transcriptional remodelling program driving key anabolic nodes.

A) Global trends of gene expression in RM and methionine supplemented MM. Boxplot shows gene expression levels of transcripts in WT cells grown in RM, or shifted to MM plus methionine, or MM plus nonSAAs in comparison to MM set.

B) Effect of methionine on a transcriptional program in cells. The heat map shows differentially expressed genes in MM plus methionine and MM plus nonSAAs compared to MM set.

C) GO analysis of the methionine-responsive genes. The pie-chart depicts the processes grouped by GO analysis for the up-regulated transcripts between MM plus methionine and MM set. Numbers in the bracket indicate the number of genes from the query set/ total number of genes in the reference set for the given GO category.

D) Manual regrouping of the methionine responsive genes into relevant metabolic pathways, focusing on carbon metabolism and central metabolic processes. The pathway map includes individual genes in central carbon metabolism which are induced by methionine (indicating the fold-changes in gene expression). The arrows marked red indicate increased expression across the pathway.

Figure 3: Methionine sets-up a hierarchical metabolic response leading to anabolism.

A) Regrouping of the methionine responsive genes in amino acid and nucleotide metabolism onto their respective pathways. The schematic shows the methionine-responsive genes in various amino acid and nucleotide biosynthesis pathways, along with their fold-changes in gene expression. The requirement of PPP metabolites, NADPH, PLP or glutamate/glutamine for the individual steps is mapped on to the schematic. The arrows marked red are the steps induced in the presence of methionine. Note that all these regulated steps use PPP intermediates, NADPH, PLP and/or glutamine/glutamate.

B) A bird's eye-view of the amino acid biosynthesis steps regulated by methionine. Each bead (or filled circle) represents a step in the pathway (prepared according to the individual amino acid pathways shown at https://pathway.yeastgenome.org/.) A step is considered expensive (marked red) when it is either the entry or the final or involves ATP utilization or involves reduction. All the rest of the steps are considered inexpensive (marked green). Methionine responsive steps are shown with a yellow fill at the centre of the circle, for the given step. The p-value for methionine dependence of genes encoding the critical, rate-limiting or costly steps in amino acid= $3.8e^{-09}$, and for the other nodes, it is non-significant (Fisher's exact test).

C) A hierarchical organization of the methionine mediated anabolic remodelling. Methionine induces expression of genes in the PPP-GDH-PLP node, which provides precursors for the overall biosynthesis of amino acids and nucleotides.

Figure 4: Gcn4p uniquely regulates the metabolic response due to methionine.

A) Trends of gene expression in RM and methionine supplemented MM in *gcn4Δ* cells. Gene expression levels of transcripts in *gcn4Δ* cells grown in RM or shifted to MM or MM plus methionine or MM plus nonSAAs were compared to WT MM set.

B) Global transcriptional response in the absence of *GCN4*. The heat map shows differentially expressed genes (log$_2$ 1.5-fold change; $p < 10^{-4}$) between WT and *gcn4Δ* cells in the respective growth conditions.

C) GO analysis of the methionine-responsive genes in *gcn4Δ* cells. A pie chart showing the processes grouped by GO analysis for the up-regulated transcripts between MM plus methionine and MM set in *gcn4Δ* background. Numbers in the bracket indicate the number of genes from the query set/ total number of genes in the reference set for the given GO category.

D) Contribution of *GCN4* to the methionine-response. The expression level of methionine-responsive transcripts in WT set (MM plus methionine versus MM) was compared in the *gcn4Δ* background. Amino acid biosynthesis related genes are marked with blue circles and ribosome biogenesis and function related genes are marked with red circle.

E) Regrouping of the *GCN4*-dependent genes onto metabolic nodes. The schematic shows the *GCN4*-dependent genes (comparison of MM plus methionine set between WT and *gcn4Δ*) in the PPP, amino acid and nucleotide biosynthesis pathways, along with fold-changes in gene expression. The arrows marked blue in the PPP pathway are the steps down-regulated in *gcn4Δ* cells. Key rate-limiting steps in the pathway are marked by asterisk.

Figure 5: Methionine stimulates key transamination and reductive biosynthetic processes.

A) Snz1p, Gnd2p and Gdh1p amounts in WT or *gcn4Δ* cells. WT and *gcn4Δ* cells expressing FLAG-tagged Snz1p or Gnd2p or Gdh1p were shifted from RM to MM or MM plus methionine and amounts of these proteins were detected by Western blotting. A representative blot is shown in each case.

B) NADP-dependent glutamate dehydrogenase activity and methionine availability. Crude extracts of WT cells grown in RM and shifted to MM or MM plus methionine were analysed for intracellular biosynthetic NADP-glutamate dehydrogenase activity.

C) Relative nucleotide amounts in the presence of methionine in WT or *gcn4Δ* cells. WT and *gcn4Δ* cells grown in RM were shifted to MM (4h) with and without methionine, and the relative amounts of AMP and GMP from metabolite extracts of the respective samples were measured by LC-MS/MS.

In all panels data indicate mean±SD. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 6: Methionine stimulates key transamination and reductive biosynthetic processes in amino acid biosynthesis

A) A schematic showing the experimental design of ^{15}N pulse-labelling experiment. Cells were shifted to MM with and without methionine, maintained for 3h, and ^{15}N-ammonium sulfate was pulsed into the medium, and the indicated, labelled metabolites were measured. Also see Table S2.

B) Methionine increases nitrogen assimilation in amino acids. Label incorporation in newly synthesized amino acids in WT and *gcn4Δ* cells was measured, as shown in the panel A. For all the labelled moieties, fractional abundance of the label was calculated. Also see Table S2.

In all panels data indicate mean±SD. *p < 0.05, **p < 0.01.

Figure 7: Methionine increases nucleotide biosynthesis in a Gcn4p dependent manner.

A) A schematic showing carbon and nitrogen inputs in nucleotide biosynthesis, and their coupling to the PPP/NADPH metabolism.

B) Methionine increases nitrogen assimilation in nucleotides. The WT and *gcn4Δ* cells
treated and pulse-labelled with ^{15}N ammonium sulfate as illustrated in the top panel. For all
the labelled moieties, fractional increase of the incorporated label was calculated, to measure
newly synthesized AMP and GMP. (also see Figure S8 for CMP and UMP).

C) Methionine enhances carbon flux into AMP biosynthesis. An experimental set-up similar
to the Panel B was employed, using ^{13}C-lactate for carbon labelling. Label incorporation into
nucleotides (from +1 to +5) was accounted for calculations. (note: GMP could not be
estimated because of MS/MS signal interference from unknown compounds in the metabolite
extract).
750
E) A model illustrating how methionine triggers cell proliferation. Methionine activates the
GCN4-mediated ISR, promoting the synthesis of PPP metabolites, PLP, NADPH and
glutamate (up-regulated genes in the pathways are shown in blue), which directly feed into
nitrogen metabolism. As a result, methionine activates biosynthesis of amino acids and
nucleotides, allowing the cells to grow in amino acid limiting medium.

In all panels data indicate mean±SD. ns: non-significant difference, **p < 0.01, ***p < 0.001.

References:

Alam MT, Zelezniak A, Mülleder M, Shliaha P, Schwarz R, Capuano F, Vowinckel J, Radmanesfahar E, Krüger A, Calvani E, Michel S, Börno S, Christen S, Patil KR, Timmermann B, Lilley KS, Ralser M. 2016. The metabolic background is a global player in *Saccharomyces* gene expression epistasis. *Nat Microbiol* **1**:15030. doi:10.1038/nmicrobiol.2015.30.

Barton MD, Delneri D, Oliver SG, Rattray M, Bergman CM. 2010. Evolutionary systems biology of amino acid biosynthetic cost in yeast. *PLOS ONE* **5**: e11935. doi:10.1371/journal.pone.0011935

Bi M, Naczki C, Koritzinsky M, Fels D, Blais J, Hu N, Harding H, Novoa I, Varia M, Raleigh J, Scheuner D, Kaufman RJ, Bell J, Ron D, Wouters BG, Koumenis C. 2005. ER stress-regulated translation increases tolerance to extreme hypoxia and promotes tumor growth. *EMBO J* **24**:3470-81. doi:10.1038/sj.emboj.7600777.

Boer VM, Amini S, Botstein D. 2008. Influence of genotype and nutrition on survival and metabolism of starving yeast. *Proc Natl Acad Sci U S A* **105**:6930-5. doi:10.1073/pnas.0802601105.

Boer VM, Crutchfield CA, Bradley PH, Botstein D, Rabinowitz JD. 2010. Growth-limiting intracellular metabolites in yeast growing under diverse nutrient limitations. *Mol Biol Cell* **21**:198-211. doi:10.1091/mbc.E09-07-0597.

Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD. 1998. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: A useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**:115-32. doi:10.1002/(SICI)1097-0061(19980130)14:2<115::AID-YEA204>3.0.CO;2-2.

Brauer MJ, Huttenhower C, Airoldi EM, Rosenstein R, Matese JC, Gresham D, Boer VM, Troyanskaya OG, Botstein D. 2008. Coordination of growth rate, cell cycle, stress

response, and metabolic activity in yeast. *Mol Biol Cell* **19**:352-67.

doi:10.1091/mbc.E07-08-0779.

Breillout F, Antoine E, Poupon MF. 1990. Methionine dependency of malignant tumors: A

possible approach for therapy. *J Natl Cancer Inst* **82**:1628-32.

doi:10.1093/jnci/82.20.1628.

Broach JR. 2012. Nutritional control of growth and development in yeast. *Genetics* **192**:73-

105. doi:10.1534/genetics.111.135731.

Cai L, Sutter BM, Li B, Tu BP. 2011. Acetyl-CoA induces cell growth and proliferation by

promoting the acetylation of histones at growth genes. *Mol Cell* **42**:426-37.

doi:10.1016/j.molcel.2011.05.004.

Cairns RA, Harris IS, Mak TW. 2011. Regulation of cancer cell metabolism. *Nat Rev Cancer*

11:85-95. doi:10.1038/nrc2981.

Campbell K, Vowinckel J, Keller MA, Ralser M. 2016. Methionine metabolism alters

oxidative stress resistance *via* the pentose phosphate pathway. *Antioxid Redox Signal*

24:543-7. doi:10.1089/ars.2015.6516.

Carmona-Saez P, Chagoyen M, Tirado F, Carazo JM, Pascual-Montano A. 2007.

GENECODIS: A web-based tool for finding significant concurrent annotations in gene

lists. *Genome Biol* **8**:R3.

Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, Holland TA, Keseler

IM, Kothari A, Kubo A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley

S, Subhraveti P, Weaver DS, Weerasinghe D, Zhang P, Karp PD. 2014. The MetaCyc

database of metabolic pathways and enzymes and the BioCyc collection of

pathway/genome databases. *Nucleic Acids Res* **42**:D459-71. doi:10.1093/nar/gkt1103.

Cavuoto P, Fenech MF. 2012. A review of methionine dependency and the role of

methionine restriction in cancer growth control and life-span extension. *Cancer Treat*

Rev **38**:726-36. doi:10.1016/j.ctrv.2012.01.004.

Cellarier E, Durando X, Vasson MP, Farges MC, Demiden A, Maurizis JC, Madelmont JC,

Chollet P. 2003. Methionine dependency and cancer treatment. *Cancer Treat Rev*

29:489-99. doi:10.1016/S0305-7372(03)00118-X.

Clarke CJ, Berg TJ, Birch J, Ennis D, Mitchell L, Cloix C, Campbell A, Sumpton D, Nixon

C, Campbell K, Bridgeman VL, Vermeulen PB, Foo S, Kostaras E, Jones JL, Haywood

L, Pulleine E, Yin H, Strathdee D, Sansom O, Blyth K, McNeish I, Zanivan S, Reynolds

AR, Norman JC. 2016. The initiator methionine tRNA drives secretion of type II

collagen from stromal fibroblasts to promote tumor growth and angiogenesis. *Curr Biol*

26:755-65. doi:10.1016/j.cub.2016.01.045.

Collart MA, Oliviero S. 2001. Preparation of yeast RNA. In *Curr Protoc Mol Biol*. Hoboken,

NJ, USA: John Wiley & Sons, Inc. doi:10.1002/0471142727.mb1312s23.

Comerford SA, Huang Z, Du X, Wang Y, Cai L, Witkiewicz AK, Walters H, Tantawy MN,

Fu A, Manning HC, Horton JD, Hammer RE, McKnight SL, Tu BP. 2014. Acetate

dependence of tumors. *Cell* **159**:1591-1602. doi:10.1016/j.cell.2014.11.020.

Corbacho I, Teixidó F, Velázquez R, Hernández LM, Olivero I. 2011. Standard YPD, even

supplemented with extra nutrients, does not always compensate growth defects of

Saccharomyces cerevisiae auxotrophic strains. *Antonie Van Leeuwenhoek* **99**:591-600.

doi:10.1007/s10482-010-9530-5.

DeBerardinis RJ, Sayed N, Ditsworth D, Thompson CB. 2008. Brick by brick: Metabolism

and tumor cell growth. *Curr Opin Genet Dev* **18**:54-61. doi:10.1016/j.gde.2008.02.003.

Dechant R, Peter M. 2008. Nutrient signals driving cell growth. *Curr Opin Cell Biol* **20**:678-

87. doi:10.1016/j.ceb.2008.09.009.

Doherty D. 1970. [119] L-Glutamate dehydrogenases (yeast). In , 850-56. doi:10.1016/0076-

6879(71)17294-1.

Dong YX, Sueda S, Nikawa JI, Kondo H. 2004. Characterization of the products of the genes

SNO1 and *SNZ1* involved in pyridoxine synthesis in *Saccharomyces cerevisiae*. *Eur J*

Biochem **271**: 745-52. doi:10.1111/j.1432-1033.2003.03973.x.

Dwight SS, Harris MA, Dolinski K, Ball CA, Binkley G, Christie KR, Fisk DG, Issel-Tarver

L, Schroeder M, Sherlock G, Sethuraman A, Weng S, Botstein D, Cherry JM. 2002.

Saccharomyces Genome Database (SGD) provides secondary gene annotation using the

Gene Ontology (GO). *Nucleic Acids Res* **30**:69-72. doi:10.1093/nar/30.1.69.

Eliot AC, Kirsch JF. 2004. Pyridoxal phosphate enzymes: Mechanistic, structural, and

evolutionary considerations. *Annu Rev Biochem* **73**:383-415.

doi:10.1146/annurev.biochem.73.011303.074021.

Faubert B, Li KY, Cai L, Hensley CT, Kim J, Zacharias LG, Yang C, Do QN, Doucette S,

Burguete D, Li H, Huet G, Yuan Q, Wigal T, Butt Y, Ni M, Torrealba J, Oliver D,

Lenkinski RE, Malloy CR, Wachsmann JW, Young JD, Kernstine K, DeBerardinis RJ.

2017. Lactate metabolism in human lung tumors. *Cell* **171**:358-371.e9.

doi:10.1016/j.cell.2017.09.019.

Fauchon M, Lagniel G, Aude JC, Lombardia L, Soularue P, Petat C, Marguerie G, Sentenac

A, Werner M, Labarre J. 2002. Sulfur sparing in the yeast proteome in response to sulfur

demand. *Mol Cell* **9**:713-23. doi:10.1016/S1097-2765(02)00500-2.

Gerashchenko MV, Gladyshev VN. 2014. Translation inhibitors cause abnormalities in

ribosome profiling experiments. *Nucleic Acids Res* **42**:e134. doi:10.1093/nar/gku671.

Gray JV, Petsko GA, Johnston GC, Ringe D, Singer RA, Werner-Washburne M. 2004.

"Sleeping Beauty": Quiescence in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev*

68:187-206. doi:10.1128/MMBR.68.2.187-206.2004.

Guo HY, Herrera H, Groce A, Hoffman RM. 1993. Expression of the biochemical defect of

methionine dependence in fresh patient tumors in primary histoculture. *Cancer Res*

53:2479-83. PMID:8495409.

Halpern BC, Clark BR, Hardy DN, Halpern RM, Smith RA. 1974. The effect of replacement

of methionine by homocystine on survival of malignant and normal adult mammalian

cells in culture. *Proc Natl Acad Sci U S A* **71**:1133-36. doi:10.1073/pnas.71.4.1133.

Hinnebusch AG. 2005. Translational regulation of *GCN4* and the general amino acid control

of yeast. *Annu Rev Microbiol* **59**:407-50. doi:10.1146/annurev.micro.59.031805.133833.

Hu X, Chao M, Wu H. 2017. Central role of lactate and proton in cancer cell resistance to

glucose deprivation and its clinical translation. *Signal Transduct Target Ther* **2**:e16047.

doi: 10.1038/sigtrans.2016.47.

Jorgensen P, Tyers M. 2004. How cells coordinate growth and division. *Curr Biol* **14**:R1014-

27. doi:10.1016/j.cub.2004.11.027.

Kaleta C, Schäuble S, Rinas U, Schuster S. 2013. Metabolic costs of amino acid and protein

production in *Escherichia coli*. *Biotechnol J* **8**:1105-14. doi:10.1002/biot.201200267.

Kennedy KM, Scarbrough PM, Ribeiro A, Richardson R, Yuan H, Sonveaux P, Landon CD,

Chi JT, Pizzo S, Schroeder T, Dewhirst MW. 2013. Catabolism of exogenous lactate

reveals it as a legitimate metabolic substrate in breast cancer. *PLOS ONE* **8**: e75154.

doi:10.1371/journal.pone.0075154.

Klosinska MM, Crutchfield CA, Bradley PH, Rabinowitz JD, Broach JR. 2011. Yeast cells

can access distinct quiescent states. *Genes Dev* **25**:336-49. doi:10.1101/gad.2011311.

Kokkinakis DM, Hoffman RM, Frenkel EP, Wick JB, Han Q, Xu M, Tan Y, Schold SC.

2001. Synergy between methionine stress and chemotherapy in the treatment of brain

tumor xenografts in athymic mice. *Cancer Res* **61**:4017-23. PMID:11358820.

Laxman S, Sutter BM, Shi L, Tu BP. 2014. Npr2 inhibits TORC1 to prevent inappropriate

utilization of glutamine for biosynthesis of nitrogen-containing metabolites. *Sci Signal*

7:ra120. doi:10.1126/scisignal.2005948.

Laxman S, Sutter BM, Wu X, Kumar S, Guo X, Trudgian DC, Mirzaei H, Tu BP. 2013.

Sulfur amino acids regulate translational capacity and metabolic homeostasis through

modulation of tRNA thiolation. *Cell* **154**:416-29. doi:10.1016/j.cell.2013.06.043.

Laxman S, Tu BP. 2010. Systems approaches for the study of metabolic cycles in yeast. *Curr*

Opin Genet Dev **20**:599-604. doi:10.1016/j.gde.2010.09.008.

Lee BC, Kaya A, Gladyshev VN. 2016. Methionine restriction and life-span control. *Ann N Y*

Acad Sci **1363**:116-24. doi:10.1111/nyas.12973.

Lee BC, Kaya A, Ma S, Kim G, Gerashchenko MV, Yim SH, Hu Z, Harshman LG,

Gladyshev VN. 2014. Methionine restriction extends lifespan of *Drosophila*

melanogaster under conditions of low amino-acid status. *Nat Commun* **5**:3592.

doi:10.1038/ncomms4592.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**:1754-60. doi:10.1093/bioinformatics/btp324.

Ljungdahl PO, Daignan-Fornier B. 2012. Regulation of amino acid, nucleotide, and

phosphate metabolism in *Saccharomyces cerevisiae*. *Genetics* **190**:885-929.

doi:10.1534/genetics.111.133306.

Lu S, Epner DE. 2000. Molecular mechanisms of cell cycle block by methionine restriction

in human prostate cancer cells. *Nutr Cancer* **38**:123-30.

doi:10.1207/S15327914NC381_17.

Mampel J, Schröder H, Haefner S, Sauer U. 2005. Single-gene knockout of a novel

regulatory element confers ethionine resistance and elevates methionine production in

Corynebacterium glutamicum. *Appl Microbiol Biotechnol* **68**:228-36.

doi:10.1007/s00253-005-1893-6.

Mariño, G, Pietrocola F, Eisenberg T, Kong Y, Malik SA, Andryushkova A, Schroeder S,

Pendl T, Harger A, Niso-Santano M, Zamzami N, Scoazec M, Durand S, Enot DP,

Fernández ÁF, Martins I, Kepp O, Senovilla L, Bauvy C, Morselli E, Vacchelli E,

Bennetzen M, Magnes C, Sinner F, Pieber T, López-Otín C, Maiuri MC, Codogno P,

Andersen JS, Hill JA, Madeo F, Kroemer G. 2014. Regulation of autophagy by cytosolic

acetyl-coenzyme A. *Mol Cell* **53**:710-25. doi:10.1016/j.molcel.2014.01.016.

Natarajan K, Meyer MR, Jackson BM, Slade D, Roberts C, Hinnebusch AG, Marton MJ.

2001. Transcriptional profiling shows that Gcn4p is a master regulator of gene

expression during amino acid starvation in yeast. *Mol Cell Biol* **21**:4347-68.

doi:10.1128/MCB.21.13.4347-4368.2001.

Nelson DL, Cox MM. 2017. Lehninger Principles of Biochemistry (7[th] ed.). New York:W. H.

Freeman. p. 533-74.

Nogales-Cadenas R, Carmona-Saez P, Vazquez M, Vicente C, Yang X, Tirado F, Carazo JM,

Pascual-Montano A. 2009. GeneCodis: interpreting gene lists through enrichment

analysis and integration of diverse biological information. *Nucleic Acids Res* **37**:W317-

22. doi:10.1093/nar/gkp416.

Palam LR, Gore J, Craven KE, Wilson JL, Korc M. 2015. Integrated stress response is

critical for gemcitabine resistance in pancreatic ductal adenocarcinoma. *Cell Death Dis*

6:e1913. doi:10.1038/cddis.2015.264.

Patra KC, Hay N. 2014. The pentose phosphate pathway and cancer. *Trends Biochem Sci*

39:347-54. doi:10.1016/j.tibs.2014.06.005.

Pietrocola F, Galluzzi L, Bravo-San Pedro JM, Madeo F, Kroemer G. 2015. Acetyl coenzyme

A: A central metabolite and second messenger. *Cell Metab* **21**:805-21.

doi:10.1016/j.cmet.2015.05.014.

Poirson-Bichat F, Gonçalves RA, Miccoli L, Dutrillaux B, Poupon MF. 2000. Methionine

depletion enhances the antitumoral efficacy of cytotoxic agents in drug-resistant human

tumor xenografts. *Clin Cancer Res* **6**:643-53. PMID:10690550.

Pronk JT. 2002. Auxotrophic yeast strains in fundamental and applied research. *Appl Environ*

Microbiol **68**:2095-2100. doi:10.1128/AEM.68.5.2095-2100.2002.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: A bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139-

40. doi:10.1093/bioinformatics/btp616.

Sherman F. 1991. Getting started with yeast. *Methods Enzymol* **194**:3-21. doi:10.1016/0076-

6879(91)94004-V.

Shi L, Tu BP. 2013. Acetyl-CoA induces transcription of the key G1 cyclin *CLN3* to promote

entry into the cell division cycle in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*

110:7318-23. doi:10.1073/pnas.1302490110.

Slavov N, Botstein D. 2011. Coupling among growth rate response, metabolic cycle, and cell

division cycle in yeast. *Mol Biol Cell* **22**:1997-2009. doi:10.1091/mbc.E11-02-0132.

Slavov N, Botstein D. 2013. Decoupling nutrient signaling from growth rate causes aerobic

glycolysis and deregulation of cell size and gene expression. *Mol Biol Cell* **24**:157-68.

doi:10.1091/mbc.E12-09-0670.

Srinivasan R, Chandraprakash D, Krishnamurthi R, Singh P, Scolari VF, Krishna S,

Seshasayee AS. 2013. Genomic analysis reveals epistatic silencing of "expensive" genes

in *Escherichia coli* K-12. *Mol Biosyst* **9**:2021-33. doi:10.1039/c3mb70035f.

Stern PH, Hoffman RM. 1986. Enhanced *in vitro* selective toxicity of chemotherapeutic

agents for human cancer cells based on a metabolic defect. *J Natl Cancer Inst* **76**:629-

39. doi:10.1093/jnci/76.4.629.

Sugimura T, Birnbaum SM, Winitz M, Greenstein JP. 1959. Quantitative nutritional studies

with water-soluble, chemically defined diets. VIII. The forced feeding of diets each

lacking in one essential amino acid. *Arch Biochem Biophys* **81**:448-55.

doi:10.1016/0003-9861(59)90225-5.

Sutter BM, Wu X, Laxman S, Tu BP. 2013. Methionine inhibits autophagy and promotes

growth by inducing the SAM-responsive methylation of PP2A. *Cell* **154**:403-15.

doi:10.1016/j.cell.2013.06.041.

Tabas-Madrid D, Nogales-Cadenas R, Pascual-Montano A. 2012. GeneCodis3: a non-

redundant and modular enrichment analysis tool for functional genomics. *Nucleic Acids*

Res **40**:W478-83. doi:10.1093/nar/gks402.

Thomas D, Cherest H, Surdin-Kerjan Y. 1991. Identification of the structural gene for

glucose-6-phosphate dehydrogenase in yeast. Inactivation leads to a nutritional

requirement for organic sulfur. *EMBO J* **10**:547-53. PMID:2001672.

Thomas D, Surdin-Kerjan Y. 1997. Metabolism of sulfur amino acids in *Saccharomyces*

cerevisiae. Microbiol Mol Biol Rev **61**:503-32. PMID:9409150.

Tong X, Zhao F, Thompson CB. 2009. The molecular determinants of *de novo* nucleotide

biosynthesis in cancer cells. *Curr Opin Genet Dev* **19**:32-37.

doi:10.1016/j.gde.2009.01.002.

Troen AM, French EE, Roberts JF, Selhub J, Ordovas JM, Parnell LD, Lai CQ. 2007.

Lifespan modification by glucose and methionine in *Drosophila melanogaster* fed a

chemically defined diet. *Age (Dordr)* **29**:29-39. doi:10.1007/s11357-006-9018-4.

Tu BP, Kudlicki A, Rowicka M, McKnight SL. 2005. Logic of the yeast metabolic cycle:

Temporal compartmentalization of cellular processes. *Science* **310**:1152-58.

doi:10.1126/science.1120499.

Tu BP, Mohler RE, Liu JC, Dombek KM, Young ET, Synovec RE, McKnight SL. 2007.

Cyclic changes in metabolic state during the life of a yeast cell. *Proc Natl Acad Sci U S*

A **104**:16886-91. doi:10.1073/pnas.0708365104.

Unger MW, Hartwell LH. 1976. Control of cell division in *Saccharomyces cerevisiae* by

methionyl-tRNA. *Proc Natl Acad Sci U S A* **73**:1664-68. doi:10.1073/pnas.73.5.1664.

van Dijken JP, Bauer J, Brambilla L, Duboc P, Francois JM, Gancedo C, Giuseppin ML,

Heijnen JJ, Hoare M, Lange HC, Madden EA, Niederberger P, Nielsen J, Parrou JL,

Petit T, Porro D, Reuss M, van Riel N, Rizzi M, Steensma HY, Verrips CT, Vindeløv J,

Pronk JT. 2000. An interlaboratory comparison of physiological and genetic properties

of four *Saccharomyces cerevisiae* strains. *Enzyme Microb Technol* **26**:706-14.

doi:10.1016/S0141-0229(00)00162-9.

Vander Heiden MG, Cantley LC, Thompson CB. 2009. Understanding the Warburg effect:

The metabolic requirements of cell proliferation. *Science* **324**:1029-33.

doi:10.1126/science.1160809.

Vattem KM, Wek RC. 2004. Reinitiation involving upstream ORFs regulates *ATF4* mRNA

translation in mammalian cells. *Proc Natl Acad Sci U S A* **101**:11269-74.

doi:10.1073/pnas.0400541101.

Warburg O. 1956. On the origin of cancer cells. *Science* **123**:309-14.

doi:10.1126/science.123.3191.309.

Warner JR, Vilardell J, Sohn JH. 2001. Economics of ribosome biosynthesis. *Cold Spring*

Harb Symp Quant Biol **66**:567-74. doi:10.1101/sqb.2001.66.567.

Warner JR. 1999. The economics of ribosome biosynthesis in yeast. *Trends Biochem Sci*

24:437-40. doi:10.1016/S0968-0004(99)01460-7.

Wu X, Tu BP. 2011. Selective regulation of autophagy by the Iml1-Npr2-Npr3 complex in

the absence of nitrogen starvation. *Mol Biol Cell* **22**:4124-33. doi:10.1091/mbc.E11-06-

0525.

Xu YF, Létisse F, Absalan F, Lu W, Kuznetsova E, Brown G, Caudy AA, Yakunin AF,

Broach JR, Rabinowitz JD. 2013. Nucleotide degradation and ribose salvage in yeast.

Mol Syst Biol **9**:665. doi:10.1038/msb.2013.21.

Ye J, Kumanova M, Hart LS, Sloane K, Zhang H, De Panis DN, Bobrovnikova-Marjon E,

Diehl JA, Ron D, Koumenis C. 2010. The *GCN2-ATF4* pathway is critical for tumour

cell survival and proliferation in response to nutrient deprivation. *EMBO J* **29**:2082-96.

doi:10.1038/emboj.2010.81.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7