# The genome of the biting midge *Culicoides sonorensis* and gene expression analyses of vector competence for Bluetongue virus

Ramiro Morales-Hojas[1,2,*], Malcolm Hinsley[3], Irina M. Armean[3], Rhiannon Silk[1], Lara E. Harrup[1], Asier Gonzalez-Uriarte[4], Eva Veronesi[1,5], Lahcen Campbell[3], Dana Nayduch[6], Christopher Saski[7], Walter J. Tabachnick[8], Paul Kersey[3], Simon Carpenter[1] and Mark Fife[1]

[1] The Pirbright Institute, Ash Road, Woking, Surrey GU24 0NF, UK

[2] Rothamsted Insect Survey, Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK

[3] EMBL-European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SD, UK

[4] Bioinformatics group, Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK

[5] National Centre for Vector Entomology, Institute of Parasitology, Vetsuisse Faculty, University of Zürich, Zürich, Switzerland

[6] USDA-ARS, Center for Grain and Animal Health Research, Arthropod Borne Animal Diseases Research Unit, 1515 College Avenue, Manhattan, KS 66502, USA

[7] Clemson University Genomics Institute, Department of Genetics and Biochemistry, BRC #310, 105 Collins Street, Clemson, SC 29634, USA

[8] Florida Medical Entomology Laboratory, Department of Entomology and Nematology, University of Florida, IFAS, 200 9th St., SE, Vero Beach, FL 32962 USA

* Corresponding author: Ramiro.morales-hojas@rothamsted.ac.uk

**Abstract**

**Background:** The use of the new genomic technologies has led to major advances in control of several arboviruses of medical importance. However, the development of tools and resources available for vectors of non-zoonotic arboviruses remains neglected. Biting midges of the genus *Culicoides* transmit some of the most important arboviruses of wildlife and livestock worldwide, with a global impact on economic productivity, health and welfare. The absence of a suitable reference genome has hindered genomic analyses to date in this important group of vectors. In the present study, the genome of *Culicoides sonorensis,* a vector of bluetongue virus (BTV) in the USA, has been sequenced to be used as reference genome for transcriptomic analyses of vector competence for BTV.

**Results:** The genome has a size of 197.4 Mb and has been assembled in 7,974 scaffolds. Its annotation using the transcriptomic data has identified 15,629 genes. Gene expression analyses of *C. sonorensis* females infected with BTV have identified 165 genes that were differentially expressed between vector competent and refractory females. Two candidate genes, *glutathione S-transferase* (*gst*) and the antiviral helicase *ski2*, were previously recognized as involved in vector competence for BTV in *C. sonorensis* (*gst*) and repressing dsRNA virus propagation (*ski2*).

**Conclusions:** The genome of *C. sonorensis* has enabled preliminary analyses of the genetic profiles of vector competent and refractory individuals. The *C. sonorensis* genome and transcriptomes generated in this study provide suitable platforms for future research on arbovirus transmission. They will be a valuable source of comparative data for other important Dipteran vector families including mosquitoes (Culicidae) and sandflies (Psychodidae).

47 **Keywords:** Culicoides, biting midges, bluetongue virus, genome, genomics,

48 transcriptomics, vector competence

## Background

Arboviruses (arthropod-borne viruses) are a taxonomically diverse group that include some of the most important emerging and re-emerging pathogens of wildlife, livestock and human beings worldwide [1-3]. Among arbovirus vectors, the majority of recent genomic studies have been carried out on the mosquitoes *Aedes* (*Stegomyia*) *aegypti* (L.) and *Aedes* (*Stegomyia*) *albopictus* (Skuse) due to their involvement in human to human transmission of a wide-range of arboviruses, their relative ease of colonization and recent technical advances in genome sequencing and annotation technologies [4]. While these studies have led to major advances in control of the arboviruses these species transmit and our understanding of what drives susceptibility to infection [5-7], tools and resources for use with many other vector groups remain neglected.

To date, no genomic analyses have been carried out for vectors of non-zoonotic arboviruses that are pathogenic to livestock and wildlife. Among the most important of these vector groups are *Culicoides* biting midges (Diptera: Ceratopogonidae), species of which transmit internationally important arboviruses of ruminants, equines and deer [8-10]. In recent years, unprecedented outbreaks of *Culicoides*-borne arboviruses such as bluetongue virus (BTV) have inflicted huge economic losses on the livestock sector of Europe (e.g. estimate of US\$ 1.4 billion in France in 2007) through clinical disease and accompanying restrictions against the movement of livestock imposed to limit virus spread [11, 12]. Globally, BTV outbreaks initiate non-tariff trade barriers restricting the movement of livestock and livestock germplasm, and cause a decrease in ruminant productivity in regions as diverse as India and the USA [13]. Globally, the economic impact of bluetongue has been estimated to US\$ 3 billion [12].

*Culicoides* are notoriously difficult to culture under laboratory conditions and their

small size of approximately 1.5mm body length renders them far from ideal subjects for

transcriptomic and genome manipulation studies [14]. Only one of 14 confirmed vector

species of *Culicoides* is currently colonized, *Culicoides sonorensis* Wirth and Jones [8, 15].

*Culicoides sonorensis* colonies have already provided significant insights into the genetic

basis of vector competence for BTV [16], a *de novo* transcriptome [17], and have been

used to construct a physical map of the *C. sonorensis* genome which consists of four

chromosomes [18, 19].

Susceptibility to infection and transmission of arboviruses by *Culicoides* is determined

in part by the heritability of barriers to virus dissemination following ingestion of the

bloodmeal [13, 20, 21]. Experimental evidence of a midgut infection barrier, a midgut

escape barrier and a haemocoel dissemination barrier in *C. sonorensis* (formerly *C.*

*variipennis* or *C. v. sonorensis* [22]) have been defined [23]. There is no evidence to date

of salivary gland barriers preventing transmission in any species of *Culicoides* [24], as

has been inferred in several species of mosquitoes[20]. Previous studies of the genetic

basis for vector competence of *C. sonorensis* for BTV, using maternal inheritance

properties, identified a 90 kd protein and used antibodies to this protein to isolate and

characterize a cDNA clone encoding a glutathione S-transferase class delta enzyme [25].

Differential gene regulation in response to BTV infection, however, has not been

investigated to date.

In this study, we sequence, *de novo* assemble, annotate and explore the first full genome

of *C. sonorensis* and then use transcriptomic analyses both to improve gene prediction

within the genome build, and to elucidate differential gene regulation associated with

*Culicoides* competence for a BTV serotype 1 strain. The full genome sequence and

transcriptome analyses are important resources for further studies of this phylogenetic

group which is separated from the other major Dipteran vector families by at least 220

million years [26]. In addition to being a valuable resource for comparative study of

vector phylogenomics, the provision of *Culicoides* genomes is of interest as a group that

demonstrates several unique features, including a significant capacity for long-distance

dispersal by semi-passive flight and the ability to reach huge population density under

suitable conditions. In addition, their close association with livestock raises questions

concerning both host preference and vector competence for arboviruses. Many of these

questions can only be addressed by understanding genetic diversity within populations

which will be enhanced by the provision of comparative genomic data, but also in the

long term by yielding targets for transgenic modification. The present genome provides

a resource to facilitate all these studies.

**Results**

**Genome sequencing, construction and annotation**

Assembly of the Illumina reads resulted in a genome of 197,417,436 bp assembled in

7,974 scaffolds. The N50 of the genome assembly was 89,502 and the proportion of Ns

was of 2.63% (Table 1). Analysis with Repeat Modeller [27] identified a total of 793

repeat regions encompassing ~14% of the genome. When low-complexity regions are

included, the total repeat content is 29.7% including 2 Mb of Type II transposons and 5

Mb of Type I transposons (mostly LINE elements). The GC content of the *C. sonorensis*

genome is 28% in the complete assembly and 34% in the exons (Table 1). This is low

when compared to the closest fully sequenced relative to *C. sonorensis* (*Belgica*

*antarctica*: total GC content = 39%) and a range of other Dipteran genomes (Table 1).

Analyses on the completeness of the assembly identified 98.79% (100% including

partial matches) of 248 Core Eukaryotic Genes Mapping Approach (CEGMA) genes and

88.33% (95.88% including partial genes) of 2,675 Benchmarking Universal Single-Copy

Orthologs (BUSCO) arthropod genes. These methods define gene sets that are widely

conserved in single copy and whose can therefore be used to ascertain the

completeness and non-redundancy of a genome sequence. Annotation of the genome

resulted in 15,629 protein-coding genes and 21,336 transcripts, which is comparable to

the number of protein-coding genes in other species of Diptera (Table 1). The mean

gene, intron and exon lengths in *C. sonorensis* are 5,040 bp, 828 bp, and 480 bp,

respectively. The average number of introns per gene in *C. sonorensis* is 6.7. A total of

2,991 genes have been estimated to be alternatively spliced and the maximum number

of splice variants per gene (19) is comparable to *A. aegypti* and *An. gambiae.*

Homology to genes annotated in other species was detected for 93% of the genes

annotated in *C. sonorensis*, with an average of 19 homologous genes and nine

orthologues identified per gene; these numbers are comparable to the other species

used in the analysis. Over 90% of the *C. sonorensis* genes were linked to their

homologues via the ancestral nodes of Arthropoda (6,609 genes), Neoptera (6,361

genes) or Diptera (1,225 genes).

Of 149 high-level categories in the GO ("GO Slim"), 137 were present in *C. sonorensis* as

opposed to 140 categories represented in the genome of *An. gambiae* and 144 in the

model Dipteran, *D. melanogaster*. GO terms with large numbers of annotated proteins

compared to *An. gambiae* and *D. melanogaster* include: ion binding (2,869 genes),

oxidoreductase activity (703 genes), kinase activity (367 genes), hydrolase activity

acting on glycosyl bonds (112 genes), methyltransferase activity (111 genes) and tRNA

metabolic process (101 genes).

To identify tandemly arrayed genes, we looked for genes that contained common

domain architectures (as defined by the set of InterPro-defined domains found within

each gene) and were located within 100,000 nucleotides of each other. 1,281 genes

were found in such arrays. The longest such array contained 10 copies of a gene

containing a chitin-binding domain. The longest array of a protein with this architecture

comprises just two copies in *D. melanogaster*, and there are no such arrays in *A. aegypti*

or *An. gambiae*, although there are various longer arrays of proteins containing this

domain alongside others in these species. The second longest such array contains 9

copies of a gene encoding a protein with arrestin and immunoglobulin domains; the

homologous proteins are found in 10 tandem copies in *A. aegypti* and *D. melanogaster*,

and four tandem copies in *An. gambiae*.

**Transcriptome sequencing and analysis**

Paired-end sequencing of transcriptomes resulted in an average of 50 million reads

with a length of 50 bp for each sample sequenced. Phred scores exceeded 34 in all cases.

The transcriptomes from the different conditions were merged with Cuffmerge prior to

a comparative analysis of their gene expression profiles. The number of genes on the

merged transcriptome is 17,263 and the number of transcripts is 35,813. Of the genes

present in the transcriptome, 15,630 had already been annotated in the genome. 1,816

transcripts, mapping to 1,635 loci, were novel. The proportion of gene loci with a single

transcript is 62.7% (10,826), but 93.3% among the genes that had not been annotated

on the genome. BLAST searches against NCBI's Arthropod non-redundant (nr) protein

database resulted in 92.4% of the isoforms (33,095) and 90% of the genes matching to a

database entry (15,268 of the genes had one or more hits).

**Differential expression of genes between competent and refractory *Culicoides***

***sonorensis***

Eighty-five (56%) *C. sonorensis* in replicate one and 53 (37%) in replicate two produced

$C_q$ values of ≤ 27 (Additional file 2: Figure S1). Sixty-five (44%) *C. sonorensis* in replicate

one and 92 (63%) in replicate two produced $C_q$ values of > 27 or no $C_q$ value. Replicate

one had a significantly greater proportion of disseminated infections than replicate two

(Fishers exact test; df=1, *P* = 0.0007). Three and nine samples within those produced

from the first and second replicates demonstrated failed RNA extraction from bodies

and were discarded. Transcriptome studies were therefore conducted using 147 and

136 female *C. sonorensis*, from the two biological replicates of the infection study,

respectively.

Gene expression profiles demonstrated a trend towards upregulation of expression in

refractory *C. sonorensis* when compared to competent individuals (Fig. 1a). A total of

165 genes demonstrated significant differential expression (DE) (Fig. 1b; Additional file

1: Table S1) of which 123 (75%) had been annotated and 42 (25%) represented novel

genes (Additional file 1: Table S2). A total of 94 (57%) were upregulated in refractory

individuals, while 71 were downregulated (43%) (Additional file 1: Table S1). 120 of

the 165 genes showing DE had one or more BLAST hits in the non-redundant (nr)

protein database from NCBI (Additional file 1: Table S2) and 88% of these

corresponded closely to genes from other members of the suborder Nematocera. Of the

165 genes, 63 were annotated with gene ontology terms. The distribution of these terms

has a mean maximum depth of six (Additional file 2: Figure S2). However, enrichment

analysis showed no significant results.

*Ski2* (XLOC_006435) and *glutathione S-transferase-1* (CSON011559) showed an increase

in expression levels in the females that were resistant to full dissemination, the log2 fold

change being -1.10 ($P$ = 0.00035; $q$-value = 0.00381) and -1.07 ($P$ = 0.00055, q-value =

0.00555), respectively (Fig. 2; Additional file 1: Table S1). In contrast, the *Toll*

(XLOC_000522) gene and the gene matching *glutathione s transferase* (CSON010973) were

expressed at higher levels in the samples that were susceptible to a fully disseminated

BTV infection (log2 fold change 1.21, $P$ = 0.00005 and $q$-value = 0.00072) and (log2 fold

change 2.07, $P$ = 0.00005 and $q$-value = 0.00072) (Fig. 2; Additional file 1: Table S1).

The direction and degree of change in expression levels of the four genes identified as

potentially influencing *C. sonorensis* competence for BTV (*Ski2*, *gst-1*, *gst* and *Toll*) was

confirmed using RT-qPCR and the same samples. Housekeeping genes used as reference

were *CytB5*, *rpl13* and *rps8*, which showed the highest level of stability of those tested

(Additional file 1: Table S3 and Table S4). The direction of up and down regulation of

the genes and the log-2 fold change was similar in all cases as that detected in the

RNAseq data (Additional file 1: Table S5).

**Molecular evolution of ski2 antiviral helicase and glutathione s transferase-1**

The most significant BLAST hit for the *C. sonorensis ski2* was the gene from *Cx.*

*quinquefasciatus* (XP_001845019.1 = CPIJ003293). There are 37 genes identified as

orthologous to *Cx. quinquefasciatus ski2* in 36 species in VectorBase. 35 are 1-to-1

orthologues and there are two orthologous genes in *An. maculatus* (Additional file 1:

Table S6a). The evolutionary relationship of this protein is consistent with species-level

phylogeny (Fig. 3a). In the *gst-1* gene from *C. sonorensis*, the most similar BLAST hit is

that of *C. variipennis* (synonym of *C. sonorensis*; AAB94639.1) [25]. The second closest

was the *gst-1* copy of *An. gambiae* (AGAP004164), for which 61 orthologues have been

identified in 31 other species in VectorBase (Additional file 1: Table S6b). The

phylogeny of the *gst-1* protein recovers the two main clades in Diptera, Brachycera and

Nematocera, but there is little resolution within these clades (Fig. 3b).

**Immune response related genes in *C. sonorensis***

All *D. melanogaster* genes from FlyBase that had InterPro hits containing Toll, Imd and

JAK/STAT associated to their names and all GO terms matching to the Toll, Imd and

JAK/STAT pathway terms were used as reference to identify the homologues in *C.*

*sonorensis* (Additional file 1: Table S7, and S8; Additional file 2: Figure S3). 36 of 42

genes identified in the Toll pathway of *D. melanogaster* were also identified in *C.*

*sonorensis*; eight of the 11 genes of the Imd signalling pathway in *D. melanogaster* were

identified in the *C. sonorensis* genome and 10 of the 13 *D. melanogaster* JAK/STAT

pathway genes were identified. Nineteen genes of the Toll pathway are duplicated in

this species, in comparison with 13 in *M. domestica* and *Ae. aegypti*, which are the

species with the second largest number of duplicated genes (13) (Additional file 1:

Table S9a). In the Imd pathway, seven genes are duplicated in comparison to five genes

in *Cx. quinquefasciatus* and *A. aeqypti* (Chi square $P = 0.009$; Additional file 1: Table S9b).

In the JAK/STAT pathway the number of duplicated genes is within that observed in

other species (Chi square $P = 0.77$; Additional file 1: Table S9c). However, the mean

number of copies per gene in these pathways was not significantly different among

Nematoceran species for Toll and the Imd pathways (single factor ANOVA with *d.f.* = 1,

11 for Toll: $P$ = 0.4; Imd: $P$ = 0.45), but was significantly different in the JAK/STAT

pathway ($P$ = 0.03).

Additional searches for immune related genes in *C. sonorensis* were carried out using

BLASTp with *A. aegypti* and *Cx. quinquefasciatus* gene sequences from the ImmunoDB

database ([http://cegg.unige.ch/Insecta/immunodb](http://cegg.unige.ch/Insecta/immunodb)) as queries (Additional file 1: Table

S10 and S11). Four annotated genes in *C. sonorensis* were identified as potential anti-

microbial peptides (AMPs) belonging to the gene subfamilies Attacin, Cecropin,

Defensin and Diptericin (Additional file 1: Table S10a); however, the analyses did not

identify more than a single gene within each of the AMP subfamilies. Eight potential

Toll-like receptors were identified using BLASTp although there were three instances in

which the most significant identities were different depending on the species used to

provide input data (Additional file 1: Table S11), reflecting the large difference in

sequence between the homologous genes in *A. aegypti* and *Cx. quinquefasciatus.* All five

genes belonging to the Toll pathway (Additional file 1: Table S11) and all genes of the

JAK/STAT signal transduction pathway were identified in *C. sonorensis* (Additional file

1: Table S11); while of the Imd pathway, one gene was identified in five of the eight

subfamilies (Additional file 1: Table S11).

The Compara approach reconciles the gene tree (derived from sequence similarity

metrics) with the species tree to identify likely orthologous relationships; but requires

an initial clustering step (which can lead to the inclusion of sequences into the wrong

trees). The only instance in which the two approaches identify a different *C. sonorensis*

gene and the orthologue of a gene in the above pathways is that of *Pelle* (which activates

the Toll Receptor), for which BLAST identifies CSON012655 as its homologue, and

Compara identifies CSON013584 and CSON013585. There were other cases in which

BLAST identifies a potential homologue while Compara does not (e.g. TNF-receptor-

associated factor 6, Traf6, or tube) and genes Imd and Fadd, neither of the two

approaches identify a homologue in *C. sonorensis*.

**Discussion**

This study has produced the first *de novo* genome assembly of the Dipteran family

Ceratopogonidae, which includes important vectors of emerging and re-emerging

veterinary arboviruses [8, 28]. The closest relative to this group for which a genome of

comparable quality is available is the non-biting midge, *Polypedilum vanderplanki*

Hinton (Family: Chironomidae) [29], which diverged from the Ceratopogonidae some

220 million years ago. This study provides a primary resource for comparative studies

with other arbovirus vector species (e.g. mosquitoes [30]). It also will facilitate efforts to

produce transgenic *Culicoides* to control *Culicoides*-borne pathogens, although the

limitations on colony production for the other epidemiologically relevant species of

*Culicoides* remains a major constraining factor [14, 31].

The genome assembly of *C. sonorensis* is not wholly contiguous, however, its general

features are consistent with similarly sized genomes from other species. Thus, features

such as the repeat content, gene count and level of alternative splicing are within the

ranges observed in *Ae. aegypti*, *An. gambiae* and *D. melanogaster*. Coverage of well-

conserved gene families is complete and comparable with other published assemblies

(98.79% of CEGMA genes and 88.33% of 2,675 BUSCO genes). The gene set has been

annotated with a comparable number of GO Slim terms in *An. gambiae*, with the

molecular functions of ion binding, oxidoreductase activity, kinase activity, hydrolase

activity acting on glycosyl bonds, methyltransferase activity all highly represented, as is

the tRNA metabolic process. In addition, the genes of the immune-related pathways

have been identified and annotated, revealing complete or nearly-complete orthologous

pathways. Overall, these observations indicate that the present assembly and

annotation of the *C. sonorensis* genome represents a reliable resource to use for some

genetic studies. Nevertheless, it is still highly fragmented and the use of long-read

sequencing technologies like PacBio would help improving the scaffolding. In addition,

the use of this genomic resource by the community will help improve the annotation of

the genes. This is the case for many of the genomes of vector species that have been

neglected over the years.

A striking feature of the *C. sonorensis* genome is its GC content, which is among the

lowest reported for Dipteran species (although close to identical to the non-biting

midge *P. vanderplanki* [29] and other non-Dipteran arthropods like the pea aphid,

*Acyrthosiphon pisum* Harris [32]). This is not just a feature of the repeated content, but

extends into the protein-coding genes (which have a higher GC content than the non-

coding regions, but less than in comparable species: Table 1). Recent comparative

analyses of codon usage in Diptera and Hymenoptera showed a significant association

between codon bias and high GC content in Diptera, but low GC content in Hymenoptera

[33]. Thus, the low GC content observed in *C. sonorensis* could reflect a codon bias in

comparison to other Diptera species. Functional investigations of GC content have also

revealed that it has an impact on genome functioning and species ecology in microbes

[34], vertebrates [35] and plants [36]. It is also known that the GC content has an impact

in the efficiency of sequencing technologies [37]. Regions with low GC content have less

coverage than more GC balanced ones. This characteristic of the *C. sonorensis* genome

should be accounted for in future genomic studies, and appropriate experimental

designs that take into account the low GC content should be used [e.g. 38].

Studies investigating gene expression during infection with BTV identified four

candidate genes whose regulation was correlated with vector competence, despite only

using a single time-point at eight days post-infection and pools of *C. sonorensis*

exhibiting different degrees of virus dissemination. Interestingly, the elevated

expression of *gst-1* in refractory *Culicoides* observed in this study is consistent with a

previously proposed genetic mechanism for *C. sonorensis* vector competence for BTV

[13, 15, 24]. These studies identified a locus controlling vector competence in the AA

colony, the origin colony of PIR-s-3 colony used in the present study.  The former study

used a maternally inherited 90kd protein consistent with genetic studies that suggested

maternal inheritance of the competence controlling factor identified in unfertilized eggs

of refractory females to isolate a clone that upon sequencing was identified as *gst-1* [25].

Though the current study evaluated gene expression after exposure to BTV and the

previous study compared genetically selected resistant and susceptible families using

individuals from these families who were not exposed to BTV, both independently

identified involvement of *gst-1*. Thus, the differential expression of the gene in the

current study is confirmation of its involvement in BTV infection in AA colony, although

its role requires functional validation.

Future studies of vector competence in *C. sonorensis* must characterize genes that

facilitate BTV competence and differentiate these from DE genes that are the

consequence of BTV infection. Studies of other *C. sonorensis* colonies from different

populations and other populations of other *Culicoides* vectors and non-vector species

are needed to determine variation in mechanisms that control competence and factors

that be different from AA and PIR-s-3 colonies. Once controlling genes are identified it

will be essential to assess the effect of polymorphisms, the various functions of the

identified genes in the absence of the pathogen to evaluate factors that influence their

frequencies, and compare these processes between different species of *Culicoides* [23].

The present paper presents the genome of *C. sonorensis* and this has been used in

transcriptomic analyses to provide information that will help facilitate the transmission

of viruses by vectors and provide new avenues of research to understand vector

competence for BTV. We present a list of candidate genes that will be further explored

to show their involvement in the transmission of BTV by *C. sonorensis*. Furthermore, the

results also provide evidence for the involvement of genes belonging similar functional

families in response to virus infection in different species belonging to different families

of Diptera. These results show the importance of comparative analyses to interrogate

the evolution of host-pathogen interactions, and the *C. sonorensis* genome will facilitate

such studies.

**Conclusions**

Here we present the first annotated genome of the biting midge, *C. sonorensis*, a vector

of economically important viruses of livestock, such as Bluetongue virus (BTV) and

Schmallenberg virus. This genome has been fundamental in gaining information about

the genetics of BTV transmission. Thus, gene expression comparison between females

susceptible to a full infection by BTV and those females refractory to full BTV infection,

identified 165 genes that are candidates to be involved in vector competence in this

species. Of these genes, the *gst-1* gene was identified previously to be involved in BTV

transmission, while the antiviral helicase *ski2* involved in suppressing the replication of

dsRNA viruses. Overall, the publication of the genome and transcriptomes will lead to

advances in the control of economically relevant arboviruses of livestock. In addition,

this genome fills a phylogenetic gap of more than 200 million years, and will be a useful

source of comparative data for studies of vector competence across Diptera.

**Methods**

**Insects used and their infection with bluetongue virus**

All *C. sonorensis* originated from the 'AA' colony which was originally produced in 1955

at the Kerrville, Texas laboratory of the US Department of Agriculture [39]. Since 1969,

this colony has been maintained at The Pirbright Institute without any additional

outbreeding [40]. Within this period, two selection bottlenecks of ≤10 individuals were

performed to increase susceptibility to BTV serotype 4 and African horse sickness virus

(AHSV) serotype 9 infection (Mellor, Pers Comm). Following the second selection in the

1990's [41], the colony strain was renamed as PIR-s-3. No attempt was made to further

reduce heterozygosity prior to using individuals from the PIR-s-3 line in this study. All

*C. sonorensis* used in genomic and transcriptomic studies were 3-4 days old and had not

been fed sucrose following emergence from pupae.

The virus used to infect *C. sonorensis* was a western topotype strain of BTV serotype 1

(GIB2007/01) [42]. Virus stock was mixed 1:1 with defibrinated horse blood (TCS

Biosciences, Buckingham, UK) with 6.2 $\text{Log}_{10}$ $\text{TCID}_{50}$/ml being the final infectious dose.

All blood-feeding of *C. sonorensis* was conducted using a Hemotek membrane-based

system (Discovery Workshops, Accrington, UK).

***Full genome sequencing***: *C. sonorensis* adults were separated by sex under a

stereomicroscope and stored in 95% ethanol at room temperature prior to use. A total

of 375 male (three pools of 100 and one pool of 75 individuals) and 150 females (one

pool of 100 and one pool of 50 individuals) were homogenized twice for 30 seconds in

75 µl of phosphate buffered saline using a Tissuelyser™ (Qiagen, UK) at a frequency of

25 Hz/s and a 3mm stainless-steel ball bearing (DeJay Distribution Ltd, UK) within 2ml

screw-topped tubes. Genomic DNA was extracted from these pools separately and was

pooled afterwards according to sex.

***Transcriptome sequencing*** *Vector competent vs refractory:* A total of 150 and 145 *C.*

*sonorensis* in two replicates were fed on the blood: BTV-1 suspension and survived the

extrinsic incubation period of eight days at 25 ᵒC. During this period of incubation, 10%

(*w/*v) sucrose solution was offered via a cotton wool pad. Each individual was then

decapitated using disposable needles and the head and remainder of the carcass of each

were stored separately in RNAlater™ (Thermofisher Scientific, UK) at -20 ᵒC.

Homogenization was carried out on individual heads and remaining carcasses in 100µl

Schneider's *Drosophila* media (Gibco™, Thermofisher Scientific, UK), but was otherwise

identical as for full genome sequencing. *Additional transcriptomes:* A further 100 *C.*

*sonorensis* that had fed on horse blood at three days post-emergence and then stored

without access to sucrose for 8 days and 100 *C. sonorensis* that had been offered 10%

(*w/*v) sucrose solution from 3-8 days following emergence, were also used to create

transcriptomes. In this case individuals were stored whole in RNAlater™ at -20 ᵒC and

homogenized as for full genome sequencing in a pool of 100 for each of the two

treatments.

**DNA/RNA extraction**

***Whole genome sequencing (WGS)***: Genomic DNA extraction was conducted using gravity-flow anion-exchange tips (Qiagen's Genomics Tip 20/G) with extraction of DNA up to 150kb in size and a maximum of 20µg of product. A user-developed protocol was followed ([https://www.qiagen.com/gb/resources/resourcedetail?id=b45c3cc3-7f2b-4f4a-aa37-21d814ed3730&lang=en](https://www.qiagen.com/gb/resources/resourcedetail?id=b45c3cc3-7f2b-4f4a-aa37-21d814ed3730&lang=en)), initially mixing the 75µl sample with 1.8 ml of lysis buffer. DNA extracted from the pools of male and female specimens were pooled by sex and the concentration of the resulting samples was evaluated using a Qubit® 2.0 Fluorometer (ThermoFisher Scientific, UK) with the Qubit® DNA BR Assay Kit (ThermoFisher Scientific, UK). The integrity of the genomic DNA was visualized using a 5 µl sample on a 1% (*w/v*) agarose gel.

***BTV Quantification:*** The ProtoScript II Reverse Transcriptase kit (New England Biolabs) was used to quantify BTV in individual decapitated heads. A reaction mixture of 5 µl of homogenized sample, 2 µl of Oligo d(T)$_{18}$ primer and 1 µl of nuclease free water (NFW) was used. The reaction mixture was incubated at 65 °C for 5 minutes and then 10 µl of ProtoScript II reaction mix and 2 µl of ProtoScript enzyme mix were added. The cDNA synthesis reaction was carried out at 42 °C for one hour with a final incubation at 80 °C for five minutes and a reaction with no enzyme was included as a control. The resulting cDNA was used as template in an RT-qPCR for the detection of BTV. The reactions included 5 µl of cDNA, 10 µl of PrimerDesign master mix with SYBR green dye, 20 pmol of each BTV specific primer BTVuni 291-311F 5' GCTTTTGAGGTGTACGTGAAC 3' and BTVuni 381-357R 5' TCTCCCTTGAAACTCTATAATTACG 3' [43] and 4 µl NFW. Controls with no template and with no reverse transcriptase were included. To test that negative reactions were not

due to a lack of cDNA in the sample, RT-qPCR for each sample were performed with *C.*

*sonorensis* specific primers for the Vacuolar ATPase gene (Vac-ATPase forward 5'

GCTGCTGCTGCCATCATTTT 3' and Vac-ATPase reverse 5' CCGGTCGCATCACTGACATA

3'). All reactions were performed in duplicate.

For the purposes of this study, *vector competent* individuals possess disseminated

infections that include replication in the head capsule and quantification threshold ($C_q$)

values ($\leq 27$) that have been demonstrated to allow isolation of BTV-1 [42]. *Refractory*

*individuals* refer to those without detectable BTV after a suitable incubation period,

those with BTV midgut infections that had not disseminated within the insect and those

where the process of dissemination had not been completed (i.e. those with haemocel

dissemination barriers, midgut escape and infection barriers) and those still retaining

inactivated BTV following the infective bloodmeal [24]. Pools of vector competent and

refractory *C. sonorensis* were created from decapitated bodies linked to the infection

status of the head as above.

***Transcriptome sequencing***: TRIzol® (ThermoFisher Scientific, UK) was added to each

homogenized pool of *C. sonorensis* to a final volume of 400 μl. Homogenized samples

were incubated at room temperature for 5 minutes after which 80 μl of chloroform

were added. Samples were inverted for 15 seconds and left at room temperature for 2-3

minutes. The samples were centrifuged at 18000 *x g* for 15 minutes at 4 °C and the clear

upper aqueous layer transferred to a new microcentrifuge tube. RNA was precipitated

with 200 μl of isopropanol incubated at room temperature for 10 minutes. The RNA was

centrifuged at 18000 *x g* for 10 minutes at 4 °C and the supernatant removed. The pellet

was washed with 500 μl of 75% ethanol and centrifuged for 5 minutes at 18000 *x g* at 4

ᵒC. The supernatant was discarded and the pellet was left to dry for 10-15 minutes

before being resuspended in 30 µl NFW at 55 ᵒC for 10 minutes. The integrity of the

RNA was analyzed using a Bioanalyser 2100 and the concentration was evaluated using

a Qubit® 2.0 Fluorometer (ThermoFisher Scientific, UK) with the Qubit® DNA BR Assay

Kit (ThermoFisher Scientific, UK).

**Genome sequencing and *de novo* assembly**

8 – 9 µg of high molecular weight (HMW) genomic DNA were used to sequence the

genome of male and female *C. sonorensis* separately. Two paired-end (PE) libraries of

insert sizes 200 bp and 400 bp and a mate-pair (MP) library with an average fragment

size of 4.4 kb were sequenced with 150 bp (200 bp and 4.4 kb libraries) and 100 bp

(400 bp library) paired-end from both female and male pools were run on Illumina

HiSeq 2000 and 2500 sequencing systems. Library construction and sequencing was

performed at The Earlham Institute (Norwich, UK). Read pair quality was assessed with

FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Results

suggested that 6 bp should be removed from the start of each read, and in the case of

MP data the last 40-50 bp had unexpectedly high levels of k-mer duplication. Both PE

and MP reads were filtered to limit their size to 100bp after removal of the first 6 bp.

MP data was processed with NextClip [44] to select reads which contained both ends of

the original fragment. This created a dataset of reads with variable length, which were

subsequently limited to 100bp.

PE and MP libraries were initially assembled using Velvet [45] with a k-mer length of

91. The N50 of the assembly was maximized using an expected coverage of 27, a

coverage cut-off of six, and with scaffolding option set to 'no'. This initial assembly was

filtered to remove short (<4kb) and repeated contigs (> 90% sequence ID to a larger

contig) using a BLAT and a custom script to count identical bases, considering any

overlaps between matches. The assembly was then scaffolded using SSPACE [46] and

the 400bp PE library, using three passes of decreasing stringency. Any remaining

contigs smaller than 500bp were removed (9913 total). The resulting assembly was

then scaffolded a second time using the 4.4kb MP library with SSPACE (Figure 1). This

assembly was assessed using FRCbam [47] and Reapr [48], and the output from Reapr

was then processed with GapFiller [49]. The final assembly was assessed using FRCbam

to confirm that an improvement in the reported error rate had occurred. Data (reads,

annotation and assembly) have been deposited in the ENA database under the accession

number PRJEB19938. The genome used in this present study is the latest, v1.

**Gene model prediction and annotation**

Prediction and annotation of the genome assembly was conducted using MAKER [50].

The genome annotation was done using the proteomes of nine species of *Anopheles* (*An.*

*albimanus* Wiedemann, *An. arabiensis* Patton, *An. christyi* (Newstead & Carter, 1911), *An.*

*dirus* Peyton & Harrison, *An. epiroticus* Linton & Harbach, *An. funestus* Giles, *An. minimus*

Theobald, *An. quadrimaculatus* Say and *An. stephensi* Liston) together with the

proteomes of *A. aegypti, Belgica antarctica* Jacobs*, Culex quinquefasciatus* Say*,*

*Drosophila melanogaster* Mg., *Lutzomyia longipalpis* (Lutz & Neiva, 1912) and

*Phlebotomus papatasi* (Scopoli). In addition, the transcriptome data from the BTV

infection experiments reported in this study and transcriptome data from previous

studies on *C. sonorensis* [17, 51] were used. SNAP [52] and AUGUSTUS [53] were used to

predict *ab initio* gene models. SNAP was trained using 500 models and AUGUSTUS using

the best 1000 models extracted from the initial MAKER output, as recommended in the

software documentation. Coverage of conserved proteins was assessed using the

CEGMA [54] and BUSCO [55] pipelines. Analysis of Gene Ontology annotation was

performed with Blast2GO [56].

The orthology of the annotated genes was established using the Ensembl Compara Gene

Trees pipeline [57]. Genes were functionally annotated using InterProScan, which

computationally identifies the presence of domain, motif or family signatures within a

protein sequence and infers functional descriptors (taken from the Gene Ontology

(GO)).

**Transcriptome sequencing and analyses**

In total, eight transcriptomes were sequenced: two biological replicates each for i) BTV-

competent; ii) BTV-refractory; iii) blood-fed; and, iv) sucrose-fed. Library construction

and sequencing was performed at Edinburgh Genomics (The University of Edinburgh,

Scotland). A total of 2.5 µg of RNA from each sample was used to construct TrueSeq

libraries, which were sequenced using 50 base paired-end (PE) reads on an Illumina

HiSeq 2500. Each library was run in two lanes.

The quality of the raw sequence data was analyzed in FastQC v 0.11.2. Reads were

aligned to the genome assembly with TopHat v2.0.6 [58] using the gene model

annotations generated from the genome as reference. The resulting bam files were

assembled with Cufflinks v 2.2.1 [59] and merged into a single transcriptome with

Cuffmerge. Quantification of gene and transcript expression and comparison of the

expression levels was performed with Cuffquant and Cuffdiff, respectively. The

comparative analysis of expression profiles between conditions used grouped replicates

of each condition (two biological + two lane runs). Differential expression analyses were

performed with the R package CummeRbund [60]. Differentially expressed (DE) genes

with a significant change in expression level were identified using an $\alpha$ value of 0.05,

which establishes the filtering value of the multiple-testing corrected $q$-values.

Identification and functional classification of the DE genes was done using BLAST and

InterProScan searches in Blast2Go version 4.0.2. Enrichment and gene set enrichment

analyses were performed on the sets of DE genes between the different conditions using

the interface provided in Blast2Go to the Fisher's Exact Test implemented in FatiGO [61]

and the Gene Set Enrichment Analysis (GSEA) package [62]. To perform the GSEA, all

genes were ranked per the logarithmic fold-change, removing the infinite values.

Transcriptome data have been deposited in the ENA database under the accession

numbers ERR2171964-ERR2171979.

**563     Validation of differentially expressed genes**

Quantitative reverse transcription PCR (RT-qPCR) was used to validate the change in

expression of four selected genes between the vector competent and refractory pools of

*C. sonorensis*. These were selected on the basis of previous identification in vector

competence studies of *C. sonorensis* or by having a direct functional link with antiviral

response. The genes were: a gene sharing 73% identity with the antiviral helicase *ski2*

of *Cx. quinquefasciatus (ski2)*; a gene sharing 63% identity with *glutathione S-transferase*

*(gst)*, a gene sharing 96% identity with *glutathione S-transferase-1* (*gst-1*) and a gene

sharing 49% identity with a gene encoding a *Toll* protein.

Expression levels were normalized against three reference genes whose stability was

ranked using three approaches, BestKeeper [63], geNorm [64], and NormFinder [65]

(Additional file 1: Table S3). Gene-specific primers and probes were designed using

Primer3Plus [66], to meet the following conditions: amplicon 80-120bp, Tm ~60 °C, GC

~40%, 3' GC clamp, minimal hairpin loops/primer dimers (Additional file 1: Table S4).

Primers were compared against the *C. sonorensis* genome using BLAST to ensure

specificity to the target region. Amplification efficiency was tested by RT-qPCR with

hydrolysis probes by generating a standard curve using triplicate technical replicates of

five-fold serial dilutions of RNA template from teneral female *C. sonorensis* specimens.

All RNA templates were treated with Turbo DNA-Free™ Kit (ThermoFisher Scientific,

UK) prior to RT-qPCR following the manufacture recommended protocol. Starting

template total RNA concentration was evaluated using a Qubit® 3.0 Fluorometer

(ThermoFisher Scientific, UK) with the Qubit® RNA HS Assay Kit (ThermoFisher

Scientific, UK).

cDNA synthesis and quantitative amplification was done in one reaction using the

Superscript® III One-Step RT-qPCR System with Platinum® Taq DNA polymerase

(ThermoFisher Scientific, UK). Reactions were performed in a total volume of 25 µl

containing 5.5 µl NFW, 12.5 µl 2x reaction mix, 0.5 µl SuperScript III/PlatinumTaq Mix,

0.5 µl ROX reference dye (1:10 dilution), 0.5 µl 10 µM forward primer, 0.5 µl 10 µM

reverse primer, 1.0 µl hydrolysis probe and 4 µl RNA template. Reaction cycles

consisted of the cDNA synthesis at 50 °C for 5 minutes, followed by a denaturation step

at 94 °C for 2 minutes and amplification for 40 cycles at 95 °C for 3s each and 56 °C for

30s. PCR amplification efficiency of each primer-probe-target combinations was

calculated via the linear regression of $C_q$ as the log2 of the relative RNA template

concentration using the ggplot2 package [67] in R. Primers which exhibited

amplification efficiency between 90-110% with an $R^2$ value of ≥0.98 were considered

valid. Relative expression levels for the gene of interest between vector competent and

refractory *C. sonorensis* were calculated using the ΔΔCq method [68].

**Molecular evolution analyses**

Orthologues of the *ski2* and *gst-1* genes in other insect species were identified using

Blast2Go. The most significant BLAST hit from *A. aegypti*, *Cx. quinquefasciatus* or *An.*

*gambiae* was then used to identify the orthologues in other vector species in VectorBase

[69]. The protein sequences identified were aligned using the mode expresso of T-

Coffee [70]. Poorly aligned regions (positions with score 0-6) were trimmed and the

resulting alignments were used to construct a phylogenetic tree. Phylogenetic analyses

were performed in MrBayes v3.2 [71] and the models of evolution that best fitted our

alignments were identified using ProtTest 3.4.2 [72]. Two independent analyses of two

million generations with four chains (one cold and three heated) were conducted. Trees

and parameters were sampled every 50th generation and the first 10,000 were

discarded. The remaining trees were used to estimate the consensus and Bayesian

posterior probabilities of each branch. The s*ki2* from *Saccharomyces cerevisiae* Meyen

ex E.C. Hansen and the *gst-1* from *Pediculus humanus* L. were used as outgroups.

**Declarations**

**Ethics approval and consent to participate**

Not applicable.

**Availability of data and material**

All sequencing data, genomic and transcriptomic, generated in this study has been

deposited at the European Nucleotide Archive (ENA) database under the accession

number PRJEB19938.

**Competing interests**

The authors declare that they have no competing interests.

**Funding**

This work was supported by the Biotechnology and Biological Sciences Research

Council (BBSRC) grant BB/J016721/1 and the core Strategic Programme Grant to The

Pirbright Institute.

**Author's contributions**

MF, SC and PK conceived the research project. RMH, MH, IMA, RS, LEH, AGU, EV and LC

have performed the laboratory experiments and bioinformatic analyses. DN and CS have

helped with the annotation of the immune-related genes. RMH, MF and SC have written

the manuscript and PK, WT, ND and CS have provided insightful revision. All authors

read and approved the final manuscript.

**Acknowledgements**

We thank Eric Denison and James Barber in the insectary for their supply of reagents,

cell lines and insects for this study.

**References**

1.    Weaver SC, Reisen WK: Present and future arboviral threats. Antiviral Res. 2010;

85:328-345.

2.    Mlakar J, Korva M, Tul N, Popovic M, Poljsak-Prijatelj M, Mraz J, Kolenc M, Rus

KR, Vipotnik TV, Vodusek VF *et al*: Zika Virus Associated with Microcephaly. New

Engl J Med. 2016; 374:951-958.

3.    Pialoux G, Gauzere BA, Jaureguiberry S, Strobel M: Chikungunya, an epidemic

arbovirosis. Lancet Infect Dis. 2007; 7:319-327.

4.    Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS,

Machol I, Lander ES, Aiden AP *et al*: De novo assembly of the *Aedes aegypti*

genome using Hi-C yields chromosome-length scaffolds. Sci. 2017; 356:92-95.

5.    Alphey L: Genetic Control of Mosquitoes. Annu Rev Entomol. 2014; 59:205-224.

6.    Franz AWE, Sanchez-Vargas I, Adelman ZN, Blair CD, Beaty BJ, James AA, Olson

KE: Engineering RNA interference-based resistance to dengue virus type 2 in

genetically modified *Aedes aegypti*. Proc Natl Acad Sci U S A. 2006; 103:4198-

4203.

7.    Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu ZJ, Loftus B, Xi ZY, Megy K,

Grabherr M *et al*: Genome sequence of Aedes aegypti, a major arbovirus vector.

Sci. 2007; 316:1718-1723.

8.    Purse BV, Carpenter S, Venter GJ, Bellis G, Mullens BA: Bionomics of Temperate

and Tropical *Culicoides* Midges: Knowledge Gaps and Consequences for

Transmission of *Culicoides*-Borne Viruses. Annu Rev Entomol. 2015; 60:373-392.

9.    Carpenter S, Mellor PS, Fall AG, Garros C, Venter GJ: African Horse Sickness Virus:

History, Transmission, and Current Status. Annu Rev Entomol. 2017; 62:343-

358.

10.   Savini G, Afonso A, Mellor P, Aradaib I, Yadin H, Sanaa M, Wilson W, Monaco F,

Domingo M: Epizootic haemorragic disease. Res Vet Sci. 2011; 91:1-17.

11.   Carpenter S, Wilson A, Mellor PS: *Culicoides* and the emergence of bluetongue

virus in northern Europe. Trends Microbiol. 2009; 17:172-178.

12.   Rushton J, Lyons N: Economic impact of Bluetongue: a review of the effects on

production. Vet Ital. 2015; 51:401-406.

13.   Tabachnick WJ: *Culicoides variipennis* and bluetongue-virus epidemiology in the

United States. Annu Rev Entomol. 1996; 41:23-43.

14.   Nayduch D, Cohnstaedt LW, Saski C, Lawson D, Kersey P, Fife M, Carpenter S:

675           Studying *Culicoides* vectors of BTV in the post-genomic era: Resources,

676           bottlenecks to progress and future directions. Virus Res. 2014; 182:43-49.

677    15.   Jones RH, Foster NM: Oral infection of *Culicoides variipennis* with bluetongue

678           virus - development of susceptible and resistant lines from a colony population. J

679           Med Entomol. 1974; 11:316-323.

680    16.   Tabachnick WJ: Genetic control of oral susceptibility to infection of *Culicoides*

681           *variipennis* with bluetongue virus. Am J Trop Med Hyg. 1991; 45:666-671.

682    17.   Nayduch D, Lee MB, Saski CA: The Reference Transcriptome of the Adult Female

683           Biting Midge (*Culicoides sonorensis*) and Differential Gene Expression Profiling

684           during Teneral, Blood, and Sucrose Feeding Conditions. PLoS One. 2014; 9.

685    18.   Nunamaker RA, Brown SE, Murphy KE, Tabachnick WJ, Knudson DL: Physical

686           mapping of the genome of the biting midge, *Culicoides variipennis*. Mol Biol Cell.

687           1998; 9:452A-452A.

688    19.   Nunamaker RA, Brown SE, McHolland LE, Tabachnick WJ, Knudson DL: First-

689           generation physical map of the *Culicoides variipennis* (Diptera : Ceratopogonidae)

690           genome. J Med Entomol. 1999; 36:771-775.

691    20.   Mellor PS, Boorman J, Baylis M: *Culicoides* biting midges: Their role as arbovirus

692           vectors. Annu Rev Entomol. 2000; 45:307-340.

693    21.   Tabachnick WJ: Nature, Nurture and Evolution of Intra-Species Variation in

694           Mosquito Arbovirus Transmission Competence. Int J Env Res Public Health.

695           2013; 10:249-277.

696    22.   Holbrook FR, Tabachnick WJ, Schmidtmann ET, McKinnon CN, Bobian RJ, Grogan

697           WL: Sympatry in the *Culicoides variipennis* complex (Diptera : Ceratopogonidae):

698           a taxonomic reassessment. J Med Entomol. 2000; 37:65-76.

23.    Carpenter S, Veronesi E, Mullens B, Venter G: Vector competence of Culicoides

for arboviruses: three major periods of research, their influence on current

studies and future directions. Revue Scientifique Et Technique-Office

International Des Epizooties. 2015; 34:97-112.

24.    Fu H, Leake CJ, Mertens PPC, Mellor PS: The barriers to bluetongue virus

infection, dissemination and transmission in the vector, *Culicoides variipennis*

(Diptera : Ceratopogonidae). Arch Virol. 1999; 144:747-761.

25.    Abdallah MA, Pollenz RS, Droog FN, Nunamaker RA, Tabachnick WJ, Murphy KE:

Isolation and characterization of a cDNA clone coding for a glutathione S-

transferase class delta enzyme from the biting midge Culicoides variipennis

sonorensis Wirth and Jones. Biochem Genet. 2000; 38:377-390.

26.    Wiegmann BM, Trautwein MD, Winkler IS, Barr NB, Kim JW, Lambkin C, Bertone

MA, Cassel BK, Bayless KM, Heimberg AM *et al*: Episodic radiations in the fly tree

of life. Proc Natl Acad Sci U S A. 2011; 108:5690-5695.

27.    Smit AFA, Hubley R: RepeatModeler Open-1.0. 2008-2015

<http://www.repeatmasker.org  >) 2015.

28.    Carpenter S, Groschup MH, Garros C, Felippe-Bauer ML, Purse BV: *Culicoides*

biting midges, arboviruses and public health in Europe. Antiviral Res. 2013;

100:102-113.

29.    Gusev O, Suetsugu Y, Cornette R, Kawashima T, Logacheva MD, Kondrashov AS,

Penin AA, Hatanaka R, Kikuta S, Shimura S *et al*: Comparative genome

sequencing reveals genomic signature of extreme desiccation tolerance in the

anhydrobiotic midge. Nature Communications. 2014; 5.

30.    Severson DW, Behura SK: Mosquito Genomics: Progress and Challenges. Annu

Rev Entomol. 2012; 57:143-166.

31.    Mills MK, Nayduch D, Michel K: Inducing RNA interference in the arbovirus

vector, Culicoides sonorensis. Insect Mol Biol. 2015; 24:105-114.

32.    Richards S, Gibbs RA, Gerardo NM, Moran N, Nakabachi A, Stern D, Tagu D,

Wilson ACC, Muzny D, Kovar C *et al*: Genome Sequence of the Pea Aphid

*Acyrthosiphon pisum*. Plos Biology. 2010; 8:24.

33.    Behura SK, Severson DW: Comparative Analysis of Codon Usage Bias and Codon

Context Patterns between Dipteran and Hymenopteran Sequenced Genomes.

PLoS ONE. 2012; 7.

34.    Foerstner KU, von Mering C, Hooper SD, Bork P: Environments shape the

nucleotide composition of genomes. EMBO Rep. 2005; 6:1208-1213.

35.    Galtier N, Piganeau G, Mouchiroud D, Duret L: GC-content evolution in

mammalian genomes: The biased gene conversion hypothesis. Genetics. 2001;

159:907-911.

36.    Smarda P, Bures P, Horova L, Leitch IJ, Mucina L, Pacini E, Tichy L, Grulich V,

Rotreklova O: Ecological and evolutionary significance of genomic GC content

diversity in monocots. Proc Natl Acad Sci U S A. 2014; 111:E4096-E4102.

37.    Chen YC, Liu T, Yu CH, Chiang TY, Hwang CC: Effects of GC bias in next-

generation-sequencing data on de novo genome assembly. PLoS One. 2013;

8:e62856.

38.    Oyola SO, Otto TD, Gu Y, Maslen G, Manske M, Campino S, Turner DJ, Macinnis B,

Kwiatkowski DP, Swerdlow HP *et al*: Optimizing Illumina next-generation

sequencing library preparation for extremely AT-biased genomes. BMC

Genomics. 2012; 13:1.

39.    Jones RH: The laboratory colonization of *Culicoides variipennis* (Coq). J Econ

Entomol. 1957; 50:107-108.

40.    Boorman J: Maintenance of laboratory colonies of *Culicoides variipennis* (Coq),

*Culicoides nubeculosus* (Mg) and *Culicoides riethi* Kieffer (Diptera,

Ceratopogonidae). Bull Entomol Res. 1974; 64:371-377.

41.    Mellor PS, Rawlings P, Baylis M, Wellby MP: Effect of temperature on African

horse sickness virus infection in Culicoides. Arch Virol. 1998:155-163.

42.    Veronesi E, Antony F, Gubbins S, Golding N, Blackwell A, Mertens PPC, Brownlie J,

Darpel KE, Mellor PS, Carpenter S: Measurement of the Infection and

Dissemination of Bluetongue Virus in Culicoides Biting Midges Using a Semi-

Quantitative RT-PCR Assay and Isolation of Infectious Virus. PLoS One. 2013; 8:8.

43.    Shaw AE, Monaghan P, Alpar HO, Anthony S, Darpel KE, Batten CA, Guercio A,

Alimena G, Vitale M, Bankowska K *et al*: Development and initial evaluation of a

real-time RT-PCR assay to detect bluetongue virus genome segment 1. J Virol

Methods. 2007; 145:115-126.

44.    Leggett RM, Clavijo BJ, Clissold L, Clark MD, Caccamo M: NextClip: an analysis and

read preparation tool for Nextera Long Mate Pair libraries. Bioinformatics. 2014;

30:566-568.

45.    Zerbino DR, Birney E: Velvet: Algorithms for de novo short read assembly using

de Bruijn graphs. Genome Res. 2008; 18:821-829.

46.    Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W: Scaffolding pre-

assembled contigs using SSPACE. Bioinformatics. 2011; 27:578-579.

47.    Vezzi F, Narzisi G, Mishra B: Reevaluating Assembly Evaluations with Feature

Response Curves: GAGE and Assemblathons. PLoS One. 2012; 7.

48.    Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD: REAPR: a

universal tool for genome assembly evaluation. Genome Biol. 2013; 14.

49. Boetzer M, Pirovano W: Toward almost closed genomes with GapFiller. Genome

Biol. 2012; 13.

50. Holt C, Yandell M: MAKER2: an annotation pipeline and genome-database

management tool for second-generation genome projects. BMC Bioinformatics.

2011; 12.

51. Nayduch D, Lee MB, Saski CA: Gene discovery and differential expression

analysis of humoral immune response elements in female *Culicoides sonorensis*

(Diptera: Ceratopogonidae). Parasites & Vectors. 2014; 7.

52. Stanke M, Schoffmann O, Morgenstern B, Waack S: Gene prediction in eukaryotes

with a generalized hidden Markov model that uses hints from external sources.

BMC Bioinformatics. 2006; 7.

53. Stanke M, Steinkamp R, Waack S, Morgenstern B: AUGUSTUS: a web server for

gene finding in eukaryotes. Nucleic Acids Res. 2004; 32:W309-W312.

54. Parra G, Bradnam K, Korf I: CEGMA: a pipeline to accurately annotate core genes

in eukaryotic genornes. Bioinformatics. 2007; 23:1061-1067.

55. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM: BUSCO:

assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics. 2015; 31:3210-3212.

56. Gotz S, Garcia-Gomez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M,

Talon M, Dopazo J, Conesa A: High-throughput functional annotation and data

mining with the Blast2GO suite. Nucleic Acids Res. 2008; 36:3420-3435.

57. Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle

SMJ, Amode R, Brent S *et al*: Ensembl comparative genomics resources (vol 2016,

bav096, 2016). Database-the Journal of Biological Databases and Curation. 2016.

58. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL: TopHat2: accurate

alignment of transcriptomes in the presence of insertions, deletions and gene

fusions. Genome Biol. 2013; 14.

59. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,

Rinn JL, Pachter L: Differential gene and transcript expression analysis of RNA-

seq experiments with TopHat and Cufflinks. Nat Protoc. 2012; 7:562-578.

60. L G, C T, D K: cummeRbund: Analysis, exploration, manipulation, and

visualization of Cufflinks high-throughput sequencing data. R package version

2180. 2013.

61. Al-Shahrour F, Diaz-Uriarte R, Dopazo J: FatiGO: a web tool for finding significant

associations of Gene Ontology terms with groups of genes. Bioinformatics. 2004;

20:578-580.

62. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,

Paulovich A, Pomeroy SL, Golub TR, Lander ES *et al*: Gene set enrichment

analysis: A knowledge-based approach for interpreting genome-wide expression

profiles. Proc Natl Acad Sci U S A. 2005; 102:15545-15550.

63. Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP: Determination of stable

housekeeping genes, differentially regulated target genes and sample integrity:

BestKeeper - Excel-based tool using pair-wise correlations. BiotL. 2004; 26:509-

515.

64. Hoerndli FJ, Toigo M, Schild A, Gotz J, Day PJ: Reference genes identified in SH-

SY5Y cells using custom-made gene arrays with validation by quantitative

polymerase chain reaction. AnBio. 2004; 335:30-41.

65. Ohl F, Jung M, Xu CL, Stephan C, Rabien A, Burkhardt M, Nitsche A, Kristiansen G,

Loening SA, Radonic A *et al*: Gene expression studies in prostate cancer tissue:

which reference gene should be selected for normalization? Journal of Molecular

Medicine-Jmm. 2005; 83:1014-1024.

66.    Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JAM:

Primer3Plus, an enhanced web interface to Primer3. Nucleic Acids Res. 2007;

35:W71-W74.

67.    Wickham H: **ggplot2: Elegant Graphics for Data Analysis**. New York, USA;

2009.

68.    Pfaffl MW: A new mathematical model for relative quantification in real-time RT-

PCR. Nucleic Acids Res. 2001; 29.

69.    Lawson D, Arensburger P, Atkinson P, Besansky NJ, Bruggner RV, Butler R,

Campbell KS, Christophides GK, Christley S, Dialynas E *et al*: VectorBase: a data

resource for invertebrate vector genomics. Nucleic Acids Res. 2009; 37:D583-

D587.

70.    Notredame C, Higgins DG, Heringa J: T-Coffee: A novel method for fast and

accurate multiple sequence alignment. J Mol Biol. 2000; 302:205-217.

71.    Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B,

Liu L, Suchard MA, Huelsenbeck JP: MrBayes 3.2: Efficient Bayesian Phylogenetic

Inference and Model Choice Across a Large Model Space. Syst Biol. 2012; 61:539-

542.

72.    Darriba D, Taboada GL, Doallo R, Posada D: ProtTest 3: fast selection of best-fit

models of protein evolution. Bioinformatics. 2011; 27:1164-1165.

 **Figure Legends**

 **Figure 1.** Differential gene expression analyses between females susceptible to full BTV

 infection and refractory females. **a** Scatterplot sof the pairwise comparison of the gene

 expression levels between the two phenotypes. The average gene expression across all

 genes is displayed by the blue line. **b** Volcano plot displaying fold changes in expression

 of the 165 differentially expressed genes between *Culicoides sonorensis* that are

 susceptible to infection and those that are refractory to full infection with BTV.

 **Figure 2.** Expression bar plots for candidate genes identified as differentially expressed

 at significant level between vector competent (orange bars) and refractory (blue bars)

 samples to full BTV infection. *Toll* corresponds to the *Toll*-like gene, *ski2* is the antiviral

 helicase *ski2*, *gst* is *glutathione s transferase*, and *gst-1* is the *glutathione s transferase-1*.

 **Figure 3.** Bayesian phylogenies of *ski2* (a) and *gst-1* (b) genes. Tip labels correspond to

 the Ensembl Identifier followed by the species name (in the case of *C. sonorensis*, the

 number that precedes correspond to the RNA transcript); colouring corresponds to the

 Nematocera in red and Brachycera in green. Labels on branches show the posterior

 probability values.

 **Supporting Information Legends**

 **Supplementary Material Tables**

 **Table S1.** Differentially expressed genes between vector competent and vector

 refractory *C. sonorensis* females identified using CummeRbund. The table shows the

 fpkm values for each phenotype, the log2 fold change of expression levels and the

 significance level of the change. Genes that are not annotated in the *C. sonorensis*

 genome are denoted with the name XLOC_XXXXXX. Genes belonging to an enriched GO

 term are shown in bold.

**Table S2.** BLAST results for the differentially expressed genes between vector

competent and refractory transcriptomes showing the top hit as determined by BLAST.

**Table S3.** Ranking of the expression stability of the reference genes tested for use in the

quantitative reverse transcription PCR (RT-qPCR). Three different tests were used to

rank the expression stability of the genes, BestKeeper, gNorm and NormFinder. *r*-

correlation coefficient, Cq – quantification cycle, SE – standard error, SD – standard

deviation.

**Table S4.** Primers forward (F), reverse (R) and hydrolysis probe (P) sequences used for

RT-qPCR analysis of the expression levels of *ski2*, *gst*, *gst-1* and *Toll*.

**Table S5.** Difference of expression levels in four genes (*Toll*, *gst*, *gst-1* and *ski2*) between

vector-competent and vector-refractory females assessed by RNAseq and RT-qPCR. *

denote a difference in expression between refractory and competent females with a *P* <

0.05 (RT-qPCR significance estimated using a Welch Two Sample t-test with d.f. = 1).

**Table S6.** Orthology analysis of two of genes *Ski2* **(a)** and *gst-1* **(b)**.

**Table S7.** Immune pathways genes identified in *C. sonorensis* using the Ensembl

Compara pipeline and the *D. melanogaster* genes as reference. **a)** Toll pathway; **b)** Imd

pathway; **c)** Jak/Stat pathway.

**Table S8.** Homologue immune genes in several invertebrate species identified using *D.*

*melanogaster* genes as reference all with InterPro and GO terms associated with Toll

**(a)**, Imd **(b)** and Jak/Stat **(c)**, in the Ensembl Compara pipeline.

**Table S9.** Number of copies of the different genes identified in several Diptera species

as homologues to the *D. melanogaster* genes containing InterPro and GO terms related

to Toll **(a)**, Imd **(b)** and Jak/Stat **(c)** pathways. Ae - *A. aegypti;* Ag - *An. gambiae;* Ba - *B.*

*antarctica;* Cq - *Cx. quinquefasciatus;* Cs - *C. sonorensis;* Dm - *D. melanogaster;* Gm - *G.*

*morsitans;* Ll - *L. longipalpis;* Lc - *L. cuprina;* Md - *M. domestica;* Pp - *P. papatasi;* Sc - *S.*

*calcitrans.*

**Table S10.** Immune related genes identified in *C. sonorensis* using BLASTp top hits with

the gene copies from *A. aegypti* and *Cx. quinquefasciatus* available in the ImmunoDB

database (http://cegg.unige.ch/Insecta/immunodb) as query. a) Anti-Microbial

Peptides; b) Toll receptors; c) Toll path; d) Jak/Stat path; e). Imd path. * - Top BLASTp

hits different found when *A. aegypti* and *Cx. quinquefasciatus* were used as query.

**Table S11.** Details of the BLASTp results using *A. aegypti* (a) and *Cx. quinquefasciatus*

(b) as query to identify the *C. sonorensis* genes of the immune pathways.

**Supplementary Material Figures**

**Figure S1.** Distribution of $C_q$ values from RT-qPCR used to detect BTV-1 virus infection

in the two virus-feeding experiments (in blue and orange) of *C. sonorensis*. The vertical,

dashed line corresponds to a $C_q$ value of 27 used to separate vector competent from

refractory females.

**Figure S2.** Classification and functional distribution of the genes differentially

expressed between vector competent and refractory *C. sonorensis* per the Gene

Ontology level 6. Blue: Molecular Function; Green: Cellular Component; Pink: Biological

Process.

**Figure S3.** Number of genes in the Toll (a), Imd (b) and Jak/Stat (c) pathways that have

been duplicated or lost in different species of Diptera as identified using the Ensembl

Compara pipeline.

**Table 1.** A comparison of genome characteristics of the *C. sonorensis* genome with other selected Diptera species.

| Parameter | *Culicoides sonorensis* | *Aedes aegypti* L3 | *Aedes aegypti* L5 | *Anopheles gambiae* AGAM P4 | *Drosophila melanogaster* BDGP6 | *Belgica antarctica* GCA_000775305.1 |
|---|---|---|---|---|---|---|
| **Genome size (Mb)** | 197.4 | 1,384.1 | 1,362.0 | 278.3 | 143.7 | 89.7 |
| **Number of contigs** | 15,810 | 36,205 | 4,901 | 16,825 | 2,442 | 22,492 |
| **Contig N50** | 30,774 | 82,500 | 10,861,303 | 85,555 | 19,488,218 | 13,551 |
| **Contig N90** | 5,567 | 15.283 | 74,389 | 5,600 | 666,663 | 1,865 |
| **Number of scaffolds** | 7,974 | 4,757 | 4,616 | 10,523 | 1,870 | 4,997 |
| **Scaffold N50** | 89,077 | 1,547,048 | 409,777,670 | 49,364,325 | 25,286,936 | 98,164 |
| **Scaffold N90** | 14,215 | 324,062 | 83,687 | 2,684 | 23,513,712 | 18,418 |
| **Ns (%)** | 2.8 | 5.3 | 0.0 | 7.6 | 0.8 | 0.9 |
| **Maximum contig length** | 552,397 | 685,587 | 71,953,859 | 808,130 | 27,905,041 | 131,402 |
| **Repeats + low complexity regions** | 29.7% | 73% | 78% | 25% | 23% | 4% |

| | | | | | | |
|---|---|---|---|---|---|---|
| **GC content** | 28.4% | 38.3% | 38.2% | 44.3% | 42.0% | 38.9% |
| **GC exons** | 34.3% | 46.8% | 46.2% | 54.6% | 49.1% | 47.1% |
| **Protein-coding genes** | 15,629 | 15,796 | 14,626 | 13,008 | 13,918 | 13,510 |
| **Mean protein-coding gene length** | 5,040 bp | 18,126 bp | 46,920 bp | 6,442 bp | 6,960 bp | 2,555 bp |
| **Mean exon length (protein coding genes)** | 480 bp | 460 bp | 494 bp | 438 bp | 538 bp | 325 bp |
| **Mean intron length (protein-coding genes)** | 632 bp | 23,265 bp | 10,456 bp | 1,008 bp | 1,147 bp | 213 bp |
| **Mean introns per transcript (in protein-coding genes)** | 6.7 | 5.1 | 7.9 | 5.5 | 6.9 | 5.3 |
| **Maximum splice isoforms/protein-coding gene** | 19 | 20 | 50 | 20 | 75 | n.a. |

Contigs refer to the contiguous sequences assembled without Ns; scaffolds refer to the final top level sequence assembly used for

annotation (they contain Ns).

**Fig. 1**

**a**

**b**

**Fig. 2**

926
927

929   **Fig. 3**
930
931   **a)**
932

933   **b)**

934
935