Development of PTSD-like behavior in adult mice after observing an acute traumatic event

Ray X. Lee,[1,2]* Greg J. Stephens,[2,3] Bernd Kuhn[1]

Affiliations:

[1] Optical Neuroimaging Unit, Okinawa Institute of Science and Technology (OIST) Graduate University, Okinawa, Japan.

[2] Biological Physics Theory Unit, OIST Graduate University, Okinawa, Japan.

[3] Department of Physics and Astronomy, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands.

* Correspondence to: rayxin.lee@oist.jp

Abstract: For post-traumatic stress disorder (PTSD) in humans, it remains a major conceptual challenge how stress reactions emerge after a protracted symptom-free period. Here, we study the behavioral development in mice isolated after observing a traumatic event inflicted upon their pair-housed partners. Fine-scale behavioral analysis showed that anxiety-related reactions gradually increased more in traumatized mice during isolation than in mice isolated without traumatic experience. These increases in anxiety-related reactions were independent of fear memory. Interestingly, instead of social avoidance from anxiety, traumatized mice showed long-term reduction of social interest to strangers. These behaviors were not observed if the traumatic event happened to a stranger mouse. Additionally, traumatized mice showed rebound responses to their partners after 4 weeks of separation. This suggests that affectional bonding as the single common factor underlies the otherwise independent development of PTSD-like behaviors in this paradigm. This conclusion is also supported by the strengthening of an anterior cingulate cortex (ACC)-centered network revealed by diffusion tensor imaging.

Introduction

Post-traumatic stress disorder (PTSD) is one of the most prevalent mental health disorders [1,2]. Although the mechanisms underlying PTSD are not well understood [3], a core belief shared across various hypotheses is to view PTSD as an association between trauma-induced states such as fear memory, increased anxiety, and social withdrawal [4–8]. In this collective model (Figure 1A), behaviors slowly change and influence each other during the stress incubation period until a new, abnormal equilibrium is reached. This new equilibrium is described by PTSD.

To identify the basic psychopathological mechanisms during stress incubation, it is beneficial to use an experimental assay with purely psychosocial manipulations on controlled subjective experiences and a homogeneous genetic background. Laboratory rodents proved to be a reliable model system for such a purpose. Behavioral studies revealed that laboratory rodents engage in advanced social behaviors and show similar psychological states as humans [9–12]. Specifically, laboratory rodents have been used to study stress behavior and the pharmacology of stress [13–16]. Also, a delay period before showing substantial stress reactions was reported in the context of rodent models simulating human PTSD [17–21]. Stress reactions have been studied with a wide variety of aversive stimuli, ranging from acute physical stress [22,23] to a prolonged period of repeated social defeat through witnessing [19,24]. A time-dependent amplification of fear expression [25] has been proposed to explain generalized avoidance, including associative fear conditioning or non-associative fear sensitization, through stress incubation in rodents [18]. The variability of emergent phenotype following trauma was proposed to be an interaction of incubated stress and fear conditioning [21].

Here, we combined behavioral scenarios, analytic methods, and psychological hypotheses to study the development of stress incubation by psychobehavioral experiments in mice. Additionally, we analyzed morphological changes of the brain due to stress incubation by diffusion tensor imaging (DTI). We propose that a single factor, affectional bonding, plays

a dominant role during the stress incubation period towards the development to PTSD-like

behavior. In this unitary model (Figure 1B), a single common factor underlies the otherwise

diverse and independent development of post-traumatic behaviors in mice.

Results

Emergence of long-term effects after trauma induction

To identify post-traumatic behavioral development with minimal physical impact and

peri-traumatic contribution, we developed the following assay of an acute witnessing trauma

(Figure 2A): Partnership between the male focal mouse and its male partner was established

by housing them together for 3 weeks (Day-21–Day0). During this pair-housing, the mice slept

together in a single nest which they built, and no aggressive interaction was observed. On

Day0 (trauma induction), the focal mouse observed its partner being attacked by 5 different

aggressor mice in succession (aggressive encounters) and stayed together with the attacked

partner between each aggressive encounter (stress infiltration and resting). Importantly, the

focal mouse only experienced the traumatic event by witnessing the attacks and by social

communication, but not through any direct physical threat, such as attack bites and pursuits

from either the aggressors or its partner. After the last aggressive encounter, the focal mouse

(Partner-observing mouse) was socially isolated for 4 weeks (Day0–Day28). Behavior was

tested on Days -7, 1, 7, and 28.

To differentiate behavioral consequences from the trauma induction and the effects of

isolation, adaptation to the tests, and aging, a control group of mice were isolated from their

partners on Day0 without trauma induction (No-scenario-isolated mice; Figure 2B). We found

that Partner-observing mice (n = 47) built nests with significantly higher walls than those

constructed by No-scenario-isolated mice (n = 47) after isolation (Figure 2C). Partner-

observing mice also increased their body masses in the late phase of the study (Figure 2D).

Both observations indicate a long-term effect of the trauma. These differences could be

alternatively explained, e.g. having better emotional stability and therefore nicer nests and

better appetites *v.s.* manifesting protective responses of anxiety by hiding behind higher nest

walls and engaging in anxiety-induced binge eating [26,27]. However, the corticosterone

concentrations in blood plasma [28] confirms increased stress in the Partner-observing mice:

Compared with No-scenario-isolated mice, Partner-observing mice showed higher baseline

plasma corticosterone level after trauma, which reached statistical significance on Day28

(Figure 2E). To confirm the stress reactions suggested by the baseline plasma corticosterone

level, in this experiment, we also isolated the attacked mice and collected their blood samples

after trauma (Attacked mice; n=5, note that 2 out of 5 mice died on Days 4 and 5, supposedly

due to a serious depression-like anorexia as no obvious physical trauma occurred; interestingly,

such losses were not observed in attacked mice which were subsequently group-housed). The

tendency of higher baseline plasma corticosterone level was also observed in Attacked mice

which had a more obvious baseline corticosterone increment during the early phase (Figure

2E), supporting the interpretation that the observed behavioral differences are behavioral

phenotypes of stress.

To investigate morphological changes in the brain, we used DTI of brains collected on

Day28 to analyze brain-wide microstructural differences (Figure 3; Supplemental Figures 1 to

4): Instead of a structural change in the hypothalamus [29], significant differences mainly

occurred in both neocortex and hippocampus. For Partner-observing mice, compared with No-

scenario-isolated mice, their DTI-based fractional anisotropy (FA; Figure 3A) was lower in

anterior cerebral cortical areas including the primary somatosensory cortex, anterior cingulate

cortex, and orbital cortex, but higher in posterior cerebral cortical areas including the

retrosplenial cortex and the primary visual cortex. It was also higher in the hippocampus.

Different measures of DTI-based water diffusivities [mean diffusivity (MD), axial diffusivity

(AD), and radial diffusivity (RD); Figure 3, B to D] were generally higher in the cerebral cortex

and hippocampus of Partner-observing mice than in No-scenario-isolated ones. Interestingly,

we observed an obvious asymmetry of the brain areas: Gray matters including the cerebral

cortex and hippocampus showed differences in the right brain hemisphere, while white matters

including the corpus callosum, anterior commissure, and cingulum bundle showed differences

in the left brain hemisphere. The asymmetry of diffusivity between the right and left brain

hemispheres were significantly enhanced in the amygdala (the right more than the left) and

areas of the lateral cerebral cortical subnetwork including dorsolateral orbital cortex and insular

cortex (the left more than the right) for Partner-observing mice. These observations support

the idea that the emergence of long-term impacts after trauma induction arise from a higher-

order process of cognition and emotion.

Evidence of stress development revealed by fine-scale behavioral analysis

To measure emotionality of Partner-observing mice in different phases after trauma, we

investigated the spontaneous behaviors of Partner-observing mice by using standard behavioral

testing setups. Although the generally believed validity to identify rodent emotionality in

standard behavioral tests has been criticized [30], spontaneous behavior in a steady environment

is arguably a representation of internal states such as emotion [31].

Here, we focus on anxiety-like reactions in mice. In humans, anxious behaviors are

shown when facing unexpected events or uncertain situations [32]. Behavioral tests for rodent

anxiety-like reactions were therefore designed to evaluate their stress reaction against their

willingness to explore (e.g. light-dark box and elevated plus-maze) or their activity under

conditions with a gradient of uncertainty (e.g. open field). We first examined spontaneous

behaviors in the light-dark box test (Figure 4A; n = 8 mice for each group), where the stressor

was a natural aversion to brightly lit areas [33]. Surprisingly, while the time spent in the light

area did not differ significantly between No-scenario-isolated and Partner-observing mice on

Day1, Partner-observing mice spent more time in the light area than No-scenario-isolated mice

on Days 7 and 28 (Figure 4A, left panels).

This result raised the following questions: (i) Did this behavioral difference start to

develop immediately after the traumatic event or only after a delay? (ii) What comparative

emotionality does this behavioral difference indicate?

To answer the first question (i), we examined the positions of the mice in the light-dark

box on a fine-scale. Emotion dynamically maps internal states and external stimuli onto a set

of corresponding behaviors [34], which should be reflected in the probability distribution of

different parameters shown during these behaviors. Based on spatial symmetry, we analyzed

T(x), the cumulative probability distribution of time that the mouse spent at positions along the

axis of the light-dark box (Figure 4A, top-right scheme and the equation, and middle panels for

the results). We calculated significance levels [presented as -log of p-values, -log(p)] of T(x)

between Partner-observing and No-scenario-isolated populations by computing position-

dependent population means and applying a two-tailed, two-sample Student's t-test (Figure 4A,

bottom-middle panels). Already on Day1, Partner-observing mice showed differences in their

spatial distribution, as they spent less time than No-scenario-isolated mice at the far end of the

dark area. This tendency increased with time: On Day 7, Partner-observing mice spent more

time in the light area close to the door compared to No-scenario-isolated, and then on Day 28

additionally on the far side of the light area. Collapsed -log(p) distributions reveal the overall

gradual increase in spatial preference differences (Figure 4A, right panel). Additionally,

Partner-observing mice maintained a higher locomotor speed compared to the gradually

decreasing speed of No-scenario-isolated mice (Figure 4B) and showed more transfers between

the boxes and shorter latencies until their first transfers from Day 1 (Figure 4C). These results

indicate an onset and gradual increase of behavioral differences immediately following the

trauma.

Regarding the second question (ii), these differences may reflect the hypo-sensitivity

to sensory harm and hyper-activity, such as the characteristic of somatic anxiety observed in

PTSD patients [35–37]; however, they may also simply indicate that Partner-observing mice

became braver and more excited. Reactions of cognitive anxiety are expected to be opposite

of those shown in somatic anxiety [35]. To determine whether the differences could be attributed

to chronic anxiety, we examined their behaviors in the elevated plus-maze test (Figure 5A; n =

8 mice for each group). In this test, stressors included fear of falling and exposure. After first

exposure on Day-7 and separation, mice spent only a fraction of the time in the open arms of

the maze, but with no significant difference between No-scenario-isolated and Partner-

observing mice (Figure 5A, left panels). However, Partner-observing mice spent increasingly

more time in the far end of the closed arms (Figure 5A, middle panels) and moved more slowly

in the elevated plus-maze after trauma induction (Figure 5B) with longer periods of freezing

and fewer entries to the central platform from the closed arms (Figure 5C).

Together, this indicates a higher anxiety level and supports the hypothesis that

behavioral differences are the result of chronic anxiety. Gradually increasing differences

between No-scenario-isolated and Partner-observing mice (Figure 5, A and B) further confirm

that stress incubation of anxiety caused the observed differences.

Development of different psychological components untangled in standard behavioral tests

How different psychological components develop during stress incubation is still

unknown. From now on through the rest of the study, two questions will be addressed: (i) How

does a trauma-induced psychological component affect trauma-independent psychological

components? (ii) How does chronic anxeity, as one psychological component, increase during

stress incubation?

To answer these questions, we sequentially identified different psychological

components observable in mouse behaviors, demonstrated their associations in stress

development, determined what cause-effect system could account for the observed processes,

and validate the conceptual framework by testing its explanatory and predictive power on a

wide range of control, rescuing, and physiological conditions.

To explore the developmental process of different psychological components, we first

investigated components revealed in the light-dark box and the elevated-plus maze. In both

tests, key incubation features were consistently more obvious within stressor-free zones (dark

area and closed arms) than within stressor zones (light area and open arms). To quantitatively

investigate the differences between zones, we analyzed locomotor speed, as an independent

behavioral index of mouse position, separately in stressor and stressor-free zones (Figure 6).

As expected, two different behavioral patterns developed in the two zones: While speed

differences consistently increased in stressor-free zones (Figure 6, B and D), speed differences

in stressor zones only showed acute increases on Day1 (Figure 6, A and C). These results

suggest that different psychological components were measured in the two zones.

Therefore, we asked what psychological basis could account for the two behavioral

features separately measured in the stressor and stressor-free zones. In humans, fear is a

response to a known threat with a magnitude that increases with the certainty of the threat,

whereas anxiety is a response to uncertainty with a magnitude that increases with the

uncertainty of a situation [32]. Together with the presented demonstration of anxiety incubation

(Figures 4 and 5), we hypothesized that behavioral differences in stressor-free zones are the

result of anxiety reactions to uncertainty, while acute stress reactions in stressor zones are the

result of lasting fear responses from trauma induction. We examined mouse locomotor speed

in uncertain and secure environments separately, using the open field test (Figure 7, A to C; n

= 8 mice for each group) and the locomotor activity test (Figure 7D; n = 8 mice for each group).

In the open field test, mice were exposed to a dimly lit environment with a gradient of spatial

uncertainty from the field center (high uncertainty) to the field boundaries (low uncertainty).

In the locomotor activity test, mice were habituated in a dark environment with familiar cues

of bedding materials from their homecages indicative of less environmental stress. Compared

to No-scenario-isolated mice, Partner-observing mice moved faster in the center region of the open field (Figure 7A), which was not observed in the periphery (Figure 7B). The difference between groups increased gradually towards the center but not in the periphery (Figure 7, A and B). The avoidance of spatial uncertainty by Partner-observing mice was also reflected in the shorter time they spent near the center (Figure 7C, left panel) and showed a shorter latency to the first rearing during the test (Figure 7C, right panel), which together suggests anxiety, represented in a high tension between high uncertainty avoidance and high exploratory motivation, as the causal emotional state. In contrast, Partner-observing mice moved significantly slower on Day1 in the locomotor activity test and recovered on Days 7 and 28 (Figure 7D), suggesting an acute post-traumatic impact under low stress condition. These results support the hypothesized psychological basis of anxiety and fear that account for the two behavioral features with separate developments.

Notably, these observations of mouse spontaneous behaviors in the light-dark box test, elevated plus-maze test, open field test, and locomotor activity test (Figures 6 and 7) indicated a previously unexpected association between different post-traumatic stress reactions: The results revealed that the two developmental patterns displayed a consistent uncertainty dependence. In addition, the patterns arose together in the acute phase, but the fear pattern vanished while the anxiety pattern continued to increase in the later phase. These results suggest that delayed anxiety should not simply be considered an expanded, sustained response of fear memory or a stress reaction induced by fear memory, because fear reactions vanish soon after trauma induction. This finding refutes a general belief [1–7,18,38,39].

Comparison of emotionality and sociality differences suggests a unitary model

Although we have demonstrated that the incubated anxiety and acute non-associative fear memory developed independently in our paradigm, this indication does not distinguish a

unitary model from a collective model (Figure 1A): In a unitary model, the incubated anxiety

and acute non-associative fear memory are independently mediated by the trauma-induced state

of a common psychological component, and the incubated anxiety does not induce generalized

behavioral changes in the late phase. In a collective model, incubated anxiety and acute non-

associative fear memory can be independently and directly induced by the traumatic event,

while the incubated anxiety can also further induce generalized behavioral changes in the late

phase. To test both models against each other, we study the relation between emotionality,

specifically anxiety, and sociality in our paradigm.

Emotional responses simplify and speed up animal reactions to complex external cues

and are critical in corresponding social interactions [40]. If a sociality difference occurs and it is

induced by incubated anxiety, as in the case of the collective model, the growing pattern of

incubated anxiety should temporally precede or co-emerge with the corresponding growth

pattern of sociality difference. Additionally, the sociality difference should be anxiety-

correlated, and therefore should not occur without incubated anxiety. In the unitary model, if

the sociality difference develops independently after trauma induction and is independent of

the incubated anxiety, then the sociality difference can precede the incubated anxiety and

should be independent of incubated anxiety.

To test potential effects of anxiety on sociality, we examined mouse social motivation

in a two-session social test (Figure 8A): A non-social session followed by a social session (n =

5 mice for each group). The social stimulus was either a female or a male stranger mouse.

During the social session, Partner-observing mice spent less time in social approaches

of nose poking toward both female and male strangers, starting from Day1, and remained less

social compared to No-scenario-isolated mice during the post-traumatic period (Figure 8B).

The time they spent in the interaction zone around the social target, however, did not differ

significantly from that of No-scenario-isolated mice (Figure 8C). Both, Partner-observing and

No-scenario-isolated mice, spent only a short but similar time on nose poking during the non-

social session through the recordings on different days, with no significant difference between

Partner-observing and No-scenario-isolated populations (Figure 8D), indicating that the

observed difference of nose poking time was socially specific. In addition, less social

vocalization was recorded during the female stranger test for Partner-observing mice (Figure

8E). These results indicated a general and chronic reduction of social interest developing after

the trauma (Supplemental Video 1). This reduction of social interest differs from social

avoidance due to anxiety because the social target is not avoided but just ignored [24,41].

We also analyzed locomotor speed of mice in empty sessions as an indicator of

emotionality previous to social sessions, and find a notable drop of spontaneous locomotor

speed starting on Days 7 and 28 (Figure 8F). Interestingly, onset of chronic reduction of social

interest (starting from Day1) preceded the drop of locomotor speed. These results suggest that

the changes of emotionality and sociality arise independently after trauma induction.

Relationship-dependent behaviors suggesting affectional mechanism

Two possibilities remain that could still explain a collective model: The sociality

difference could be a higher-order consequence of incubated anxiety (Figures 4 to 7) or it could

also be a parallel mechanism induced directly by the traumatic event. However, the following

experiments reject these possibilities (Figure 9):

First, we show that affectional bonding is the key trauma-associated psychological

factor that commonly mediated the development of both emotionality and sociality (Figure 10):

As Partner-observing mice experienced the traumatic event purely psychosocially,

partnership between mice served as a candidate variable with which to alter the emotional

impact and coping strategy [10,42,43]. Thus, we analyzed a second group of mice (Stranger-

observing mice; Figure 9A) as each of them observed different stranger mice being attacked

by different aggressors and stayed together with each stranger between aggressive encounters

before isolation.

Stranger-observing mice did not experience any physical stress from strangers or

aggressors. Two notable behavioral differences during trauma induction were observed: While

tail rattling during aggressive encounters and hiding under bedding material with the partner

during resting were observed in 83% (n = 39 out of 47 mice) and 100% (n = 47 out of 47 mice)

Partner-observing mice (Supplemental Video 2), respectively, no such behaviors were shown

by Stranger-observing mice (n = 0 out of 20 mice; n = 0 out of 20 mice). In Partner-observing

mice, the frequency of tail rattles dropped progressively during aggressive encounters (Figure

10A), suggesting that it represents a rapid phasic psychological reaction during trauma

induction. These results suggest that social relationship rapidly modulated emotional impact

and social reactions during trauma induction. Moreover, chronic behavioral differences were

not observed in Stranger-observing mice (Figure 10, B to F), which further excluded potential

effects from salient, non-specific environmental manipulation (e.g. rotation through aggressors'

homecages) and sensory shock (e.g. olfactory cues from urine and vocalization indicating fear)

in trauma induction. Taken together, the results reveal that social relationship constitutes a

critical common psychological factor that determined post-traumatic psychobehavioral

development.

Considering a unitary model with social relationship as the key factor, an important

question remained: Why is stress incubated, but not attenuated? As traumatic consequences

were modulated by social relationship, we hypothesized that the traumatic event resulted in

gradually increasing separation distress through partnership memory during social isolation.

This is analog to human patients tending to recall relatively vivid memories of the past

relationship after a traumatic relationship dissolution [44]. Social isolation generally disrupts

mouse capacities for social recognition and social memory measured in curiosity-based

behavioral tests [45,46]. We examined the persistence of social memory in a partner-revisiting test

on Day28 (Figure 10G; n = 5 mice for each group). Social stimuli were the previous partner

and a stranger mouse, both immobilized to allow enough social cues to be attractive, but no

active interaction with the focal mouse. To avoid possible influences of social cues from socially defeated mice, stranger mice used to test No-scenario-isolated and Stranger-observing mice were partners of Partner-observing mice. Strikingly, Partner-observing mice spent three times as much time allogrooming or pushing their previous partners as did No-scenario-isolated and Stranger-observing mice (Figure 10, H and Supplemental Video 3), which was opposite to the lower contacts with familiar social targets in a three-chamber social novelty test where curiosity to novelty dominates the behavior [46]. The preference of Partner-observing mice to their previous partners, together with the findings of chronic anxiety (Figures 4 and 5) and reduction of social interest (Figure 8) after separation, suggests affectional bonding as a key mechanism of stress incubation in this paradigm.

Rescue of stress development by social buffering

Next, we observed the development of sociality differences without the development of emotionality differences when testing the necessary and sufficient conditions in our paradigm (Figure 11):

Two important conditions in our behavioral paradigm were the increased number of mice during trauma induction (visual exposure to 5 different aggressors) and the forced social isolation after trauma, of which the potential effects should further be identified. We first compared behavioral developments of No-scenario-isolated mice with a third control group of mice (Stranger-exposed mice; Figure 9B) where the aggressor mice for trauma induction were replaced by non-aggressive strangers. As the expectation (Figure 10), Stranger-exposed mice did not show the developments of behavioral difference (Figure 11), confirming that social aggression is necessary to induce stress developments.

Conceptualization of social support as a "stress buffer" have been proposed to explain the positive association between responsive social resources in a small social network and adverse effects of stressful events [47]. To further characterize social effects in post-traumatic

period, we examined the impact of social support on behavioral developments by a fourth control group of mice (Partner-buffered mice; Figure 9C) as each of them was kept pair-housed with its attacked partner after trauma induction. Noteworthily, developments of the differences in emotionality of stress did not occur in Partner-buffered mice (Figure 9, A to C), but importantly, developments that reflected their sociality showed the differences of Partner-observing mice (Figure 9, D and E). This observation manifested the sufficiency of socially isolated condition for anxiety incubation but not for the reduction of sociality differences. Their similarity with Partner-observing mice in a longer time spent with partners extrapolates that the partner revisiting test is a memory-based assay to measure social bonding in mice. The observation also strengthens the position that emotionality and sociality during stress incubation were separately regulated.

Since Partner-buffered mice experienced the same traumatic scenario as Partner-observing mice, questions remain regarding the psychological mechanism for the partial rescue of stress development in Partner-buffered mice. Because stress reduction through social buffering depends on both social contact and social communication which can be altered by emotional contagion based on different social experiences [10–12,48], and the stress-buffering resources of Partner-buffered mice experienced direct trauma, we hypothesized that, in social buffering, the development of emotionality difference is regulated by the social availability, while the development of sociality difference is regulated by the social context. To address these issues, we conducted a fifth control group of mice where each of them was kept pair-housed with a stranger after trauma induction (Stranger-buffered mice; Figure 9D), and a sixth control group of mice where each of them was provided with toys during social isolation after trauma induction (Environment-enriched mice; Figure 9E).

In the Stranger-buffered paradigm, we found aggressive attacks toward defeated strangers by all focal mice (n=5 out of 5) during pair-housing after trauma. Similar behavior was observed when Partner-observing mice, but not their defeated partners, were group-housed

after trauma induction (non-recorded observation). While the observations importantly illustrate partnership-specific aggressive behavior, we tested the behaviors of Stranger-buffered mice only for those which were pair-housed with a non-defeated stranger. For these Stranger-buffered mice, they did not displayed the developments of both emotionality and sociality differences (Figure 12), with only an acute effect of positions in the light-dark box test (Figure 12A).

Notably, although Environment-enriched mice showed similar nose poking times in the female stranger test (Figure 12B) suggesting a rescue of sociality difference, they showed the behavioral differences of Partner-observing mice in the light-dark box test and the non-social session of female stranger test, which even had a stronger difference during the early phase (Figure 12, A and C). The data corroborate the hypothesized mechanism of social buffering in rescuing stress incubation and additionally emphasize the differential effects between social and physical environmental enrichment.

Changes in brain connectome suggest neural substrate of unitary model

Based on our behavioral results, as well as analyses of physiological conditions (Figures 2E and 3), we next asked if a corresponding neural framework could account for the unitary model deduced from our psychobehavioral experiments. The major neuromodulatory system of serotonin has long been linked to broad neural controls of physiological, sensorimotor, emotional, motivational, cognitive, and social modulations [49]. Fluoxetine, as a selective serotonin reuptake inhibitor, is widely used in prevalent disorders, including PTSD, regarding its supposed importance on inconclusive physiological roles [50]. We investigated a seventh control group of mice (FLX-treated mice; Figure 9F) as each of them was injected with fluoxetine daily after trauma. Although FLX-treated mice showed less behavioral difference in the early post-traumatic phase (Figure 12, A and D) and did not display a reduction of social interest (Figure 12B), incubated anxiety kept developing and reached significance in the late

phase (Figure 12, A and D). These results exposed the limitation of a global pharmacological

treatment for stress incubation.

To further explore the physiological basis, beyond the DTI-based microstructures of

the brain, we investigated long-range connections of the hippocampus, as the region showing

the most notable differences from various measures (Figure 3). Compared with No-scenario-

isolated mice, among all brain regions, DTI-based fiber tracking highlighted the significant

increases of structural connections between the right hippocampus (CA3 and dentate gyrus)

and right dorsolateral entorhinal cortex (Figure 13A), an area also showed difference of its

mean diffusivity (Figure 3A) in Partner-observing mice. While the entorhinal cortex is the

main interface between the hippocampus and neocortex, we further tracked its long-range

connections.   The right dorsolateral entorhinal cortex showed significantly increased

connections with the right perirhinal cortex, but had significantly decreased connections with

the right piriform cortex (Figure 13A). Interestingly, the piriform and perirhinal cortices are

the two core parahippocampal structures involve in the kindling phenomenon, the daily

progressive increase in response severity of both electrographic and behavioral seizure activity

[51,52] supposedly linked to fear conditioning in rat PTSD models [53,54]. We found a decrease of

the connections between the piriform and perirhinal cortices (Figure 13A). Trough further

tracking, we exposed a trauma-induced changes of DTI-based connectivity framework: the

perirhinal cortex-entorhinal cortex-hippocampus, retrosplenial cortex-hippocampus, piriform

cortex-amyglala, and amyglala-hypothalamus systems were centered at the anterior cingulate

cortex (ACC; Figure 13B), in agreement with a unitary model. These results emphasize the

importance of a treatment on the specific clinical target in the full mechanistic picture

according to a unitary model that explains, integrates, and predicts multiple mechanisms

underlying highly diverse behavioral consequences.

Discussion

We present a novel approach to interpret previously challenging and inconclusive

behavioral development and use it to study the internal psychological development of stress

incubation in laboratory mice.   These results demonstrate a system-level view of

experimentally disentangled components, processes, and determinants in post-traumatic stress

development. While our data suggest that laboratory mice developed diverse post-traumatic

reactions with comparable psychological features as observed in human, we found little support

for the collective model view that a complex association among trauma-induced psychological

states govern the developmental process [1–7,18,38,39] (Figure 14A). Instead, we propose a unitary

model where the alteration of a single cognitive source is sufficient for triggering changes of

diverse and otherwise independent psychological states (Figure 14B). Indeed, we found that

the strengthening of an ACC-centered network supports the proposed unitary model for the

diversity of behavioral phenotypes.

Here we summarize our results and interpretations:

Did mice display stress reactions after observing a traumatic event inflicted upon their

pair-housed partners? Yes. Higher nest wall (Figure 2C), higher baseline corticosterone levels

(Figure 2E), more time spent in the further end of close arms (Figure 5A), less exploration from

close arms to center (Figure 5C, left panel), longer freezing time (Figure 5C, right panel), and

slower locomotion (Figure 5B) in the elevated plus-maze test, more time spent in the center

region (Figure 7C, left panel), and shorter latency to the first rearing (Figure 7C, right panel)

in the open field test, and slower locomotion in the locomotor activity test (Figure 7D) are all

classically interpreted as stress reactions. However, we also note that increased body mass

(Figure 2D), more time spent in light area (Figure 4A), faster locomotion (Figure 4B), more

transfers (Figure 4B, right panel), and shorter latency to the first transfer (Figure 4C, right panel)

in the light-dark box test are controversial in classic readouts, an aspect we address in the next

point.

Do standard behavioral tests of mouse anxiety completely and exclusively measure

mouse anxiety? No. We argue that mice display a mixture of multiple emotional aspects in

the standard behavioral tests with classic analyses. First, we showed that an emotional aspect

can yield opposite behavioral phenotypes in classic analyses of standard behavioral tests

(Figures 4 and 5, e.g. somatic and cognitive reactions of chronic anxiety). Second, we showed

that multiple emotional aspects can couple in a single behavioral test and are mixed in

corresponding classic analyses (Figure 6, e.g. distinct developmental patterns of fear and

anxiety). Classically, the emotion is inferred from specific behavioral analyses (behavior-

inferred emotion approach). Instead, we first considered a complete and exclusive definition

of the focused emotion. Then, we quantified how much of the overall behaviors matches with

the essential property inferred from the provided definition (emotion-inferred behavior

approach). Our interpretation of mouse anxiety was validated in four behavioral tests (Figures

6 and 7).

Did fear memory support the development of chronic anxiety? No. We observed the

continuous growth of anxiety when fear reactions had long vanished (Figures 4 to 7).

Did the social deficits reflect increased anxiety? No. The time course of social deficits

led that of increased anxiety (Figures 4, 5, and 8), and social buffering with partner mice

selectively rescued increased anxiety but not social deficits (Figure 11).

Did the trauma affect multiple behavioral aspects in parallel? No. There is a single

factor, affective bonding, commonly mediating diverse behavioral developments. Alternation

of a single social cognitive factor, social relationship, eliminated behavioral differences from

mice with no traumatic experience (Figures 10 and 11). Also, among all rescuing controls,

including social, environmental, or pharmacological approach (Figure 9, C to F), social

buffering with stranger mice showed the best rescuing effects on diverse behavioral aspects

(Figure 12).

Is there a neural substrate for the proposed psychological model?   Yes.   An

enhancement of an ACC-centered network was shown in DTI-based brain-wide connectome

analysis (Figures 3 and 13).

Detection and identification of animal emotionality

We argue that classical behavioral analyses are too coarse to capture intricate emotional

states. The traditional approach to identify animal emotionality is to test if animals show the

behaviors specified by the experimental test assumed when it was designed (behavior-inferred

emotion approach) [55]. However, it is commonly observed that significant, robust, but

inconsistent behavioral outcomes are shown in different tests with the same logic and

assumptions [30]. Even if the results are consistent with the inferred behaviors, they can still be

alternatively explained [56]. The ambiguity reaches deeply into the history of widely used

behavioral tests and therefore have resulted in a considerable amount of inconclusive and

seemingly paradoxical results [57–63]. These limitations caused by circular arguments and

alternative interpretations should be due to the psychological and environmental factors beyond

the reductive condition considered in an experimental design. Classically, researchers

addressed the difficulties by an effort to show a proof-of-concept [9]. Here, we propose the

emotion-inferred behavior approach to identify animal emotionality from standard behavioral

tests which give inconclusive results when analyzed and interpreted in the traditional way.

We also introduced a simple and powerful analysis of standard behavioral tests, which

allowed us to discover otherwise obscured behavioral details [64]. Explaining subtle behavioral

differences is challenging without both richer measurements and quantitative analyses [65]. The

fine-scale analysis developed in our study solves this difficulty, while its utility was validated

by comparing it with standard analyses which in most cases prove inconclusive; that is, for the

analysis of stress incubation.

Biological mechanisms of mouse PTSD

Behavioral paradigms of laboratory rodents that simulate PTSD were established to

expose the mechanistic and causal insights of long-term fear memory following acute physical

stress [22,23] and learned depression after repeated social defeat, either directly [66] or through

witnessing [19,24]. The diversity of PTSD models in mice is even highlighted by a paradigm of

trauma-free pharmacologically-induced memory impairments in mice, which was recognized

as a PTSD model and further identified the corresponding pathophysiological mechanism [14].

The phenomenon of stress incubation was observed and discussed in a few of these

animal PTSD studies. Early inhibition of PTH2R (parathyroid hormone 2 receptor)-mediated

TIP39 (tuberoinfundibular peptide of 39 residues) signaling in the medial amygdalar nucleus

was reported to enhance the increasing fear memory following a single scrambled footshock in

mice [20], although a system-level mechanism of stress incubation was limited by the sole

observation of freezing time in the study. Differentiating the effects from directly physical

impact, stress incubation of multiple phenotypes was more obvious in mice experiencing

observational stress but not in those experiencing direct attacks [19]. However, an identification

of post-traumatic stress incubation was challenged by significant effects of the prolonged peri-

traumatic stress development from the repeated trauma induction for more than a week [66].

Our behavioral paradigm overcomes these limitations in identifying stress incubation

process. With the behavioral foundation of our mouse paradigm, we propose that stress

incubation is a process caused by the activity-induced changes of global networks from ACC

after its mediation of traumatic experience. While researches on animal emotion usually

emphasize their ability to show human behavioral and physiological conditions (face validity),

to reproduce pharmacological effects in human (predictive validity), and to share the same

biological processes as human (construct validity) [15], a few studies have taken the possibility

of human-unique, animal-unique, and human-animal-sharing emotions under consideration [67].

We demonstrated human-animal-sharing emotions by the emotion-inferred behavior approach,

and showed in our paradigm that affective bonding, instead of fear memory, explains stress

incubation in mice.

Potential insights and applications to human PTSD

Although our study focused on stress incubation in mice, the observed psychological

homology between mice and humans suggests that our paradigm can be used as simple model

of psychobehavioral development in humans. Complex developmental trajectories of human

PTSD symptoms were demonstrated with associated physiological and environmental

regulators [8], yet the combination of psychological therapy and pharmacotherapy does not

provide a more efficacious treatment than psychological therapy alone for PTSD development

[68,69]. More than leading toward a better efficacy and lower risk in conducting psychological

treatments, our research provides a foundation to test pharmacotherapies in a system-level view

that integrates multiple mechanisms underlying highly diverse behavioral consequences

(Figure 14). The brain heavily integrates not only external but also internal causes [70,71,80,81,72–

79]. Amygdala, medial prefrontal cortex, and hippocampus are the brain regions traditionally

focused on in human PTSD research [82], with reports emphasizing the morphology of the right

hippocampus [83,84]. Found on highly controlled conditions in experiments that overcome major

difficulties in human studies, our data in laboratory mice provide an unexpected insight of

PTSD development, toward brain physiology on how the potentially slow and global change

of brain dynamics arises from an altered dynamic in social bonding circuits through their

interconnected nodes (Figures 13 and 14). The efficacy of pharmacotherapies could therefore

be improved.

While much remains to be considered before clinical applications, including a better

connection between our mouse results and human PTSD criteria as well as a corresponding

study in humans, our study establishes a solid basis to uncover psychological theories for

therapeutic strategies [3]. Compared with the classic view of PTSD development as a process of

complex associations [85], we propose to consider human PTSD development as a relatively

simple process with a unitary origin. While the core factor in a unitary model of human PTSD

is not necessarily social bonding, we suggest a special focus on affective bonding as the core

factor for patient diagnosed with witnessing PTSD.

We expect that behavioral signs of PTSD development could also be detected in

humans already shortly after the traumatic event. The fine-scale behavioral analyses we

introduced here provides a simple, non-invasive analytic tool to capture informative behavioral

details and is not limited to laboratory animals. It opens a new window for early detection and

prediction with the potential to prevent a development into PTSD.

Materials and Methods:

Study design

Sample size. No statistical methods were used to predetermine sample sizes. Animal numbers

were determined based on previous studies [9,86].

Rules for stopping data collection. Partner mice were expected to get minor injuries from

aggressor mice during aggressive encounters, which normally should be through attack bites

on the dorsal side of posterior trunk [86]. The aggressive encounter and all further experiments

were terminated once (i) the partner mouse showed severe bleeding or ataxia, or (ii) the

aggressor mice showed abnormal attack bites on any other body part. Partner mice that reached

the criteria (i) were euthanized. Aggressor mice that reached the criteria (ii) were terminated

its usage in any further experiments. If any aggressive sign (sideways threat, tail rattle, pursuit,

and attack bite) was shown by the partner mouse, all further experiments with the partner mouse,

aggressor mouse, and observer mouse were terminated.

Data inclusion/exclusion criteria. No data were excluded.

Outliers. No outliers were defined.

Selection of endpoints. The endpoints were prospectively selected. The criteria were approved by the Institutional Animal Care and Use Committee (IACUC) in the animal facility at the Okinawa Institute of Science and Technology (OIST) Graduate University, accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC).

Replicates. All behavioral tests were conducted in quintuplicate to octuplicate sampling replicates. All behavioral tests were conducted in single experimental cohorts. All other records were conducted in quadruplicate to septuplicate experimental cohorts.

Research objectives. The goal of this work is to identify diverse psychological aspects, temporal patterns, and associations of behavioral development in mice after a single trauma induction. Pre-specified hypotheses stated that (i) internal psychological development is already represented in behavioral details during early post-traumatic phase, while (ii) an association among trauma-induced states such as fear memory, increased anxiety, and social withdrawal explains PTSD-like behavioral development. Our data support the first pre-specified hypothesis, but not the second. All other hypotheses were suggested after initiation of the data analyses.

Research subjects or units of investigation. 8-week-old male C57BL/6J mice, 16-week-old female C57BL/6J mice, and 20-week-old or older male Slc:ICR mice were used.

Experimental design. We approached the research goal by developing a novel mouse model of psychosocial trauma under highly controlled conditions (psychosocial manipulations, subjective experiences, and genetic background) and by applying fine-scale analysis to standard behavioral tests. To induce acute witnessing trauma, a pair-housed mouse observed how its partner got bullied by a larger, aggressive mouse on the day of trauma induction. After this trauma, the observer mouse was isolated and developed behavioral differences compared to control mice in the ensuing weeks. The control groups included (i) mice isolated without experiencing trauma induction, (ii) mice isolated after observing how a stranger mouse got

bullied by a larger, aggressive mouse, (iii) mice isolated after exposed to a non-aggressive

stranger mouse, (iv) mice which were pair-housed with their defeat partners after observing

how its partner got bullied by a larger, aggressive mouse, (v) mice which were pair-housed

with strangers after observing how its partner got bullied by a larger, aggressive mouse, (vi)

mice isolated with their environment enriched, and (vii) mice isolated with daily injections of

fluoxetine. The behavioral tests included the light-dark box test, elevated plus-maze test, open

field test, locomotor activity test, active social contact test to a female stranger, active social

contact test to a male stranger, and partner-revisiting test. The non-behavioral tests included

the body mass measurement, nest wall height measurement, baseline corticosterone

concentration test, and ex vivo diffusion tensor imaging.

Randomization. Mice were from different litters. Mice were randomly paired. A focal mouse

was randomly selected from each pair of mice. Mice were randomly allocated into

experimental groups. Testing order among groups was counterbalanced. Strangers and

aggressors were randomly assigned.

Blinding. The investigator was blinded to behavioral outcomes but not to group allocation

during data collection and/or analysis.

Overview

All experimental procedures were approved by IACUC in the animal facility at the

OIST Graduate University, accredited by AAALAC. In total, 467 male C57BL/6J mice (CLEA

Japan, Inc.), 49 female C57BL/6J mice (CLEA Japan, Inc.), and 33 male Slc:ICR mice (Japan

SLC, Inc.; retired from used for breeding) were used in this study. In CLEA Japan, nursing

females were individually housed (CL-0103-2; 165×234×118 mm), while pups were separated

on P21 according to gender and housed ≤15 mice per cage (CL-0104-2; 206×317×125 mm).

Pups were re-arranged on P28 according to their weights and housed ≤13 mice per cage (CL-

0104-2). Mice were shipped in boxes each with 10 – 30 mice to the OIST Animal Facility. In

the OIST Animal Facility, mice were housed in 380×180×160-mm transparent holding cages

(Sealsafe Plus Mouse DGM - Digital Ready IVC; Tecniplast Inc., Quebec, Canada) bedded

with 100% pulp (FUJ9298101; Oriental Yeast Co., Ltd., Tokyo, Japan) under a 12-hr dark/light

cycle (350-lux white light) at a controlled temperature of 22.7 – 22.9 ºC, humidity of 51 – 53%,

and differential pressure of -14 – -8 Pa with food and water available ad libitum. Circadian

time (CT) is defined to start at mid-light period and described in a 24-hr format, i.e. light off at

CT 6:00.

The experimenter and caretakers wore laboratory jumpsuits, lab boots, latex gloves,

face masks, and hair nets when handling mice and performing experiments. Handling of mice

during the dark cycle was done under dim red light and mice were transported in a lightproof

carrier within the animal facility. For mice in experimental and control groups tested on the

same dates, the testing order was alternated. Surfaces of experimental apparatuses were wiped

with 70% ethanol in water and dry paper tissues after testing each mouse to remove olfactory

cues. Each mouse was only used for one behavioral test (in total 4 records with intervals of 6

– 21 days) to avoid confounded results due to cross-testing and to minimize measurement

effects on its psychological development [41].

Pre-traumatic period (Day-21 to Day0)

To establish partnerships between mice, a male C57BL/6J mouse (focal mouse; 8 weeks)

was pair-housed with another male C57BL/6J mouse (partner mouse; 8 weeks) for 3 weeks

(Day-21 to Day0, with trauma induction on Day0). The partner was initially marked by ear

punching. The holding cage was replaced once per week, with the last change 3 days before

the traumatic event (Day-3).

To establish the territory of an aggressor mouse in its homecage, an Slc:ICR mouse

(aggressor mouse; ≥20 weeks) was pair-housed with a female C57BL/6J mouse (female mouse;

16 weeks) for 3 weeks (Day-21 to Day0). The holding cage was replaced with a clean one once a week, with the last change one week before the traumatic event (Day-7).

Aggression level of aggressors was screened on Days -5, -3, -1 through intruder encounters [87] toward different screening mice to determine appropriate aggressors to be used for trauma induction on Day0. Aggression screening was carried out in the behavior testing room at 22.4 – 23.0 ºC, 53 – 58% humidity, -4 – -3 Pa differential pressure, and 57.1 dB(C) ambient noise level during the light period (CT 4:00 – 6:00) with 350-lux white light. After the female and pups with the aggressor were taken out of their homecage and kept in a clean holding cage in the behavior testing room, a 3-min aggression screening was started after a male C57BL/6J mouse (screening mouse; 10 weeks) was brought into the homecage of the aggressor, followed by covering the cage with a transparent acrylic lid. During screening, the aggressor freely interacted with the screening mouse. The aggressor was brought back to the holding room after the screening mouse was taken away from the aggressor's homecage and the female and pups were brought back to its homecage right after screening. Aggressors were selected for trauma induction on Day0 if they showed biting attacks on all of these screening days and the latencies to the initial bites on Day-3 and Day-1 were less than 20 s.

Trauma induction (Day0)

The following experimental assay emotionally introduced an acute traumatic experience in mice through a social process. The setup was the aggressor's homecage, divided into an 80×180-mm auditorium zone and a 300×180-mm battle arena by the insertion of a stainless-steel mash with 8×8-mm lattices. The cage was covered with a transparent acrylic lid. The behavioral procedure was carried out in the behavior testing room during CT 4:00 – 6:00, with 3 – 5 experiments done in parallel.

After the female and pups with the aggressor were taken out of their homecage, a divider was inserted into the aggressor's homecage, allowing the aggressor to freely behave in

the battle arena, but not to enter the auditorium zone. A 5-min aggression encounter session started after the focal mouse was brought to the auditorium zone and its partner to the battle arena. Tail rattling counts of the focal mouse during aggressive encounter were recorded by experimenter. The aggressive encounter session was followed by a 5-min stress infiltration session, in which the partner was brought to the focal mouse in the auditorium zone, while the aggressor remained in the battle arena. Right after the stress infiltration session, both focal mouse and its partner were brought back to their homecage in the behavior testing room for a 10-min resting period. The procedure was repeated 5 times with different aggressors. During each resting session, the aggressor stayed in its homecage without the divider before its next intruder encounter. Each aggressor had 3 – 5 encounters with resting periods of 10 – 30 min. After the 5th aggression encounter session, the focal mouse was placed back in its homecage where the nest had been razed, and brought back to the holding room. Partners from different pairs were brought to a new holding cage and housed in groups of 3 – 5 per cage. Right after the last intruder encounter for each aggressor, the female and pups were brought back to the homecage and returned to the holding room together with the aggressor.

Post-traumatic period (Day0 to Day28)

To investigate the behavior of focal mice after trauma induction (now called Partner-observing mice, Figure 2A), they were housed individually for 4 weeks after the procedure (Day0 to Day28). No environmental enrichment was provided, except to the Environment-enriched mice, and the holding cage was not changed during social isolation.

Control experiments

To differentiate behavioral consequences of the emotionally traumatic experience from consequences of social isolation, a control group of mice had their partners taken away and their nests razed during body weighing on Day0 without trauma induction (No-scenario-isolated mice, Figure 2B).

To test the influence of social relationship on the emotionally traumatic experience, a

control group of mice witnessed the traumatic events toward stranger mice of the same strain,

gender, and age instead (Stranger-observing mice, Figure 9A). In each iteration of the

aggression encounter, stress infiltration, and resting period, a different stranger mouse was

presented.

To identify the impacts of aggression during trauma induction, a control group of mice

experienced exposure to strangers of the same strain, gender, and age, instead of the aggressor

mice for trauma induction (Stranger-exposed mice, Figure 9B).

To examine the potential reversal effects of social support on the emotionally traumatic

experience, a control group of mice was kept pair-housed with their attacked partners after

trauma induction (Partner-buffered mice, Figure 9C).

To further test the critical component of social relationship in the potential social

support reversal, a control group of mice was kept pair-housed but with a stranger mouse after

trauma induction (Stranger-buffered mice, Figure 9D).

To characterize potential reversal effects through environmental factors besides social

factors, a control group of mice was housed individually with environmental enrichment,

provided with a pair of InnoDome™ and InnoWheel™ (Bio-Serv, Inc., Flemington, NJ, USA)

and a Gummy Bone (Petite, Green; Bio-Serv, Inc.), after trauma induction (Environment-

enriched mice, Figure 9E).

To demonstrate predictive validity of potential treatment on stress by an antidepressant,

a control group of mice was intraperitoneally injected with fluoxetine (2 µl/g of 10 mg/ml

fluoxetine hydrochloride dissolved in saline, i.e. 20 mg/kg; F132-50MG; Sigma-Aldrich, Inc.,

Saint Louis, MO, USA) once per day at CT 1:00 – 2:00 after trauma induction (FLX-treated

mice, Figure 9F).

Body mass and nest wall height

In the holding room, body masses of all individuals were recorded on Days -7, 0, 1, 7,

28, while the heights of nest walls built by each individual were recorded on Days 1, 7, 28.

The height of the nest wall was measured with 5-mm resolution using a transparent acrylic

ruler, while the mouse was weighed with 10-mg resolution on a balance. Mice were placed

back in their homecages right after recording.

Light-dark box test

The light-dark box test is an experimental assay to measure anxiety in rodents [88],

designed to evaluate their natural aversion to brightly lit areas against their temptation to

explore. The light-dark box setup consisted of two connected 200×200×250-mm non-

transparent PVC boxes, separated by a wall with a 50×30-mm door. The boxes were covered

with lids with white and infrared LED light illumination for the light and dark areas,

respectively, and CCD cameras in the centers (4-chamber system; O'Hara & Co., Ltd., Tokyo,

Japan). The floors of the boxes were white, while the walls of the boxes were white for the

light area and black for the dark area. Uniform illumination in the light area was 550 lux.

Behavioral tests were carried out on Days -7, 1, 7, and 28 in the behavior testing room at 22.7

– 23.0 ℃, 51 – 54% humidity, -11 – -9 Pa differential pressure, and 53.6 dB(C) ambient noise

level during dark period (CT 6:00 – 8:00).

After habituation for 10 min individually in the homecage in the behavior testing room

in darkness, the focal mouse was transferred to the dark area through a 50×50-mm side door.

A 5-min behavior record was started right after the side door of dark area was closed and the

door between light and dark areas was opened. Locomotion was recorded 2-dimensionally at

15 Hz from top-view with CCD video cameras. Right after recording, the mouse was returned

to its homecage, and brought back to the holding room.

Elevated plus-maze test

The elevated plus-maze test is an experimental assay to measure anxiety in rodents [89],

designed to evaluate their natural fear of falling and exposure against their temptation to

explore. The elevated plus-maze setup consisted of a gray PVC platform raised 500 mm above

the ground (single maze system; O'Hara & Co., Ltd.). The platform was composed of a 50×50-

mm square central platform, two opposing 50×250-mm open arms, and two opposing 50×250-

mm closed arms with 150-mm semi-transparent walls. Each of the two open arms emanated at

90° to each of the two closed arms, and vice versa. The apparatus was installed in a soundproof

box with white fluorescent lamp illumination (20 lux) and ventilators. Behavioral tests were

carried out on Days -7, 1, 7, 28 in the behavior testing room at 22.8 – 23.0 ºC, 53 – 56%

humidity, -13 – -11 Pa differential pressure, and 52.1 dB(C) ambient noise level during dark

period (CT 8:00 – 10:00).

After habituation for 10 min individually in the homecage in the behavior testing room

in darkness, the focal mouse was brought to the central platform of the elevated plus-maze,

facing the open arm on the opposite side from the door of the soundproof box. A 5-min

behavior recording was started right after the door of the soundproof box was closed.

Locomotion was recorded 2-dimensionally at 15 Hz from top-view with a CCD video camera

installed above the center of the central platform. Delineated entrances to open and closed

arms were defined at 50 mm from the center of the central platform. Right after recording, the

mouse was placed back in its homecage, and brought back to the holding room.

Open field test

The open field test is an experimental assay to measure anxiety in rodents [90], designed

to evaluate their spontaneous activity under a gradient of spatial uncertainty (high in the field

center and low along the walls and at the corners of the field). The open field setup consisted

of a 400×400×300-mm non-transparent gray PVC box with no cover, installed in a soundproof

box with white LED light illumination and ventilators (2-chamber system; O'Hara & Co., Ltd.).

Behavioral tests were carried out on Days -7, 1, 7, 28 in the behavior testing room at 22.8 –

23.0 ºC, 53 – 56% humidity, -13 – -11 Pa differential pressure, and 56.7 dB(C) ambient noise level during dark period (CT 8:00 – 10:00).

After habituation for 10 min individually in the homecage in the behavior testing room in darkness, the focal mouse was brought to the center of the open field arena under 20-lux uniform illumination, facing the wall on the opposite side from the door of the soundproof box. A 5-min behavior recording was started right after the door of the soundproof box was closed. Locomotion was recorded 2-dimensionally at 15 Hz from top-view with a CCD video camera installed above the center of the open field arena. Vertical activity of exploratory rearing behavior was recorded by the blocking of invisible infrared beams created and detected by photocell emitters and receptors, respectively, positioned 60 mm high on the walls of the open field box. A delineated center region was defined as the central 220×220 mm area. Right after recording, the mouse was placed back in its homecage, and returned to the holding room.

Locomotor activity test

The locomotor activity test is an experimental assay to measure spontaneous activity of rodents in an environment without an experimentally designed stressor. The locomotor activity setup consisted of a 200×200×250 mm non-transparent covered PVC box with infrared LED illumination and a CCD camera in the center (the dark area of the light-dark box setup, while the door between the light and dark areas was closed and fixed). The floor of the box was embedded with bedding material from the homecage of the focal mouse, while the walls of the box were black. Behavioral test was carried out on Days -7, 1, 7, 28 in the behavior testing room at 22.7 – 23.0 ºC, 51 – 54% humidity, -11 – -9 Pa differential pressure, and 53.6 dB(C) ambient noise level during dark period (CT 6:00 – 8:00).

After habituation for 30 min individually in the behavior testing box, a 1-hr behavior recording was started. The behavior testing box was not covered completely in order to allow air circulation. Locomotion was recorded 2-dimensionally at 15 Hz from top-view with the

CCD video camera. Right after recording, the mouse was returned to its homecage, and
brought back to the holding room.

Active social contact test

The active social contact test (also known as "social interaction test", but to be
distinguished with the one-session test using an open field with a social target freely behaving
in the field [91] or the one-session test placing a social target-containing cylinder into the center
of the testing subject's homecage for social instigation [92]) is a 2-session experimental assay to
measure social motivation in rodents [93]. The setup consists of a 400×400×300-mm non-
transparent gray PVC box with no cover, installed in a soundproof box with 20-lux white LED
illumination and ventilators. A 60×100×300-mm stainless-steel chamber with wire grid sides
was placed in the center of the wall on the opposite side from the door of the soundproof box.
The wire grid had 8×8 mm lattices at a height of 10 – 60 mm from the bottom. An ultrasound
microphone (CM16/CMPA; Avisoft Bioacoustics, Glienicke, Germany) with an acoustic
recording system (UltraSoundGate; Avisoft Bioacoustics) was hung outside the chamber, 100
mm above the ground. Behavioral tests were carried out on Days -7, 1, 7, 28 in the behavior
testing room at 22.8 – 23.0 ºC, 53 – 56% humidity, -13 – -11 Pa differential pressure, and 56.7
dB(C) ambient noise level during dark period (CT 8:00 – 10:00).

The social target used for active social contact tests was either a male or a female
C57BL/6J mouse (18 weeks), pair-housed with a partner of the same strain, gender, and age
for more than 2 weeks before the tests. The social target was adapted to the experimental
protocol one day before the tests in the behavior testing room during dark period (CT 8:00 –
9:00): After habituation for 5 min individually in the homecage in the soundproof box under
20-lux uniform illumination, the social target was brought into the chamber in the open field
arena under 20 lux uniform illumination. A male C57BL/6J mouse (11–16 weeks; from
partners of No-scenario-isolated mice in previous experiment) was then brought to the open
field arena for a 2.5-min spontaneous exploration and interaction with the social target. The

social target was then brought back to its homecage in the soundproof box under 20-lux uniform light for a 5-min rest. The social interaction procedure was repeated with a different male C57BL/6J mouse right afterward. After the social target had interacted with 4 different mice, it was returned to its homecage and brought back to the holding room.

On testing days, after 10-min habituation individually in its homecage in the behavior testing room in darkness, the first session of the active social contact test started by placing the focal mouse at the center of the open field arena under 20-lux uniform light, facing the empty chamber. A 2.5-min behavior recording started right after the door of the soundproof box was closed. Locomotion was recorded 2-dimensionally at 15 Hz from top-view with a CCD video camera installed above the center of the open field arena. Ultrasonic vocalization was recorded at 250 kHz. In the second session of the active social contact test, which followed the first session, the social target was brought into the chamber. Another 2.5-min behavior recording started as soon as the door of the soundproof box was closed. Right afterward, the focal mouse was returned to its homecage and brought back to the holding room.

The focal mouse experienced active social contact tests with different social targets on different recording days (Days -7, 1, 7, 28), while different focal mice were tested with the same social target on the same recording day (5 – 10 records). The social target remained in its homecage in a soundproof box under 20-lux uniform illumination before and between each test. A delineated interaction zone was taken as the region within 80 mm of the edges of the chamber. Social approaches of the focal mouse poking its nose toward the social target were recorded manually using the event recording software, tanaMove ver0.09 (http://www.mgrl-lab.jp/tanaMove.html).

Partner-revisiting test

The partner-revisiting test is a memory-based experimental assay to measure social bonding in rodents (sharing similar concept of "familiar v.s. novel social target recognition",

but to be distinguished with the three-chamber paradigm test [94]). The partner-revisiting setup

was the uncovered homecage of the focal mouse, installed in a soundproof box with white LED

illumination and ventilators (O'Hara & Co., Ltd., Tokyo, Japan). The long sides of the

homecage were parallel to the door of soundproof box. The partner-revisiting test was carried

out on Day28 in the behavior testing room at 22.8 – 23.0 ºC, 53 – 56% humidity, -13 – -11 Pa

differential pressure, and 56.7 dB(C) ambient noise level during light period (CT 4:00 – 6:00)

with 350-lux light intensity.

The previously separated partner of the focal mouse, being a social target in the test,

was initially sedated with 3%v/v isoflurane in oxygen, and then anesthetized by intraperitoneal

(i.p.) injection of a mixture of medetomidine (domitor, 3%v/v in saline, 0.3 mg/kg), midazolam

(dormicum, 8%v/v in saline, 4 mg/kg), and butorphanol (vetorphale, 10%v/v in saline,5 mg/kg).

Also, a stranger mouse (15 weeks; a separated partner of a Partner-observing or Buffered mouse

for testing a No-scenario-isolated, Stranger-observing, or Stranger-exposed mouse, and *vice*

versa) was anesthetized as an alternative social target. Both anesthetized mice were kept on a

heating pad at 34ºC (B00O5X4LQ2; GEX Co., Ltd., Osaka, Japan) to maintain their body

temperatures before the test.

The focal mouse was brought to a clean, uncovered holding cage in the soundproof box

under 50-lux uniform illumination for 5-min habituation, while its homecage was placed in

another soundproof box under 50-lux uniform light. During habituation of the focal mouse,

the anesthetized social targets were injected with atipamezole hydrochloride (antisedan; 6%v/v

in saline for 0.3 mg/kg, i.p.) to induce recovery from anesthesia. During the waking-up period,

the social targets were still immobilized and not able to actively interact with the focal mouse

during the following recording, but showed enough social cues to be attractive for the focal

mouse. The immobilized social targets were then placed in the homecage of the focal mouse

with their nose pointing toward the center of the short side of the wall (10 mm of nose-to-wall

distance) with their bellies facing the door of the soundproof box. After habituation, the focal

mouse was brought to the center of its homecage, facing the long side of the homecage wall on

the opposite side from the door of soundproof box. A 10-min behavior record started right

after the door of the soundproof box was closed. Locomotion was recorded 2-dimensionally

at 15 Hz from top-view with a CCD video camera installed above the center of the homecage.

Right after recording, social targets were taken out of the focal mouse's homecage and the focal

mouse was brought back to the holding room.

Social contacts including sniffing, allogrooming, and pushing of the focal mouse

toward each of the social targets were recorded manually using the event recording software,

tanaMove ver0.09 (http://www.mgrl-lab.jp/tanaMove.html).

Baseline plasma corticosterone concentration test

The baseline plasma corticosterone (CORT) concentration test is a competitive-

inhibition enzyme-linked immunosorbent assay (ELISA) to measure physiological stress level

in rodents, designed to quantitatively determinate CORT concentrations in blood plasma. The

sample collection was carried out on Days -7, 1, 7, 28 in the behavior testing room at 22.4 –

23.0 ºC, 53 – 58% humidity, -4 – -3 Pa differential pressure, and 57.1 dB(C) ambient noise

level during CT 4:00 – 6:00 with 350-lux white light.

After habituation for 30 min individually in the homecage in the behavior testing room,

the mouse was initially sedated with 3%v/v isoflurane in oxygen. Six drops of blood from the

facial vein pricked by a 18G needle were collected in a EDTA-lined tube [K2 EDTA (K2E)

Plus Blood Collection Tubes, BD Vacutainer; Becton, Dickinson and Company (BD), Franklin

Lakes, NJ, USA] and kept on ice. Right after collection, the mouse was returned to its

homecage, and brought back to the holding room. Whole blood samples were then centrifuged

(MX-300; Tomy Seiko Co., Ltd., Tokyo, Japan) at 3,000 rpm for 15 min at 4°C. Plasma

supernatant was decanted and kept at -80°C until the measurement on Day 29.

CORT concentrations in blood plasma were tested with Mouse Corticosterone (CORT)

ELISA Kit (MBS703441, 96-Strip-Wells; MyBioSource, Inc., San Diego, USA; stored at 4°C

before use) on Day 29. All reagents [assay plate (96 wells, pre-coated with goat-anti-rabbit

antibody), standards (0, 0.1, 0.4, 1.6, 5, and 20 ng/ml of CORT), rabbit-anti-CORT antibody,

HRP-conjugated CORT, concentrated wash buffer (20x phosphate-buffered saline (PBS)),

3,3',5,5'-tetramethylbenzidine (TMB) color developing agent (substrates A and B), and TMB

stop solution] and samples were brought to room temperature for 30 min before use. Collected

plasma samples after thawing were centrifuged again (MX-300; Tomy Seiko Co.) at 3,000 rpm

for 15 min at 4°C. 20 µl of standard or sample was added per well, assayed in duplicate, with

blank wells set without any solution. After 20 µl of HRP-conjugated CORT was added to each

well except to the blank wells, 20 µl of rabbit-anti-CORT antibody was added to each well and

mixed. After incubation for 1 hour at 37°C, each well was aspirated and washed, repeated for

3 times, by filling each well with 200 µl of wash buffer (diluted to 1x PBS) using a squirt bottle,

standing for 10 s, and completely removing liquid at each step. After the last wash and the

removal of any remaining wash buffer by decanting, the plate was inverted and blotted against

clean paper towels. After TMB color developing agent (20 µl of substrate A and 20 µl of

substrate B) was added to each well, mixed, and incubated for 15 min at 37°C in dark, 20µl of

TMB stop solution was added to each well and mixed by gently tapping the plate. The optical

density (O.D.) of each well was determined, within 10 min, using a microplate reader

(Multiskan GO; Thermo Fisher Scientific, Inc., Waltham, MA, USA) measuring absorbance at

450 nm, with correction wavelength set at 600–630 nm.

CORT concentrations were calculated from the O.D. results using custom scripts

written in MATLAB R2015b (MathWorks). The duplicate O.D. readings for each standard and

sample was averaged and subtracted the average O.D. of the blanks, $X = \langle O.D. \rangle -$

$\langle O.D. \rangle_{blank}$. A standard curve was determined by a four parameter logistic (4PL) regression

fitting the equation $\rho_{CORT}(X_{standard}) = d + \frac{a-d}{1+(\frac{X_{standard}}{c})^b}$, where ρ_{CORT} is the CORT

concentration, a is the minimum asymptote, b is the Hill's slope, c is the inflection point, and

d is the maximum asymptote. CORT concentrations of the samples were calculated from the

fitted 4PL equation with respected to X_{sample}.

Ex vivo diffusion tensor imaging

Ex vivo diffusion tensor imaging (DTI) is a magnetic resonance imaging (MRI)

technique to determinate structural information about tissues [95], designed to measure the

restricted diffusion of water in tissue. The sample collection was carried out on Day 28 in the

necropsy room at 22.4 - 22.5 ºC, 53 - 54 % humidity, and 10 - 12 Pa differential pressure during

CT 4:00 - 6:00 with 750-lux white light.

After mice were brought individually in their homecages to the necropsy room, they

were initially sedated with 3%v/v isoflurane in oxygen, then deeply anaesthetized with a

ketamine-xylazine mixture (>30 μl/g body weight of 100 mg/ml ketamine and 20 mg/ml

xylazine), and perfused transcardially. The perfusates, in a two-step procedure, were (i) 20 ml

of ice cold 1x phosphate-buffered saline (PBS), and (ii) 20 ml of ice cold 4% paraformaldehyde,

0.2% sodium meta-periodate, and 1.4% lysine in PBS. Mouse skull including the brain was

removed and stored in the perfusate (ii) at 4 ºC for 2 weeks. Each skull with the brain was then

transferred into 2 mM gadolinium with diethylenetriaminepentaacetic acid (Gd-DTPA) and

0.5% azide in PBS for 2 weeks.

Isolated fixed brains within the skulls were positioned in an acrylic tube filled with

fluorinert (Sumitomo 3M Ltd., Tokyo, Japan) to minimize the signal intensity attributable to

the medium surrounding the brain during MRI scanning. All MRI was performed with an 11.7-

T MRI system (BioSpec 117/11; Bruker Biospec, Ettlingen, Germany) using ParaVision 6.0.1

software (Bruker Biospec, Ettlingen, Germany) for data acquisition. The inner diameter of the

integrated transmitting and receiving coil (Bruker Biospec, Ettlingen, Germany) was 35 mm

for the ex vivo MRI. DTI data were acquired by using a 3-D diffusion-weighted spin-echo

imaging sequence, with repetition time (TR) = 267 ms, echo time (TE) = 18.5 ms, b-value =

2,000 s/mm2, and 30 non-collinear directions. Five T2-weighted measurements were acquired

together with DTI, for one every 6 diffusion measurements. The acquisition matrix was

216×216×168 over a 27.0×27.0×21.0 mm3 field of view, resulting in a native isotropic image

resolution of 125 μm. Total acquisition time was 96 hr.

MRI data was processed using custom scripts written in MATLAB R2015b

(MathWorks). All 30 DTI and 5 T2 3-D images were masked by thresholding at the half of

mean values of diffusion weights for each voxel and omiting clusters smaller than 10 voxels.

After diffusion tensor of each voxel was estimated by solving the Stejskal-Tanner equation

through linear regression [96], the 3 eigenvalues (λ_1, λ_2, and λ_3) with respect to the 3 axes of the

diffusion ellipsoid (the longest, middle, and shortest axes, respectively) were calculated by

eigenvalue decomposition of the diffusion tensor. Four focused DTI-based measures [97] are the

mean diffusivity (MD) that represents membrane density

945
$$MD = \langle\lambda\rangle = \frac{\lambda_1 + \lambda_2 + \lambda_3}{3},$$

946   axial diffusivity (AD) that represents neurite organization

947
$$AD = \lambda_1,$$

948   radial diffusivity (RD) that represents myelination

949
$$MD = \frac{\lambda_2 + \lambda_3}{2},$$

950   and fractional anisotropy (FA) that represents average microstructural integrity

951
$$FA = \sqrt{\frac{3 \times [(\lambda_1 - \langle\lambda\rangle)^2 + (\lambda_2 - \langle\lambda\rangle)^2 + (\lambda_3 - \langle\lambda\rangle)^2]}{2 \times (\lambda_1{}^2 + \lambda_2{}^2 + \lambda_3{}^2)}}.$$

After the registration of these 3-D DTI-based brain maps to a template brain atlas

(DSURQE Atlas; https://wiki.mouseimaging.ca/display/MICePub/Mouse+Brain+Atlases),

mean values of these DTI-based quantities were identified in a total of 244 brain regions ($)

for each individual. Based on the FA maps, DTI-based long-range fiber tracking from a focal

seed region ($B) was calculated with 2 samplings, distance of forward fiber in one step = 50

μm, and thresholds of minimal fiber length = 750 μm, maximal fiber length = 75,000 μm,

maximal fiber deviation angle = 57.3º, and minimal FA for keeping tracking = 0.4.

Video image process

Video image data was processed using custom scripts written in MATLAB R2015b

(MathWorks). Each video frame from a recorded AVI video file was read as a 2-dimensional

matrix with an 8-bit gray scale. Each of these matrices was then divided by a background

matrix read from a TIF image file of the background taken before bringing the test mouse to

the setup. The centroid of the area with non-one values in each matrix ratio was taken as the

position of the mouse at this specific time point. Speed was calculated as the distance between

temporally adjacent positions multiplied by 15 (15-Hz recording). Freezing periods were

sorted out if the area of the mouse body between temporally adjacent frames was less than 20

mm^2.

Audio signal process

Audio signal data was processed with custom scripts written in MATLAB R2015b

(MathWorks). Each recorded WAV audio file was read and transformed into a spectrogram

using fast Fourier transform with non-overlapping 0.4-ms time windows. To identify the time

segments with ultrasonic vocalization signals, recordings were thresholded at a power spectral

density (PSD) ≥-75 dB/Hz, and time segments with averaged PSD between 0–50 kHz higher

than that between 50–120 kHz were removed. The duration of remaining time segments was

calculated.

Statistical analysis

Numerical data were analyzed with custom scripts written in MATLAB R2015b

(MathWorks). Statistical significance of the difference between 2 mean values was estimated

with two-tailed, two-sample Student's t-test, except of DTI quantities which were estimated

with one-tailed, two-sample Student's t-test. Statistical significance of the difference between

2 median values (vocalization analysis; Figure 8F) was estimated using one-tailed Mann–

Whitney U test.

To capture fine-scale behavioral details of location within the light-dark box and the

elevated plus-maze (Figures 4 and 5), we computed T(x), the cumulative probability of finding

position $\leq$x, for each individual (light traces) for all measured locations (a collection of

locations from all mice for the statistics). We then show the average across the control group

(bold blue trace) and the Partner-observing group (bold red trace). We compared the averages

of each group with a two-tailed, two-sample Student's t-test and plot the resulting p-values,

presented as -log(p), the negative logarithm of p-values. We also show the box plot (the

minimum, lower quartile, median, upper quartile, and maximum) of -log(p) values collapsed

across all measured locations. To capture the fine- scale behavioral details of speed, we

followed a similar procedure as above, but with U(v), the cumulative distribution function of

finding speed $\leq$v.

To evaluate the uncertainty of the percentage for each tail rattle count (Figure 10A), we

created 10,000 bootstrapped data sets where each sample was randomly picked with

replacement from the original data set. Each bootstrapped data set had the same sample size as

the original data set. The standard error was taken as the standard deviation of the bootstrapped

percentages for a tail rattle count. A similar procedure was carried out to evaluate the standard

error of mean for the percentage of time spent in each behavior in the partner-revisiting test

(Figures 10H and 11E), where each sample of the bootstrapped data sets was a set of the

percentages of the three classified behaviors (partner concern, stranger concern, and non-social activity/sniffing at social targets) from a mouse record. Standard errors of means for other results were estimated with the formula $\sigma/\sqrt{n}$, where σ is the sample standard deviation and n is the sample size.

Acknowledgments: We thank Tsuyoshi Koide (NIG, Japan), Aki Takahashi (University of Tsukuba), Teruhiro Okuyama (MIT), Charles Yokoyama (RIKEN BSI), Ai Koizumi (CiNet), and Tsai-Wen Chen (National Yang-Ming University) for comments, Tadashi Yamamoto (OIST) for sharing behavior testing setups, and Steven D. Aird (OIST) for editing the manuscript.

Competing interests: The authors declare no competing financial interests.

Funding: We acknowledge a JSPS KAKENHI Grant (JP 16J10077) by DC1 Student Research Fellowship to R.X.L. and OIST internal funding to B.K. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions: R.X.L. conceptualized the study, designed the research, performed experiments, analyzed data, interpreted results, wrote the original draft, and revised the article; G.J.S. advised on data analysis and revised the article; B.K. advised on research design and revised the article.

References:

1.	DSM-III. Diagnostic and statistical manual of mental disorders (DSM-III). *Am. Psychiatr. Assoc.*

2.	DSM-5. Diagnostic and statistical manual of mental disorders (DSM-5). *Am. Psychiatr. Assoc.* 947 (2013). doi:10.1176/appi.books.9780890425596

3.	Schnyder, U. & Cloitre, M. *Evidence based treatments for trauma-related psychological disorders : a practical guide for clinicians.*

4.   Ehlers, A. & Clark, D. M. A cognitive model of posttraumatic stress disorder. *Behav.*

Res. Ther. **38,** 319–345 (2000).

5.   Siegmund, A. & Wotjak, C. T. Toward an Animal Model of Posttraumatic Stress

Disorder. *Ann. N. Y. Acad. Sci.* **1071,** 324–334 (2006).

6.   Hayes, J. P., VanElzakker, M. B. & Shin, L. M. Emotion and cognition interactions in

PTSD: a review of neurocognitive and neuroimaging studies. *Front. Integr. Neurosci.* **6,**

89 (2012).

7.   Zoladz, P. R. & Diamond, D. M. Current status on behavioral and biological markers of

PTSD: A search for clarity in a conflicting literature. *Neurosci. Biobehav. Rev.* **37,** 860–

895 (2013).

8.   Bryant, R. A., O'Donnell, M. L., Creamer, M., McFarlane, A. C. & Silove, D. A

Multisite Analysis of the Fluctuating Course of Posttraumatic Stress Disorder. *JAMA*

Psychiatry **70,** 839 (2013).

9.   Ben-Ami Bartal, I., Decety, J. & Mason, P. Empathy and pro-social behavior in rats.

Science **334,** 1427–30 (2011).

10.   Ben-Ami Bartal, I., Rodgers, D. A., Bernardez Sarria, M. S., Decety, J. & Mason, P.

Pro-social behavior in rats is modulated by social experience. *Elife* **3,** (2014).

11.   Burkett, J. P. *et al.* Oxytocin-dependent consolation behavior in rodents. *Science* **351,**

375–8 (2016).

12.   Smith, M. L., Hostetler, C. M., Heinricher, M. M. & Ryabinin, A. E. Social transfer of

pain in mice. *Sci. Adv.* **2,** e1600855–e1600855 (2016).

13.   Cryan, J. F. & Holmes, A. The ascent of mouse: advances in modelling human

depression and anxiety. *Nat. Rev. Drug Discov.* **4,** 775–790 (2005).

14.   Kaouane, N. *et al.* Glucocorticoids can induce PTSD-like memory impairments in mice.

Science **335,** 1510–3 (2012).

15.   Calhoon, G. G. & Tye, K. M. Resolving the neural circuits of anxiety. *Nat. Neurosci.*

18, 1394–1404 (2015).

16.   Tovote, P., Fadok, J. P. & Lüthi, A. Neuronal circuits for fear and anxiety. *Nat. Rev.*

Neurosci. **16,** 317–331 (2015).

17.   Davis, M. Sensitization of the acoustic startle reflex by footshock. *Behav. Neurosci.* **103,**

495–503 (1989).

18.   Pamplona, F. A. *et al.* Prolonged fear incubation leads to generalized avoidance behavior

in mice. *J. Psychiatr. Res.* **45,** 354–360 (2011).

19.   Warren, B. L. *et al.* Neurobiological Sequelae of Witnessing Stressful Events in Adult

Mice. *Biol. Psychiatry* **73,** 7–14 (2013).

20.   Tsuda, M. C., Yeung, H.-M., Kuo, J. & Usdin, T. B. Incubation of Fear Is Regulated by

TIP39 Peptide Signaling in the Medial Nucleus of the Amygdala. *J. Neurosci.* **35,**

12152–61 (2015).

21.   Sillivan, S. E. *et al.* Susceptibility and Resilience to Posttraumatic Stress Disorder–like

Behaviors in Inbred Mice. *Biol. Psychiatry* **82,** 924–933 (2017).

22.   Balogh, S. A., Radcliffe, R. A., Logue, S. F. & Wehner, J. M. Contextual and cued fear

conditioning in C57BL/6J and DBA/2J mice: Context discrimination and the effects of

retention interval. *Behav. Neurosci.* **116,** 947–957 (2002).

23.   Philbert, J. *et al.* Acute inescapable stress exposure induces long-term sleep disturbances

and avoidance behavior: A mouse model of post-traumatic stress disorder (PTSD).

Behav. Brain Res. **221,** 149–154 (2011).

24.   Sial, O. K., Warren, B. L., Alcantara, L. F., Parise, E. M. & Bolaños-Guzmán, C. A.

Vicarious social defeat stress: Bridging the gap between physical and emotional stress.

J. Neurosci. Methods **258,** 94–103 (2016).

25. Houston, F. P., Stevenson, G. D., McNaughton, B. L. & Barnes, C. A. Effects of age on
the generalization and incubation of memory in the F344 rat. *Learn. Mem.* **6,** 111–9
(1999).

26. Goto, T. *et al.* Subchronic and mild social defeat stress accelerates food intake and body
weight gain with polydipsia-like features in mice. *Behav. Brain Res.* **270,** 339–348
(2014).

27. Otabi, H., Goto, T., Okayama, T., Kohari, D. & Toyoda, A. The acute social defeat stress
and nest-building test paradigm: A potential new method to screen drugs for depressive-
like symptoms. *Behav. Processes* **135,** 71–75 (2017).

28. Ben-Ami Bartal, I. *et al.* Anxiolytic Treatment Impairs Helping Behavior in Rats. *Front.*
Psychol. **7,** 850 (2016).

29. DeMorrow, S., DeMorrow & Sharon. Role of the Hypothalamic–Pituitary–Adrenal Axis
in Health and Disease. *Int. J. Mol. Sci.* **19,** 986 (2018).

30. Ramos, A. Animal models of anxiety: do I need multiple tests? *Trends Pharmacol. Sci.*
29, 493–8 (2008).

31. Archer, J. Tests for emotionality in rats and mice: A review. *Anim. Behav.* **21,** 205–235
(1973).

32. Grupe, D. W. & Nitschke, J. B. Uncertainty and anticipation in anxiety: an integrated
neurobiological and psychological perspective. *Nat. Rev. Neurosci.* **14,** 488–501 (2013).

33. Kumar, V., Bhat, Z. A. & Kumar, D. Animal models of anxiety: A comprehensive
review. *J. Pharmacol. Toxicol. Methods* **68,** 175–183 (2013).

34. Pessoa, L. On the relationship between emotion and cognition. *Nat. Rev. Neurosci.* **9,**
148–158 (2008).

35. Cloninger, C. R. Anxiety and Theories of Emotion. *Handb. Anxiety* **2,** 1–29 (1988).

36.   Adler, L. A., Kunz, M., Chua, H. C., Rotrosen, J. & Resnick, S. G. Attention-
1100   Deficit/Hyperactivity Disorder in adult patients with Posttraumatic stress disorder
1101   (PTSD): Is ADHD a vulnerability factor? *J. Atten. Disord.* **8,** 11–16 (2004).

1102   37.   Mostoufi, S. *et al.* Pain sensitivity in posttraumatic stress disorder and other anxiety
1103   disorders: a preliminary case control study. *Ann. Gen. Psychiatry* **13,** 31 (2014).

1104   38.   Andrews, B., Brewin, C. R., Philpott, R. & Stewart, L. Delayed-Onset Posttraumatic
1105   Stress Disorder: A Systematic Review of the Evidence. *Am. J. Psychiatry* **164,** 1319–
1106   1326 (2007).

1107   39.   Bryant, R. A. *et al.* Acute and Chronic Posttraumatic Stress Symptoms in the Emergence
1108   of Posttraumatic Stress Disorder. *JAMA Psychiatry* **74,** 135 (2017).

1109   40.   Anderson, D. J. Circuit modules linking internal states and social behaviour in flies and
1110   mice. *Nat. Rev. Neurosci.* **17,** 692–704 (2016).

1111   41.   Krishnan, V. *et al.* Molecular adaptations underlying susceptibility and resistance to
1112   social defeat in brain reward regions. *Cell* **131,** 391–404 (2007).

1113   42.   Sandi, C. & Haller, J. Stress and the social brain: behavioural effects and neurobiological
1114   mechanisms. *Nat. Rev. Neurosci.* **16,** 290–304 (2015).

1115   43.   Yang, T. *et al.* Social Control of Hypothalamus-Mediated Male Aggression. *Neuron* **95,**
1116   955–970.e4 (2017).

1117   44.   Chung, M. C. *et al.* Coping with post-traumatic stress symptoms following relationship
1118   dissolution. *Stress Heal.* **19,** 27–36 (2003).

1119   45.   Kogan, J. H., Franklandand, P. W. & Silva, A. J. Long-term memory underlying
1120   hippocampus-dependent social recognition in mice. *Hippocampus* **10,** 47–56 (2000).

1121   46.   Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S. & Tonegawa, S. Ventral CA1 neurons
1122   store social memory. *Science* **353,** 1536–1541 (2016).

1123   47.   Cohen, S. & Wills, T. A. *Stress, Social Support, and the Buffering Hypothesis.*
1124   *Psychologkal Bulletin* **98,** (1985).

1125   48.   Yu, Y.-Q., Barry, D. M., Hao, Y., Liu, X.-T. & Chen, Z.-F. Molecular and neural basis

1126   of contagious itch behavior in mice. *Science* **355,** 1072–1076 (2017).

1127   49.   Li, Y. *et al.* Serotonin neurons in the dorsal raphe nucleus encode reward signals. *Nat.*

1128   *Commun.* **7,** 10503 (2016).

1129   50.   Fischer, A. G. & Ullsperger, M. An Update on the Role of Serotonin and its Interplay

1130   with Dopamine for Reward. *Front. Hum. Neurosci.* **11,** 484 (2017).

1131   51.   McIntyre, D. C. & Kelly, M. E. The Parahippocampal Cortices and Kindling. *Ann. N. Y.*

1132   *Acad. Sci.* **911,** 343–354 (2006).

1133   52.   McIntyre, D. C. The Kindling Phenomenon. *Model. Seizures Epilepsy* 351–363 (2006).

1134   doi:10.1016/B978-012088554-1/50030-X

1135   53.   Rau, V., DeCola, J. P. & Fanselow, M. S. Stress-induced enhancement of fear learning:

1136   An animal model of posttraumatic stress disorder. *Neurosci. Biobehav. Rev.* **29,** 1207–

1137   1223 (2005).

1138   54.   Knox, D. *et al.* Single prolonged stress disrupts retention of extinguished fear in rats.

1139   *Learn. Mem.* **19,** 43–9 (2012).

1140   55.   Walf, A. A. & Frye, C. A. The use of the elevated plus maze as an assay of anxiety-

1141   related behavior in rodents. *Nat. Protoc.* **2,** 322–328 (2007).

1142   56.   Garcia, A. M. B., Martinez, R. C. R. & Morato, S. Preference for the light compartment

1143   of a light/dark cage does not affect rat exploratory behavior in the elevated plus-maze.

1144   *Psychol. & Neurosci.* **1,** 73–80 (2008).

1145   57.   Hascoët, M., Bourin, M. & Nic Dhonnchadha, B. Á. The mouse ligth-dark paradigm: A

1146   review. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **25,** 141–166 (2001).

1147   58.   Crusio, W. E. in *Behavioral Genetics of the Mouse* (eds. Crusio, W. E., Sluyter, F.,

1148   Gerlai, R. T. & Pietropaolo, S.) 148–154 (Cambridge University Press, 2013).

1149   doi:10.1017/CBO9781139541022.016

59. Kulesskaya, N. & Voikar, V. Assessment of mouse anxiety-like behavior in the light–
dark box and open-field arena: Role of equipment and procedure. *Physiol. Behav.* **133,**
30–38 (2014).

60. Ennaceur, A. Tests of unconditioned anxiety — Pitfalls and disappointments. *Physiol.*
Behav. **135,** 55–71 (2014).

61. Carobrez, A. P. & Bertoglio, L. J. Ethological and temporal analyses of anxiety-like
behavior: The elevated plus-maze model 20 years on. *Neurosci. Biobehav. Rev.* **29,**
1193–1205 (2005).

62. Engin, E. & Treit, D. The effects of intra-cerebral drug infusions on animals'
unconditioned fear reactions: A systematic review. *Prog. Neuro-Psychopharmacology*
Biol. Psychiatry **32,** 1399–1419 (2008).

63. Henriques-Alves, A. M. & Queiroz, C. M. Ethological Evaluation of the Effects of
Social Defeat Stress in Mice: Beyond the Social Interaction Ratio. *Front. Behav.*
Neurosci. **9,** 364 (2016).

64. Hånell, A. & Marklund, N. Structured evaluation of rodent behavioral tests used in drug
discovery research. *Front. Behav. Neurosci.* **8,** 252 (2014).

65. Anderson, D. J. & Perona, P. Toward a Science of Computational Ethology. *Neuron* **84,**
18–31 (2014).

66. Golden, S. A., Covington, H. E., Berton, O. & Russo, S. J. A standardized protocol for
repeated social defeat stress in mice. *Nat. Protoc.* **6,** 1183–1191 (2011).

67. Anderson, D. J. & Adolphs, R. A framework for studying emotions across species. *Cell*
157, 187–200 (2014).

68. Mataix-Cols, D. *et al.* D-Cycloserine Augmentation of Exposure-Based Cognitive
Behavior Therapy for Anxiety, Obsessive-Compulsive, and Posttraumatic Stress
Disorders. *JAMA Psychiatry* **74,** 501 (2017).

69.   Hetrick, S. E., Purcell, R., Garner, B. & Parslow, R. Combined pharmacotherapy and

1176   psychological therapies for post traumatic stress disorder (PTSD). *Cochrane Database*

1177   *Syst. Rev.* (2010). doi:10.1002/14651858.CD007316.pub2

1178   70.   Funamizu, A., Kuhn, B. & Doya, K. Neural substrate of dynamic Bayesian inference in

1179   the cerebral cortex. *Nat. Neurosci.* **19,** 1682–1689 (2016).

1180   71.   Larkum, M. A cellular mechanism for cortical associations: an organizing principle for

1181   the cerebral cortex. *Trends Neurosci.* **36,** 141–151 (2013).

1182   72.   Lee, R. X., Huang, J.-J., Huang, C., Tsai, M.-L. & Yen, C.-T. Plasticity of cerebellar

1183   Purkinje cells in behavioral training of body balance control. *Front. Syst. Neurosci.* **9,**

1184   1–15 (2015).

1185   73.   Lee, R. X., Huang, J. J., Huang, C., Tsai, M. L. & Yen, C. T. Collateral projections from

1186   vestibular nuclear and inferior olivary neurons to lobules I/II and IX/X of the rat

1187   cerebellar vermis: A double retrograde labeling study. *Eur. J. Neurosci.* **40,** 2811–2821

1188   (2014).

1189   74.   Donaldson, G. P., Melanie Lee, S. & Mazmanian, S. K. Gut biogeography of the

1190   bacterial microbiota. *Nat. Publ. Gr.* **14,** (2015).

1191   75.   Tononi, G., Boly, M., Massimini, M. & Koch, C. Integrated information theory: from

1192   consciousness to its physical substrate. *Nat. Rev. Neurosci.* **17,** 450–461 (2016).

1193   76.   Murugan, M. *et al.* Combined Social and Spatial Coding in a Descending Projection

1194   from the Prefrontal Cortex. *Cell* **171,** 1663–1677.e16 (2017).

1195   77.   Zelikowsky, M. *et al.* The Neuropeptide Tac2 Controls a Distributed Brain State Induced

1196   by Chronic Social Isolation Stress. *Cell* **173,** 1265–1279.e19 (2018).

1197   78.   Kohl, J. *et al.* Functional circuit architecture underlying parental behaviour. *Nature* **556,**

1198   326–331 (2018).

1199   79.   Remedios, R. *et al.* Social behaviour shapes hypothalamic neural ensemble

1200   representations of conspecific sex. *Nature* **550,** 388–392 (2017).

80.　Matias, S., Lottem, E., Dugué, G. P. & Mainen, Z. F. Activity patterns of serotonin
neurons underlying cognitive flexibility. *Elife* **6,** e20552 (2017).

81.　Roome, C. J. & Kuhn, B. Simultaneous dendritic voltage and calcium imaging and
somatic recording from Purkinje neurons in awake mice. *Nat. Commun.* **9,** 3388 (2018).

82.　Shin, L. M., Rauch, S. L. & Pitman, R. K. Amygdala, Medial Prefrontal Cortex, and
Hippocampal Function in PTSD. *Ann. N. Y. Acad. Sci.* **1071,** 67–79 (2006).

83.　Gilbertson, M. W. *et al.* Smaller hippocampal volume predicts pathologic vulnerability
to psychological trauma. *Nat. Neurosci.* **5,** 1242–1247 (2002).

84.　Pavić, L. *et al.* Smaller right hippocampus in war veterans with posttraumatic stress
disorder. *Psychiatry Res. Neuroimaging* **154,** 191–198 (2007).

85.　McFarlane, A. C. The long-term costs of traumatic stress: intertwined physical and
psychological consequences. *World Psychiatry* **9,** 3–10 (2010).

86.　Takahashi, A. *et al.* Glutamate input in the dorsal raphe nucleus as a determinant of
escalated aggression in male mice. *J. Neurosci.* **35,** 6452–63 (2015).

87.　Miczek, K. A. & O'Donnell, J. M. Intruder-evoked aggression in isolated and
nonisolated mice: Effects of psychomotor stimulants and l-Dopa. *Psychopharmacology*
(Berl). **57,** 47–55 (1978).

88.　Crawley, J. & Goodwin, F. K. Preliminary report of a simple animal behavior model for
the anxiolytic effects of benzodiazepines. *Pharmacol. Biochem. Behav.* **13,** 167–170
(1980).

89.　Pellow, S., Chopin, P., File, S. E. & Briley, M. Validation of open : closed arm entries
in an elevated plus-maze as a measure of anxiety in the rat. *J. Neurosci. Methods* **14,**
149–167 (1985).

90.　Hall, C. & Ballachey, E. L. A study of the rat's behavior in a field. A contribution to
method in comparative psychology. *Univ. Calif. Publ. Psychol.* 1–12 (1932).

91.   Arakawa, T. *et al.* A male-specific QTL for social interaction behavior in mice mapped

1227   with automated pattern detection by a hidden Markov model incorporated into newly

1228   developed freeware. *J. Neurosci. Methods* **234,** 127–134 (2014).

1229   92.   Tsuda, M. C. & Ogawa, S. Long-Lasting Consequences of Neonatal Maternal

1230   Separation on Social Behaviors in Ovariectomized Female Mice. *PLoS One* **7,** e33028

1231   (2012).

1232   93.   Berton, O. *et al.* Essential role of BDNF in the mesolimbic dopamine pathway in social

1233   defeat stress. *Science* **311,** 864–8 (2006).

1234   94.   Nadler, J. J. *et al.* Automated apparatus for quantitation of social approach behaviors in

1235   mice. *Genes, Brain Behav.* **3,** 303–314 (2004).

1236   95.   Basser, P. J., Mattiello, J. & LeBihan, D. MR diffusion tensor spectroscopy and imaging.

1237   *Biophys. J.* **66,** 259–267 (1994).

1238   96.   Hrabe, J., Kaur, G. & Guilfoyle, D. N. Principles and limitations of NMR diffusion

1239   measurements. *J. Med. Phys.* **32,** 34–42 (2007).

1240   97.   Mori, S. (Susumu). *Introduction to diffusion tensor imaging*. (Elsevier, 2007).

1241

Figure 1. **Two conceptual models of the post-traumatic stress incubation process. (A)** In the collective model, different psychological elements (Node #, a type of emotion or cognition) influence each other during different phases of stress incubation. **(B)** In the unitary model a single common factor underlies the development of post-traumatic behaviors. Orange and red boxes represent medium and strong trauma-induced differences, respectively. It is assumed that each psychological element can be connected to an associated neural substrate with trauma-induced dynamic changes.

1249

Figure 2. Paradigm induces long-term and delayed effects on multiple behaviors and physical conditions. (A) Paradigm with acute psychosocial trauma induction in mice. Focal mouse (dark gray; the Partner-observing mouse), partner mouse (light gray P), aggressor mouse (white), focal mouse's homecage (black), aggressor's homecage (brown), and wire-meshed divider (dashed line). **(B)** Experiments of No-scenario-isolated mice which had no experience of trauma induction identified the scenario effect in the behavioral paradigm. **(C)** Nest wall heights show long-lasting significant differences after trauma induction. **(D)** Body mass shows a significant increase 28 days after trauma induction. **(E)** Baseline plasma corticosterone level increased after trauma induction for both Partner-observing mice and their partners, the Attacked mice. Error bars indicate standard errors of the means; n.s., p≥0.05; *, 0.01≤p<0.05; ***, p<0.001; two-tailed, two-sample Student's t-test.

Figure 3. Diffusion tensor imaging (DTI) reveals chronic effect of acute observational stress on microstructures in the brain. Graphs show all brain structures with significant differences between No-scenario-isolated mice (n=4) and Partner-observing mice (n=4). **(A)** Fractional anisotropy (FA) of DTI-based water diffusivity suggests the changes of average microstructural integrity in multiple areas of the cerebral cortex. lOC, lateral orbital cortex; SInh, non-homunculus region of the primary sensory cortex; SIfl, forelimb region of the primary sensory cortex; SIjaw, jaw region of the primary sensory cortex; vACC, ventral region of the anterior cingulate cortex; CA2sr, stratum radiatum of the hippocampal cornu ammonis (CA) 2 area; RSC, the retrosplenial cortex; VI, the primary visual cortex; (R), the right area; (L), the left area. **(B)** DTI-based mean water diffusivity (MD) suggests the changes of membrane density in multiple areas of the entorhinal cortex-hippocampus system and the straitening of structural hemispheric specializations in the amygdala-insular cortex system. dlOC, dorsolateral orbital cortex; InC, the insular cortex; CoA, the cortical amygdalar nucleus; MeA, medial amygdalar nucleus; CA3sl, stratum lucidum of the hippocampal CA3 area; dlEC, dorsolateral entorhinal cortex. **(C)** DTI-based axial water diffusivity (AD) suggests the changes of neurite organization in multiple areas of the cerebral cortex and white matter mainly in the right hemisphere. OP, olfactory peduncle; PLC, prelimbic cortex; mOC, medial orbital cortex; CL, claustrum; MS, medial septal complex; mPOA, medial preoptic area; CB, cingulum bundle; CA3sr, stratum radiatum of the hippocampal CA3 area; CA3sl, stratum lucidum of the hippocampal CA3 area;CA3so, stratum oriens of the hippocampal CA3 area; CA3ipl, inner pyramidal layer of the hippocampal CA3 area; CA3opl, outer pyramidal layer of the hippocampal CA3 area. **(D)** DTI-based radial water diffusivity (RD) suggests the changes of myelination in multiple areas of the cerebral cortex in the right hemisphere and the white matter in the left hemisphere. vOC, ventral orbital cortex; ACal, anterior limb of the anterior commissure; CC, corpus callosum; CA1so, stratum oriens of the hippocampal CA1 area. *, $0.01 \leq p < 0.05$; **, $0.001 \leq p < 0.01$.

1288

Figure 4. Fine-scale behavioral analysis in light-dark box test detects the gradually developing process of the behavioral difference. (A) Light-dark box test quantified through the cumulative position probability T(x) along the light-dark axis (left; total time = 300 s). On average, Partner-observing mice (red) spent more time in the light area than No-scenario-isolated mice (blue) during the late post-traumatic period [T(0), bottom-left]. Spatially fine-scale behavioral analysis reveals significant differences between Partner-observing and No-scenario-isolated populations already in the early post-traumatic period (middle). For each position, we compute the mean T(x) across the No-scenario-isolated and Partner-observing populations and compute statistical significance through a two-population Student's t-test. These differences gradually increased, as evidenced by significance distributions collapsed across all positions (right; box plots show the minima, lower quartiles, medians, upper quartiles,

and maxima). **(B)** We similarly quantified speed using the fine-scale cumulative distribution

$U(v)$ of having speed $\leq v$ and we show the statistical analysis of population differences in $U(v)$.

Cumulative distribution functions of locomotion speed ($U(v)$ of having speed $\leq v$) and

corresponding significance distributions provide an additional independent behavioral index

that showed a gradually increasing differences of higher speed in Partner-observing mice. **(C)**

Higher transfer counts and shorter latency to the first transfers in Partner-observing mice

suggest their higher activity and exploratory motivation, respectively.

1307

Figure 5. Behavioral testing in the elevated plus-maze demonstrates that stress incubation of anxiety caused the observed differences. **(A)** Partner-observing and No-scenario-isolated mice did not differ significantly in the time they spent in opened arms (total time = 300 s); however, spatial distributions show differences in preferred location between Partner-observing and No-scenario-isolated mice in the closed arms, which increased with time. **(B)** Cumulative distribution functions of locomotion speed and corresponding significance distributions show a gradually increasing differences of lower speeds in Partner-observing mice. **(C)** Less exploration from close arms to platform center and longer freezing time in Partner-observing mice suggest their stronger stress reactions.

1317

Figure 6. **Gradually increasing anxiety and acute fear reaction were untangled in standard behavioral tests as different psychological components with distinctive developments.** Locomotion speed shows acute difference only in stressor zones [open arms **(A)** and light area **(C)**], but incubated differences in stressor-free zones [dark area **(B)** and closed arms **(D)**].

1323

Figure 7. Behavioral testing in the open field test and locomotor activity test confirms distinctive psychological substrates and their corresponding development patterns. (A) Anxiety is evident in the open field test through a delayed onset of locomotor speed differences in the center region with higher spatial uncertainty. **(B)** The differences of locomotor speed observed in the center region did not occur in the periphery region with lower spatial uncertainty. **(C)** Less time spent in the central region by Partner-observing mice suggests their avoidance of a region with high special uncertainty, while shorter latency to their first rearing indicates their higher exploratory motivation. **(D)** In the locomotor activity test without stressors, acute effects of activity reduction recovered in the later post-traumatic period.

1333

Figure 8. Acute psychosocial trauma decreases social interest. (A) The active social interaction test with consecutive non-social and social phases. During the social phase, motivation for social contact toward a stranger mouse (light gray S) was evaluated as the time spent for social approaches of nose poking and the time spent in the delineated interaction zone. **(B)** Partner-observing mice made fewer nose poking to both female (left) and male (right)

strangers. **(C)** There was no significant difference in the time spent in the interaction zone

during the social phase, suggesting a decrease of social interest instead of an active social

avoidance. **(D)** There was no significant difference in the time spent of nose poking during the

non-social phase, confirming that the observed differences of nose poking time stemmed from

a specifically social root. **(E)** Spectrogram of short but conspicuous ultrasonic vocalizations

emphasizes a specific behavioral repertoire during the social session in the female stranger test

of a No-scenario-isolated mouse on Day1. More vocalization was recorded from No-scenario-

isolated mice than Partner-observing mice on Days 1 and 7. Reduced ultrasonic vocalization

during the social session of the female stranger test in Partner-observing mice attests to

diminished social communication. Note that data points greater than 0.05% are not visible in

the right panel which zoom in the data of the middle panel to emphasize the data distributions

in the range of 0–0.05%. Data points and median, one-tailed Mann-Whitney U test; PSD, power

spectral density. **(F)** Partner-observing mice showed a delayed onset of reduction in

locomotion speed during the non-social phase.

Figure 9. Control experiments investigate psychosocial and physiological factors that determined and recued the behavioral developments. (A) Experiments of Stranger-observing mice which had witnessing experience of trauma that happened to strangers, rather than to their pair-housed partners, identified the relationship effect in the behavioral paradigm. (B) Experiments of Stranger-exposed mice which had experience of social interactions without witnessing stress from strangers identified the aggression effect in the behavioral paradigm. (C) Experiments of Partner-buffered mice which were pair-housed with their partners after trauma induction identified the social transfer effect of coexperiencing trauma in the behavioral paradigm. (D) Experiments of Stranger-buffered mice which were pair-housed with strangers after trauma induction identified the isolation effect in the behavioral paradigm. (E) Experiments of Environment-enriched mice which were provided with toys after trauma induction identified the social rescue effect in the behavioral paradigm. (F) Experiments of

FLX-treated mice which were treated with fluoxetine after trauma induction identified the

pharmacological rescue effect in the behavioral paradigm.

Figure 10. Social relationship in trauma induction determines stress development. (A)

Social relationship increased emotional impact, evidenced by tail rattling behavior during

trauma induction. **(B–F)** No significant chronic difference was found in Stranger-observing

mice. **(G)** The partner-revisiting test. A stranger mouse (light gray S) and the previously pair-

housed partner (light gray P), both immobilized, were presented as social targets. Pink

rectangle, heating pad. **(H)** Partner-observing mice showed significantly longer allogrooming

or pushing their partners (yellow, % of time spent in partner concern behavior) than either No-

scenario-isolated or Stranger-observing mice. Standard errors were calculated from

bootstrapped data.

1378

Figure 11. Control experiments identify necessity of relationship-dependent vicarious defeat and sufficiency of social isolation for anxiety incubation. (A) Positions in the light-dark box test indicate that Partner-buffered and Stranger-exposed mice did not develop the chronic stress reactions. (B) Locomotor speeds indicate that Partner-buffered and Stranger-exposed mice did not develop the acute stress reactions. (C) Body masses indicate that Partner-buffered and Stranger-exposed mice did not develop the physiological differences. (D) Nose poking times in the female stranger test indicate that Partner-buffered mice did, but Stranger-exposed mice did not develop the social differences. (E) Partner-buffered mice and Partner-observing mice spent longer time on the behaviors reflecting partner concern.

Figure 12. Control experiments uncover that social buffering with non-defeated
strangers rescued the full phenotypic dimensions of stress development. (A) Positions in
the light-dark box test indicate that Environment-enriched and FLX-treated mice did, but
Stranger-buffered mice did not develop the behavioral differences in the late phase. **(B)** Nose
poking times in the female stranger test indicate that Stranger-buffered, Environment-enriched,
and FLX-treated mice did not develop the social differences. **(C)** Environment-enriched mice
showed of reduction in locomotion speed during the non-social phase. **(D)** Positions in the
elevated plus-maze test indicate that Stranger-buffered mice were better rescued than FLX-
treated mice.

1398

Figure 13. DTI-based long-range connections suggest neural substrate of unitary view for stress development. (A) DTI-based network-wise fiber tracking reveals specific chronic changes of structural connectivity in the brain. AMY, the amygdala; HTh, the hypothalamus; DG, the hippocampal dentate gyrus; PC, the piriform cortex; PRC, the perirhinal cortex; APir, the amygdalopiriform transition area. Note that the brain regions were in the right brain hemisphere. **(B)** Trauma-induced structural changes of the underlying brain connectome uncovered an enhancement of a unitary network centered at the anterior cingulate cortex.

1406

Figure 14. Foregoing data demonstrate a unitary psychopathological model with its

corresponding neural substrate for the stress development paradigm in this study. Our

data show little support for a collective view speculated in previous hypotheses **(A)**, but support

a unitary view **(B)** where diverse PTSD-like reactions independently developed via a common

cognitive source ("Affective bonding"). This unitary view suggests a psychological substrate

in system-level that integrates multiple mechanisms underlying highly diverse behavioral

consequences, as a specific clinical target in the full mechanistic picture.

Day28 Diffusion tensor imaging:
Fractional anisotropy (FA)

1414

Supplemental Figure 1. Brain-wide microstructural changes measured by DTI fractional

anisotropy. Po, Partner-observing mice; Nsi, No-scenario-isolated mice; -log(p), statistical

significance through a two-population Student's t-test; A, anterior; P, posterior; D, dorsal; V,

ventral; L, left; R, right; lOC, lateral orbital cortex; SInh, non-homunculus region of the

primary sensory cortex; SIfl, forelimb region of the primary sensory cortex; SIjaw, jaw region

of the primary sensory cortex; vACC, ventral region of the anterior cingulate cortex; CA2sr,

stratum radiatum of the hippocampal cornu ammonis (CA) 2 area; RSC, the retrosplenial cortex;

VI, the primary visual cortex.

1423

Supplemental Figure 2. Brain-wide microstructural changes measured by DTI mean diffusivity. dlOC, dorsolateral orbital cortex; InC, the insular cortex; CoA, the cortical amygdalar nucleus; MeA, medial amygdalar nucleus; CA3sl, stratum lucidum of the hippocampal CA3 area; dlEC, dorsolateral entorhinal cortex.

1428

Supplemental Figure 3. Brain-wide microstructural changes measured by DTI axial diffusivity. OP, olfactory peduncle; PLC, prelimbic cortex; mOC, medial orbital cortex; CL, claustrum; MS, medial septal complex; mPOA, medial preoptic area; CB, cingulum bundle; CA3sr, stratum radiatum of the hippocampal CA3 area; CA3sl, stratum lucidum of the hippocampal CA3 area;CA3so, stratum oriens of the hippocampal CA3 area; CA3ipl, inner pyramidal layer of the hippocampal CA3 area; CA3opl, outer pyramidal layer of the hippocampal CA3 area.

1436

Supplemental Figure 4. Brain-wide microstructural changes measured by DTI radial diffusivity. vOC, ventral orbital cortex; ACaI, anterior limb of the anterior commissure; CC, corpus callosum; CA1so, stratum oriens of the hippocampal CA1 area.

Supplemental Video 1. Social apathy is an observed behavioral characteristic of Partner-

observing mice. Examples of 30-s recordings during the social session in the female stranger

test on Day1. First scene, No-scenario-isolated mouse; Second scene, Partner-observing mouse.

Supplement Video 2. Illustration of behavioral characteristics that were specific to observer

mice during trauma induction when their partners were attacked. First scene, tail rattling during

aggressive encounter; Second scene, tail rattling during aggressive encounter (4x slower);

Third scene, hiding under bedding material with the partner during resting. These behaviors

were not observed if a stranger mouse got attacked.

Supplemental Video 3. Illustration of rebound reaction of a Partner-observing mouse to its

previously pair-housed partner during the partner-revisiting test.