Host range and genetic plasticity explain the co-existence of integrative and extrachromosomal mobile genetic elements

Jean Cury[1,2], Pedro H. Oliveira[1,2,*], Fernando de la Cruz[3], Eduardo P.C. Rocha[1,2]

Affiliations.

[1]Microbial Evolutionary Genomi cs, Institut Pasteur, 28, rue Dr Roux, Paris, 75015, France,
[2]CNRS, UMR3525, 28, rue Dr Roux, Paris, 75015, France,
[3] Instituto de Biomedicina y Biotecnología de Cantabria (IBBTEC),

Universidad de Cantabria, Calle Albert Einstein 22, 39011 Santander, Spain

*Currently at: Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA

Classification: Biological Sciences/Evolution

Corresponding author: Jean Cury (jean.cury@normalesup.org)

Abstract

Self-transmissible mobile genetic elements drive horizontal gene transfer between
prokaryotes. Some of these elements integrate in the chromosome, whereas others
replicate autonomously as plasmids. Recent works showed the existence of few differences,
and occasional interconversion, between the two types of elements. Here, we enquired on
why evolutionary processes have maintained the two types of mobile genetic elements by
comparing integrative and conjugative elements (ICE) with extrachromosomal ones
(conjugative plasmids) of the highly abundant MPF_T conjugative type. Plasmids encode
more replicases, partition systems, and antibiotic resistance genes, whereas ICEs encode
more integrases and metabolism-associated genes. Plasmids are more variable in size, have
more DNA repeats, and exchange genes more frequently. On the other hand, ICEs are more
frequently transferred between distant taxa, and this drives the conversion of plasmids into
ICEs after transfer to distantly related hosts. Hence, differential plasticity and
transmissibility explain the occurrence of both integrative and extra-chromosomal elements
in microbial populations.

Keywords: Mobile genetic elements, horizontal gene transfer, molecular evolution,
microbial genomics, conjugation

Introduction

The genomes of Prokaryotes have mobile genetic elements (MGEs) integrated in the
chromosome or replicating as extrachromosomal elements. These MGEs usually encode
non-essential but ecologically important traits (1, 2). Extra-chromosomal elements, such as
conjugative plasmids (CPs) and lytic phages, replicate autonomously in the cell using
specialized replicases to recruit the bacterial DNA replication machinery (or to use their
own). Plasmids and extra-chromosomal prophages can also increase their stability in cellular
lineages using partition systems, for proper segregation during bacterial replication (3),
resolution systems, to prevent accumulation of multimers (4), and restriction-modification
or toxin-antitoxin systems, for post-segregation killing of their hosts (5). Alternatively, many
MGEs integrate into the chromosome. This is the case of the vast majority of known
prophages, of most conjugative elements (ICEs), and of many elements with poorly
characterized mechanisms of genetic mobility (*e.g.*, many pathogenicity islands)(6–8). The
integrated elements are replicated along with the host chromosome and require an
additional step of excision before being transferred between cells. The existence of both
integrative and extra-chromosomal elements was a fruitful source of controversy in the
dawn of molecular biology, eventually leading to the discovery of the molecular
mechanisms allowing both states (9, 10). Yet, a complementary question does not seem to
have been addressed in the literature: Why are there both types of elements? What are the
relative benefits and disadvantages of the integrated and extrachromosomal MGE?

To address these questions, we analyzed the differences and similarities between ICEs and
CPs. We focused on these elements because both forms are frequently found in bacteria,
they can be easily detected in genomes, and the mechanism of conjugation is well known.
Conjugative elements have a crucial role in spreading antibiotic resistance and virulence
genes among bacterial pathogens (11–14). Recently, several works suggested that the line
separating integrative ICEs and CPs could be thinner than anticipated (15), because some
ICEs encode plasmid-associated functions like replication (16) or partition (17), some
plasmids encode integrases (18), and ICEs and CPs are intermingled in the phylogenetic tree
of conjugative systems (19). Finally, both forms – ICEs and CPs – are found throughout the
bacterial kingdom, but their relative frequency depends on the taxa and on the mechanisms

of conjugation (7). These differences suggest that conjugative elements endure diverse

selective pressures for being integrative or extrachromosomal depending on unknown

environmental, genetic, or physiological variables.

We thought that key differences in the biology of integrative and extrachromosomal

elements might provide them with different types of advantages. ICEs require an additional

step of integration/excision during transfer, which may take time and requires genetic

regulation. Their integration in the chromosome may affect the latter's organization and

structure, and these collateral effects might depend on the size of the element. On the

other hand, ICEs replicate as part of chromosomes and could thus be lost from the cell at

lower rates than plasmids. Furthermore, plasmids must recruit the host replication

machinery, which may render elements incompatible and is known to constrain their host

range: many plasmids are able to conjugate into distantly related hosts, but are unable to

replicate there (20–22). We thus hypothesize that ICEs might be favored when transfers

occur between distant hosts, whereas plasmids might provide more genetic plasticity

because their size is not constrained by chromosomal organization.

Here, we study conjugative elements of the type MPF_T. This is the most frequent and best-

studied type of conjugative systems (19), and the only one for which we can identify

hundreds of elements of each of the forms (ICEs and CPs). We restricted our analysis to

genera containing both CPs and ICEs, to avoid, as much as possible, taxonomical biases. We

first describe the content of both types of elements and highlight their differences and

similarities. Next, we quantify their genetic similarity and the extent of their gene

exchanges. Finally, we show that chromosomal integration facilitates the colonization of

novel taxa by a conjugative element.

 Results

 Functional and genetic differences between ICEs and CPs

We analyzed a set of 151 ICEs and 136 CPs of the same genera and of type MPF_T, most of

which were from Proteobacteria (96.9%). Both ICEs and CPs were found to be AT-richer than

their host chromosomes, which is a common feature in MGEs and horizontally transferred

genes (23). However, the difference was three times smaller for ICEs (Fig. 1A), presumably

because they replicate with the chromosome or remain a longer time in the same host. The

average size of CPs is slightly larger (75kb vs 59kb), and the median slightly smaller (46kb vs

52kb) than that of ICEs. In contrast, CPs have more diverse sizes than ICEs (Fig. 1B), showing

a coefficient of variation twice as large (1.05 vs 0.49). The size of the conjugative elements

depends on the size of the bacterial genome (after discounting the size of their conjugative

elements), this effect being much stronger for CPs (Fig. 1C). CPs also have four times higher

density of large DNA repeats than ICEs (Fig. 1D). These results suggest that CPs diversify

faster than ICEs.

Fig. 1: Comparison between 136 CPs (blue) and 151 ICEs (yellow) in composition and sizes.
A. CPs are AT richer than ICEs relative to their hosts (Wilcoxon rank sum test, p-value<10^{-3}).
B. ICEs and CPs have different distributions of size (same test, p-value<0.05). Median sizes:
ICEs (52.5 kb) > CPs (46.1 kb). Averages: ICEs (59 kb) < CPs (74.6 kb) **C.** Size of the element
as a function of the genome size of its host (decreased by the size of the mobile element
itself). Shaded regions indicate the 95% confidence interval. The Y-axis is identical to the one

in panel B. **D**. Density of repeats is higher in CPs than in ICEs (0.30 vs 0.078 repeats per kb,
same test, p-value < 10^{-10}).

HGT concentrates in a few hotspots in bacterial chromosomes, presumably to minimize
disruption in their organization (24). We used HTg50, a measure of the concentration of
HGT in chromosomes that corresponds to the minimal number of spots required to account
for 50% of horizontally transferred genes (24), to test if chromosomes with fewer
integration hotspots had more plasmids. Indeed, there is a negative association between
the number of plasmids, weighted by their size, and the chromosomes' HTg50 (Spearman
ρ=-0.35, p-value=0.0016, Fig. S1).

We then quantified the differences between ICEs and CPs in terms of functions associated
with their biology, with a focus on stabilization functions. Relaxases are part of the rolling
circle replication initiator proteins and some have been shown to act as replicases(16, 25) or
site-specific recombinases (26, 27). Since all conjugative elements encode a relaxase, by
definition, they may also have these functions. In the following, we focused on typical
plasmid replication initiator proteins (more than 95% of them are involved in theta-
replication, and none is matched by the protein profiles of relaxases), and serine or tyrosine
recombinases as integrases. Expectedly, ICEs showed higher frequency of integrases, while
CPs had more frequently identifiable partition and replication systems. Some ICEs encode
partition systems (11%) and many encode a replicase (40%), while 37% of CPs encode at
least one tyrosine or serine recombinase (Fig. 2A). These results further illustrate a
continuum between the two types of elements: about half of the elements (40% and 48%,
ICEs and CPs, respectively) have functions usually associated with the other type and may
(rarely) lack functions typically associated with its own type (Fig. 2B). We identified plasmid
incompatibility systems of diverse types, whereas no ICE could not be typed in the current
scheme (Fig. S2).

Fig. 2: Comparisons of the functions carried by ICEs and CPs. **A.** Elements encoding (+) or lacking (-) replication or partition systems ("Rep or Par") or an integrase. **B.** Accessory functions over-represented (yellow) or under-represented (blue) in ICEs relative to CPs. Colored bars: significantly different from zero, p-value < 0.05 Fisher exact test with Bonferroni-Holm correction for multiple tests. Grey bars otherwise.

We then made similar analyses for functions usually regarded as accessory or unrelated to the biology of MGEs (Fig. 2B). ICEs were more likely to carry restriction-modification systems (x2.8) than CPs (but not orphan methylases), suggesting that ICEs endure stronger selective pressure for stabilization in the genome. In contrast, they were significantly less likely to carry antibiotic resistance genes or integrons. They also had fewer identifiable entry-exclusion systems, which may reflect the ability of ICEs to tolerate the presence of multiple similar elements in the cell (28). The classification of genes in the four major functional categories of the EggNOG database, showed that ICEs had relatively more genes encoding metabolic and cellular processes. We have previously shown that genes of unknown or poorly characterized function were over-represented in ICEs relative to their host chromosome (29). The frequency of these genes is even higher in plasmids (61% vs 46%). Hence, both types of elements have many functions in common, but their relative frequency often differs significantly.

Genetic similarities between ICEs and CPs

The results of the previous section, together with previously published studies (see Introduction), suggest that ICEs and CPs either share a common history or often exchange genes (or both). We detailed the relationships of homology between ICEs and CPs using the weighted Gene Repertoire Relatedness (wGRR), which measures the frequency of bi-directional best hits between two elements weighted by their sequence similarity (see Methods). We clustered the matrix of wGRR using the Louvain algorithm (30), and found six well-distinguished groups (Fig. 3). Two groups (1 and 6) are only constituted of CPs, two are composed of more than 90% of ICEs (3 and 5) and two have a mix of both types of elements (2 and 4) (Fig.3, top bar). Bacterial species are scattered between groups, showing that they are not the key determinant of the clustering. Some groups are only from γ-proteobacteria, but others include bacteria from different classes. Groups where elements are from the same taxonomic classes tend to have either CPs or ICEs, whereas the others have mixtures of both elements. Group 4, includes many ICEs and CPs, where all ICEs have integrases while more than half of the CPs lack both replication and partition systems (Fig. 4). This group includes almost only elements from ε-proteobacteria that may have specificities that we were not able to take into account. In contrast, almost all ICEs of groups 2 and 3 encode an integrase and all CPs have partition or replication systems.

Fig. 3: Heatmap of the wGRR scores, ordered after the size of Louvain's group (depicted on the left bar). The top bar represents the proportion of ICEs (yellow) and CPs (blue) for each

group. The bottom bar assigns a color corresponding to the host's species or class (γ-
proteobacteria in red, β-proteobacteria in green, α-proteobacteria in blue, ε-
proteobacteria in orange, Fusobacteria and Acidobacteria in grey).

Fig. 4: Mosaic plot representing the frequency of key functions of conjugative elements in

terms of the Louvain's groups (Fig. 3). The width of the bar is proportional to the number of

elements in a given Louvain's group (see the number of elements of each group on the top

of the bars) and the areas reflect the proportion of the elements with the function. The

colors represent the type of function, and their tint represent the part of ICEs (lighter) and

CPs (darker).

We controlled for the effect of the MPF genes in the previous clustering analysis by re-doing

it without these genes (Fig. S3). This produced the same number of groups - N1 to N6 - and

90% of the elements of the former groups were classed in the same novel groups (Fig. S4).

The only qualitatively significant difference between the two analyses concerned the group

2 for which 36% of the elements were now classed in groups N4 or N6. Overall, these

controls confirm that ICEs and CPs can be grouped together, and apart from other elements

of the same type. The grouping is not caused by sequence similarity between conjugative

systems. Instead, it probably reflects either within group genetic exchanges between ICEs

and CPs, or interconversions of the two types of elements.

Genetic exchanges: CPs become ICEs for broader host range

The clustering of ICEs and CPs could be explained by genetic transfer between them. To

address this question, we searched for pairs of conjugative elements with low wGRR (<30%)

but some highly similar homologs (>80% sequence identity). This identified several cases of

recent transfer of a single or a few genes between elements (Fig. 5A). In agreement with our

observations of higher genetic plasticity in CPs, most transfers took place between these

types of elements (Fig. 5.B-D).

Fig. 5: 2D histogram of the wGRR score of pairs of conjugative elements as a function of the

protein identity of their homologues. **A**. Distribution for the entire dataset. wGRR values are

correlated with protein identity of the elements' homologs ($\rho_{ICE-ICE} = 0.90$, $\rho_{CP-CP} = 0.83$). The

blue rectangle zooms on a region where the pairs of elements are very different (GRR<30%),

yet they encode at least one very similar protein (identity > 80%). The dashed line separates

the elements where protein identity is higher than wGRR x 2/3. **B**. Zoom for ICE-ICE

comparisons. **C**. ICE-CP comparisons. **D**. CP-CP comparisons.

We hypothesized that ICEs could hold an advantage over CPs to colonize novel hosts,

because replication restricts plasmid host range. To test this hypothesis, we analyzed the

wGRR between pairs of ICEs and pairs of CPs in function of the phylogenetic distance

between their bacterial hosts. This showed similar patterns for the two types of elements,

with the notable exception that there are no pairs of highly similar plasmids (wGRR>50%) in

distant hosts (more than 0.1 substitutions/position, *e.g.,* the average distance in the tree

between *E. coli* and *P. aeruginosa*). In contrast, a third of all ICEs (n=50) are in these conditions (Fig. 6A, Fig. S5). The same analysis after removing the MPF genes shows wGRR values shifted to lower wGRR values for all elements, but qualitatively similar trends (Fig. S6). This suggests a major difference in the ability of ICEs and CPs to be stably maintained after their transfer into a distant host.

Fig. 6: wGRR as function of the host phylogenetic distance **A**. 2D histogram of the distribution of the wGRR score as a function of the phylogenetic distance for pairs of CPs (left) and pairs of ICEs (right). The bottom row corresponds to all pairs with distance lower than 10^{-5} (including those in the same host). Elements in the black rectangle are depicted in a phylogenetic tree in Fig. S5. **B**. Proportion of ICEs more distinct in terms of tri-nucleotides from their host than CPs. Bins are larger than in A. to increase the power of the statistical analysis. Color code: ICEs (Red) or CPs (Blue) are more distinct from the host in terms of tri-nucleotide composition than the other element. Grey: not significant (binomial test, p-value $> 10^{-2}$). White: no elements in the bin.

We then analyzed the pairs ICE-CP. We found few pairs of highly similar ICEs and CPs in closely related hosts (bottom right corner of Fig. 6, n=8 for wGRR>50% and d < 10^{-2}), suggesting that interconversion between these elements remains rare within a clade. A larger number of ICE-CP pairs were very similar but present in distant hosts (n=38, Fig. 6). The most parsimonious explanation for these observations, is the recent transfer of one of the elements (ICE or CP) to a distant bacterial host. We identified the latter element based on the differences in terms of tri-nucleotide composition between the elements and the host chromosomes (defined as pvalue in (31)). We then computed for each ICE-CP pair the difference between the pvalue of the pair ICE-host and that of the pair CP-host (see Methods). In agreement to our observation that ICEs have broader host ranges, these

differences indicate that ICEs are relatively more distant from the host chromosome for

pairs with high wGRR in distant hosts than for the rest of the pairs (Wilcoxon rank sum test,

p-value$<10^{-20}$, Fig. 6B, and Fig. S7). The rarity of ICE-CP pairs in closely related hosts, their

abundance in distant hosts, and the identification that ICEs are the most compositionally

atypical relative to the host in the latter, suggest that successful transfer of CPs to distant

hosts is favored when they integrate the chromosome and become ICEs.

Discussion

In this study, we quantified the differences between ICEs and CPs to evaluate the claims that they are essentially equivalent MGEs (15, 16, 29). We found that numerous CPs have integrases (although these may serve for dimer resolution and not integration (32)), numerous ICEs encode replication and partition functions, the elements often cluster them together, and they exchange genetic information. Furthermore, relaxases – present in both ICEs and CPs – have been shown to act as integrases or replication initiators (33). Hence, ICEs and CPs constitute closely related elements sharing many functions beyond those related to conjugation. Yet, there are also some clear differences between them (Fig. 7). First, genes encoding plasmid replicases and partition systems are more frequent in plasmids, and tyrosine and serine recombinases are more frequent in ICEs. Interestingly, we could not attribute incompatibility groups to ICEs, suggesting that the replication module is rarely exchanged between ICEs and CPs. Second, the frequency of certain accessory traits is different: plasmids are more likely to encode antibiotic resistance genes whereas ICEs encode more metabolism-related genes, even if this could result from biases in the database towards nosocomial pathogens. Finally, the %GC relative to the host, the number of repeats, the patterns of gene variation and exchange, and the host range are quantitatively different in the two types of elements. After integrating all this information, we propose that in spite of their similarities each type has specific advantages.

Fig. 7: To integrate or not to integrate? ICEs and CPs have similar conjugative systems but persist in different ways (resp. integrating the chromosome or replicating independently).

ICEs over-represent integrases, whereas plasmids over-represent replication and partition

functions (although both can have all of these functions). ICEs seem at an advantage when

conjugating to distant hosts, presumably because they integrate the chromosome. On the

other hand, plasmids have wider distribution of size and exchange more genetic

information.

Even if there are some known families of large (>200kb) ICEs (34, 35), most of these

elements in our dataset have a narrower variation in size than CPs. This suggests that CPs

are more flexible than ICEs in terms of the amount of genetic information they can carry and

in their ability to accommodate novel information. CPs also exchange genes more

frequently. Mechanistically, the rate of recombination between plasmids may be higher

because they encode more repeats, integrons, and transposable elements. Plasmid copy

number, when high, may also contribute to increase recombination rates. Interestingly,

recombination mediated by transposable elements has been shown to be drive the

evolution of certain plasmids(36, 37), and to accelerate the reduction of plasmid cost thus

stabilizing the element after horizontal transfer (38). The restrictions in size variation of ICEs

are probably not due to the mechanism of integration or excision because such reactions

occur between very distant recombination sites (39). Instead, very large ICEs may disrupt

chromosome organization by affecting the distribution of motifs, changing chromosome

folding domains, or unbalancing the sizes of replichores (40). Repeat-mediated

recombination leads to replicon rearrangements, and may lead to stronger counter-

selection of DNA repeats in ICEs than in CPs, further restricting their size variation and their

flux of gene exchange. Interestingly, the size of plasmids varies much more steeply with

genome size than the size of ICEs, suggesting that CPs may play a particularly important role

in the evolution of large bacterial genomes of Proteobacteria, which have higher rates of

genetic exchanges (41), and often contain mega-plasmids (42).

Some plasmids are known to be broad-host range and adapt to novel hosts, especially if

they carry adaptive traits that compensate for the initially poor intrinsic persistence of the

element (43, 44). However, within the large phylogenetic span considered in this work,

MPF_T ICEs have broader host ranges than CPs. Actually, the first known ICE, Tn916 (not

MPF$_T$, thus not included in this study), became notorious due to its ability to spread

antibiotic resistance between distant phyla (45).

Finally, we show novel evidence for interconversion between the two types of elements,

which had been proposed based on the phylogeny of the conjugative system (7). This is

consistent with the clustering of ICEs and CPs in terms of gene repertoires, even when

removing the conjugation system from the analysis, in certain groups and not in others.

Also, the transition of CPs to ICEs is the best explanation for the observations of the

presence of pairs of similar elements ICE-CP in distant hosts.

Other traits may provide advantages to ICEs or CPs. The ability of plasmids to modify their

copy number may accelerate adaptive evolutionary processes, such as the acquisition of

antibiotic resistance (46). On the other hand, ICEs might be more stably maintained in

lineages because they replicate within the chromosome. Finally, the carriage of ICEs and CPs

may have different costs. The cost of plasmids has been extensively studied and is strongly

dependent on the traits they encode (47). Much less is known about the cost of ICEs; several

reports suggest that they lead to low fitness costs when conjugation is not expressed, but

their fitness cost varies much more between elements during transfer (14). Direct

comparisons of the cost of carriage of ICE and CPs carrying similar traits are unavailable.

Further work will be needed to test these hypotheses.

Occasional transfers between CPs and ICEs allow them to access the other elements' gene

pool. These events may create elements with traits of ICEs and of CPs, as observed in more

than a third of all conjugative elements, and lead to their clustering in the wGRR group 5.

Additionally, they facilitate the interconversion of one type of element into the other.

Many elements are mobilizable but not able to conjugate independently (42, 48). These

elements often encode a relaxase that recognizes the element's origin of transfer and is able

to interact with a T4SS from an autonomously conjugative element to transfer to other cells.

Many of the disadvantages of conjugative plasmids and ICEs are similar to those of

mobilizable plasmids and integrative mobilizable elements, whether they encode a relaxase

or not. Notably, the former must be replicated in the extrachromosomal state, and the

latter integrate the genome where they must not disrupt genome organization. Patterns

observed in conjugative elements are thus likely to be applicable to mobilizable ones.

These results may also be relevant to understand lysogeny by temperate phages. The vast

majority of known prophages are integrated in the chromosome, but some replicate like

plasmids (49, 50). Considering that prophages share some of the constraints of conjugative

elements, they are likely to be under similar trade-offs. However, phages are under

additional constraints. Notably, their genome size is much less variable than that of

conjugative elements, because it must be packaged into the virion (51), and this may render

the extrachromosomal prophages less advantageous in terms of accumulating novel genes.

This could explain why most prophages are integrative whereas conjugative systems are

more evenly split between integrative and extrachromosomal elements.

Material and Methods

Data

Conjugative systems of type T (MPF_T) were searched in the set of complete bacterial genomes from NCBI RefSeq (http://ftp.ncbi.nih.gov/genomes/refseq/bacteria/, last accessed in November 2016). We analyzed 5562 complete genomes from 2268 species, including 4345 plasmids and 6001 chromosomes. The classification of the replicon in plasmid or chromosome was taken from the information available in the GenBank file. Our method to delimit ICEs is based on comparative genomics of closely related strains. Hence, we restricted our search for conjugative systems to the species for which we had at least five genomes completely sequenced (164 species, 2990 genomes).

Detection of conjugative systems and delimitation of ICEs

Conjugative systems were detected using the CONJscan module of MacSyFinder (52), with protein profiles and definitions of the MPF type T, published previously (53). ICEs were delimited with the same methodology, as developed in a previous work (29). Briefly, we identified the core genomes of the species. The region between two consecutive genes of the core genome defined an interval in each chromosome. We then defined spots as the sets of intervals in the chromosome flanked by genes of the same two families of the core genome (24). We then identified the intervals and the spots with conjugative systems. The information on the sets of gene families of the spots with ICEs (i.e., the spot pan-genome) was used to delimit the element boundaries (script available at https://gitlab.pasteur.fr/gem/spot_ICE). This methodology was shown to be accurate at the gene level (precise nucleotide boundaries are not identifiable by this method, see (29)).

Functional analyses

Partition systems, replication systems, entry-exclusion systems and restriction modification systems were annotated with HMM profiles, as described in our previous work (29, 54). Integrases were annotated with the PFAM profile PF00589 for the Tyrosine recombinases and the combination of PFAM profiles PF00239 and PF07508 for Serine recombinases. DDE Transposases were detected with Macsyfinder (52) with models used previously (55). Antibiotic resistance genes were detected with ResFams profiles (core version v1.1) (56)

using the --cut_ga option. We determined the functional categories of genes using their

annotation as provided by their best hits to the protein profiles of the EggNOG database for

bacteria (version 4.5, bactNOG) (57). Genes not annotated by the EggNOG profiles were

classed as "Unknown" and included in the "Poorly characterized" group. The HMM profiles

were used to search the genomes with HMMER 3.1b2 (58), and we retrieved the hits with

an e-value lower than 10^{-3} and with alignments covering at least 50% of the profile.

Integrons were detected using IntegronFinder version 1.5.1 with the --local_max option for

higher accuracy (59). Repeats (direct and inverted) were detected with Repseek (version

6.6) (60) using the option -p 0.001 which set the p-value for determining the minimum seed

length.

Statistics

We tested the over-representation of a given function or group of functions using Fisher's

exact tests on contingency tables. For partition, replication and integration, the contingency

table was made by splitting replicons in those encoding or not encoding the function and

between ICEs and CPs. The use of presence/absence data instead of the absolute counts

was made because the presence of at least one occurrence of a system is sufficient to have

the function and because the counts were always low. For the other functions, the

contingency table was made by splitting the proteins of the element in those annotated for

a given function and the remaining ones. This allowed to take into account the differences in

the number of genes between elements. The Fisher-exact tests were considered as

significant after sequential Holm-Bonferroni correction, with a family-wise error rate of 5%

(the probability of making at least one false rejection in the multiple tests, the type I error).

From the contingency table, we computed the relative ratio (or relative risk) of having a

given function more often in ICEs than in CPs. The relative ratio is computed as follow:

$RR = \frac{ICEwF/N_{ICE}}{CPwF/N_{CP}}$ where *ICEwF* is the number of ICE (or proteins in ICEs) with the given

function, and N_{ICE}, the total number of ICE (or proteins in ICEs), and likewise for CP. The

term $ICEwF/N_{ICE}$ is an estimation of the probability of an ICE (or a protein in an ICE) to

carry a given a function.

Phylogenetic distances

Phylogenetic distances were extract from the Proteobacterial tree of the Core-genome. To build the tree, we identified the genes present in at least 90% of the 2897 genomes of Proteobacteria larger than 1 Mb that were available in GenBank RefSeq in November 2016. A list of orthologs was identified as reciprocal best hits using end-gap free global alignment. Hits with less than 37% similarity in amino acid sequence and more than 20% difference in protein length were discarded. We then identified the protein families with relations of orthology in at least 90% of the genomes. They represent 341 protein families. We made multiple alignments of each protein family with MAFFT v.7.205 (with default options) (61) and removed poorly aligned regions with BMGE (with default options) (62). Genes missing in a genome were replaced by stretches of "–" in each multiple alignment, which has been shown to have little impact in phylogeny reconstruction (63). The tree of the concatenate alignment was computed with FastTree version 2.1 under the LG model (64). We chose the LG model because it was the one that minimized the AIC.

Distance to the host

We used the differences in tri-nucleotide composition to compute the genetic distance between the mobile element and its host chromosome, as previously proposed (65). The analysis of ICEs was done by comparing the element with the chromosome after the removal of its sequence from the latter. Briefly, we computed the trinucleotide relative abundance ($x_{ijk} \forall i, j, k \in \{A, T, C, G\}$) for the chromosomes (in windows of 5 kb) and for the conjugative elements (entire replicon), which is given by: $x_{ijk} = f_{ijk}/f_i f_j f_k$, with f the frequency of a given k-mer in the sequence (31). We first computed the Mahalanobis distance between each window and the host chromosome as follow:

$$D = \sqrt{(w - h)^T H^{-1}(w - h)}$$

with w, the vector of tri-nucleotide abundances (x_{ijk}) in a given window, and h, the mean of the vector of x_{ijk} (*i.e.*, the average tri-nucleotide abundance in the chromosome). H is the covariance matrix of the tri-nucleotide relative abundances. The inverse of the covariance matrix (H^{-1}) downweights frequent trinucleotides, like the tri-nucleotides corresponding to start codons, which are common to conjugative elements and chromosome and could bias

the distance. We computed the Mahalanobis distance between conjugative elements and

their hosts' chromosomes (same formula as above, but *w* is now for a conjugative element

instead of a chromosome window). We then computed the probability (p-value) that the

measured distance between a conjugative element and the host's chromosome is the same

as any fragment of the host's chromosome.

We compared ICEs and CPs in relation to their compositional distance to the host. For this,

we made the null hypothesis that the proportion of ICEs having a p-value lower than CPs

follows a binomial distribution whose expected proportion is that of the entire dataset (the

proportion of ICEs having a p-value lower than CP), precisely:

$H_0 = N(pvalue_{ICE} < pvalue_{CP})/N_{Comparisons}$, where $N_{Comparisons}$ is the total number of

ICE-CP pairs, *i.e.* $151 \times 136 = 20536$.

Network analysis of gene repertoire relatedness

We built a network describing the relations of homology between the elements. The nodes

in the network are conjugative elements and they are linked if they share a given relation of

homology. More precisely, the relationship between two elements was quantified with the

weighted Gene Repertoire Relatedness score (wGRR). This score represents the number of

homologous proteins between two elements, weighted by their sequence identity, as

described in (29). The formula is:

$$wGRR_{A,B} = \sum_i \frac{id(A_i, B_i)}{min(A,B)} \, if \, f \overset{\leftrightarrow}{evalue}(A_i, B_i) < 10^{-5}$$

Where (A_i, B_i) is the i^{th} pair of homologous protein between element A and element B,

id(A$_i$,B$_i$) is the sequence identity of their alignment, min(A,B) is the number of proteins of

the element with fewest proteins (A or B). The sequence identity was computed with blastp

v.2.2.15 (default parameters)(66) and kept all bi-directional best hits with an e-value lower

than 10^{-5}.

The network was built based on the wGRR matrix. Its representation was made using the

Fruchterman-Reingold force-directed algorithm as implemented in the NetworkX v1.11

python library. The groups were made using the Louvain algorithm (30). We controlled for

the consistency of the heuristic used to assess that the group found are not form a local

optimal. We performed 100 clustering, which led to the same classification in 95% of the

time.

Incompatibility typing

456 We determined the incompatibility group of replicons using the method of PlasmidFinder

457 (67). We used BLASTN (66) to search the replicons for sequences matching the set of 116

458 probes used by PlasmidFinder. We kept the hits with a coverage above 60% and sequence

459 identity above 80%, as recommended by the authors. Around 3% of the elements had

460 multiple incompatibility types attributed.

Acknowledgements

This work was supported by an European Research Council grant [EVOMOBILOME

n°281605], and a grant from the Agence National de la Recherche [MAGISBAC, ANR-14-

CE10-0007]. Work in FdlC lab was supported by grants BFU2014-55534-C2-1-P and

BFU2014-62190-EXP from the Spanish Ministry of Economy and Competitiveness. We

thank Alan Grossman and Marie Touchon for comments and suggestions, and Aude

Bernheim for providing the phylogenetic tree of proteobacteria. J.C. is a member of the

'Ecole Doctorale Frontière du Vivant (FdV) – Programme Bettencourt'.

References

1. Frost LS, Leplae R, Summers AO, Toussaint A (2005) Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 3(9):722–32.

2. Ochman H, Lawrence JG, Groisman EA (2000) Lateral gene transfer and the nature of bacterial innovation. *Nature* 405(6784):299–304.

3. Ebersbach G, Gerdes K (2005) Plasmid segregation mechanisms. *Annu Rev Genet* 39:453–79.

4. Summers DK (1991) The kinetics of plasmid loss. *Trends Biotechnol* 9(8):273–278.

5. Kobayashi I (2001) Behavior of restriction-modification systems as selfish mobile elements and their impact on genome evolution. *Nucleic Acids Res* 29(18):3742–3756.

6. Dobrindt U, Hochhut B, Hentschel U, Hacker J (2004) Genomic islands in pathogenic and environmental microorganisms. *Nat Rev Microbiol* 2(5):414–24.

7. Guglielmini J, Quintais L, Garcillán-Barcia MP, de la Cruz F, Rocha EPC (2011) The repertoire of ICE in prokaryotes underscores the unity, diversity, and ubiquity of conjugation. *PLoS Genet* 7(8):e1002222.

8. Canchaya C, Proux C, Fournous G, Bruttin A, Brussow H (2003) Prophage Genomics. *Microbiol Mol Biol Rev* 67(2):238–276.

9. Lederberg J (1998) Plasmid (1952 – 1997). *Plasmid* 9:1–9.

10. Jacob F, Schaeffer P, Wollman EL (1960) Episomic Element in Bacteria. *Symp. Soc. Gen. Microbiol*, pp 67–91.

11. Johnson CM, Grossman AD (2015) Integrative and Conjugative Elements (ICEs): What They Do and How They Work. *Annu Rev Genet* 49(1):annurev-genet-112414-055018.

12. Bellanger X, Payot S, Leblond-bourget N, Guédon G (2014) Conjugative and mobilizable genomic islands in bacteria: evolution and diversity. *FEMS Microbiol Rev*

38:720–760.

13. Carraro N, Burrus V (2014) Biology of Three ICE Families: SXT/R391, ICEBs1, and
ICESt1/ICESt3. *Microbiol Spectr* 2(6):1–20.

14. Delavat F, Miyazaki R, Carraro N, Pradervand N, van der Meer JR (2017) The hidden
life of integrative and conjugative elements. *FEMS Microbiol Rev* 41(4):512–537.

15. Carraro N, Burrus V (2015) The dualistic nature of integrative and conjugative
elements. *Mob Genet Elements* 5(6):98–102.

16. Lee C a, Babic A, Grossman AD (2010) Autonomous plasmid-like replication of a
conjugative transposon. *Mol Microbiol* 75(2):268–79.

17. Carraro N, Poulin D, Burrus V (2015) Replication and Active Partition of Integrative
and Conjugative Elements (ICEs) of the SXT/R391 Family: The Line between ICEs and
Conjugative Plasmids Is Getting Thinner. *PLOS Genet* 11(6):e1005298.

18. Nunes-Düby SE, Kwon HJ, Tirumalai RS, Ellenberger T, Landy a (1998) Similarities and
differences among 105 members of the Int family of site-specific recombinases.
Nucleic Acids Res 26(2):391–406.

19. Guglielmini J, de la Cruz F, Rocha EPC (2013) Evolution of conjugation and type IV
secretion systems. *Mol Biol Evol* 30(2):315–31.

20. Guiney DG (1982) Host range of conjugation and replication functions of the
Escherichia coli sex plasmid Flac. Comparison with the broad host-range plasmid RK2.
J Mol Biol 162(3):699–703.

21. Zhong Z, Helinski D, Toukdarian A (2005) Plasmid host-range: Restrictions to F
replication in Pseudomonas. *Plasmid* 54(1):48–56.

22. Klümper U, et al. (2015) Broad host range plasmids can invade an unexpectedly
diverse fraction of a soil bacterial community. *ISME J* 9:934–945.

23. Rocha EPC, Danchin A (2002) Base composition bias might result from competition for
metabolic resources. *Trends Genet* 18(6):291–4.

24. Oliveira PH, Touchon M, Cury J, Rocha EPC (2017) The chromosomal organization of
horizontal gene transfer in Bacteria. *Nat Commun* 8(841):1–10.

25. Carraro N, et al. (2016) Plasmid-like replication of a minimal streptococcal integrative
and conjugative element. *Microbiol (United Kingdom)* 162(4):622–632.

26. Francia MV, Clewell DB (2002) Transfer origins in the conjugative Enterococcus
faecalis plasmids pAD1 and pAM373: Identification of the pAD1 nic site, a specific
relaxase and a possible TraG-like protein. *Mol Microbiol* 45(2):375–395.

27. César CE, Machón C, De La Cruz F, Llosa M (2006) A new domain of conjugative
relaxase TrwC responsible for efficient oriT-specific recombination on minimal target
sequences. *Mol Microbiol* 62(4):984–996.

28. Garcillán-Barcia MP, de la Cruz F (2008) Why is entry exclusion an essential feature of
conjugative plasmids? *Plasmid* 60(1):1–18.

29. Cury J, Touchon M, Rocha EPC (2017) Integrative and conjugative elements and their
hosts: composition, distribution and organization. *Nucleic Acids Res* (17):1–14.

30. Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E (2008) Fast unfolding of
communities in large networks. *J Stat Mech*. doi:10.1088/1742-
5468/2008/10/P10008.

31. Suzuki H, Yano H, Brown CJ, Top EM (2010) Predicting plasmid promiscuity based on
genomic signature. *J Bacteriol* 192(22):6045–55.

32. Carnoy C, Roten C-A (2009) The dif/Xer recombination systems in proteobacteria.
PLoS One 4(9):e6531.

33. Wawrzyniak P, Płucienniczak G, Bartosik D (2017) The Different Faces of Rolling-Circle
Replication and Its Multifunctional Initiator Proteins. *Front Microbiol* 8(November):1–
13.

34. Sullivan JT, Ronson CW (1998) Evolution of rhizobia by acquisition of a 500-kb
symbiosis island that integrates into a phe-tRNA gene. *Proc Natl Acad Sci*

95(April):5145–5149.

35.   Kers JA, et al. (2005) A large, mobile pathogenicity island confers plant pathogenicity
on Streptomyces species. *Mol Microbiol* 55(4):1025–1033.

36.   Hall JPJ, Williams D, Paterson S, Harrison E, Brockhurst MA (2017) Positive selection
inhibits gene mobilization and transfer in soil bacterial communities. *Nat Ecol Evol*
1(9):1348–1353.

37.   He S, et al. (2016) Mechanisms of Evolution in High-Consequence Drug Resistance.
MBio 7(6):e01987-16.

38.   Porse A, Schønning K, Munck C, Sommer MOA (2016) Survival and Evolution of a
Large Multidrug Resistance Plasmid in New Clinical Bacterial Hosts. *Mol Biol Evol*
33(11):2860–2873.

39.   Wu LJ, Errington J, Rossignol M, Cornet F, Boccard F (2002) A large dispersed
chromosomal region required for chromosome segregation in sporulating cells of
Bacillus subtilis. *EMBO J* 21(15):4001–11.

40.   Touchon M, Rocha EPC (2016) Coevolution of the organization and structure of
prokaryotic genomes. *Cold Spring Harb Perspect Biol* 8(1):1–18.

41.   Oliveira PH, Touchon M, Rocha EPC (2016) Regulation of genetic flux between
bacteria by restriction-modification systems. *Proc Natl Acad Sci U S A* 113(20):5658–
63.

42.   Smillie C, Garcillán-Barcia MP, Francia MV, Rocha EPC, de la Cruz F (2010) Mobility of
plasmids. *Microbiol Mol Biol Rev* 74(3):434–52.

43.   De Gelder L, Williams JJ, Ponciano JM, Sota M, Top EM (2008) Adaptive plasmid
evolution results in host-range expansion of a broad-host-range plasmid. *Genetics*
178(4):2179–2190.

44.   Loftie-Eaton W, et al. (2017) Compensatory mutations improve general
permissiveness to antibiotic resistance plasmids. *Nat Ecol Evol* 1:1354–1363.

45.   Clewell DB, Flannagan SE, Jaworski DD (1995) Unconstrained bacterial promiscuity:
the Tn916–Tn1545 family of conjugative transposons. *Trends Microbiol* 3(6):229–236.

46.   San Millan A, Escudero JA, Gifford DR, Mazel D, Maclean RC (2016) Multicopy
plasmids potentiate the evolution of antibiotic resistance in bacteria. *Nat Ecol Evol*
1(10):1–8.

47.   San Millan A, MacLean RC (2017) Fitness Costs of Plasmids: a Limit to Plasmid
Transmission. *Microbiol Spectr* 5(5):1–12.

48.   Guédon G, Libante V, Coluzzi C, Payot S, Leblond-Bourget N (2017) The obscure world
of integrative and mobilizable elements, highly widespread elements that pirate
bacterial conjugative systems. *Genes (Basel)* 8(11). doi:10.3390/genes8110337.

49.   Ravin N V. (2011) N15: The linear phage-plasmid. *Plasmid* 65(2):102–109.

50.   Łobocka MB, et al. (2004) Genome of Bacteriophage P1. *J Bacteriol* 186(21):7032–
7068.

51.   Touchon M, Moura de Sousa JA, Rocha EP (2017) Embracing the enemy: The
diversification of microbial gene repertoires by phage-mediated horizontal gene
transfer. *Curr Opin Microbiol* 38:66–73.

52.   Abby SS, Néron B, Ménager H, Touchon M, Rocha EPC (2014) MacSyFinder: A
Program to Mine Genomes for Molecular Systems with an Application to CRISPR-Cas
Systems. *PLoS One* 9(10):e110726.

53.   Guglielmini J, et al. (2014) Key components of the eight classes of type IV secretion
systems involved in bacterial conjugation or protein secretion. *Nucleic Acids*
Res:gku194-.

54.   Oliveira PH, Touchon M, Rocha EPC (2014) The interplay of restriction-modification
systems with mobile genetic elements and their prokaryotic hosts. *Nucleic Acids Res*
42(21):1–14.

55.   Touchon M, et al. (2014) The Genomic Diversification of the Whole Acinetobacter

Genus: Origins, Mechanisms, and Consequences. *Genome Biol Evol* 6(10):2866–2882.

56.   Gibson MK, Forsberg KJ, Dantas G (2014) Improved annotation of antibiotic resistance

determinants reveals microbial resistomes cluster by ecology. *ISME J* 9(1):207–216.

57.   Huerta-Cepas J, et al. (2016) eggNOG 4.5: a hierarchical orthology framework with

improved functional annotations for eukaryotic, prokaryotic and viral sequences.

Nucleic Acids Res 44(D1):D286-93.

58.   Eddy SR (2011) Accelerated Profile HMM Searches. *PLoS Comput Biol* 7(10):e1002195.

59.   Cury J, Jové T, Touchon M, Néron B, Rocha EP (2016) Identification and analysis of

integrons and cassette arrays in bacterial genomes. *Nucleic Acids Res* 44(10):4539–50.

60.   Achaz G, Boyer F, Rocha EPC, Viari A, Coissac E (2007) Repseek, a tool to retrieve

approximate repeats from large DNA sequences. *Bioinformatics* 23(1):119–121.

61.   Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. *Mol Biol Evol* 30(4):772–780.

62.   Criscuolo A, Gribaldo S (2010) BMGE (Block Mapping and Gathering with Entropy): a

new software for selection of phylogenetic informative regions from multiple

sequence alignments. *BMC Evol Biol* 10:210.

63.   Filipski A, Murillo O, Freydenzon A, Tamura K, Kumar S (2014) Prospects for building

large timetrees using molecular data with incomplete gene coverage among species.

Mol Biol Evol 31(9):2542–2550.

64.   Price MN, Dehal PS, Arkin AP (2009) Fasttree: Computing large minimum evolution

trees with profiles instead of a distance matrix. *Mol Biol Evol* 26(7):1641–1650.

65.   Suzuki H, Sota M, Brown CJ, Top EM (2008) Using Mahalanobis distance to compare

genomic signatures between bacterial plasmids and chromosomes. *Nucleic Acids Res*

36(22):e147.

66.   Altschul SF, et al. (1997) Gapped BLAST and PSI-BLAST: A new generation of protein

database search programs. *Nucleic Acids Res* 25(17):3389–3402.

67.   Carattoli A, et al. (2014) In Silico detection and typing of plasmids using plasmidfinder

626   and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58(7):3895–

627   3903.