Analysis of mycobacterial infection-induced changes to host lipid metabolism in a zebrafish infection model reveals a conserved role for LDLR in infection susceptibility

Matt D. Johansen[1, 2], Elinor Hortle[2], Joshua A. Kasparian[1, 2], Alejandro Romero[3], Beatriz Novoa[3], Antonio Figueras[3], Warwick J. Britton[2, 4, 5], Kumudika de Silva[1, 5], Auriol C. Purdie[1, 5], Stefan H. Oehlers[2, 4, 5 *]

[1] Sydney School of Veterinary Science, Faculty of Science, The University of Sydney, Camden, NSW, Australia

[2] Tuberculosis Research Program, Centenary Institute, Camperdown, NSW, Australia

[3] Institute of Marine Research (IIM), Spanish National Research Council (CSIC), Vigo, Spain

[4] Sydney Medical School, The University of Sydney, Newtown, NSW, Australia

[5] Marie Bashir Institute, The University of Sydney, Newtown, NSW, Australia

[*] Corresponding author: stefan.oehlers@sydney.edu.au

Website: http://sydney.edu.au/medicine/people/academics/profiles/stefan.oehlers.php

Keywords: zebrafish, mycobacterium, pathogenesis, lipid, metabolism

Abstract

Changes to lipid metabolism are well-characterised consequences of tuberculosis infection but their functional relevance are not clearly elucidated. The zebrafish-*Mycobacterium marinum* infection model recapitulates many aspects of human-*M. tuberculosis* pathogenesis but has not been widely used to study the role of infection-induced lipid metabolism. We find infection-induced alterations in host Low Density Lipoprotein metabolism are conserved in the zebrafish model of mycobacterial pathogenesis. Depletion of LDLR, a key lipid metabolism node, decreased *M. marinum* burden, and corrected infection-induced altered lipid metabolism resulting in decreased LDL and the reduced rate of macrophage transformation into foam cells. Our results demonstrate a conserved role for infection-

induced alterations to host lipid metabolism, and specifically the LDL-LDLR axis, in

mycobacterial pathogenesis.

Introduction

Mycobacterial manipulation of host lipid metabolism and the transformation of macrophages

into foam cells are important pathways in the pathogenesis of *Mycobacterium tuberculosis*

(*Mtb*) [1]. Mycobacteria drive the transformation of infected macrophages into foam cells

through the imbalance of low density lipoprotein (LDL) influx and efflux mechanisms [2].

Given the significant proportion of the mycobacterial genome that is allocated to lipid

metabolism, and the utilisation of lipids such as cholesterol for growth *in vitro*, the abundance

of neutral lipids contained within the granuloma microenvironment may supply a carbon

source for intracellular mycobacterial survival and persistence [3].

Dyslipidaemia and foam cell differentiation are known to contribute to disease pathology and

prognosis in human *Mtb* infection where foamy macrophages may inhibit lymphocyte access

to infected macrophages, increase the build-up of caseum at the center of granulomas, and

provide lipids utilized during evasion of phagocytosis [4, 5]. However, the involvement of

foam cells and dyslipidemia in the pathogenesis of non-tubercular mycobacterial infections

such as *Mycobacterium avium* subspecies *paratuberculosis* (MAP) or *Mycobacterium*

marinum has not been explored. Our group has identified infection-induced changes to host

lipid metabolism genes in MAP-infected cattle and aberrant host lipid metabolism gene

expression and intracellular cholesterol accumulation within MAP-infected macrophages [6,

7]. Here, we have turned to the natural host-pathogen pairing of the zebrafish-*M. marinum*

platform to investigate the conservation and consequences of altered host lipid metabolism as

a conserved motif across mycobacterial infections.

Zebrafish (*Danio rerio*) have a functional innate immune system from early in embryogenesis with a high degree of homology to mammals and are adaptable to the study of early host-pathogen interactions [8]. *M. marinum* is a natural pathogen of fish and amphibian species and genomic analysis has identified *M. marinum* as the closest genetic relative of the *Mtb* complex, sharing 85% of orthologous regions with *Mtb* [9]. Zebrafish models of *M. marinum* infection have provided unique insight into host-pathogen interplay responsible for granuloma formation, exquisitely reproducing the cellular structure of the human-*M. tuberculosis* granulomas [10-12]. As such, the zebrafish-*M. marinum* model is an important platform for investigating immunopathogenesis of the mycobacterial granuloma, providing fundamental insights into the role of macrophage epithelial transdifferentiation in granuloma maturation [10].

The aim of this study was to explore the biological significance of infection-induced lipid metabolism in a model non-tubercular mycobacterial infection. Our findings demonstrate the key lipid metabolism node Low Density Lipoprotein Receptor (LDLR) mediates the uptake of lipids into the mycobacterial granuloma niche and aids the transformation of macrophages into foam cells.

Results

Infection-induced altered host dyslipidemia is recapitulated in the zebrafish-*M. marinum* infection model

Previous reports have identified hypercholesterolemia as a correlate of disease pathology during human *Mtb* infection [5]. To determine whether these findings were conserved within the zebrafish-*M. marinum* platform, we infected zebrafish embryos with *M. marinum* then

analysed the total embryo LDL and HDL content at 3 and 5 dpi, corresponding to an early

infection timepoint and the onset of granuloma necrosis, respectively. Calculation of

lipoprotein content revealed a skewing towards LDL demonstrating conservation of the

infection-induced dyslipidemia phenotype in our zebrafish embryo model (Figure 1A).

Knockdown of *Ldlra* reduces *M. marinum* burden in zebrafish embryos

Infection of macrophages with various species of mycobacteria, including in the zebrafish-*M.*

marinum system, increases *LDLR* transcription [6, 13]. Given the infection-induced changes

in the ligand LDL in our model, we tested the functional relevance of LDLR by morpholino

knockdown of the zebrafish *ldlra* gene [14]. Knockdown of Ldlra had no effect on embryo

development and editing of the *ldlra* transcript was observed (Figures 1B and 1C). Oil Red O

staining for lipids in Ldlra-depleted embryos revealed marked vascular lipid accumulation

consistent with previous reports of Ldlra depletion being atherogenic (Figure 1D) [14].

Following Ldlra-depletion, embryos were infected with fluorescent *M. marinum* and bacterial

burden was analysed by fluorometry. Zebrafish embryos depleted of Ldlra displayed a

significantly decreased bacterial burden at 3 and 5 dpi when compared to control embryos

(Figure 2).

Knockdown of Ldlra corrects infection-induced altered host lipid metabolism

Host lipid metabolism is a tightly controlled biological pathway, responsible for addressing

the cellular lipid requirements through the regulation of influx and efflux mechanisms. Of the

many surface receptors involved in lipid metabolism, LDLR is arguably the most important,

playing a key role in driving the uptake of circulating LDL [2]. We hypothesised knockdown

of Ldlra would thus restore homeostatic lipid metabolism.

Following *M. marinum* infection, expression of the LDLR-network and lipid efflux gene

ABCA1 was markedly increased (Figure 3A). Depletion of Ldlra significantly reduced the

magnitude of infection-induced *abca1* transcription (Figure 3A). Next, we hypothesised that

depletion of LDLR would prevent infection-induced hypercholesterolemia. Following *M.*

marinum infection, Ldlra-deficient embryos displayed a decreased LDL:HDL ratio compared

to infected controls at 5 dpi (Figure 3B). Together, these data demonstrate conservation and

correction of an infection-induced lipid metabolism network in our model.

Next, we hypothesised that the combination of reduced potential for LDL uptake and

correction of total cholesterol levels would reduce granuloma lipid accumulation within

granuloma foam cells in Ldlra-depleted embryos. Granuloma Oil Red O lipid staining density

was reduced in infected Ldlra-deficient embryos compared to infected controls demonstrating

a functional role for LDLR in directing bacterial-beneficial lipid accumulation during

mycobacterial infection (Figure 4).

Discussion

The elevation of host lipids following mycobacterial infection has been well documented in

both human and murine *Mtb* infections [5, 15]. Recently, we have shown that elevated serum

cholesterol is associated with histopathological lesions in sheep and cattle infected with MAP

[16]. The results from the current study are in agreement with previous findings and suggest

that the induction of host hypercholesterolemia is a conserved motif of pathogenic

mycobacteria. Our findings in this study demonstrate 1) infection-induced altered host lipid

metabolism is a conserved feature of non-tuberculous mycobacterial infection, and 2) host

expression of LDLR is co-opted by mycobacterial infection to increase susceptibility to

mycobacterial infection. Furthermore, our results demonstrated that *LDLR* knockdown

significantly altered the availability of circulating lipoproteins and decreased Oil Red O
staining density, providing strong evidence to suggest that the abundance of neutral lipids and
therefore foam cells, was largely decreased as a consequence of *LDLR* depletion.

Host lipid metabolism is a complex pathway involving many transcription factors, surface
molecules and feedback mechanisms. In the absence of key components, compensatory
pathways are utilised to achieve cellular homeostasis. Expression of *abca1* was increased
following *M. marinum* infection, in agreement with previous studies identifying the
upregulation of *ABCA1* by mycobacterial infection across multiple host species [17].
Interestingly, there was a notably lower expression in *ldlra*-deficient embryos, presumably
due to the decreased efflux of lipids from mycobacteria-infected macrophages as a
consequence of the altered lipoprotein abundance.

Collectively, this study provides evidence of conserved infection-induced altered host lipid
metabolism in the zebrafish-*M. marinum* model, and highlights a crucial role of *LDLR* in
infection susceptibility across host and pathogen species.

Materials and Methods

Zebrafish breeding and ethics statement

Adult zebrafish were housed at the Garvan Institute of Medical Research Biological Testing
Facility (St Vincent's Hospital AEC Approval 1511) and housed for infection experiments at
the Centenary Institute (Sydney Local Health District AEC Approval 2016-037). Zebrafish
embryos were obtained by natural spawning and embryos were raised at 28°C in E3 media .
All experiments and procedures were completed in accordance with Sydney Local Health
District animal ethics guidelines for zebrafish embryo research.

Morpholino design and microinjection

Splice-blocking morpholino oligonucleotide for *LDLR* (gene variant *ldlra*) was designed

from Ensembl transcript sequences (http://www.ensembl.org/index.html) and purchased from

Gene Tools. The *LDLR* morpholino (5'-ATCACATTTCATTTCTTACAGCAGT-3') was

targeted at the exon 4/intron 4 and 5 splice junction that was expected to cause the excision of

exon 4. The control morpholino (5'-CCTCTTACCTCAGTTACAATTTATA-3') is a

negative control targeting a human beta-globin intron known to cause little phenotype

variations in zebrafish.

Embryos at the 1-4 cell stage were injected with 2 nL of morpholino into the yolk. Embryos

were raised at 28°C in E3 media.

Molecular analysis of *ldlra* knockdown

RNA was extracted in TRIZOL (Thermofisher) by precipitation and cDNA was synthesised

with an Applied Biosystems High Capacity cDNA reverse transcription kit (Thermofisher).

PCR was performed with *ldlra*-specific primers (5'-AGAGCTGGAAATGTGACGGA-3'

and 3'-CTCATCTGGACGGCATGTTG-5') and visualised by gel electrophoresis.

Oil Red O staining

Oil Red O lipid staining on whole mount embryos was completed as previously described

[18]. Briefly, embryos were individually imaged for bacterial distribution by fluorescent

microscopy, fixed, and stained in Oil Red O (0.5% w/v in propylene glycol). Oil Red O

staining intensity at sites of infection were quantified in ImageJ and calculated as the pixel

density above background tissue.

Infection of zebrafish embryos

At 30 hours post-fertilisation, zebrafish embryos were anaesthetised with 160 µg/mL tricaine (Sigma-Aldrich) and infected with approximately 200 CFU M strain *M. marinum* via caudal vein injection. A dose of approximately 1000 CFU Δ*ESX1 M. marinum* was injected to match parental strain burden at 5 dpi.

Lipoprotein assay

Lipoprotein abundance was measured using the HDL & LDL/VLDL Cholesterol Assay Kit (Cell Biolabs) as per manufacturer's instructions and based on previously published methods [14]. Briefly, 10 embryos per treatment group were terminally anaesthetised and placed into BHT solution (10 µg/mL, Sigma-Aldrich) and homogenised. Homogenates were centrifuged at 4°C at 2000 x *g* for 5 mins to pellet debris. The supernatant was mixed with an equal volume of precipitation solution to separate the high density lipoprotein (HDL) and low density lipoprotein (LDL) fractions. Each lipoprotein fraction was mixed with an equal volume of reaction mix and quantified by fluorescent plate reader. A standard curve was included in each run to calculate the concentration of lipoproteins within each treatment.

Imaging

Live zebrafish embryos were anaesthetized in M-222 (Tricaine) and mounted in 3% methylcellulose for imaging on a Leica M205FA fluorescence stereomicroscope. Histological sections were imaged on a Leica DM6000B. Further image manipulation and/or bacterial quantification was carried out with Image J Software Version 1.51j.

Quantification of *M. marinum* burden by fluorescent pixel count

Infection burden was measured as the number of pixels in each embryo above background

fluorescence in ImageJ (National Institutes of Health) and pixels counted using the 'Analyse

particles' function [19].

Quantitative PCR

Quantitative PCR (qPCR) reactions were completed as previously described [20] using

Mx3000P real-time PCR systems (Stratagene). Gene-specific primers were generated for

abca1 (5'-AGCTGCTGGTCGAGATCATA-3' and 3'- CTGTCTCACTGCATGCTTGG-5')

for comparison against housekeeping gene *actb* (5'-CCTTCCAGCAGATGTGGATT-3' and

3'-CACCTTCACCGTTCCAGTTT-5').   Samples were run in duplicate with reaction

specificity confirmed based on the post-amplification dissociation curve.

Statistical analysis

Graphpad Prism was used to perform statistical testing as indicated.

Acknowledgements

Drs Anneliese Ashurst and Gayathri Nagalingam for training assistance with laboratory

equipment; Dr Kristina Jahn and Sydney Cytometry for assistance with imaging equipment.

Special thanks goes to the animal house staff at the Garvan Institute of Medical Research and

CSIC.

This work was supported by the Australian National Health and Medical Research Council

(APP1099912 and APP1053407 to S.H.O.); Meat and Livestock Australia (P.PSH. 0813 to

A.C.P. and K. dS); the Marie Bashir Institute for Infectious Diseases and Biosecurity (grant

to S.H.O., A.C.P. and K. dS); the Kenyon Family Foundation Inflammation Award (grant to

S.H.O.); the University of Sydney (fellowship to S.H.O.); Consellería de Economía, Emprego

e Industria (GAIN), Xunta de Galicia (grant IN607B 2016/12 to Institute of Marine Research

(IIM-CSIC)).

Author contributions

M.D.J. Planned experiments, performed experiments, analysed data, wrote the paper; E.H.

Planned experiments, performed experiments; J.A.K. Performed experiments; A.R.

Performed experiments, contributed reagents; B.N. Performed experiments, contributed

reagents; A.F. Contributed reagents; W.J.B. Analysed data, contributed reagents; K.dS.

Planned experiments, analysed data, contributed reagents; A.C.P. Planned experiments,

analysed data, contributed reagents; S.H.O. Planned experiments, performed experiments,

analysed data, contributed reagents, wrote the paper.

References

1. Lovewell, R. R., Sassetti, C. M. & VanderVen, B. C. (2016) Chewing the fat: lipid
metabolism and homeostasis during M. tuberculosis infection, *Curr Opin Microbiol.* **29**, 30-
6.
2. Cardona, P., Allain, S., Altare, F., Kim, M. & Russell, D. G. (2009) Foamy macrophages
and the progression of the human tuberculosis granuloma, *Nature Immunology.* **10**, 943-948.
3. Pandey, A. K. & Sassetti, C. M. (2008) Mycobacterial persistence requires the utilization
of host cholesterol, *Proceedings of the National Academy of Sciences of the United States of
America.* **105**, 4376-4380.
4. Russell, D. G., Cardona, P. J., Kim, M. J., Allain, S. & Altare, F. (2009) Foamy
macrophages and the progression of the human tuberculosis granuloma, *Nat Immunol.* **10**,
943-8.
5. Dong, Z., Shi, J., Dorhoi, A., Zhang, J., Soodeen-Lalloo, A. K., Chen, W., Yin, H., Sha,
W., Li, W., Zheng, R., Liu, Z., Yang, H., Qin, L., Wang, J., Huang, X., Wu, C., Kaufmann, S.
H. E. & Feng, Y. (2017) Hemostasis and Lipoprotein Indices Signify Exacerbated Lung
Injury in Tuberculosis With Diabetes Comorbidity, *Chest*.
6. Johansen, M. D., de Silva, K., Plain, K. M., Whittington, R. J. & Purdie, A. C. (submitted)
Mycobacterium avium subspecies *paratuberculosis* is able to manipulate host lipid
metabolism and accumulate cholesterol within macrophages, *Pathogens and Disease*.
7. Thirunavukkarasu, S., de Silva, K., Plain, K. M. & Whittington, R. J. (2014) Role of host-
and pathogen-associated lipids in directing the immune response in mycobacterial infections,

with emphasis on Mycobacterium avium subsp. paratuberculosis, *Critical reviews in*
microbiology, 1-13.
8. Traver, D., Herbomel, P., Patton, E. E., Murphey, R. D., Yoder, J. A., Litman, G. W.,
Catic, A., Amemiya, C. T., Zon, L. I. & Trede, N. S. (2003) The Zebrafish as a Model
Organism to Study Development of the Immune System, *Advances in Immunology.* **81**, 253-
330.
9. Stinear, T. P., Seemann, T., Harrison, P. F., Jenkin, G. A., Davies, J. K., Johnson, P. D. R.,
Abdellah, Z., Arrowsmith, C., Chillingworth, T., Churcher, C., Clarke, K., Cronin, A., Davis,
P., Goodhead, I., Holroyd, N., Jagels, K., Lord, A., Moule, S., Mungall, K., Norbertczak, H.,
Quail, M. A., Rabbinowitsch, E., Walker, D., White, B., Whitehead, S., Small, P. L. C.,
Brosch, R., Ramakrishnan, L., Fischbach, M. A., Parkhill, J. & Cole, S. T. (2008) Insights
from the complete genome sequence of *Mycobacterium marinum* on the evolution of
Mycobacterium tuberculosis, *Genome Research.* **18**, 729-741.
10. Cronan, M. R., Beerman, R. W., Rosenberg, A. F., Saelens, J. W., Johnson, M. G.,
Oehlers, S. H., Sisk, D. M., Jurcic Smith, K. L., Medvitz, N. A., Miller, S. E., Trinh, L. A.,
Fraser, S. E., Madden, J. F., Turner, J., Stout, J. E., Lee, S. & Tobin, D. M. (2016)
Macrophage Epithelial Reprogramming Underlies Mycobacterial Granuloma Formation and
Promotes Infection, *Immunity.* **45**, 861-876.
11. Oehlers, S. H., Cronan, M. R., Scott, N. R., Thomas, M. I., Okuda, K. S., Walton, E. M.,
Beerman, R. W., Crosier, P. S. & Tobin, D. M. (2015) Interception of host angiogenic
signalling limits mycobacterial growth, *Nature.* **517**, 612-5.
12. Oehlers, S. H., Cronan, M. R., Beerman, R. W., Johnson, M. G., Huang, J., Kontos, C.
D., Stout, J. E. & Tobin, D. M. (2017) Infection-induced vascular permeability aids
mycobacterial growth, *J Infect Dis.* **215**, 813-7.
13. Kim, M. J., Wainwright, H. C., Locketz, M., Bekker, L. G., Walther, G. B., Dittrich, C.,
Visser, A., Wang, W., Hsu, F. F., Wiehart, U., Tsenova, L., Kaplan, G. & Russell, D. G.
(2010) Caseation of human tuberculosis granulomas correlates with elevated host lipid
metabolism, *EMBO Molecular Medicine.* **2**, 258-74.
14. O'Hare, E. A., Wang, X., Montasser, M. E., Chang, Y. C., Mitchell, B. D. & Zaghloul, N.
A. (2014) Disruption of ldlr causes increased LDL-c and vascular lipid accumulation in a
zebrafish model of hypercholesterolemia, *Journal of lipid research.* **55**, 2242-2253.
15. Martens, G. W., Arikan, M. C., Jinhee, L., Ren, F., Vallerskog, T. & Kornfeld, H. (2008)
Hypercholesterolemia Impairs Immunity to Tuberculosis, *Infection and Immunity.* **76**, 3464-
3472.
16. Johansen, M. D., De Silva, K., Plain, K. M., Begg, D. J., Whittington, R. J. & Purdie, A.
C. (submitted) Sheep and cattle exposed to *Mycobacterium avium* subspecies
paratuberculosis exhibit altered total serum cholesterol profiles during the early stages of
infection, *Veterinary Immunology and Immunopathology*.
17. Long, J., Roy, R. B., Zhang, Y. J. J., Antrobus, R., Du, Y. X., Smith, D. L., Weekes, M.
P. & Javid, B. (2016) Plasma Membrane Profiling Reveals Upregulation of ABCA1 by
Infected Macrophages Leading to Restriction of Mycobacterial Growth, *Frontiers in*
Microbiology. **7**, 1086-1097.
18. Passeri, M. J., Cinaroglu, A., Gao, C. & Sadler, K. C. (2009) Hepatic steatosis in
response to acute alcohol exposure in zebrafish requires sterol regulatory element binding
protein activation, *Hepatology.* **49**, 443-52.
19. Matty, M. A., Oehlers, S. H. & Tobin, D. M. (2016) Live Imaging of Host-Pathogen
Interactions in Zebrafish Larvae, *Methods Mol Biol.* **1451**, 207-23.
20. Taylor, D. L., Zhong, L., Begg, D. J., de Silva, K. & Whittington, R. J. (2008) Toll-like
receptor genes are differentially expressed at the sites of infection during the progression of

Johne's disease in outbred sheep, *Veterinary Immunology and Immunopathology.* **124**, 132-
151.

Figure Legends

Figure 1: LDL is increased by infection and validation of Ldlra knockdown.

A, Quantification of LDL concentration in wild-type zebrafish embryos at 5 days following

M. marinum infection. Each data point represents a pooled sample of 10 embryos. Scale bars

represent 500 µm. Statistical tests were performed using unpaired Student's t-tests.

Experiments were completed with 4 independent biological replicates. *B,* Normal

morphology and phenotypes at 5 days post-fertilisation as visible by light microscopy. *C*,

Molecular validation of *ldlra* morpholino efficacy at 3 and 5 days post-treatment

demonstrating depletion of native 464 bp amplicon and appearance of aberrant 378 bp

amplicon. *D*, Oil Red O staining demonstrates visible lipid abundance within the vasculature

of Ldlra-deficient embryos. White arrows represent expansive lipid accumulation within the

vasculature.

Figure 2: Knockdown of Ldlra reduces *M. marinum* bacterial burden..

A, Representative images of *M. marinum* bacterial burden at 5 days-post infection. Scale bars

represent 500 µm. *B,* Quantification of *M. marinum* bacterial burden at 3 days post infection.

C, Quantification of *M. marinum* burden at 5 days post infection. Error bars represent

standard deviations, each data point represents a single embryo. Analysis of *M. marinum*

bacterial burden was unmatched T tests. Experiments were completed with 4 independent

biological replicates for each panel.

Figure 3: Mycobacterial infection-induced modulation of host lipid metabolism is corrected

in Ldlra-deficient embryos.

A, *abca1* transcript fold change expression at 5 days post-infection measured by qPCR,

normalised to control uninfected sample. Each column represents two independent biological

replicates analysed with technical replicates. *B*, Quantification of LDL concentration in

Ldlra-deficient embryos at 5 days-post infection. Each data point represents a pooled sample

of 10 embryos completed with four independent replicates.

Gene expression analysis was completed using one-way ANOVA. Analysis of LDL

concentration and Oil Red O density units was completed using unpaired Student's *t*-tests.

Figure 4: Infection-induced granuloma lipid accumulation is reduced in Ldlra-deficient

embryos.

A, Representative images of Oil Red O staining density at 5 days post-infection. Red circles

indicate areas of embryos with largest foci of infection. Scale bar represents 500 µm. *B,*

Quantification of Oil Red O density units at 3 days post-infection with *M. marinum. C,*

Quantification of granuloma Oil Red O staining density units at 5 days post-infection. Each

data point represents an individual granuloma. Error bars represent standard deviation.

Statistical tests were performed using unpaired Student's t-tests.