The *Chlamydia* type III effector TarP alters the dynamics and organization of host cell focal adhesions

António T. Pedrosa[b,c,1], Korinn N. Murphy[a,b], Ana T. Nogueira[b,c,2], Amanda J. Brinkworth[a], Tristan R. Thwaites[c,3], Jesse Aaron[d], Teng-Leong Chew[d], and Rey A. Carabeo[a]

[a]Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE; [b]School of Molecular Biosciences, College of Veterinary Medicine, Washington State University, Pullman, WA; [c]Bacteriology Section, Programme in Microbiology, Institute of Medical Sciences, University of Aberdeen, Aberdeen, UK; [d]Advanced Imaging Center, Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA

Corresponding author:
Rey A. Carabeo, Ph.D.
Department of Pathology and Microbiology
University of Nebraska Medical Center
Omaha, NE 68198-5900
(402) 836-9778
Email: rey.carabeo@unmc.edu

Current address:
[1]Molecular Cell Biology Laboratory, NHLBI, NIH, Bethesda, MD
[2]Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC
[3]Cell and Gene Therapy Catapult, London, UK

Keywords: *Chlamydia*, Focal adhesion, Pathogenesis, Cell adhesion, iPALM

Abstract

The human pathogen *Chlamydia trachomatis* targets epithelial cells lining the genital mucosa.

We observed that infection of various cell types, including fibroblasts and epithelial cells resulted

in the formation of unusually stable focal adhesions that resisted disassembly induced by the

myosin II inhibitor, blebbistatin. Super-resolution microscopy revealed in infected cells the

vertical displacement of paxillin and FAK from the signaling layer of focal adhesions; while

vinculin remained in its normal position within the force transduction layer. The candidate type III

effector TarP which localized to focal adhesions during infection and when expressed

ectopically, was sufficient to mimic both the reorganization and blebbistatin-resistant

phenotypes. These effects of TarP, including its localization to focal adhesions required

interaction with the host protein vinculin through a specific domain at the C-terminus of TarP.

The consequence of *Chlamydia*-stabilized focal adhesions was restricted cell motility and

enhanced attachment to the extracellular matrix. Thus, via a novel mechanism, *Chlamydia*

inserts TarP within focal adhesions to alter their organization and dynamics.

Introduction

Bacterial infection of mucosal epithelial cells triggers the antimicrobial defense strategy of cell exfoliation and apoptosis induction (reviewed in: Kim *et al.*, 2010). The controlled extrusion of damaged host cells and colonizing pathogens requires the degradation of cell adhesion factors. In epithelial cells, focal adhesions and hemidesmosomes are primarily responsible for attachment to the extracellular matrix, and their assembly and turnover are exquisitely regulated at multiple levels, by kinases, phosphatases, protein-protein interactions, internalization of components, and degradation (Borradori and Sonnenberg, 1999; Geiger *et al.*, 2001; Rosenblatt *et al.*, 2001; Zaidel-Bar *et al.*, 2007). Disruption of one or more of these regulatory processes alters the adhesion dynamics and properties of the cells.

One strategy employed by bacteria to neutralize exfoliation relies on the precise targeting of one or more components of the focal adhesion proteome. The best-characterized example is that of *Shigella*, which neutralizes epithelial extrusion to colonize the epithelium efficiently (Kim *et al.*, 2009). It does so by delivering the OspE effector by the type III secretion system (T3SS). This protein reinforces host cell adherence to the basement membrane by interacting with integrin-linked kinase (ILK), a serine/threonine kinase that is part of the focal adhesome (Kim *et al.*, 2009; Zaidel-Bar *et al.*, 2007). A consequence of the OspE-ILK interaction is an increased surface expression of β1-integrin, which in turn promotes focal adhesion (FA) assembly. In addition, the OspE-ILK complex stabilizes the focal adhesions (FAs) by reducing phosphorylation of focal adhesion kinase (FAK) at a functionally important Tyr397 residue and of paxillin. Inhibition of both phosphorylation events has been shown to induce FA disassembly (Kim *et al.*, 2009). Interestingly, some EPEC and EHEC strains, as well as *Citrobacter rodentium* possess the effector EspO, which shares strong homology with OspE (reviewed in Vossenkämper, Macdonald and Marchès, 2011; Morita-Ishihara *et al.*, 2013). As such, it is conceivable that these pathogens also reinforce adherence of the infected epithelial cells to secure an infectious foothold. The EspZ effector of EPEC and EHEC has been shown to reduce cell death and detachment *in vitro* (Shames *et al.*, 2010). EspZ binds the transmembrane glycoprotein CD98 and enhances its effect on β1-integrin signalling and cell survival via activation of FAK (Shames *et al.*, 2010).It is possible that EspO and EspZ may cooperate to confer enhanced adhesion of the host epithelial cells to the extracellular matrix. Finally, through interaction with human carcino-embryonic antigen-related cell adhesion molecules (CEACAM), bacterial pathogens such as *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Moraxella catarrhalis*, and *Haemophilus influenzae* can activate β1-integrin signalling and inhibit epithelial cell detachment (reviewed in: Kim *et al.*, 2010). Despite

numerous examples of pathogens manipulating host cell adhesion, the details of these

mechanisms remain uncharacterized.

Chlamydiae are obligate intracellular pathogens that are distinguished by their biphasic

developmental cycle that alters between the infectious elementary body (EB), and the

replicative, but non-infectious reticulate body (RB). At late time points, the non-infectious RBs

convert back to EBs to produce infectious particles for the next round of infection. The entire

intracellular growth cycle of *Chlamydia* takes ~48–96 h and occurs within a membrane-bound

inclusion, and most of it is spent in the non-infectious RB form. Thus, it is essential that the

adhesion of the infected cells to the epithelium is sustained during chlamydial development to

enable the differentiation of the non-infectious RBs to the infectious and stable elementary

bodies (EBs) (reviewed in AbdelRahman and Belland, 2005). This means that *Chlamydia* must

evade a host of anti-microbial defenses, including epithelial extrusion.

Previous works by Kumar and Valdivia (2008) and Heymann *et al.*, (2013) described the

loss of motility of *Chlamydia*-infected epithelial cells. Heymann *et al.*, (2013) attributed this to the

chlamydial inhibition of Golgi polarization that occurs at >24 h post-infection, leading to loss of

directional migration. In this report, we offer an alternate and possibly complementary

mechanism of FA stabilization, which could lead to an increase of host-cell adhesion to the

extracellular matrix (ECM), thus culminating in previously reported loss of motility (Kumar and

Valdivia, 2008; Heymann *et al.*, 2013). Using quantitative confocal and live-cell imaging and

super-resolution microscopy, we describe the various *Chlamydia* infection-dependent changes

that occur to FAs, such as increased numbers, enhanced stability, and altered organization. We

provide evidence implicating the T3SS effector TarP, and its interaction with the focal adhesion

protein vinculin. We show that vinculin and its binding motif in TarP is required for the

localization of the effector to focal adhesions, and their resistance to blebbistatin-induced

disassembly. TarP localization to focal adhesions is also required for the displacement of the

focal adhesion kinase and paxillin from their normal position within the integrin signaling layer.

We also show that TarP alone was sufficient to restrict cell motility. Overall, the results indicate

that *Chlamydia* has a dedicated mechanism of modulating focal adhesion dynamics, which may

be linked to the maintenance of *Chlamydia* infection in a high-turnover tissue site.

RESULTS

Chlamydia infection enhances FA numbers

Cos7 cells were infected and 24 h post-infection (hpi), cells were fixed and prepared for indirect

immunofluorescence imaging of paxillin-positive FAs. As shown in **Figure 1**, cells infected with

C. trachomatis serovar L2, serovar D, serovar B, *C. caviae* GPIC, and *C. muridarum* (MoPn)

consistently had greater numbers of FAs than mock-infected cells. We further explored the

apparent infection-dependent increase in FA numbers using serovar L2, and observed

enhanced FA numbers at 8 hpi that increased by 24 hpi. Next, we asked if the process is

pathogen-directed. Specifically, we investigated if this effect on FAs required de novo protein

synthesis by *Chlamydia*. Cos7 cells were infected with CtrL2 for the specified duration followed

by treatment by the bacterial translation inhibitor, chloramphenicol (Cm). We observed that

while an 8-h protein synthesis inhibition was not sufficient to prevent the effects on FA number,

the 24-h Cm treatment reduced FA numbers of infected cells to the level of mock-infected

control (**Figure 2**). The results indicate that the latter phase of focal adhesion alterations

requires either the *de novo* synthesis of new proteins by *Chlamydia* or replenishment of

effectors packaged in the metabolically quiescent EBs. These effectors are delivered by the

type III secretion system early in infection, prior to differentiation to the vegetative reticulate

body form, when they gain the ability for macromolecular synthesis.

A marked difference was the increased numbers of FAs at the interior relative to the cell

periphery. Focal adhesion maturation is associated with movement away from the cell periphery

and towards the center (Smilenov, et al. 1999). FAs at the interior of the cell either mature to

become stable fibrillar adhesions to promote cell attachment or disassemble during migration

(Smilenov, et al. 1999 ; Dumbauld *et al.*, 2010; reviewed in: Nagano *et al.*, 2012). We speculated

that the increased numbers of interior FAs arose from infection-dependent stabilization. To

assess stability, we took advantage of the enhanced turnover of FAs in the presence of the

myosin II-specific inhibitor, blebbistatin (Straight *et al.*, 2003). FA stability is dependent on

tension within and between focal adhesions (Chrzanowska-Wodnicka and Burridge, 1996;

Pasapera *et al.*, 2010; Carisey *et al.*, 2013). This tension is largely provided by the contractile

action of the molecular motor myosin II on stress fibers (SF); and its inhibition by blebbistatin

consistently leads to FA disassembly and altered motility (Wang *et al.*, 2008; Dumbauld *et al.*,

2010, Liu *et al.*, 2010). We evaluated the relative resistance of FAs to a 60-min treatment with

10 μM blebbistatin in the context of infection with CtrL2 at 20 hpi. As shown in **Figure 3** of

representative samples, SFs in the mock-infected cell disassembled, with the simultaneous

disappearance of FAs marked with paxillin. In contrast, FAs in the CtrL2-infected cells

remained. Infection did not prevent blebbistatin-induced disassembly of stress fibers. Taken

together, the results point to a dedicated mechanism in *Chlamydia trachomatis* and perhaps

other chlamydial species to stabilize focal adhesions. In addition, focal adhesion stabilization in

infected cells did not require stress fibers.

The type III effector TarP localizes to focal adhesions in a vinculin-dependent manner

The chlamydial type III effector TarP has been implicated in the invasion process during

infection of non-phagocytic cells. Specifically, TarP translocation by the elementary bodies

contributes to the actin remodeling that is required for uptake of the pathogen (Clifton *et al.*,

2005; Jewett *et al.*, 2006; Lane *et al.*, 2008; Thwaites *et al.*, 2014). Interestingly, TarP was also

reported to have a role in increased resistance of infected cells to apoptosis, raising the

possibility that this effector has post-invasion function (Mehlitz *et al.*, 2010). Consistent with this

idea is the continuous presence of the protein throughout infection (Clifton *et al.*, 2004). In

addition, there was sustained presence of TarP translocated to the cytosol as indicated by its

reactivity to the anti-phosphotyrosine 4G10 antibody under conditions that prevented further

synthesis of this protein, i.e. Cm treatment (**Figure S1**). During invasion, TarP localizes to sites

of chlamydial adhesion at the plasma membrane (Clifton *et al.*, 2004). If TarP has a role post-

invasion, we expect TarP to be found at sites where it exerts its function. Immunostaining with a

rabbit polyclonal antibody to C. *trachomatis* serovar L2 (CtrL2) TarP of infected mouse

embryonic fibroblasts (MEFs) revealed specific staining of focal adhesions (FA), in addition to

punctae within inclusions, which are likely to be the bacteria (**Figure 4**). Uninfected cells

consistently exhibited diffused background immunofluorescence signal, illustrating specificity of

the antibody to TarP localized to FAs.

We then sought to determine if ectopically expressed TarP would yield a similar

subcellular localization to FAs. TarP and its deletion derivatives shown in Figure 5A were fused

to the fluorescent protein mTurquoise2, and ectopically expressed in MEFs . TarP has multiple

domains that resemble motifs for protein-protein interaction and signaling, including a repeated

50-amino acid domains that is tyrosine-phosphorylated by Src family kinases and the Abl

kinase, actin-binding domains, a leucine-aspartate (LD) domain recognized by the focal

adhesion kinase (FAK), and vinculin-binding domains (VBD) (Clifton *et al.*, 2005; Jewett *et al.*,

2006, 2010; Lutter *et al.* 2010; Mehlitz *et al.* 2010; Jiwani *et al.*, 2013; Thwaites *et al.*, 2014,

2015; Braun *et al.*, 2019*)* . All have been demonstrated in invasion-related actin-recruitment

assays to be functional(Lane *et al.*, 2008; Thwaites *et al.*, 2014, 2015) . We created various C-

terminal deletion constructs of TarP fused to mTurquoise for heterologous expression in MEFs,

as well as a TarP derivative lacking the proline-rich domain (PRD) to minimize non-specific

aggregation of the protein. Transfected cells were counterstained with a monoclonal antibody to

paxillin to visualize FAs. As shown in **Figure 5B**, colocalization of TarP with the focal adhesion

marker paxillin required the LD and VBD motifs. Note that paxillin localization was observed for

both full-length TarP and TarP ∆PRD (**Figure S2**)

We previously reported that the LD and VBD domains were recognized by FAK and

vinculin, respectively (Thwaites *et al*., 2014, 2015). They are distinct non-overlapping domains

that interacted with their respective binding partners independently. Therefore, we evaluated if

the loss of either of the binding partner (e.g. FAK or vinculin) would result in the loss of FA

localization. To address the functional relevance of these interactions, albeit in a post-invasion

context, the LDVBD construct was expressed in wild type, *vcl-/-* (vinculin), or *ptk2-/-* (FAK) MEF

knockout mutants. We observed FA localization of mTurquoise-LDVBD in wild type MEFs, but

not in *vcl-/-* MEFs (**Figure 5B**). The loss of FAK did not affect the focal adhesion localization of

LDVBD (**Figure 5C**). Together the data indicated that TarP localization to FAs required the host

protein vinculin, likely through its interaction with the VBD domain. FAK was dispensable in this

regard. The TarP-positive subcellular structures were verified as FAs based on β1-integrin

staining using a monoclonal antibody specific to the conformationally active form of the receptor

(**Figure 5D**).

The ectopic expression of TarP is sufficient to increase focal adhesion numbers and

confer vinculin-dependent resistance to blebbistatin

We observed increased paxillin-positive FAs in TarP-transfected cells, provided that the TarP

construct retained the VBD domain. Images from the FA localization experiments in Cos7 cells

were re-analyzed by quantifying the number of focal adhesions per cell in cells transfected with

the mTurquoise vector alone, full-length TarP, or LDVBD (**Figure 6A**). Paxillin-positive

structures in transfected cells were counted in NIH ImageJ, and the data is illustrated as box-

whisker plots in **Figure 6B**. We observed statistically significant increases in focal adhesion

numbers in cells expressing TarP ∆PRD and LDVBD. These results suggested that the ability of

TarP to localize to FAs, which was mediated by the LDVBD domain of TarP was linked to its

effects on FA numbers.

We also evaluated the effect of ectopically expressed LDVBD on the stability of focal

adhesions in fixed cells. As illustrated in **Figure 7A**, *Chlamydia*-infected wild type MEFs

retained paxillin-marked FAs after 60 min of treatment with 10 uM blebbistatin, while mock-

infected cells lost them. In contrast, infection of the *vcl-/-* MEFs failed to inhibit blebbistatin-

induced disassembly of focal adhesions, highlighting the crucial role of the host protein vinculin

in FA stability. Using these results as reference, we investigated if the LDVBD domain was

sufficient to confer a similar level of resistance to blebbistatin-induced disassembly, and if it did

so in a vinculin-dependent manner. LDVBD transfection of wild type MEFs led to the retention of

FAs after the 60-min treatment with blebbistatin, while the *vcl-/-* MEFs lost these structures

despite LDVBD expression (Figure 7B). Therefore, we concluded that vinculin plays an

important role in TarP-dependent stabilization of FAs. However, we could not determine if the

apparent stabilizing role of vinculin is due to the FA localization of TarP or an alteration of its

activity as a result of its interaction with TarP.

Infection disrupts the stratified organization of focal adhesions, a phenotype mimicked

by the ectopic expression of TarP

Focal adhesions are organized into distinct strata termed the integrin layer, the signaling layer,

which harbors paxillin and FAK amongst others, and the force transduction layer that contains

vinculin, talin, and other mechanosensitive proteins. At the highest layer, actin and actin-

associated proteins, such as α-actinin and myosin II are found (Betzig *et al.*, 2006;

Kanchanawong *et al.*, 2010). Given the profound effect of infection and TarP ectopic expression

on FA stability, we investigated using interferometric photoactivated localization microscopy

(iPALM) their effects on FA organization. The localization of paxillin, FAK, and vinculin, all fused

to mTurquoise2 were monitored in mock-infected, CtrL2-infected, TarP full-length, or LDVBD-

transfected cells (**Figure 8**). Image analysis revealed dramatic reorganization of FAs with

regards to paxillin and FAK. In control samples, paxillin, FAK, and vinculin were found 50.5,

40.4, and 71 nm from the bottom of the cell, respectively, consistent with previous findings

(Kanchanawong *et al.*, 2010). However, paxillin and FAK shifted upwards to 176.9 and 96 nm,

respectively in infected cells, while no change in location was observed for vinculin (**Figure 8A**).

The state of infection of cells analyzed are shown in **Figure S3**.

Expression of TarP ΔPRD also caused a shift (240.2 nm) in paxillin localization (**Figure**

8B). LDVBD expression caused a noticeable shift (47 nm, with a second peak at 130 nm), but to

a lesser degree than full-length TarP. Interestingly, vinculin was not affected by either infection

or TarP ΔPRD (or LDVBD) expression, which points to the specific disruption of FA organization

by *Chlamydia*.

Cell motility is restricted in Chlamydia-infected or TarP-expressing cells

It was previously reported that *Chlamydia*-infected cells were restricted in their motility, and this

was attributed to the inability of infected cells to establish front-rear polarity due to Golgi

fragmentation induced by the pathogen. We decided to reinvestigate the loss of motility of

infected cells by focusing on focal adhesion dynamics. The decision of the cell to migrate or

adhere involves the regulation of focal adhesion stability in response to external cues, such as

chemoattractants and extracellular matrix (ECM) stiffness. First, we verified that *Chlamydia*-

infected mouse embryo fibroblasts were severely limited in their ability to migrate, relative to

mock-infected control cells (Movie S1). The manual tracking plugin of ImageJ was utilized to

obtain cell trajectory tracks for the motility assay. Cells were tracked using the position of the

nucleus over time. The coordinate data was input into ibidi's chemotaxis and migration tool to

obtain velocity and distance measurements. Velocity measurements revealed a 1.5-fold

decrease in the mean rate of migration of infected cells (**Figure 9A and 9C**).

Mouse embryonic fibroblasts were transfected to ectopically express mTurquoise2

vector only or the LDVBD domain. Cell migration was monitored in the DIC channel with

fluorescence images taken at the end of the motility assay (**Movie S1**; **Figure S4**). To quantify

motility, the cells were tracked as described above, with accompanying velocity calculations. As

shown in **Figure 9A and 9B**, transfected cells were significantly restricted in their motility

relative to mock-transfected controls. Both distance and velocity of LDVBD-transfected cells

were further restricted to those of infected cells (**Figure 9C**) indicating that inhibition of cell

migration by *C. trachomatis* could be accounted for fully by TarP overexpression. The enhanced

inhibition of migration distance and velocity in transfected cells may have been due to increased

levels of LDVBD when compared to levels present during infection.

Infection by Chlamydia trachomatis but not ectopic expression of TarP confers

resistance to detachment by mild trypsinization

Exfoliation of epithelial cells from the infected epithelium has been reported in rodent models of

ocular and genital infection; and both reports speculated the involvement of neutrophil-derived

proteases in the process (Ramsey *et al.*, 2005; Lacy *et al.*, 2011). We evaluated the resistance

of infected epithelial cells to detachment by 0.025% trypsin, and monitored for cell rounding by

time-lapse imaging at 1-min intervals for 30 min (Movie 1). Uninfected HeLa cells started

detaching by 7 min post-trypsinization, while *C. trachomatis* L2-infected cells remained attached

through the duration of imaging (30 min *C. trachomatis* post-trypsinization). To evaluate if TarP

ectopic expression would be sufficient to resist detachment, the cells were transfected for 24 h

to overexpress (LDVBD-mTurquoise2). Monolayers were imaged under fluorescence

microscopy at 0, 15, and 30-min after trypsinization. If TarP LDVBD overexpression was

sufficient to induce detachment resistance, we would expect an enrichment in remaining

adherent cells of those expressing LDVBD-mTurquoise2 than cells expressing mTurquoise2

only. Percentage values for both LDVBD-mTurquoise2 and vector-only samples were 11.9%

vs. 11.0% prior to trypsinization. We obtained the following for LDVBD vs. vector-only; 10.2%

vs. 10.3% (15 min) and 8.7% vs. 11.0% (30 min). No statistically significant differences were

found between LDVBD and vector only for either trypsinization time point. These results

indicated that, while TarP is able to reduce cell motility, it was not sufficient to confer resistance

to detachment by mild trypsinization. We interpret this to mean that additional changes to focal

adhesions, possibly mediated by additional chlamydial factors are required.

Discussion

Discussion

All chlamydial species, to varying extents exhibit tropism to epithelial cells, and thus likely

evolved to counteract cell extrusion associated with the normal cycle of turnover of the

epithelium, or as an anti-microbial mechanism to limit dissemination and eliminate infection. The

latter may also involve polymorphonuclear (PMN) cells, which secrete proteases to degrade

adhesion structures of epithelial cells. Thus, *Chlamydia*, with its biphasic developmental cycle

that involves a temporary loss of infectivity, is subjected to a very strong selective pressure to

acquire mechanisms to inhibit epithelial cell extrusion. A large portion of the developmental

cycle is spent in the non-infectious form, and thus, it is crucial to the survival of the pathogen to

inhibit extrusion of host epithelial cells before *Chlamydia* could convert to the infectious form.

Here, we demonstrated that adhesion of infected cells is enhanced via the action of TarP, an

effector protein conserved in the genus *Chlamydia*, and the host cell protein vinculin.

Our data collectively point to FAs as targets for modulation by *Chlamydia*. The type III

effector TarP plays a role in this modulation. The mechanism involves the vinculin-dependent

localization of TarP to FAs resulting in increased stability as indicated by their resistance to

disassembly by the myosin II inhibitor blebbistatin. We also made the novel observation of FA

reorganization in *Chlamydia*-infected or TarP-expressing cells, with paxillin and FAK displaced

from the signaling layer. A pressing question is whether the reorganization is the cause or the

effect of enhanced stability of TarP-targeted FAs. Considering the reported relative stability of

FAs in FAK-depleted cells, it is possible that the displacement of FAK by TarP could be

analogous to a loss-of-function. A more comprehensive investigation of FA disorganization by

TarP is required to define the exact mechanism of FA stabilization by this chlamydial effector.

We did not observe any change to tyrosine autophosphorylation at position 397; and we cannot

exclude the possibility that despite the apparent phosphorylation level, FAK displacement might

negatively affect interactions with signaling proteins that recognize and bind this motif, like the

Src kinase. The progression of tyrosine phosphorylation along the FAK protein by Src is crucial

to the role of FAK in disassembling FAs. Therefore, restricted FAK-Src interaction as a

consequence of FAK displacement in infected cells remains a possibility. Paxillin was similarly

displaced in both infected and TarP-expressing cells, which likely disrupts protein-protein

interactions and signaling related to paxillin within FAs. To our knowledge, FA reorganization to

the extent that we observed in infected cells has not been reported, pointing to a possible novel

mechanism of regulating FA dynamics.

While TarP, specifically the LDVBD domain was sufficient to drive FA stability to inhibit

cell motility, it was unable to mimic the resistance of infected cells to detachment by mild

trypsinization, which was meant to replicate polymorphonuclear cell (PMN)-mediated extrusion

of infected epithelial cells. This would be consistent with the involvement of additional

chlamydial factors that may mediate various aspects of focal adhesion characteristics, in

addition to numbers. Additional factors may facilitate progression of maturation, possibly to

fibrillar adhesions. Another might be the infection-dependent production of extracellular matrix

components by the host cell, including collagen. Increased deposition of collagen underneath

infected cells would likely influence FA stability by engaging integrins and inducing "outside-in"

FA-stabilizing signals. Thus, *Chlamydia* might have multiple cooperating mechanisms ensuring

the strong adhesion of its host cell, highlighting the importance of a mechanism to counteract

epithelial cell extrusion.

A pressing question is the biological significance of FA stabilization by this pathogen. We

speculate that this might be one mechanism by which *Chlamydia* neutralize extrusion of

epithelial cells. It is known that this process could limit infection dissemination. Shedding of

epithelial cells from uropathogenic *E. coli* (UPEC)-infected bladder is thought to reduce bacterial

burden, facilitating resolution of infection (Mysorekar *et al.*, 2002). The intestinal pathogen

Shigella possesses an effector, OspE that modulates epithelial cell attachment to facilitate the

pathogen's cell-to-cell spread. Indeed, loss of OspE resulted in a significant decrease in

virulence (Kim *et al.*, 2009, 2010). Epithelial cells of the genital tract or the ocular mucosa also

experience higher rates of turnover and are constantly replenished (Kuwabara, Perkins and

Cogan, 1976; Crosson, Klyce and Beuerman, 1986; Anderson, Marathe and Pudney, 2014).

Epithelial extrusion is a complex process that involves, not only promoting detachment of the

cells from the ECM, but also modulating interactions with neighboring cells via disassembly of

intercellular junctions (Rosenblatt, Raff and Cramer, 2001; Gudipaty and Rosenblatt, 2017).

Another possibility is increased resistance to apoptosis. Cell detachment is associated with

anoikis, a programmed cell death associated with loss of adherence (Frisch and Francis, 1994;

Gudipaty *et al.*, 2018). Focal adhesions provide anti-apoptotic signals (Frisch *et al.*, 1996), and

Chlamydia stabilization of these structures could promote apoptosis resistance.

An interesting question is the means by which epithelial cell extrusion is triggered. Is it

part of the normal cell turnover during tissue remodeling/renewal, or is it linked to pathogen

recognition? The shedding of bladder epithelial cells during UPEC infection requires the

expression of bacterial Type 1 pili, which is a potent pathogen-associated molecular pattern

(PAMP) that is recognized by the toll-like receptor 4 (TLR4) (Mysorekar *et al.*, 2002), raising the

intriguing possibility of a direct link between regulation of cell adhesion dynamics and pathogen

recognition. Various chlamydial species are recognized by toll-like receptors expressed on

epithelial cells e.g. TLR2, TLR3, TLR4, and TLR9 (Derbigny, Kerr and Johnson, 2005;

O'Connell *et al.*, 2006; Shaw *et al.*, 2011; Derbigny *et al.*, 2012; Pan *et al.*, 2017; Carrasco *et*

al., 2018; Kumar *et al.*, 2019), which may be indicative of epithelial detachment being linked to

pathogen recognition.

In addition to the demonstration that all tested *Chlamydia* species exhibited enhanced

FA stability, the presence of a type III effector with FA modulating function highlights the

importance of this process to *Chlamydia*; and yet, this aspect of Chlamydia-host cell interaction

is relatively understudied. The availability of a *Chlamydia* mutant lacking TarP and an efficient

method to image epithelial cell extrusion in the genital tract of animal models of infection are

critical to advancing studies in this area.

Materials and Methods

Cell culture. Cos7 (ATCC CRL-1651), NIH3T3 (kindly supplied by Hector Aguilar-Carreño

ATCC CRL-1658) and HeLa 229 (ATCC CCL-2.1) Mouse embryonic fibroblasts (MEFs) $vcl^{-/-}$

and matched HeLa 229 MEFs $vcl^{+/+}$ (Marg *et al.*, 2010) (were kindly provided by Dr. Wolfgang

Ziegler Hannover Medial School).cells were culture using Dulbecco's Modified Eagle Medium

(DMEM) (Thermofisher scientific, 11960-085). Media were supplemented with 10% fetal bovine

serum (Sigma, F0804-500ML), 2mM L-glutamine, and 10µg/ml gentamicin. The human

keratinocytes HaCaT cells (kindly supplied by Dr. Kristin M. Braun) were cultured in 3 parts

DMEM and 1 part Ham's F-12 Nutrient Mix (Thermofisher scientific 11765054), supplemented

with 10% fetal bovine serum (Sigma, F0804-500ML), 2 mM L-glutamine, 10µg/ml gentamicin,

insulin (Sigma I9278-5ML) and hydrocortisone cholera toxin EGF (HCE) cocktail. *Chlamydia*

trachomatis serovar L2 (L2/434/Bu) was propagated in HeLa 229. EBs were harvested by

discontinuous density gradient centrifugation in gastrografin (Bracco Diagnostics), as previously

described (Thwaites *et al.*, 2014).

***Chlamydia* infections.** Cells were infected with *Chlamydia trachomatis* serovar L2 (L2/434/Bu,

CtrL2) at the multiplicity of infection MOI of 5, for 20 h, and of 25, for 8 h, in ice cold serum-free

DMEM. Cells were centrifuged at 1000 rpm for 5 min at 4 °C to synchronize the infection. After

centrifugation, the inoculum was replaced with warm DMEM supplemented with 10% fetal

bovine serum, 2 mM L-glutamine, and 10µg/ml gentamicin. In parallel, a mock-infected control

was made following the same protocol but without *Chlamydia* infectious particles. Infection by

other strains/serovars was as follows. Cos7 cells were grown on glass coverslips. Cells were

infected with *Chlamydia trachomatis* serovar L2, serovar D, serovar B, *Chlamydia muridarum*

(MoPn), or *Chlamydia caviae* (GPIC) at an MOI of 5 for 24 hours. Cells were centrifuged at 500

x g for 15 minutes at 4°C to synchronize the infection. A mock-infected control was made

following the same protocol but without *Chlamydia* infectious particles. Prior to infection with

Serovar D, cells were pre-treated with 1x DEAE-Dextran for 15 minutes at room temperature.

Pre-treatment was followed by two washes with 1x HBSS and replacement with DMEM to

continue the infection.

Immunostaining. Cells were grown on fibronectin coated coverslips (Neuvitro, GG-12-

fibronectin) for the duration of the experiment. At the pre-determined, time cells were rinsed with

Hank's Balanced Salt Solution (HBSS) (Thermofisher scientific, 14025-100) and fixed using 4%

paraformaldehyde (PFA) in PBS pH 7.4 (Gibco, 14190-094) for 20 min at room temperature.

The fixed cells were then permeabilized using PBS with 0.2% Triton X-100. Subsequently,

permeabilized cells were incubated with 1% BSA (Sigma, A9418) in PBS for 30 min at room

temperature to block non-specific antigen binding. Cells were then incubated with the primary

antibodies overnight at 4ºC with rocking. The primary antibodies used in this study were rabbit

polyclonal antibody against FAK phosphorylated at tyrosine 397 (pFAK-Y397) (Abcam, ab4803),

rabbit monoclonal antibody paxillin (Abcam, ab32084), mouse monoclonal antibody vinculin

(Abcam, ab18058), rat monoclonal 9EG7 against the active form of β1-integrin (BD Biosciences,

553715), mouse monoclonal Flag-tag antibody (Cell Signalling, 8146S) mouse monoclonal

antibody *Chlamydia* LPS (Abcam, ab62708) and convalescent human sera. Afterwards cells

were incubated with appropriate fluorescently conjugated secondary antibodies and, when

specified, with DAPI (Roche, 10236276001) and Alexa flour 488 phalloidin stains, for 1 hr at

room temperature, with rocking. In this study, the following secondary antibodies were used:

goat anti-rabbit Alexa flour 488 (Thermofisher Scientific, A11008), goat anti-rabbit Alexa flour

633 (Thermofisher Scientific, A21071), goat anti-mouse Alexa flour 594 (Thermofisher Scientific,

A11005), goat anti-human Alexa flour 647 (Thermofisher scientific A-21445). Following staining,

the coverslips were mounted with Mowiol, and visualized in ZEISS LSM 710 confocal

microscope, in the Microscopy and Histology Core Facility at the University of Aberdeen, or the

Leica SP8 confocal microscope in Washington State University Integrative Physiology and

Neuroscience advance image equipment. FIJI software (Schindelin *et al.*, 2012; Schneider,

Rasband and Eliceiri, 2012) was used to generate the final images.

To localize endogenous TarP to focal adhesions, Mouse embryonic fibroblasts (MEFs)

grown on glass coverslips were infected with *Chlamydia trachomatis* L2 at an MOI of 10 for 20

hours. Cells were centrifuged at 500G for 15 minutes at 4°C to synchronize the infection. A

mock-infected control was made following the same protocol but without *Chlamydia* infectious

particles. Cells were fixed using ice cold 100% methanol for one minute. Cells were then

blocked with 5% BSA for one hour at room temperature. Focal adhesions and TarP were

visualized respectively using a primary monoclonal Talin1 antibody (Novus Biologics, NBP2-

50320) and rabbit polyclonal TarP antibody generated against the epitope (661-710 a.a.) (Li

International, Denver, CO). Samples were incubated overnight at 4°C. Immunostaining with

secondary antibodies was as described above.

Time-lapse microscopy. For live-cell imaging of FAs fibroblasts were seeded on ibidi μ-slide 8

well chambers with fibronectin coating (ibidi, 80823) at the recommended seeding density and

left overnight in a 37 ºC, 5% CO_2 incubator. The following day, cells were infected with CtrL2

with a MOI of 5. At 2 h post-infection, the cells were transfected with either Vinculin-venus

(Grashoff *et al.*, 2010) a gift from Martin Schwartz, (Addgene, 27300), paxillin-pEGFP (Laukaitis

et al., 2001) a gift from Rick Horwitz, (Addgene, 15233), or FAK-GFP (Gu *et al.*, 1999; Lane *et*

al., 2008) a gift from Kenneth Yamada, (Addgene, 50515) using Lipofectamine 3000

transfection reagent (Thermofisher Scientific, L3000008), following the manufacture instructions.

After 20 to 22 h, time lapsed images of transfected cells were obtained using a Leica SD6000

AF in TIRF mode, in Washington State University IPN advance image equipment. Images of the

GFP-tagged proteins were collected every minute for 90 min. The time lapse images were

uploaded to the Focal adhesion Analysis server (Berginski and Gomez, 2013).

Cell Motility Assay. Mouse embryonic fibroblasts (MEFs) were seeded on ibidi μ-slide live cell

imaging chambers (ibidi, 80426). Cells were infected with *Chlamydia trachomatis* serovar L2 at

an MOI of 10 by rocking at 4°C for one hour. Infected cells were imaged starting at 20 hours

post-infection. A mock-infected control was made following the same protocol but without

Chlamydia infectious particles. Cells were transfected with N1-mTurquoise2 empty vector

control or TarP$^{829-1006}$-mTurquoise2 (LDVBD) using electroporation, seeded into an ibidi μ-slide,

and imaged starting at 22 hours post electroporation. Cells were imaged in live cell imaging

solution (Thermofisher scientific, A14291DJ) supplemented with 5% fetal bovine serum within a

37°C, 5% CO_2 controlled environment. Time-lapse DIC images were obtained using a Leica

SD6000 AF microscope every ten minutes for ten hours. To minimize the risk of phototoxicity,

we restricted image acquisition of the fluorescent mTurquoise2 channel for our transfected cells

to the last frame alone. We utilized a similar individual cell tracking data analysis approach as

described in (Pijuan *et al.*, 2019). Cell motility was tracked using the manual tracking function in

ImageJ. Each individual cell was tracked using the position of the nucleus over time. We

maximized the time of analysis for each experiment based on the number of cells that remained

within a trackable field of view over the imaging span. The coordinate data from the manual

tracking function was uploaded to ibidi's chemotaxis and migration tool. Measurements were

taken from spatially calibrated images with a (pixel/μm) scale contained within the meta data.

The x/y calibration was set to 0.800001 based on the microscope's settings contained within the

file's meta data. The statistics function was used to determine the velocity and euclidean

(straight-line) distance traveled for each cell.

***De novo* protein inhibition.** Cos7 cells were cultured as previously described. Prior to

infection, cells and EB particles were treated with 60 μg/ml of chloramphenicol (Sigma C0378)

for 30 min. Cells and EBs were kept in chloramphenicol supplemented DMEM until fixation.

Cells were fixed at 8 or 20 h post-infection and were immunostained as described above.

Western blot. Cos7 cells were plated in 6 well plates with duplicate wells and incubated at 37°C

and 5% CO2, until 80% confluency. Cells were infected, as previously described, with CtrL2 for

0 min, 8 or 24 hrs with an MOI of 200. Proteins were harvested using ice cold RIPA buffer

(Millipore 20-188) supplemented with phosphatase (Sigma 4906845001) and protease inhibitors

(Sigma 5892970001). Cells were scraped and incubated for 30 min on ice. The lysates were

centrifuged at 13,000 x g for 20 min at 4°C. Supernatants were diluted in Laemmli buffer (Biorad

161-0747) and kept at -20°C before analysis. Samples were resolved on 10% acrylamide SDS-

PAGE. Proteins were transferred to a nitrocellulose membrane (Bio-rad 1620115).

Immunoblotting was performed by blocking membranes with 5% BSA in TBS-T overnight at 4°C

and incubation using antibodies against TarP (a generous gift from Dr. Raphael Valdivia, Duke

University) and β-tubulin HRP conjugated (Abcam ab21058). The secondary antibody used was

anti-mouse HRP conjugated (DAKO P0161). Immobilin chemiluminescence kit (Millipore,

WBKLS0500) was used to develop the blot.

Cloning and transfection of TarP contructs. A summary of the primers used in this study is

provided in Table S1. Initially TarP Full-length, TarP LDVBD, TarP LD, and TarP VBD were

PCR amplified from CtrL2 genomic DNA using the primers combination 1-2, 5-2, 5-6 and 4-2,

respectively. A *Bam*H1 (reverse primer) and *Kpn*I (forward primer) restriction sites were used for

fusion with the N1-mTurquoise2 plasmid. The TarP ΔPRD was obtained using the 7-8 primer

pair for PCR amplification from TarP Full-length-mTurquoise2 fusion plasmid. The primers were

created to amplify the whole TarP Full-length-mTurquoise2 except the nucleotides that

constitute the proline rich domain (PRD) 625-650. The resulting PCR product was recombined

using in-Fusion HD cloning plus CE (Clontech, 638916) to create a functional circular plasmid.

TarP ΔLDVBD was PCR amplified from the TarP ΔPRD-mTurquoise2 plasmid using the primers

pair 1-3. The same restriction enzymes were used to clone these fragments into N1-

mTurquoise2. Transformations using restriction enzymes recombination were made into

chemically competent Top10 (invitrogen) *E. coli*, and vectors sequence was verified using

sequencing (Eurofins) The construct pFH-TarP ΔPRD and pFH-TarP ΔLDVBD used for super-

resolution experiments was PCR amplified from TarP ΔPRD-mTurquoise2 using primers

combination 9-10 and 9-17, respectively. To use homology cloning the vector backbone 1436

pcDNA3-Flag-HA, kindly provided by William Sellers (Addgene 10792), was linearized by PCR

using the primers pair 11-12 and 11-18, creating homology overhang regions to the TarP ΔPRD

and TarP ΔLDVBD, respectively. TarP LDVBD was amplified from CtrL2 genomic DNA using

the primer pairs 13-14. To use homology cloning the vector backbone 1436 pcDNA3-Flag-HA

was linearized by PCR using the primer pair 15-16. Fragments and vector backbone were

recombined using in-Fusion HD cloning plus CE (Clontech, 638916) to create a functional

circular plasmid. Transformations using homology recombination were made into chemically

competent Stellar (Clontech) *E. coli*, and vectors sequence was verified using sequencing

(Eurofins). The pcDNA3-Flag-Apex-Nes was a gift from Alice Ting (Addgene, 49386). The N1-

mTurquoise2 (Addgene, 54843). Transfections were done as described above. For iPALM

experiments 1µg of DNA and 2µl of sheared salmon sperm DNA were mixed together in 15µl of

Opti-MEM (Thermofisher scientific, 31985062), and kept on ice for 15 min. $1x10^6$ Cos7 cells

were resuspended in 200µl of cold Opti-MEM, mixed with the DNA solution and kept on ice for

30 seconds. Cells and DNA suspension were transferred to a 4mm gap cuvette (BioRad,

1652088) and electroporated using BioRad Gene Pulser XCell using the following settings:

190V; 950uF; infinity. After electroporation 1.5ml of warm growth media was added. 400µl of cell

solution was added to a 6 well plate well containing 1.5ml of warm growth media and the gold

fiducial coverslip. Cells were incubated 37°C, 4% CO_2 for 4 h to adhere to the gold fiducial

coverslip. Cells were washed to remove dead cells debris and further incubated for 20 h.

iPALM imaging and analysis. The principle of instrumentation for iPALM imaging and analysis

were performed as previously described (Shtengel *et al.*, 2009; Kanchanawong *et al.*, 2010)

with the following modifications. After 24°C of transfection cells plated in gold fiducial coverslip

were fixed with 0.8% PFA and 0.1% glutaraldehyde (Sigma G7526-10ML) solution (in PBS) for

10 min. After fixation cells were washed 3 times with PBS and quenched using 1% NaBH4

(Sigma, 452882-25G) solution (in PBS) for 7 min. Cells were then washed again 3 times with

PBS. After washing cells were immunostained (when necessary) and/or processed for iPALM

imaging as previously described (Kanchanawong *et al.*, 2010). The vertical coordinates relative

to the golden fiducial markers are indicating by a color scale from red (0 nm) to purple (250 nm).

Trypsin assay. Cells were plated in 24 well plates and incubated at 37°C and 5% CO2, until

80-90% confluency. Afterwards, cells were infected with CtrL2 with a multiplicity of infection of 5

for 20 h. Cells were then treated with 0.01% trypsin diluted in serum-free DMEM media at 37°C,

for 0, 10, 20, 30, or 35 min. Cells were fixed with 4% PFA, carefully washed with PBS and

stained with DAPI to count the number of remaining cells as well as to visualize *Chlamydia*

inclusions. Images were taken using Nikon eclipse TE2000-U. Detachment assays were

performed as above, and transfection procedures were as described.

Acknowledgments

We would like to thank Drs. Martin Schwartz, Rick Horwitz, Kenneth Yamada, William Sellers,

Alice Ting, and Michael Davidson for various constructs obtained through Addgene; Dr. Raphael

Valdivia (Duke University Medical Center) for the generous gift of the TarP antibody; Dr. Lisa

Rucks (University of Nebraska Medical Center) for the *C. trachomatis* serovar D strain; the

services provided by the WSU Integrated Physiology and Neuroscience imaging facilities; and

Satya Khuon (AIC Janelia) for technical help and advice. This project was supported by grants

from the National Institute for Food and Agriculture (#1010265), National Institutes of Health

(AI065545), and start-up funds from the WSU College of Veterinary Medicine to RAC. ATP and

ATN were recipients of the Fundação para a Ciência e Tecnologia, SFRH/BD/76741/2011 and

SFRH/BD/86670/2012, respectively. TRT is a Medical Research Council studentship awardee.

KNM was supported by the NIH Protein Biotechnology Training Grant T32 GM008336 awarded

to WSU. AJB was supported by the Infectious Diseases and Microbial Immunology Training

Grant T32 AI007025 awarded to WSU. iPALM data used in this publication was produced in

collaboration with the Advanced Imaging Center, a facility jointly supported by the Gordon and

Betty Moore Foundation and Howard Hughes Medical Institute at the Janelia Research

Campus.

Author Contributions

Conceptualization, RAC; Methodology, ATP, JA, and RAC; Software, JA; Formal Analysis, ATP; Investigation, ATP, ATN, KNM, JA, AJB, and TRT; Writing – Original Draft, ATP, and RAC; Writing – Review & Editing, RAC, KNM and AJB; Funding Acquisition RAC; Resources, JA, TLC and RAC; Supervision, TLC and RAC.

Conflict of Interest

The authors have declared that no competing or financial interests exist.

References

AbdelRahman, Y. M. and Belland, R. J. (2005) 'The chlamydial developmental cycle', *FEMS*

Microbiology Reviews, 29(5), pp. 949–959. doi: 10.1016/j.femsre.2005.03.002.

Anderson, D. J., Marathe, J. and Pudney, J. (2014) 'The Structure of the Human Vaginal

Stratum Corneum and its Role in Immune Defense', *American Journal of Reproductive*

Immunology, 71(6), pp. 618–623. doi: 10.1111/aji.12230.

Berginski, M. E. and Gomez, S. M. (2013) 'The Focal Adhesion Analysis Server: a web tool for

analyzing focal adhesion dynamics.', *F1000Research*, 2, p. 68. doi: 10.12688/f1000research.2-

68.v1.

Betzig, E., Patterson, G. H., Sougrat, R., Lindwasser, O. W., Olenych, S., Bonifacino, J. S.,

Davidson, M. W., Lippincott-Schwartz, J. and Hess, H. F. (2006) 'Imaging Intracellular

Fluorescent Proteins at Nanometer Resolution', *Science*, 313(5793), pp. 1642–1645. doi:

10.1126/science.1127344.

Borradori, L. and Sonnenberg, A. (1999) 'Structure and function of hemidesmosomes: more

than simple adhesion complexes.', *The Journal of investigative dermatology*, 112(4), pp. 411–8.

doi: 10.1046/j.1523-1747.1999.00546.x.

Braun, C., Alcázar-Román, A. R., Laska, A., Mölleken, K., Fleig, U. and Hegemann, J. H.

(2019) 'CPn0572, the *C. pneumoniae* ortholog of TarP, reorganizes the actin cytoskeleton via a

newly identified F-actin binding domain and recruitment of vinculin.', *PloS one*. Edited by D. M.

Ojcius, 14(1), p. e0210403. doi: 10.1371/journal.pone.0210403.

Carisey, A., Tsang, R., Greiner, A. M., Nijenhuis, N., Heath, N., Nazgiewicz, A., Kemkemer,

R., Derby, B., Spatz, J. and Ballestrem, C. (2013) 'Vinculin regulates the recruitment and

release of core focal adhesion proteins in a force-dependent manner.', *Current biology : CB*.

Elsevier, 23(4), pp. 271–81. doi: 10.1016/j.cub.2013.01.009.

Carrasco, S. E., Hu, S., Imai, D. M., Kumar, R., Sandusky, G. E., Yang, X. F. and Derbigny,

W. A. (2018) 'Toll-like receptor 3 (TLR3) promotes the resolution of *Chlamydia muridarum*

genital tract infection in congenic C57BL/6N mice.', *PloS one*. Public Library of Science, 13(4),

p. e0195165. doi: 10.1371/journal.pone.0195165.

Chrzanowska-Wodnicka, M. and Burridge, K. (1996) 'Rho-stimulated contractility drives the

formation of stress fibers and focal adhesions', *The Journal of Cell Biology*, 133(6), pp. 1403–

1415. doi: 10.1083/jcb.133.6.1403.

Clifton, D. R., Fields, K. A., Grieshaber, S. S., Dooley, C. A., Fischer, E. R., Mead, D. J.,

Carabeo, R. A. and Hackstadt, T. (2004) 'A chlamydial type III translocated protein is tyrosine-

phosphorylated at the site of entry and associated with recruitment of actin.', *Proceedings of the*

National Academy of Sciences of the United States of America, 101(27), pp. 10166–71. doi:

10.1073/pnas.0402829101.

Clifton, D. R., Dooley, C. A., Grieshaber, S. S., Carabeo, R. A., Fields, K. A. and Hackstadt,

T. (2005) 'Tyrosine Phosphorylation of the Chlamydial Effector Protein Tarp Is Species Specific

and Not Required for Recruitment of Actin', *Infection and Immunity*, 73(7), pp. 3860–3868. doi:

10.1128/IAI.73.7.3860-3868.2005.

Crosson, C. E., Klyce, S. D. and Beuerman, R. W. (1986) 'Epithelial wound closure in the

rabbit cornea. A biphasic process.', *Investigative ophthalmology & visual science*, 27(4), pp.

464–73. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3957565 (Accessed: 27 November

2019).

Derbigny, W. A., Shobe, L. R., Kamran, J. C., Toomey, K. S. and Ofner, S. (2012)

'Identifying a role for Toll-like receptor 3 in the innate immune response to *Chlamydia*

muridarum infection in murine oviduct epithelial cells', *Infection and Immunity*. Edited by R. P.

Morrison, 80(1), pp. 245–65. doi: 10.1128/IAI.05549-11.

Derbigny, W. A., Kerr, M. S. and Johnson, R. M. (2005) 'Pattern recognition molecules

activated by *Chlamydia muridarum* infection of cloned murine oviduct epithelial cell lines.',

Journal of immunology (Baltimore, Md. : 1950), 175(9), pp. 6065–75. doi:

10.4049/jimmunol.175.9.6065.

Dumbauld, D. W., Michael, K. E., Hanks, S. K. and García, A. J. (2010) 'Focal adhesion

kinase-dependent regulation of adhesive forces involves vinculin recruitment to focal

adhesions', *Biology of the Cell*, 102(4), pp. 203–213. doi: 10.1042/BC20090104.

Frisch, S. M., Vuori, K., Ruoslahti, E. and Chan-Hui, P. Y. (1996) 'Control of adhesion-

dependent cell survival by focal adhesion kinase', *The Journal of Cell Biology*, 134(3), pp. 793–

799. doi: 10.1083/jcb.134.3.793.

Frisch, S. M. and Francis, H. (1994) 'Disruption of epithelial cell-matrix interactions induces

apoptosis', *The Journal of Cell Biology*, 124(4), pp. 619–626. doi: 10.1083/jcb.124.4.619.

Geiger, B., Bershadsky, A., Pankov, R. and Yamada, K. M. (2001) 'Transmembrane crosstalk

between the extracellular matrix--cytoskeleton crosstalk.', *Nature reviews. Molecular cell*

biology, 2(11), pp. 793–805. doi: 10.1038/35099066.

Grashoff, C., Hoffman, B. D., Brenner, M. D., Zhou, R., Parsons, M., Yang, M. T., McLean,

M. A., Sligar, S. G., Chen, C. S., Ha, T. and Schwartz, M. A. (2010) 'Measuring mechanical

tension across vinculin reveals regulation of focal adhesion dynamics.', *Nature*, 466(7303), pp.

263–6. doi: 10.1038/nature09198.

Gu, J., Tamura, M., Pankov, R., Danen, E. H., Takino, T., Matsumoto, K. and Yamada, K. M.

(1999) 'Shc and FAK differentially regulate cell motility and directionality modulated by PTEN.',

The Journal of cell biology. Rockefeller University Press, 146(2), pp. 389–403. doi:

10.1083/jcb.146.2.389.

Gudipaty, S. A., Conner, C. M., Rosenblatt, J. and Montell, D. J. (2018) 'Unconventional

Ways to Live and Die: Cell Death and Survival in Development, Homeostasis, and Disease.',

Annual review of cell and developmental biology, 34(1), pp. 311–332. doi: 10.1146/annurev-

cellbio-100616-060748.

Gudipaty, S. A. and Rosenblatt, J. (2017) 'Epithelial cell extrusion: Pathways and

pathologies', *Seminars in Cell & Developmental Biology*, 67, pp. 132–140. doi:

10.1016/j.semcdb.2016.05.010.

Heymann, J., Rejman Lipinski, A., Bauer, B., Meyer, T. F. and Heuer, D. (2013) '*Chlamydia*

trachomatis infection prevents front-rear polarity of migrating HeLa cells', *Cellular Microbiology*,

15(7), pp. 1059–1069. doi: 10.1111/cmi.12114.

Jewett, T. J., Fischer, E. R., Mead, D. J. and Hackstadt, T. (2006) 'Chlamydial TARP is a

bacterial nucleator of actin', *Proceedings of the National Academy of Sciences*, 103(42), pp.

15599–15604. doi: 10.1073/pnas.0603044103.

Jewett, T. J., Miller, N. J., Dooley, C. A. and Hackstadt, T. (2010) 'The conserved Tarp actin

binding domain is important for chlamydial invasion.', *PLoS pathogens*, 6(7), p. e1000997. doi:

10.1371/journal.ppat.1000997.

Jiwani, S., Alvarado, S., Ohr, R. J., Romero, A., Nguyen, B. and Jewett, T. J. (2013)

'*Chlamydia trachomatis* Tarp Harbors Distinct G and F Actin Binding Domains That Bundle Actin

Filaments', *Journal of Bacteriology*, 195(4), pp. 708–716. doi: 10.1128/JB.01768-12.

Kanchanawong, P., Shtengel, G., Pasapera, A. M., Ramko, E. B., Davidson, M. W., Hess,

H. F. and Waterman, C. M. (2010) 'Nanoscale architecture of integrin-based cell adhesions',

Nature, 468(7323), pp. 580–584. doi: 10.1038/nature09621.

Kim, M., Ogawa, M., Fujita, Y., Yoshikawa, Y., Nagai, T., Koyama, T., Nagai, S., Lange, A.,

Fässler, R. and Sasakawa, C. (2009) 'Bacteria hijack integrin-linked kinase to stabilize focal

adhesions and block cell detachment.', *Nature*, 459(7246), pp. 578–82. doi:

10.1038/nature07952.

Kim, M., Ogawa, M., Mimuro, H. and Sasakawa, C. (2010) 'Reinforcement of epithelial cell

adhesion to basement membrane by a bacterial pathogen as a new infectious stratagem',

Virulence, 1(April 2010), pp. 52–55. doi: 10.1038/nature07952.he.

Kumar, R., Gong, H., Liu, L., Ramos-Solis, N., Seye, C. I. and Derbigny, W. A. (2019) 'TLR3

deficiency exacerbates the loss of epithelial barrier function during genital tract *Chlamydia*

muridarum infection.', *PloS one*. Edited by G. Zhong, 14(1), p. e0207422. doi:

10.1371/journal.pone.0207422.

Kumar, Y. and Valdivia, R. H. (2008) 'Reorganization of the host cytoskeleton by the

intracellular pathogen *Chlamydia trachomatis*', *Commun Integr Biol*, 1(2), pp. 175–177. doi:

10.1016/j.chom.2008.05.018.www.landesbioscience.com.

Kuwabara, T., Perkins, D. G. and Cogan, D. G. (1976) 'Sliding of the epithelium in

experimental corneal wounds.', *Investigative ophthalmology*, 15(1), pp. 4–14. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/1245378 (Accessed: 27 November 2019).

Lacy, H. M., Bowlin, A. K., Hennings, L., Scurlock, A. M., Nagarajan, U. M. and Rank, R. G.

(2011) 'Essential role for neutrophils in pathogenesis and adaptive immunity in *Chlamydia*

caviae ocular infections.', *Infection and immunity*. Edited by J. N. Weiser, 79(5), pp. 1889–97.

doi: 10.1128/IAI.01257-10.

Lane, B. J., Mutchler, C., Al Khodor, S., Grieshaber, S. S. and Carabeo, R. A. (2008)

'Chlamydial entry involves TARP binding of guanine nucleotide exchange factors.', *PLoS*

pathogens, 4(3), p. e1000014. Available at: http://www.ncbi.nlm.nih.gov/pubmed/18383626

(Accessed: 22 November 2016).

Laukaitis, C. M., Webb, D. J., Donais, K. and Horwitz, A. F. (2001) 'Differential dynamics of

alpha 5 integrin, paxillin, and alpha-actinin during formation and disassembly of adhesions in

migrating cells.', *The Journal of cell biology*, 153(7), pp. 1427–40. doi: 10.1083/jcb.153.7.1427.

Liu, Z., van Grunsven, L. A., Van Rossen, E., Schroyen, B., Timmermans, J.-P., Geerts, A.

and Reynaert, H. (2010) 'Blebbistatin inhibits contraction and accelerates migration in mouse

hepatic stellate cells.', *British journal of pharmacology*, 159(2), pp. 304–15. doi: 10.1111/j.1476-

5381.2009.00477.x.

Lutter, E. I., Bonner, C., Holland, M. J., Suchland, R. J., Stamm, W. E., Jewett, T. J.,

McClarty, G. and Hackstadt, T. (2010) 'Phylogenetic Analysis of *Chlamydia trachomatis* Tarp

and Correlation with Clinical Phenotype', *Infection and Immunity*, 78(9), pp. 3678–3688. doi:

10.1128/IAI.00515-10.

Marg, S., Winkler, U., Sestu, M., Himmel, M., Schönherr, M., Bär, J., Mann, A., Moser, M.,

Mierke, C. T., Rottner, K., Blessing, M., Hirrlinger, J. and Ziegler, W. H. (2010) 'The vinculin-

DeltaIn20/21 mouse: characteristics of a constitutive, actin-binding deficient splice variant of

vinculin.', *PloS one*. Edited by D. P. Martin, 5(7), p. e11530. doi: 10.1371/journal.pone.0011530.

Mehlitz, A., Banhart, S., Mäurer, A. P., Kaushansky, A., Gordus, A. G., Zielecki, J.,

Macbeath, G. and Meyer, T. F. (2010) 'Tarp regulates early Chlamydia-induced host cell

survival through interactions with the human adaptor protein SHC1.', *The Journal of cell biology*,

190(1), pp. 143–57. doi: 10.1083/jcb.200909095.

Morita-Ishihara, T., Miura, M., Iyoda, S., Izumiya, H., Watanabe, H., Ohnishi, M. and

Terajima, J. (2013) 'EspO1-2 regulates EspM2-mediated RhoA activity to stabilize formation of

focal adhesions in enterohemorrhagic Escherichia coli-infected host cells.', *PloS one*. Edited by

F. Börnke, 8(2), p. e55960. doi: 10.1371/journal.pone.0055960.

Mysorekar, I. U., Mulvey, M. A., Hultgren, S. J. and Gordon, J. I. (2002) 'Molecular regulation

of urothelial renewal and host defenses during infection with uropathogenic Escherichia coli.',

The Journal of biological chemistry, 277(9), pp. 7412–9. doi: 10.1074/jbc.M110560200.

Nagano, M., Hoshino, D., Koshikawa, N., Akizawa, T. and Seiki, M. (2012) 'Turnover of

Focal Adhesions and Cancer Cell Migration', *International Journal of Cell Biology*, 2012, pp. 1–

10. doi: 10.1155/2012/310616.

O'Connell, C. M., Ionova, I. A., Quayle, A. J., Visintin, A. and Ingalls, R. R. (2006)

'Localization of TLR2 and MyD88 to *Chlamydia trachomatis* inclusions. Evidence for signaling

by intracellular TLR2 during infection with an obligate intracellular pathogen.', *The Journal of*

biological chemistry, 281(3), pp. 1652–9. doi: 10.1074/jbc.M510182200.

Pan, Q., Zhang, Q., Chu, J., Pais, R., Liu, S., He, C. and Eko, F. O. (2017) '*Chlamydia*

abortus Pmp18.1 Induces IL-1β Secretion by TLR4 Activation through the MyD88, NF-κB, and

Caspase-1 Signaling Pathways.', *Frontiers in cellular and infection microbiology*, 7, p. 514. doi:

10.3389/fcimb.2017.00514.

Pasapera, A. M., Schneider, I. C., Rericha, E., Schlaepfer, D. D. and Waterman, C. M.

(2010) 'Myosin II activity regulates vinculin recruitment to focal adhesions through FAK-

mediated paxillin phosphorylation.', *The Journal of cell biology*, 188(6), pp. 877–90. doi:

10.1083/jcb.200906012.

Pijuan, J., Barceló, C., Moreno, D. F., Maiques, O., Sisó, P., Marti, R. M., Macià, A. and

Panosa, A. (2019) 'In vitro Cell Migration, Invasion, and Adhesion Assays: From Cell Imaging to

Data Analysis.', *Frontiers in cell and developmental biology*, 7, p. 107. doi:

10.3389/fcell.2019.00107.

Ramsey, K. H., Sigar, I. M., Schripsema, J. H., Shaba, N. and Cohoon, K. P. (2005)

'Expression of matrix metalloproteinases subsequent to urogenital *Chlamydia muridarum*

infection of mice.', *Infection and immunity*, 73(10), pp. 6962–73. doi: 10.1128/IAI.73.10.6962-

6973.2005.

Rosenblatt, J., Raff, M. C. and Cramer, L. P. (2001) 'An epithelial cell destined for apoptosis

signals its neighbors to extrude it by an actin- and myosin-dependent mechanism', *Current*

Biology, 11(23), pp. 1847–1857. doi: 10.1016/S0960-9822(01)00587-5.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J.,

Hartenstein, V., Eliceiri, K., Tomancak, P. and Cardona, A. (2012) 'Fiji: an open-source

platform for biological-image analysis.', *Nature methods*, 9(7), pp. 676–82. doi:

10.1038/nmeth.2019.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012) 'NIH Image to ImageJ: 25 years

of image analysis', *Nature Methods*, 9(7), pp. 671–675. doi: 10.1038/nmeth.2089.

Shames, S. R., Deng, W., Guttman, J. A., de Hoog, C. L., Li, Y., Hardwidge, P. R., Sham, H.

P., Vallance, B. A., Foster, L. J. and Finlay, B. B. (2010) 'The pathogenic E. coli type III

effector EspZ interacts with host CD98 and facilitates host cell prosurvival signalling.', *Cellular*

microbiology, 12(9), pp. 1322–39. doi: 10.1111/j.1462-5822.2010.01470.x.

Shaw, J. L. V, Wills, G. S., Lee, K.-F., Horner, P. J., McClure, M. O., Abrahams, V. M.,

Wheelhouse, N., Jabbour, H. N., Critchley, H. O. D., Entrican, G. and Horne, A. W. (2011)

'*Chlamydia trachomatis* infection increases fallopian tube PROKR2 via TLR2 and NFκB

activation resulting in a microenvironment predisposed to ectopic pregnancy.', *The American*

journal of pathology, 178(1), pp. 253–60. doi: 10.1016/j.ajpath.2010.11.019.

Shtengel, G., Galbraith, J. A., Galbraith, C. G., Lippincott-Schwartz, J., Gillette, J. M.,

Manley, S., Sougrat, R., Waterman, C. M., Kanchanawong, P., Davidson, M. W., Fetter, R.

D. and Hess, H. F. (2009) 'Interferometric fluorescent super-resolution microscopy resolves 3D

cellular ultrastructure.', *Proc Natl Acad Sci U S A*, 106(9), pp. 3125–30. doi:

10.1073/pnas.0813131106.

Smilenov, L. B., Mikhailov, A., Pelham, R. J., Marcantonio, E. E. and Gundersen, G. G.

(1999) 'Focal Adhesion Motility Revealed in Stationary Fibroblasts', *Science*, 286(5442), pp.

1172–1174. doi: 10.1126/science.286.5442.1172.

Straight, A. F., Cheung, A., Limouze, J., Chen, I., Westwood, N. J., Sellers, J. R. and

Mitchison, T. J. (2003) 'Dissecting temporal and spatial control of cytokinesis with a myosin II

Inhibitor.', *Science (New York, N.Y.)*, 299(5613), pp. 1743–7. doi: 10.1126/science.1081412.

Thwaites, T., Nogueira, A. T., Campeotto, I., Silva, A. P., Grieshaber, S. S. and Carabeo, R.

A. (2014) 'The *Chlamydia* Effector TarP Mimics the Mammalian Leucine-Aspartic Acid Motif of

Paxillin to Subvert the Focal Adhesion Kinase during Invasion', *Journal of Biological Chemistry*,

289(44), pp. 30426–30442. doi: 10.1074/jbc.M114.604876.

Thwaites, T. R., Pedrosa, A. T., Peacock, T. P. and Carabeo, R. A. (2015) 'Vinculin Interacts

with the *Chlamydia* Effector TarP Via a Tripartite Vinculin Binding Domain to Mediate Actin

Recruitment and Assembly at the Plasma Membrane', *Frontiers in Cellular and Infection*

Microbiology. Frontiers, 5, p. 88. doi: 10.3389/fcimb.2015.00088.

Vossenkämper, A., Macdonald, T. T. and Marchès, O. (2011) 'Always one step ahead: How

pathogenic bacteria use the type III secretion system to manipulate the intestinal mucosal

immune system.', *Journal of inflammation (London, England)*, 8(1), p. 11. doi: 10.1186/1476-

9255-8-11.

Wang, H. H., Tanaka, H., Qin, X., Zhao, T., Ye, L.-H., Okagaki, T., Katayama, T., Nakamura,

A., Ishikawa, R., Thatcher, S. E., Wright, G. L. and Kohama, K. (2008) 'Blebbistatin inhibits

the chemotaxis of vascular smooth muscle cells by disrupting the myosin II-actin interaction.',

American journal of physiology. Heart and circulatory physiology, 294(5), pp. H2060-8. doi:

10.1152/ajpheart.00970.2007.

Zaidel-Bar, R., Itzkovitz, S., Ma'ayan, A., Iyengar, R. and Geiger, B. (2007) 'Functional atlas

of the integrin adhesome.', *Nature cell biology*, 9(8), pp. 858–67. doi: 10.1038/ncb0807-858.

Figure legends

Figure 1. *C. trachomatis*-infected Cos7 cells exhibit increased focal adhesion numbers.

(A) Cos7 cells infected with the indicated chlamydial strain/species or mock-infected were

monitored at 24 hpi, and focal adhesions visualized by immunostaining for paxillin.

Figure 2. Infection-dependent increase in focal adhesion numbers requires *de novo*

chlamydial protein synthesis. (A) Cos7 cells infected with *C. trachomatis* serovar L2 were

mock- or chloramphenicol-treated at the start of infection for either 8 or 20 h. Focal adhesions

were visualized by immunostaining for paxillin (green). Scale bar length 10µm. (B) Focal

adhesions were counted using the particle tracker plug-in in NIH ImageJ. Analysis revealed a

lack of effect of the 8-h Cm treatment on FA numbers, while the 24-h treatment yielded a

statistically significant decreasein FA numbers. Data are represented as box-and-whisker plots.

Whiskers represent the lowest and highest data point still within 1.5 times the interquartile

range. The light blue asterisks indicate significant difference relative to the mock-infected control

(Wilcoxon rank sum test * = $p < 0.05$). The black cross shows the average for each experimental

sample.

Figure 3. Focal adhesions of *Chlamydia* infected cells are resistant to blebbistatin. Cos7

cells were mock-infected or infected with CtrL2 for 24 h or 8 h. Cells were fixed and stained for

the focal adhesion marker paxillin (green), F-actin (red) and human convalescent serum for *C.*

trachomatis (white). Cells were also mock- or pre-treated with chloramphenicol (Cm) followed by

infection of live EBs. Cm treatment was maintained for the duration of the experiment.

Blebbistatin 10 μM was introduced during the last hour of infection. Cells without blebbistatin

treatment showed clear F-actin stress fibres and paxillin-labeled focal adhesions. While both

structures were lost in mock-infected cells, infected cells retained the focal adhesions. This

characteristic was sensitive to Cm inhibition of *de novo* chlamydial protein synthesis. Scale bar

= 10µm.

Figure 4. The type III effector TarP localizes to focal adhesions during *Chlamydia*

infection. (A) CtrL2-infected MEF cells were immunostained for a rabbit polyclonal antibody to

TarP and a mouse monoclonal antibody to talin. Inclusions were visualized by staining with

DAPI. TarP localized to talin-positive FAs (arrowheads) in infected, but not in mock-infected

controls. Scale bar = 10µm.

Figure 5. The focal adhesion localization of TarP requires its LDVBD domain and the host

protein vinculin, but not FAK. (A) Representation of *C. trachomatis* effector protein TarP and

its known domains fused to mTurquoise2 fluorescent protein. (B) Wild type or vinculin-knockout

MEFs were transfected with different mTurquoise2-tagged TarP constructs (green) and imaged

by confocal microscopy to evaluate colocalization with paxillin (red) at focal adhesions.

Phalloidin was used to stain F-actin (blue). Cells were transfected for 20 h, at which time the

cells were fixed and processed for immunofluorescence staining for paxillin. (C) In a parallel

experiment, wild type and FAK1-deficient MEFs were transfected to express Flag-HA-LDVBD

and stained for flag (green). Colocalization with paxillin (red) was assessed by confocal

microscopy. (D) Ectopically expressed LDVBD localizes to β1-integrin and paxillin-positive focal

adhesions. Scale bar = 10µm.

Figure 6. The ectopic expression of TarP LDVBD is sufficient to increase focal adhesion

numbers. (A) Cos7 cells expressing different deletion derivatives of TarP or vector only were

processed for immunofluorescence with anti-paxillin antibody to visualize focal adhesions.

Representative images are shown. Scale bar = 10 µm. (B) Focal adhesion numbers were

counted using the particle counting plug-in in NIH ImageJ. Data are focal adhesion number per

cell and illustrated as box-whisker plot. Whiskers represent the lowest and highest data point

still within 1.5 times the interquartile range. For statistical analyses the Wilcoxon Rank sum test

was used to determine significance when compared to vector-only control (* = p<0.05).

Figure 7. The LDVBD domain of TarP and the host protein vinculin are required for focal

adhesion resistance to blebbistatin treatment. (A) Wild type or vinculin-knockout MEFs were

infected with C. trachomatis serovar L2. Cells at 24 hpi were mock-treated or treated for 60 min

with 10 µM of blebbistatin. The cells were then processed for immunofluorescence staining for

paxillin (green) and actin (red). Retention or loss of focal adhesions were monitored. Focal

adhesions were only resistant to blebbistatin-induced disassembly if the cell was infected and

expressing vinculin. (B) In a parallel experiment, wild type or vinculin-knockout MEFs were

transfected for 20 h with the empty vector or LDVBD-mTurquoise2 fusion protein. During the

last hour, cells were either mock- or blebbistatin-treated. Cells were processed to visualize

paxillin (red), LDVBD (green), and actin (blue; shown in composite images). LDVBD was

sufficient to confer resistance to blebbistatin-induced disassembly to focal adhesions.

Resistance also required vinculin. Scale bar = 10 µm.

Figure 8. TarP-targeted focal adhesions display altered nanoscale architecture. (A) Cos7

cells pre-transfected with paxillin-tdEos, FAK-TdEos, or Vinculin-TdEos, and grownon gold

fiducial coverslips were mock- or C. trachomatis-infected for 24 h. The cells were fixed and

processed for iPALM imaging. Representative images are shown. For each sample, multiple

panels are provided. The top panel shows the top view of area around the focal adhesion of

interest (white border). The middle panel displays a top view of the focal adhesion indicated by

the white border. The bottom panel shows the side view and corresponding z histograms. Note

the significant shifts in paxillin and FAK localization, but not vinculin. (B) Cos7 cells were co-

transfected with paxillin-tdEos and either TarP ΔPRD or LDVBD only by electroporation. The

cells were seeded on gold fiducial coverslips, and processed for iPALM at 20 h post-

transfection. Description of each panel is as above in (A). Note the significant shift in the

location of paxillin within the TarP-positive focal adhesions. The various colors indicate the

distance (z-coordinates) from the gold fiducial marker, (e.g. z = 0 nm; red). Red scale bar =

1µm. White scale bar = 200 nm.

Figure 9. The LDVBD domain of TarP is sufficient to inhibit cell migration. MEFs that were

mock-infected, Chlamydia-infected, vector-only-transfected, or LDVBD-transfected were seeded

within ibidi µ-slide live cell imaging chambers. Time-lapse imaging was performed every 10

minutes for 10 h to evaluate cell motility. (A) For analysis of the infection experiments, a 5-h

imaging window common to both mock- and *Chlamydia*-infected samples was chosen that

maximized the number of cells that remained within the field of view. Cells were tracked using

the manual tracking function in NIH ImageJ, and the cell trajectory was traced and plotted with

the starting points assigned to the origin. (B) Analysis of the transfection experiment was in a

common 10-h imaging window. Data were acquired and plotted as in (A). (C) Velocity and

Euclidean distance traveled were calculated for each cell from each experimental group. Values

were plotted as dot-plots with mean $\pm$ S.D. indicated by the bars. Statistical significance was

calculated using the Wilcoxon Rank Sum test. * indicates $p < 0.05$.

Mock-infected

Ctr L2

Ctr D (urogenital)

Ctr B (ocular)

C. muridarum MoPn

C. caviae GPIC

A

FIGURE 3

FIGURE 5

A

B

Wild type MEFs   vcl-/- MEFs

TarP Paxillin Merge   TarP Paxillin Merge

TarP ΔPRD

TarP ΔLDVBD

LDVBD only

C

Wild type MEFs   ptk2-/- MEFs

TarP Paxillin Merge   TarP Paxillin Merge

LDVBD only

D

TarP   β1-integrin   Paxillin   Merge

LDVBD only

FIGURE 6

FIGURE 7

FIGURE 8

Supplemental Table 1 - Primers used in this study

Primer number	Primer name	Primer sequence 5' to 3'
1	Fwd-TarP	GCATGGTACCATGACGAATTCTATATCAGGTG
2	Rev-VBD domain	TAATGGATCCGGGTATCCTACGGTATCAATC
3	Rev- TarP$^{1-829\Delta625-650}$	TAATGGATCCCTTGGAGGTGGTGAAGG
4	Fwd-VBD domain	TAATGGTACCATGAACAAATTCCGCAAAGAA
5	Fwd-LD domain	TAATGGTACCATGACCCCATCAACTACAACA
6	Rev-LD domain	TAATGGATCCATAGATCCCGGCTTGCC
7	FWD-TarP$\Delta^{625-650}$	ATGGCATTGTCAATGTCAACGTTGGC
8	Rev-TarP$\Delta^{625-650}$	ACATTGAAATGCCATCGTCTTCGCT
9	Fwd-TarP-FH	AACGGCCGCCAGTGTATGACGAATTCTATATCAGGTGATCAACCT
10	Rev-TarP-FH	CCCTCTAGATGCATGTTATCCTACGGTATCAATCAGTGAGCTT
11	Fwd-FH-TarP	GATACCGTAGGATAACATGCATCTAGAGGGCCCTATTCTATAG
12	Rev-FH-TarP	TATAGAATTCGTCATACACTGGCGGCCGTTACTA
13	Fwd-TarP$^{829-1006}$-FH	AACGGCCGCCAGTGTACCCCATCAACTACAACATTAAGAACG
14	Rev- TarP$^{829-1006}$-FH	CCCTCTAGATGCATGTCCTACGGTATCAATCAGTGAGCTTAG
15	Fwd-FH- TarP$^{829-1006}$	ATTGATACCGTAGGACATGCATCTAGAGGGCCCTATTCTATAG
16	Rev-FH- TarP$^{829-1006}$	TGTAGTTGATGGGGTACACTGGCGGCCGTTACTA
17	Rev-TarP$^{1-829\Delta625-650}$-FH	CCCTCTAGATGCATGTTATGGAGGTGGTGAAGGCAGTAG
18	Fwd-FH- TarP$^{1-829\Delta625-650}$	CCTTCACCACCTCCATAACATGCATCTAGAGGGCCCTATTCTATAG

Figure S1. TarP translocated by *C. trachomatis* is stable. (A) Cos7 cells were either mock-infected or *Chlamydia*-infected for the indicated times, and maintained throughout in the presence of the prokaryotic protein synthesis inhibitor chloramphenicol. Whole cell lysates were harvested to monitor by Western blot the presence and stability of translocated TarP proteins. A rabbit polyclonal anti-TarP antibody was used. Protein concentrations were determined and adjusted to ensure equal loading. β-tubulin was used as the loading control. Mock-infected and EB-only samples were added to demonstrate specificity of the anti-TarP antibody used.

Figure S2. The PRD domain of TarP is not required for focal adhesion localization upon ectopic expression. Full-length TarP and TarP ΔPRD, both fused to mTurquoise2 were expressed ectopically in wild type MEFs for 19 h. The cells were processed for immunostaining for paxillin to visualize focal adhesions. Both full-length TarP and TarP ΔPRD colocalized with paxillin-positive structures, with the forming forming larger protein aggregates (arrowheads). Smaller TarP-positive punctae are indicated by white arrows.

Figure S3. Infection disrupts the stratified organization of focal adhesions. Cos7 cells were either mock-infected or infected with *C. trachomatis* serovar L2 for 20 h. DIC images of cells analyzed by iPALM were acquired to demonstrate the infection state. Regions of interests are bounded by white lines, and inclusions indicated by arrowheads.

Figure S4. LDVBD-expressing cells exhibited restricted motility relative to the empty vector-transfected control cells. Acquisition of the fluorescent channel was limited to the final frame of the motility assay to minimize phototoxicity. Images show the final DIC and fluorescent channel captured following 10 hours of time-lapse imaging. The DIC channel includes the dot and line overlay generated via ImageJ's manual tracking function to indicate the cell's movement over time.

Movie S1. Infection inhibits cell motility. A representative video assembled from a 17-h time-lapse imaging of mock-infected MEFs shows motility of individual cells with track outlines included.

Movie S2. Infection inhibits cell motility. A representative video assembled from a 17-h time-lapse imaging of *C. trachomatis*-infected MEFs shows motility of individual cells with track outlines included.

Movie S3. LDVBD is sufficient to inhibit cell motility. MEFs transfected with the vector alone was monitored for 17 h. A representative video assembled from a series of time-lapse images shows the degree of cell motility in the vector-only control group.

Movie S4. LDVBD is sufficient to inhibit cell motility. MEFs transfected with the LDVBD-mTurquoise2 was monitored for 17 h. A representative video assembled from a series of time-lapse images shows inhibition of cell motility of LDVBD-expressing cells.

Movie S5. *Chlamydia*-infected cells are resistant to detachment by mild trypsinization. Mock-infected HeLa cells growing on glass coverslips were treated by 0.025% Trypsin + EDTA, and imaged at 60-s intervals for 30 min. Note that the cells start to round up by seven min of incubation in trypsin.

Movie S6. *Chlamydia*-infected cells are resistant to detachment by mild trypsinization. *C. trachomatis* serovar L2-infected HeLa cells growing on glass coverslips were treated by 0.025% Trypsin + EDTA at 24 h post-infection, and imaged at 60-s intervals for 30 min. In contrast to

mock-infected cells shown in Movie S5, the infected cells remained attached and spread out after 30 min of mild trypsinization, indicating a possible enhancement of adhesion of infected cells to the substrate.

FIGURE S2

Paxillin-tdEos
Mock-infected

Paxillin-tdEos
CtrL2 infected

FAK-tdEos
Mock-infected

FAK-tdEos
CtrL2 infected

Vinculin-tdEos
Mock-infected

Vinculin-tdEos
CtrL2 infected

pFH-TarP ΔPRD
Paxillin-tdEos

pFH-TarP LDVBD
Paxillin-tdEos

FIGURE S3

FIGURE S4