Metagenomics reveals diet-specific specialization in fungus gardens of grass- and dicot-cutter ants

Lily Khadempour^{1,2,3}, Huan Fan¹, Ken Keefover-Ring⁴, Camila Carlos^{1,3}, Nilson S. Nagamoto⁵, Miranda A. Dam⁶, Monica T. Pupo⁷, Cameron R. Currie^{1,3}

Affiliations:

- ¹ Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA, 53706
- ² Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA, 53706
- ³ Department of Energy Great Lakes Bioenergy Research Center, University of Wisconsin-Madison, Madison, WI, USA, 53706
- ⁴ Departments of Botany and Geography, University of Wisconsin-Madison, Madison, WI, USA, 53706
- ⁵ Department of Plant Protection, UNESP São Paulo State University, Botucatu, SP, Brazil, CEP: 18603-970
- ⁶ Department of Nutritional Sciences, University of Wisconsin-Madison, Madison, WI, USA 53706
- ⁸ School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, São Paulo, Brazil, CEP: 14040-903

Conceived of the project (L.K., C.R.C), performed computational and statistical analysis (L.K., H.F., C.C.), analyzed data (L.K.) designed sampling methodology (L.K.) performed lab work (L.K., M.A.D., K.K.), performed field work (L.K., N.S.N.) provided resources and facilities (M.T.P., C.R.C) and wrote the manuscript (L.K., H.F., K.K., C.R.C.).

Abstract

Leaf-cutter ants are dominant herbivores in the Neotropics. While most leaf-cutter ant species cut dicots to incorporate into their fungus gardens, some species specialize on grasses. Here we examine the bacterial community associated with the fungus gardens of grass- and dicot-cutter ants to elucidate the potential role of bacteria in leaf-cutter ant substrate specialization. We sequenced the metagenomes of 12 Atta fungus gardens, across four species of ants, with a total of 5.316 Gbp of sequence data. We show that community composition was significantly different between dicot- and grass-cutter ants, with grasscutter ant fungus gardens having significantly lower diversity and a significantly higher abundance of *Pantoea*, the most abundant genus overall. Reflecting this difference in community composition, the bacterial functional profiles between the fungus gardens are significantly different. Specifically, grass-cutter ant fungus garden metagenomes are particularly enriched for genes responsible for amino acid, siderophore, and terpenoid biosynthesis while dicot-cutter ant fungus gardens metagenomes are enriched in genes involved in membrane transport. Our results suggest that bacteria in leaf-cutter ant fungus gardens aid in nutrient supplementation, a function especially relevant for the fungus gardens of ants that forage grass, a plant source relatively lower in nutrient value.

Introduction

Understanding the role of microbial symbionts in aiding nutrient acquisition is fundamental to understanding the biology of herbivores. Most herbivores host microbial symbionts that serve as an interface between them and the plants that they consume. These microbes can compensate for the hosts' lack of physiological capacity to obtain energy and nutrients from plants (Hansen & Moran 2013). Herbivore microbial symbionts, often residing in the guts of animals, have been implicated in aiding plant biomass breakdown (Talbot 1977; Kudo 2009; Hess et al. 2011; Adams et al. 2011), plant defense compound remediation (Wang et al. 2010; Adams et al. 2013; Boone et al. 2013), and nutrient supplementation (Warnecke et al. 2007; Hansen & Moran 2011; LeBlanc et al. 2013). Microbial communities differ between hosts that specialize on different substrates (Muegge et al. 2011) and changes in these communities and their functional capacity are integral to their hosts' transition to utilizing novel substrates (Delsuc et al. 2013; Kohl et al. 2014; Li et al. 2014; Hammer & Bowers 2015; Kohl et al. 2016).

Leaf-cutter ants represent a paradigmatic example of the microbial mediation of herbivory. They are dominant herbivores in the Neotropics, consuming an estimated 17% of foliar biomass in the systems in which they live (Costa et al. 2008). These ants have a significant impact on their surrounding ecosystems, due to the volume of plant biomass they consume and soil that they excavate in building their underground colonies (Fowler et al. 1986; Moutinho et al. 2003; Gutiérrez & Jones 2006; Herz et al. 2007; Costa et al. 2008). Like other metazoans, leaf-cutter ants lack the capacity to break down recalcitrant plant material. Instead, they gain access to the nutrients in plant biomass by farming a fungus, Leucoagaricus gongylophorus, which serves as an external gut that enzymatically breaks down recalcitrant biomass in the leaf material that the ants forage (Nagamoto et al. 2011; Kooij et al. 2011;

Suen et al. 2011a; Grell et al. 2013; Aylward et al. 2013; Khadempour et al. 2016). Leucoagaricus gongylophorus produces gongylidia, specialized hyphal swellings that contain an abundance of sugars and lipids, that the ants consume and feed to larvae (Bass & Cherrett 1995; North et al. 1997).

Recent work has revealed that a community of bacteria reside within leaf-cutter ant fungus gardens (Suen et al. 2010; Aylward et al. 2012; Moreira-Soto et al. 2017). These communities were dominated by Gammaproteobacteria, and consistently contained strains of Pseudomonas, Enterobacter and either Rahnella or Pantoea and were consistent with communities of bacteria associated with other fungus-farming insects (Aylward et al. 2014). Some garden bacteria are vertically transmitted, as they are present in the fungus pellets that queens use to establish new fungus gardens (Moreira-Soto et al. 2017). The community consistency and their vertical transmission, suggest that the bacterial communities are important to the fitness of their hosts. One study, by Pinto-Tomás et al. (2009) showed that Pantoea and Klebsiella bacteria fix nitrogen that supplements the ant diet, which is important for a strict herbivorous system. Nevertheless, the functional role of most garden bacteria remains unknown.

While most leaf-cutter ants use dicots, at least three species of *Atta* are specialized on cutting grass, and another three species cut both grasses and dicots (Fowler *et al.* 1986). All previous studies on the microbial community in leaf-cutter ant fungus gardens have been focused on dicot-cutting ants, likely because dicot-cutters are more common and grass-cutter ants are notoriously difficult to maintain in the lab (Nagamoto *et al.* 2009). In this study, we compare the bacterial communities of fungus gardens from ants that cut grass and dicots. Given that grasses and dicots differ in terms of the cell wall composition (Ding & Himmel 2008; Popper & Tuohy 2010), plant defense compounds (Wetterer 1994; Mariaca *et al.* 1997)

and nutrient availability (Mattson 1980; Winkler & Herbst 2004), we hypothesize that the bacterial community in these fungus gardens will differ in terms of community composition and functional capacity, in response to the different composition of the substrates the ants incorporate into their gardens. To address this, we collected fungus gardens from grass- and dicot-cutter ants and obtained their metagenomes using Illumina sequencing. We analyzed the bacterial community in terms of its taxonomic composition and its functional capacity. We also conducted analyses on the fungus gardens to determine their plant composition, their nutritional composition and their plant defense compound contents.

Methods

Collection of fungus garden

Fungus gardens were collected on the campuses of the University of São Paulo (USP) in Ribeirão Preto, SP, Brazil and the State University of São Paulo (UNESP) in Botucatu, SP, Brazil. Collection dates and GPS coordinates are listed in Table 3.1. We collected fungus gardens from four species of *Atta* leaf-cutter ants: *A. bisphaerica* and *A. capiguara*, which both specialize on grass, *A. laevigata*, which cuts both grasses and dicots, and *A. sexdens*, a dicot-cutter ant (Fowler *et al.* 1986).

To collect the fungus gardens, we identified the ant species by worker morphology then followed the entrance tunnel by digging until we found a fungus garden. Care was taken to enter fungus garden chambers from the side, to avoid damaging the garden with digging tools and to avoid contamination with surrounding soil. Fungus gardens were transported to the laboratory and were aseptically transferred into 50 mL conical tubes. The majority of worker ants were removed from the fungus garden material before being transferred to the tubes. In order to further reduce the chance of soil contamination, only intact fungus garden from the central region of the fungal mass was included in the tubes. Once filled, the tubes

were frozen in liquid nitrogen and stored at -80°C. At least six 50 mL conical tubes were filled from each colony. For each colony, four tubes were used for metagenomics, one tube was used for gas chromatography, and one tube was used for iron content measurements.

DNA extraction

To target the bacteria in the fungus gardens, DNA was extracted by first using a differential centrifugation method (Aylward et al. 2012). PBS buffer with 1% tween 80 was added to the tubes and they were shaken for 30 min on a vortex. They were then kept at 4°C for 30 min so that large particles would settle. The liquid portion was decanted and passed through a 40 µm filter. The remaining leaf material from the fungus gardens was photographed after the differential centrifugation, to demonstrate the difference in leaf material consistency (Figure 1). The filtrate was centrifuged for 30 min at 4°C, after which a bacterial cell pellet was formed and the liquid was removed. This process was repeated with the original fungus garden tube. For each fungus garden, cell pellets from four tubes were combined and the DNA was extracted using the Qiagen Plant DNA Extraction Maxi Kit (Qiagen, Hilden, Germany).

DNA sequencing and assembly

All metagenomic sequencing was conducted at the Joint Genome Institute in Walnut Creek, CA. Since some of the DNA concentrations were too low for standard library prep, a low-input prep was completed for all of the samples. Sequencing was performed on an Illumina HiSeq-2500 platform (2 x 151 bp). BBDuk adapter trimming (Bushnell 2017) was used to remove known Illumina adapters. The reads were then processed using BBDuk filtering and trimming. Read ends were trimmed where quality values were less than 12. Read pairs containing more than three ambiguous bases, or with quality scores (before trimming) averaging less than three over the read, or length under 51 bp after trimming, as

well as reads matching Illumina artifact, spike-ins or phiX were discarded. Trimmed, screened, paired-end Illumina reads were assembled using the megahit assembler using with the "--k-list 23,43,63,83,103,123" option. Functional annotation and taxonomic classification were performed using the Integrated Microbial Genomes pipeline.

Plant genus richness

To determine the richness of plant substrate integrated in the fungus gardens of the ants, we used JGI's Integrated Microbial Genomes and Microbiomes (IMG) database "find gene" function to retrieve all genes annotated as *MatK* from the dataset. *MatK* is a widely used chloroplast plant DNA barcode (Hollingsworth *et al.* 2011). Retrieved *MatK* sequences for each metagenome were identified using BLAST. To ensure consistent and reliable certainty with the identified plants, we identified all sequences to the genus level. Because most of the plant biomass was removed from samples before DNA extraction only presence/absence of genera were considered, not abundance.

Bacterial taxonomic analysis

Abundance of bacterial groups (phyla and genera) were determined based on the IMG Phylogenetic Distribution tool, which is part of JGI's standard operating procedure (Huntemann et al. 2016). Briefly, IMG uses USEARCH (Edgar 2010) to compare metagenome gene sequences to all identified genomes in their database. One top USEARCH hit per gene is used to assign phylogenetic lineage. To determine relative abundance of bacterial taxonomic groups within each sample, we used the PhyloDist raw data from IMG and first removed all gene sequences that were identified as Eukaryote or Virus. We then matched the PhyloDist data to the gene counts for each gene and normalized them to the total number of genes from Bacteria and Archaea. We used the relative abundances of each phylum and genus to run an non-metric multidimensional analysis (NMDS) using a Bray-

Curtis dissimilarity index with the vegan package in the R statistical programming environment (Oksanen et al. 2013; R Core Team 2013). Also using the vegan package, we used ANOSIM and PERMANOVA to determine if groups (grass-cutters vs. dicot-cutters) were significantly different, and we used the Shannon diversity index to compare the diversity of each sample by bacterial genus. To test whether specific genera have significantly different relative abundances between grass- and dicot-cutter ant fungus gardens, we used DESeq2 in the R statistical programming environment (Love et al. 2014). Since DESeq2 requires inputs to be integers, we used number of gene copies per million genes in the metagenomes as our input (Alneberg et al.).

Bacterial functional analysis

In order to make functional comparisons of the bacteria in grass- and dicot-cutter fungus gardens, we used the Kyoto Encyclopedia of Genes and Genomes (KEGG) annotations of the metagenomes through IMG's KEGG Orthology (KO) pipeline, which is part of JGI's standard operating procedure (Huntemann et al. 2016). Briefly, genes were associated with KO terms (Kanehisa et al. 2014) based on USEARCH 6.0.294 results (Edgar 2010) and were filtered for minimum identity matches and gene sequence coverage. For an overall comparison of functional differences between the fungus gardens, we used the same ordination and statistical methods as for bacterial genus abundance. As with genus group differences, we used DESeq2 to determine what genes are significantly enriched between grass- and dicot-cutter ant fungus gardens, with number of gene copies per million genes in the metagenomes as our input (Alneberg et al.).

Iron content

Separate 50 mL tubes of fungus garden material, from the same colonies as above, were used for determination of iron content. All ants were removed from fungus garden

then the remaining material was analyzed at the UW Soil and Forage Lab in Marshfield, WI, using standard methods. Briefly, total iron content was determined by first digesting the fungus garden material in nitric acid/peroxide then analyzing by inductively coupled plasma optical emission spectroscopy (ICP-OES) (Fassel & Kniseley 1974).

Results

Metagenomic statistics

A summary of metagenome statistics is presented in (Table 2). A total of 5.316 Gbp of assembled sequence data was produced in this study, with an average of 443 Mbp per metagenome. The smallest metagenome was from the grass-cutter colony *A. capiguara* 1 at 148.7 Mbp, and the largest metagenome was from the dicot-cutter colony *A. sexdens* 2 at 812.9 Mbp. Maximum scaffold lengths ranged from 61.96 Kbp to 701.42 Kbp, with an average maximum scaffold length of 266.6 Kbp. Between 91.63% and 99.31% of reads were aligned.

Bacterial taxonomic analysis

Proteobacteria (70-99%) were the most abundant bacterial phylum detected in the fungus gardens of *Atta* spp., followed by Actinobacteria (0.13-24%) and Firmicutes (0.096-2.4%) (Supplemental Figure 1). Between fungus gardens, genus-level comparisons showed greater variability than phylum-level comparisons (Figure 2, Figure 3). Overall, *Pantoea* was the most abundant genus in all the fungus gardens (average 37%), followed by *Pseudomonas* (average 17%). The abundance of these two genera was especially pronounced in the grass-cutter ant fungus gardens, where *Pantoea* and *Pseudomonas* averaged 45% and 28%, respectively. The high relative abundance of these two genera contributed to a lower overall diversity in the grass-cutter ant gardens (Shannon diversity index of 1.20-2.44) (Figure 3). While *Pantoea* and *Pseudomonas* were still abundant in fungus gardens of the dicot-cutter ants,

A. laevigata and A. sexdens, it accounted for a lower proportion (28% and 6.5%, respectively) of the bacteria in these more diverse gardens (Shannon diversity index of 2.80-4.67). Other dominant bacterial genera included Enterobacter, Burkholderia, Erwinia, Emticicia, Serratia and Klebsiella. DESeq2 analysis revealed that six bacterial genera (Entoplasma, Flavobacterium, Mesoplasma, Pantoea, Pseudomonas, and Spiroplasma) were significantly different in relative abundance between the fungus gardens. They were all more abundant in the grass-cutter ant fungus gardens.

Bacterial functional analysis

Overall, we found significant differences in the predicted bacterial community functional profiles between grass- and dicot-cutter ant fungus gardens (Figure 4). All individual bacterial genes that were significantly different between grass- and dicot-cutter ant fungus gardens are listed in Supplemental Table 1. In total, 514 predicted bacterial genes were significantly enriched in one group or another, with 313 and 201 genes significantly enriched in grass- and dicot-cutter ant gardens, respectively (Supplemental Table 2, Supplemental Figures 4-6). Grass-cutter ant fungus gardens were enriched for amino acid biosynthesis genes for phenylalanine, tryptophan, tyrosine, histidine, arginine, lysine, cysteine, methionine, glycine, serine and threonine. They were also significantly enriched in terpenoid and siderophore biosynthesis genes (Figure 5) and had a significantly higher abundance of a gene in the nitrogen fixation pathway, nitrogenase molybdenum-iron protein beta chain (Supplementary Table 2). Dicot-cutter ant fungus gardens were particularly enriched in membrane transport genes (Figure 5).

Plant taxonomy and consistency

The incorporated plant material was markedly different in consistency between the fungus gardens. *Atta bisphaerica* and *A. capiguara* gardens both contained material that was

clearly grass, which was not mulched (Figure 1). In contrast, the leaf material in the fungus gardens of *A. laevigata* and *A. sexdens* was mulched to the point of being unrecognizable as plant material (Figure 1). We detected 68 plant species based on the *MatK* gene query in the metagenomes, from 40 genera and 15 families (Table 3). The fungus gardens of dicot-cutter ants had a significantly higher richness of plant genera than those of grass-cutter ants (ANOVA F=9.14, p=0.0128). As expected, the grass-cutter ant fungus gardens all contained grass (*Paspalum*, Poaceae). The dicot-cutter ant fungus gardens contained more genera and families of plants, which were mostly dicots, but three of these fungus gardens also contained some grass (Table 3).

Iron content

The iron content of the fungus gardens is displayed in Figure 6. The grass-cutter ant fungus gardens have lower amounts of iron than the dicot-cutter ant fungus, but this difference is not significant due to the high variability between *A. sexdens* gardens.

Discussion

Understanding how microbial symbiont communities change in relation to host substrate specialization can help inform on animal diet specialization and evolutionary transitions to utilizing novel substrates. *Atta* ants provide a relatively unique opportunity to examine a group of closely related herbivores that have transitioned from specialization on dicots to grasses. Dicots and grasses differ in terms of their cell wall composition, nutrient content and plant defense compounds. Here, using metagenomic sequencing, we examine this transition in the bacterial community in the fungus gardens of ants that are specialized on these different substrates. The results of this study demonstrate that the bacterial community differs depending on type of substrate and likely facilitates the ants' ability to

specialize on grasses, which represents a lower quality of substrate on which to grow their fungal crop.

If bacteria in fungus gardens are responsible for the breakdown of recalcitrant plant biomass, which is found in plant cell walls, we expect that the bacterial communities in the two ant groups examined here would be differentially enriched in the genes necessary for plant biomass breakdown. Grasses have a unique cell wall structure, containing (1→3),(1→4)-β-D-glucan chains and silica, neither of which are present in dicots (Popper & Tuohy 2010). In other systems specialized on grass biomass breakdown, the microbes responsible for this produce specialized enzymes (King et al. 2011) and have genomes that are adapted for this function (Wolfe et al. 2012). Since we do not observe any changes in abundance of these genes between these two systems, it is unlikely that the bacteria here are contributing to plant biomass breakdown. Indeed, recent work has implicate the fungal cultivar as the primary degrader of plant biomass in leaf-cutter ant fungus gardens (Nagamoto et al. 2011; Grell et al. 2013; Aylward et al. 2013; Khadempour et al. 2016).

Leaf-cutter ants, in general, cut an exceptionally broad diversity of plants (Mayhé-Nunes & Jaffe 1998; Solomon 2007) and thus, have the potential to encounter a myriad of plant defense compounds that are toxic to themselves and their fungal cultivar. The ants are not enriched in genes families for plant defense compound detoxification (Rane *et al.* 2016), so they must reduce the intake of these chemicals in other ways. Plant defense compound avoidance occurs in several steps. First, ants avoid cutting plants that contain plant defense compounds that are particularly toxic or abundant (Hubbell *et al.* 1984; Howard 1988; Wirth *et al.* 1997). Second, many plant defense compounds that the ants encounter are volatile chemicals (Howard 1988; Howard *et al.* 1988), and in the time that the ants cut and carry the leaf material back to their colonies, some of the volatiles will have had time to dissipate.

Finally, ants often leave leaf material in caches before they incorporate them into their fungus gardens (Hart & Ratnieks 2000; Roschard & Roces 2003), providing further opportunity for the defense compounds to evaporate. Nevertheless, some amount of volatiles can make their way into the gardens. In this study, using gas chromatography, we were able to detect eucalyptus-related compounds (eucalyptol, α -pinene, β -pinene, γ -cymene and γ -terpinene) in the fungus garden of one ant colony (γ -laevigata 1) that was observed cutting considerable amounts of eucalyptus (Supplemental methods and Supplemental Figure 2).

In order to mitigate the deleterious effects of plant defense compounds, we expect the fungal cultivar L. gongylophorus would produce enzymes to degrade them. Indeed, work by De Fine Licht et al. (2013) suggests that laccases from the fungal cultivar help detoxify plant defense compounds. Nevertheless, bacteria in the garden may also play a role in mediating plant defense compounds. The bacterial community contains the genes necessary for plant defense compound remediation, including many cytochrome P450s, gluthione S-transferases, and other genes involved in xenobiotic degradation, and aromatic compound degradation, but they are not consistently enriched in the dicot-cutter ant fungus gardens (Supplemental Table 1). We expected that since dicot-cutter ants incorporate a higher diversity of plants into their gardens (Table 3), that the diversity of bacteria would also be higher in these gardens, and that the bacteria would have a higher capacity for the degradation of these defense compounds. While we did observe a greater diversity of bacteria in the dicot-cutter ant fungus gardens (Figure 3) we did not see a significant enrichment of plant defense compound degradation genes in these gardens (Figure 5, Supplemental Table 1). However, we still cannot exclude the possibility that bacteria are taking part in this process. Since each dicot-cutter ant colony cuts a unique set of plants (Table 3), they potentially encounter a

unique set of plant defense compounds. If the bacterial community were to respond in a substrate-specific manner to different plant defense compounds, our analysis in this study would not reveal that. To elucidate the role of bacteria in plant defense compound remediation, closely controlled experiments with particular defense compounds of interest applied to bacterial cultures and to fungus gardens would be necessary.

Pinto-Tomas et al. (2009) establishes that Pantoea and Klebsiella bacteria in Central American leaf-cutter ant fungus gardens are supplementing the ant diet through nitrogen fixation. Plant material, in general, is low in nitrogen, and many herbivores supplement their diets through bacterial nitrogen fixation (Douglas 2009; Hansen & Moran 2013). Grasses are especially low in nitrogen (Mattson 1980; Winkler & Herbst 2004), so we would hypothesize that grass-cutter ant fungus gardens would be enriched in nitrogen-fixing bacteria with a corresponding enrichment of nitrogen-fixing genes. Here we show that Pantoea are more abundant in the grass-cutter ant fungus gardens, and that a nitrogenase molybdenum-iron protein beta chain gene is significantly more abundant in grass-cutter ant fungus gardens (Supplemental Table 1). Other genes that are related to nutrient acquisition are also significantly more abundant in the grass-cutter ant fungus gardens (Figure 5), such as genes in amino acid metabolism pathways. While it has been shown that nitrogen fixed by bacteria is incorporated into the bodies of ants (Pinto-Tomás et al. 2009), animals cannot simply absorb nitrogen as ammonium or nitrate, they require it to either be in the form of amino acids or other organic nitrogen-containing compounds (White 1993). The enrichment of arginine biosynthesis genes is of particular interest since the genome of Atta is deficient in genes in this pathway (Suen et al. 2011b), and no evidence has been found that the fungus provides arginine (Aylward et al. 2013; Khadempour et al. 2016).

Other categories of genes enriched in the grass-cutter ant fungus garden bacteria are those involved in metabolism of terpenoids and secondary metabolites, especially their biosynthesis. Grass-cutter ant fungus gardens are significantly enriched in 67 of these genes. This list includes seven siderophores, which are responsible for iron acquisition (Crosa 1989; Winkelmann 2002). Siderophores are costly to produce so the enrichment of these genes suggests that iron acquisition is important in this system. The grass-cutter and fungus gardens examined in this study contained lower amounts of iron than the dicot-cutter ant fungus gardens (Figure 6). Terpenoids are the most abundant secondary metabolites found in plants, and serve diverse roles (Langenheim 1994; Gershenzon & Dudareva 2007). The majority of research into the connection between plant terpenoids and animal-microbe symbioses are in regards to the detoxification of terpenes that would be deleterious to the animal host (Wang et al. 2012; Cheng et al. 2013; Adams et al. 2013; Boone et al. 2013; Raffa 2013). However, not all terpenes are toxic to all organisms (Raffa 2013), and in at least one instance they have been shown to supplement a herbivore's diet after some modification by a gut bacterium (Berasategui et al. 2017). Dicots contain higher quantities of terpenoids (Wetterer 1994; Mariaca et al. 1997). One possibility is that the bacteria in these fungus gardens are producing terpenes as a nutritional additive, especially in the grass-cutter ant fungus gardens where there are lower terpene inputs and these genes are enriched (Figure 5, Supplemental Figure 2).

It is important to consider this system, not only from the perspective of the bacteria, but also that of the ants. One may ask: if grasses are so poor in nutrients, why do the ants use them as a substrate at all? Grass-cutter ants can be categorized as facultative specialists (Shipley *et al.* 2009), as they have the capacity to use grass, a difficult substrate, as their primary fungus garden input but they are also capable of consuming a wider range of

substrates. The ants themselves are adapted to cutting grass with shorter, wider mandibles than their dicot-cutter counterparts (Fowler et al. 1986; Silva et al. 2016). They also process leaves differently - they do not mulch the material, likely because the silica contained in grasses would dull their mandibles (Massey & Hartley 2009; Silva et al. 2016). The ants' adaptation to grass-cutting, combined with the community response of the bacteria in the gardens, allow grass-cutter ants to use grass as a substrate more efficiently than dicot-cutter ants can. This has allowed grass-cutter ants to exploit a novel niche, presumably reducing the amount of interspecific competition they experience. It should be noted, however, that grass-cutter ants, if given the opportunity to use palatable dicots, actually prefer those over grasses (Nagamoto et al. 2009), probably due to their lower recalcitrance and higher nutrient content. This indicates that grass-cutter ants' fundamental niche is broader than their realized one. Not only do grass-cutters cut dicots, when they are available to them, but in this study, we detected grasses in the fungus gardens of some dicot-cutters as well (Table 3) and both A. laevigata and A. sexdens workers were observed cutting grass (personal observation). Even though there is some grass in these fungus gardens, the majority of species that these ants cut are dicots. As well, A. laevigata and A. sexdens both process their leaves in the manner consistent with other dicot-cutter ants (Fowler et al. 1986)).

Grass-cutter ants forage on a substrate that is lower in nutritional quality than their dicot-cutter counterparts. The grasses they cut are lower in nitrogen and iron. Optimal foraging theory predicts that when the quality of forage is lower, the ants should be cutting a greater diversity of plants (Rockwood & Hubbell 1987). Instead, grass-cutter ants cut a significantly lower diversity of plants than their dicot-cutter counterparts. Grass-cutter ants to do not have access to the diversity of plants necessary to compensate for their low forage quality. Instead, the bacteria in their fungus gardens can provide the necessary nutrition that

a diverse diet provides in dicot-cutter ant fungus gardens, allowing grass-cutter ant species to exploit this novel niche. Not only does our work here provide further evidence of the importance of bacteria in the leaf-cutter ant system, it provides further support that microbial symbionts are important players in novel substrate utilization by animals.

Acknowledgements

The authors would like to thank Andre Rodrigues of UNESP for his help with collections and permits. This work was funded in part by the U.S. Department of Energy Great Lakes Bioenergy Research Center (DOE Office of Science BER DE-FC02-07ER64494), National Institute of Food and Agriculture, United States Department of Agriculture, under ID number 1003779, National Institutes of Health (NIH grant U19TW009872) and São Paulo Research Foundation (FAPESP grant 2013/50954-0). Collection at the UNESP campus in Botucatu was completed under the ICMBio permit 31534-1. Collection at the USP campus in Riberão Preto was completed under the SISBIO permit 46555-5 and CNPq permit 010936/2014-9. The work conducted by the DOE Joint Genome Institute, a DOE Office of Science User Facility, is supported by the Office of Science of the DOE under Contract No. DE-AC02-05CH11231.

Table 1 Summary of collection details for leaf-cutter ant colonies used in this study

Leaf-cutter ant	Substrate	IMG genome			
colony	niche	number	Collection date	Latitude	Longitude
A. bisphaerica 1	Grass	3300013023	1-Feb-15	S22°50'47.7"	W48°26′.9″
A. bisphaerica 2	Grass	3300013025	3-Feb-15	S22°50'48.4"	W48°26'1.4"
A. bisphaerica 3	Grass	3300013022	3-Feb-15	S22°50'48.4"	W48°26'2.3"
A. capiguara 1	Grass	3300012994	2-Feb-15	S22°54'32.1"	W48°18'28.7"
A. capiguara 2	Grass	3300012996	3-Feb-15	S22°50'47.2"	W48°26'1.3"
A. capiguara 3	Grass	3300012997	3-Feb-15	S22°50'47.6"	W48°26'1.2"
A. laevigata 1	Dicot*	3300013000	20-Jan-15	S21°9'55.5"	W47°50'51.3"
A. laevigata 2	Dicot*	3300012995	17-Jan-15	S21°10'3"	W47°50'47"
A. laevigata 3	Dicot*	3300012998	19-Jan-15	S21°9'56.8"	W47°50'52.7"
A. sexdens 1	Dicot	3300012999	30-Jan-15	S21°9'50"	W47°51'26.9"
A. sexdens 2	Dicot	3300013002	30-Jan-15	S21°9'53.4"	W47°51'10.5"
A. sexdens 3	Dicot	3300013001	31-Jan-15	S21°10'2"	W47°51'5"

^{*}While A. laevigata has been described as a grass/dicot-cutter ant (Fowler et al. 1986), due to its leaf-processing behavior and fungus garden plant composition observed in this study, we consider it a dicot-cutter.

Table 2 Metagenome sequencing statistics for leaf-cutter ant fungus gardens

		Scaffold	Main	Main	Max			
		sequence	genome	genome	scaffold	Scaffolds		Protein
Leaf-cutter ant	Scaffold	total	scaffold	scaffold	length	> 50		coding
colony	total	(Mbp)	N/L50	N/L90	(Kbp)	Kbp	Aligned reads	genes
A. bisphaerica 1	628724	390.8	122506/	467228/	249.52	93	163185122	607 042
21. vispisaciica 1	020724	370.0	740	298	247.32	(2.07%)	(98.76%)	(99.39%)
A. bisphaerica 2	939707	630.9	148370/	680177/	253.54	69	148406252	910 609
21. vispoučina 2	737101	050.7	926	292	233.34	(0.88%)	(96.49%)	(99.61%
A. bisphaerica 3	285649	186.0	29401/	244722/	187.50	49	167169016	358 547
21. <i>он</i> ървает а 3	203047	100.0	972	247	107.50	(1.90%)	(98.76%)	(98.19%
A. capiguara 1	205334	148.7	16608/	178745/	273.83	37	199009346	272 096
21. tapignara 1	203334	140.7	1403	247	273.03	(2.49%)	(99.31%)	(98.99%
A. capiguara 2	345332	261.2	35330/	303130/	180.27	34	204772230	456 916
21. tupignuru 2	343332	201.2	1420	247	100.27	(1.06%)	(98.31%)	(98.70%
A. capiguara 3	573737	359.5	83958/	508567/	135.51	13	203079026	644 865
1. rapignara 3	313131	337.3	790	247	155.51	(0.29%)	(98.77%)	(98.79%
A. laevigata 1	853367	517.5	178678/	645038/	274.98	87	178461364	871 330
21. iucu gaia 1	033307	317.3	686	301	274.70	(1.53%)	(96.65%)	(99.42%
A. laevigata 2	295365	189.2	32897/	266928/	252.44	92	189659024	332 737
A. iuevigaia 2	<u> </u>	109.2	990	247	232.44	(4.23%)	(96.85%)	(96.95%
A. laevigata 3	601593	546.3	74824/	398992/	241.71	17	209535750	722 718
A. iaeu gaia 3	001373	340.3	1744	340	241.71	(0.30%)	(96.06%)	(99.01%
A. sexdens 1	674609	708.4	48220/	412923/	701.43	167	156148622	822 403
∠n. sexuens 1	0/4009	700.4	3118	341	701.43	(2.40%)	(97.92%)	(99.46%
A. sexdens 2	857038	812.9	65552/	548662/	61.96	17	150809208	1 088 71
∠1. Sexuens Z	037038	012.9	2346	328	01.90	(0.11%)	(95.32%)	(99.51%
A. sexdens 3	1006806	564.7	221493/	772141/	386.55	68	186976430	1 029 78
∠1. sexuens 3	1000000	304.7	614	277	200.33	(1.17%)	(91.63%)	(99.20%

Table 3 Plant genera detected in each fungus garden sample using MatK gene

Sample Family Genus match % Fabaceae Chamaecrista 99.3 A. bisphaerica 1 Poaceae Paspalum 99.6 Polygalaceae Polygala 99.3 A. bisphaerica 2 Poaceae Paspalum 99.4 Fabaceae Chamaecrista 99.3 A. bisphaerica 3 Fabaceae Chamaecrista 99.3 Fabaceae Polygala 99.5 Polygalaceae Polygala 99.5 Polygalaceae Polygala 99.5 Polygalaceae Polygala 99.0 A. capiguara 1 Poaceae Paspalum 99.6 A. capiguara 2 Poaceae Paspalum 99.6 Fabaceae Chamaecrista 99.3 A. capiguara 3 Fabaceae Chamaecrista 99.3 Poaceae Paspalum 99.6 Fabaceae Paspalum 99.6 A. laenigata 1 Poaceae Paspalum 99.5 Poaceae Paspalum 99.5 Poaceae Paspalum 99.5 Poaceae Paspalum 99.5 Poaceae Paspalum 99.6 A. laenigata 2 Fabaceae Centrolobium 98.2 Fabaceae Schizgolobium 100 Anacardiaceae Asteraceae Paspalum 99.5 A. laenigata 3 Fabaceae Lencaena 100 Myrtaceae Paspalum 99.8 Fabaceae Lencaena 100 Myrtaceae Paspalum 99.8 Fabaceae Lencaena 100 Myrtaceae Paspalum 99.8 Fabaceae Lencaena 100 Myrtaceae Balesia 98.8 Fabaceae Paspalum 99.9 Anacardiaceae Andira 98.6 Fabaceae Paspalum 99.9 Anacardiaceae Andira 98.6 Fabaceae Paspalum 99.9 Malvaceae Balesia 98.8 Fabaceae Loscoptergium 98.4 A. sexalens 1 Fabaceae Prengyne 100 Fabaceae Loscoptergium 98.5 Bignoniaceae Tabebria 98.1 Fabaceae Loscoptergium 98.4 A. sexalens 1 Fabaceae Prengyne 99.9 Malvaceae Balesia 98.8 Fabaceae Loscoptergium 99.9 Malvaceae Balesia 98.8 Fabaceae Loscoptergium 99.9 Malvaceae Balesia 98.3 Combretaceae Engelia 99.2 Solanaceae Centrolobium 98.2 Fabaceae Fabaceae Prengyne 99.3				Species
A. bisphaerica 1 Poaceae Poisgala Poisgala Polygalaceae Polygalaceae Polygala Polygalaceae Polygala Polygalaceae Polygala Polygalaceae Parpatum Polygalaceae Polygala Poaceae Parpatum Pooceae Polygalaceae Polygalac	Campla	Damil	Comus	
A. bisphaetica Polygalaceae Paspalum 99.6 Polygalaceae Polygala 99.3 99.3 A. bisphaetica Poaceae Paspalum 99.4 Fabaceae Zortia 100 Poaceae Paspalum 99.5 Polygalaceae Paspalum 99.5 Polygalaceae Paspalum 99.5 Polygalaceae Polygala 99.0 A. capiguara Poaceae Paspalum 99.6 A. capiguara Poaceae Paspalum 99.6 Poaceae Paspalum 99.5 Poaceae Paspalum 99.5 Poaceae Paspalum 99.5 Poaceae Propalum 99.5 Poaceae Paspalum 99.6 Paspalum 99.8 Paspalum 99.8 Paspalum 99.8 Paspalum 99.8 Paspalum 99.8 Paspalum 99.9 Paspalum	Sample			
Polygalaceae	1 high barries 1			
A. bisphaerica 2	A. vispnaenia 1			
Fabaceae Chamaecrista 99.3	1 high bassing 2			
A. bisphaerica 3	A. vispnaenica 2	•		•
Poaceae				
Polygalaceae	A. bisphaerica 3			
A. capignara 1 Poaceae Paspalum 99.7 A. capignara 2 Poaceae Paspalum 99.6 A. capignara 3 Fabaccae Chamaecrista 99.3 A. laevigata 1 Myrtaceae Perogyne 99.4 Myrtaceae Puspalum 99.5 Poaceae Puspalum 99.5 Poaceae Purochloa 100 Asteraceae Rensonia 99.6 A. laevigata 2 Fabaceae Centrolobium 98.2 Fabaceae Schizolobium 98.6 Asteraceae Ring anthus 95.4 Asteraceae Podanthus 99.5 A. laevigata 3 Fabaceae Desmodium 99.8 Asteraceae Podanthus 99.5 A. laevigata 3 Fabaceae Desmodium 99.8 Fabaceae Desmodium 99.8 Asteraceae Podanthus 99.8 Poaceae Puspalum 99.9 Antaceae Leucaphytus 99.8 P				
A. capignara 2 Poaceae Paspalum 99.6 A. capignara 3 Fabaceae Chamaecrista 99.3 Poaceae Paspalum 99.6 Fabaceae Pterogyne 99.4 Myttaceae Encalptus 99.9 Poaceae Baspalum 99.5 Poaceae Urochloa 100 Asteraceae Rensonia 99.5 Fabaceae Schizolobium 100 Anacardiaceae Schizolobium 99.6 Anacardiaceae Pachycomus 98.6 Asteraceae Poadnthus 99.5 Asteraceae Podanthus 99.5 A steraceae Podanthus 99.5 A steraceae Podanthus 99.5 A steraceae Podanthus 99.5 A laenigata 3 Fabaceae Lucaena 100 Myttaceae Paspalum 99.8 Poaceae Lucaena 100 Myttaceae Paspalum 98.1 Fabaceae Tabe	4 cationara 1		***	
Fabaceae			· · · · · · · · · · · · · · · · · · ·	
Poaceae				
Fabacea	A. capiguara 3			
A. laevigata 1				
Poaceae	4 1 1 4 4			
Poaceae	A. taevigata 1	Poaceae		99.5
A. laevigata 2 Fabaceae Centrolobium 98.2 Fabaceae Schizolobium 100 Anacardiaceae Pachycormus 98.6 Asteraceae King anthus 95.4 Asteraceae Dodanthus 99.5 A. laevigata 3 Fabaceae Desmodium 99.8 Fabaceae Desmodium 99.8 Fabaceae Desmodium 99.8 Fabaceae Desmodium 99.8 Poaceae Desmodium 99.8 Poaceae Desmodium 99.8 Poaceae Desmodium 99.8 Poaceae Eucalyptus 99.8 Poaceae Paspatum 99.9 Anacardiaceae Loxopterygum 98.4 Asteraceae Adira 98.6 Fabaceae Batesia 98.6 Fabaceae Libidibia 99.8 Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6		Poaceae	Urochloa	100
Fabaceae		Asteraceae	Rensonia	99.6
Anacardiaceae	A. laevigata 2	Fabaceae	Centrolobium	98.2
Asteraceae Ringianthus 95.4 Asteraceae Podanthus 99.5 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Myrtaceae Paspalum 99.9 Anacardiaceae Loxopterygium 98.4 Asteraceae Cymophora 98.5 Bignoniaceae Tabebuia 98.1 Fabaceae Batesia 98.8 Fabaceae Batesia 98.8 Fabaceae Bussea 100 Fabaceae Libidibia 99.8 A. sexdens 1 Fabaceae Pierogyne 100 Fabaceae Pierogyne 100 Myrtaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eucalyptus 99.6 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitgera 99.3 Fabaceae Perogyne 99.3 Fabaceae Pierogyne 99.3 Fabaceae Echipaa 100 Lecythidaceae Echipaa 100 Lecythidaceae Echipaa 100 Asteraceae Echipaa 100 Asteraceae Echipaa 100 Commelinaceae Careya 94.0 Santalaceae Perymeniopsis 99.8 Asteraceae Echipaa 100 Commelinaceae Commelina 100 Commelinaceae Desmodium 99.8 Fabaceae Desmodium 99.8		Fabaceae	Schizolobium	100
A. laevigata 3 Fabaceae		Anacar diaceae	Pachycormus	98.6
A. laevigata 3 Fabaceae Fabaceae Leucaena 100 Myrtaceae Encatyptus 99.8 Poaceae Paspalum 99.9 Desmodium 99.9 Anacardiaceae Asteraceae Gymophora 98.5 Bignoniaceae Fabaceae Bignoniaceae Batesia 98.6 Fabaceae Batesia 98.8 Fabaceae Bussea 100 Fabaceae Libidibia 99.8 Fabaceae Prerogyne 100 Fabaceae Prerogyne 100 Myrtaceae Pachira 100 Myrtaceae Encatyptus 99.6 Myrtaceae Encatyptus 99.6 Myrtaceae Engenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lyclanthes 100 Bignoniaceae Genipa 99.2 Solanaceae Lyclantee 100 Fabaceae Lyclanthes 100 Myrtaceae Encatyptus 99.6 Myrtaceae Encatyptus 99.6 Myrtaceae Encatyptus 99.8 Poaceae Scutachne 98.4 Rubiaceae Centrolohium 98.3 Combretaceae Lyclanthes 100 Ecythidaceae Centrolohium 98.2 A. sexdens 2 A. sexdens 2 Fabaceae Tipuana 100 Lecythidaceae Pherogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Pherogyne 99.3 Asteraceae Echpta 100 Asteraceae Echpta 100 Asteraceae Perymeniopsis 99.8 Asteraceae Echpta 100 Asteraceae Echpta 100 Commelinaceae Commelina 99.6 Asteraceae Perymeniopsis 99.8 Asteraceae Perymeniopsis 99.8 Asteraceae Desmodium 99.8 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Encatyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Encatyptus 98.6 Phyllanthus 100 Rubiaceae Genipa 100 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Encatyptus 98.6 Phyllanthus 100 Rubiaceae Encatyptus 98.6 Phyllanthus 100 Genipa 100 Rubiaceae Encatyptus 98.6 Phyllanthus 100 Genipa 100 Rubiaceae Genipa 100 Rubiaceae Genipa 100 Genipa 100 Rubiaceae Genipa 100 Genipa 100 Rubiaceae Genipa 100 Genipa		Asteraceae	Kingianthus	
Fabaceae Leucalpytus 99.8		Asteraceae	Podanthus	99.5
Myrtaceae Eucalyptus 99.8 Poaceae Paspalum 99.9	A. laevigata 3	Fabaceae	Desmodium	99.8
Poaceae		Fabaceae	Leucaena	100
Anacar diaceae		Myrtaceae	Eucalyptus	99.8
Asteraceae Cymophora 98.5 Bignoniaceae Tabebnia 98.1 Fabaceae Andira 98.6 Fabaceae Batesia 98.8 Fabaceae Bussea 100 Fabaceae Libidibia 99.8 A. sexdens 1 Fabaceae Pterogyne 100 Fabaceae Pabira 100 Myrtaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Compa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Centrolohium 98.2 A. sexdens 2 Fabaceae Pterogyne 99.3 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Eclipta 100 Asteraceae Perymeniopsis 99.8 Fabaceae Leucaena 100 Commelinaceae Murdannia 92.0 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100		Poaceae	Paspalum	99.9
Bignoniaceae		Anacar diaceae	Loxoptery gi um	98.4
Fabaceae		Asteraceae		
Fabaceae Batesia 98.8 Fabaceae Bussea 100 Fabaceae Bussea 100 Fabaceae Libidibia 99.8 Fabaceae Libidibia 99.8 Fabaceae Pterogyne 100 Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucafyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Migraceae Lecytanthes 100 Migraceae Lecythidaceae Centrolohium 98.2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Migraceae Eclipta 100 Asteraceae Eclipta 100 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 Malvaceae Murdannia 92.0 Malvaceae Sida 99.7 Myrtaceae Eucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucaena 100 Rubiaceae Eucaena 100 Rubiaceae Eucaena 100 Migraceae Eucaena				
Fabaceae Bussea 100 Fabaceae Libidibia 99.8 A. sexdens 1 Fabaceae Pterogyne 100 Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scuachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 Fabaceae Tipuana 100 Lecythidaceae Tipuana 100 Lecythidaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Echinacea 99.6 Asteraceae Echinacea 99.6 Asteraceae Echinacea 99.8 Asteraceae Echinacea 99.8 Asteraceae Echinacea 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100				
A. sexdens 1 Fabaceae Libidibia 99.8 A. sexdens 1 Fabaceae Pterogyne 100 Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lyclanthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 A. sexdens 2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echpta 100 Asteraceae Echpta 100 Asteraceae Felpta 100 Commelinaceae Commelina 100 Commelinaceae Desmodium <td></td> <td></td> <td></td> <td></td>				
A. sexdens 1 Fabaceae Pterogyne 100 Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 A. sexdens 2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.8 Asteraceae Desmodium 99.8 <td></td> <td>_</td> <td></td> <td></td>		_		
Fabaceae Tipuana 99.9 Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebnia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Pterogyne 99.3 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 100 Asteraceae Desymeniopsis 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Synedrella 100 Commelinaceae Synedrella 100 Malvaceae Jida 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Encalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100		_		
Malvaceae Pachira 100 Myrtaceae Eucalyptus 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lyclanthes 100 Bignoniaceae Tabebnia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Lumnitzera 93.4 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 100 Asteraceae Synedrella 100 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Synedrella 100 Malvaceae Jida 99.8 Fabaceae Leucalyptus 99.8 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100 Rubiaceae Genipa 100	A. sexdens 1	_		
Myrtaceae Eugenia 99.6 Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 Fabaceae Tipuana 100 Lecythidaceae Tabaeae Tipuana 100 Lecythidaceae Tapenaa 100 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 100 Asteraceae Desmodium 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Sida 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100			,	
Myrtaceae Eugenia 99.8 Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebnia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 Fabaceae Terogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.8 Asteraceae Echinacea 100 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 Asteraceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100				
Poaceae Scutachne 98.4 Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.8 Asteraceae Echinacea 100 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 100 Commelinaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100		2		
Rubiaceae Genipa 99.2 Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Eclipta 100 Asteraceae Perymeniopsis 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100 <td></td> <td></td> <td>* *</td> <td></td>			* *	
Solanaceae Lycianthes 100 Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 A. sexdens 2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echipta 100 Asteraceae Perymeniopsis 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Commelina 92.0 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100			_	
Bignoniaceae Tabebuia 98.3 Combretaceae Lumnitzera 93.4 Fabaceae Centrolobium 98.2 A. sexdens 2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echinacea 99.3 Asteraceae Perymeniopsis 99.8 Asteraceae Perymeniopsis 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Desmodium 99.8 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100				
Combretaceae				
A. sexdens 2 Fabaceae Centrolobium 98.2 Fabaceae Pterogyne 99.3 Fabaceae Tipuana 100 Lecythidaceae Careya 94.0 Santalaceae Phoradendron 99.6 Asteraceae Echinacea 99.3 Asteraceae Echipta 100 Asteraceae Perymeniopsis 99.8 Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100		/ ·		
A. sexdens 2Fabaceae $Pterogyne$ 99.3Fabaceae $Tipuana$ 100Lecythidaceae $Careya$ 94.0Santalaceae $Pboradendron$ 99.6Asteraceae $Echinacea$ 99.3Asteraceae $Eclipta$ 100Asteraceae $Perymeniopsis$ 99.8Asteraceae $Synedrella$ 100Commelinaceae $Commelina$ 100Commelinaceae $Murdannia$ 92.0 $A. sexdens 3$ Fabaceae $Desmodium$ 99.8Fabaceae $Leucaena$ 100Malvaceae $Sida$ 99.7Myrtaceae $Eucalyptus$ 98.6Phyllanthaceae $Phyllanthus$ 100Rubiaceae $Genipa$ 100			**	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	A sexdens?			
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	000viii0110 <u>/</u>			
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$				
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$				
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$				
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$				
Asteraceae Synedrella 100 Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100				
Commelinaceae Commelina 100 Commelinaceae Murdannia 92.0 A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100			./ /	
A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus Rubiaceae Genipa 100			./	
A. sexdens 3 Fabaceae Desmodium 99.8 Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100				
Fabaceae Leucaena 100 Malvaceae Sida 99.7 Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100	A. sexdens 3			
Myrtaceae <i>Eucalyptus</i> 98.6 Phyllanthaceae <i>Phyllanthus</i> 100 Rubiaceae <i>Genipa</i> 100				
Myrtaceae Eucalyptus 98.6 Phyllanthaceae Phyllanthus 100 Rubiaceae Genipa 100		Malvaceae	Sida	
Phyllanthaceae <i>Phyllanthus</i> 100 Rubiaceae <i>Genipa</i> 100				
,				100
Solanaceae Acnistus 99.4		Rubiaceae	Genipa	100
		Solanaceae	Acnistus	99.4

Figure captions

Figure 1 Grass- and dicot-cutter ants differ in the niches that they occupy, and the way that they cut and process leaf material. Field sites in (A) Botucatu, SP and (B) Ribeirão Preto, SP, Brazil. Fungus gardens of (C) grass- and (D) dicot-cutter ants. C. Visual inspection of leaf material from leaf-cutter ant fungus gardens demonstrates the degree of mulching that the different ants complete, with grass-cutters leaving the leaf material more intact (E -A. bisphaerica and F -A. capiguara), while dicot-cutters mulch to the point of unrecognizable leaf fragments (G -A. laevigata and H -A. sexdens).

Figure 2 NMDS plot of the relative abundance of bacterial genera in fungus gardens of grass- and dicot-cutter ants. Grass- and dicot-cutter fungus garden bacterial communities are significantly different.

Figure 3 Genus-level bacterial community analysis of leaf-cutter ant fungus gardens from grass- and dicot-cutter ants, demonstrating that dicot-cutter ant fungus gardens have a higher diversity of bacteria. A. Pie charts showing proportions of different bacterial genera in the fungus gardens. B. Shannon diversity index of bacterial genera. C. Bacterial genus richness (for genera that consist of more than 1% of the total normalize gene count).

Figure 4 NMDS plot of KO functional genes from grass- and dicot-cutter ant fungus gardens. The KO profiles are significantly different between the fungus gardens of ants cutting the different substrates.

Figure 5 Particular groups of genes are enriched in either the grass- or dicot-cutter ant fungus gardens. Grass-cutter ant fungus gardens are enriched for genes involved in metabolism of terpenoids and secondary metabolites, as well as genes involved in amino acid metabolism. In contrast, dicot-cutter ant fungus gardens are enriched for genes involved in membrane transport. Bars extending to the left (blue) represent genes that are significantly more abundant in dicot-cutter ant fungus gardens and bars extending to the right (red) represent genes that are significantly more abundant in grass-cutter ant fungus gardens.

Figure 6 Iron content of fungus gardens from this study as measured by inductively coupled plasma optical emission spectroscopy. The iron content in the grass-cutter ant fungus gardens was lower than in the dicot-cutter ant fungus gardens. This difference is not statistically significant, however, since the *A. sexdens* fungus garden iron content is highly variable.

References

- Adams AS, Aylward FO, Adams SM *et al.* (2013) Mountain pine beetles colonizing historical and naive host trees are associated with a bacterial community highly enriched in genes contributing to terpene metabolism. *Applied and Environmental Microbiology*, **79**, 3468–3475.
- Adams AS, Jordan MS, Adams SM et al. (2011) Cellulose-degrading bacteria associated with the invasive woodwasp Sirex noctilio. The ISME Journal, 5, 1323–1331.
- Alneberg J, Bengtsson-Palme J, de Bruijn I et al. (2014) Normalization of count data from the metagenomic data sets. Stockholm, Sweden.
- Aylward FO, Burnum KE, Scott JJ et al. (2012) Metagenomic and metaproteomic insights into bacterial communities in leaf-cutter ant fungus gardens. The ISME Journal, 6, 1688–1701.
- Aylward FO, Burnum-Johnson KE, Tringe SG et al. (2013) Leucoagaricus gongylophorus produces diverse enzymes for the degradation of recalcitrant plant polymers in leaf-cutter ant fungus gardens. Applied and Environmental Microbiology, 79, 3770–3778.
- Aylward FO, Suen G, Biedermann PHW et al. (2014) Convergent bacterial microbiotas in the fungal agricultural systems of insects. mBio, 5, e02077–14.
- Bass M, Cherrett JM (1995) Fungal hyphae as a source of nutrients for the leaf □ cutting ant Atta sexdens. *Physiological Entomology*, **20**, 1–6.
- Berasategui A, Salem H, Paetz C et al. (2017) Gut microbiota of the pine weevil degrades conifer diterpenes and increases insect fitness. *Molecular Ecology*, **26**, 4099–4110.
- Boone CK, Keefover-Ring K, Mapes AC *et al.* (2013) Bacteria associated with a tree-killing insect reduce concentrations of plant defense compounds. *Journal of Chemical Ecology*, **39**, 1003–1006.
- Bushnell B (2017) BBMap.
- Cheng X-Y, Tian X-L, Wang Y-S et al. (2013) Metagenomic analysis of the pinewood nematode microbiome reveals a symbiotic relationship critical for xenobiotics degradation. *Nature Publishing Group*, **3**, 237–10.
- Costa AN, Vasconcelos HL, Vieira-Neto EHM, Bruna EM (2008) Do herbivores exert top-down effects in Neotropical savannas? Estimates of biomass consumption by leaf-cutter ants. *Journal of Vegetation Science*, **19**, 849–854.
- Crosa JH (1989) Genetics and molecular biology of siderophore-mediated iron transport in bacteria. *Microbiological reviews*, **53**, 517–530.
- De Fine Licht HH, Schiøtt M, Rogowska-Wrzesinska A et al. (2013) Laccase detoxification mediates the nutritional alliance between leaf-cutting ants and fungus-garden symbionts. *Proceedings of the National Academy of Sciences*, **110**, 583–587.
- Delsuc F, Metcalf JL, Wegener Parfrey L et al. (2013) Convergence of gut microbiomes in myrmecophagous mammals. *Molecular Ecology*, **23**, 1301–1317.
- Ding S-Y, Himmel ME (2008) Anatomy and ultrastructure of maize cell walls: An example of energy plants. In: *Biomass Recalcitrance: Deconstructing the Plant Cell Wall for Bioenergy* (ed

- Himmel ME), pp. 38-60. Blackwell Publishing Ltd, Oxford, UK.
- Douglas AE (2009) The microbial dimension in insect nutritional ecology. *Functional Ecology*, **23**, 38–47.
- Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* (Oxford, England), **26**, 2460–2461.
- Fassel VA, Kniseley RN (1974) Inductively coupled plasma-optical emission spectroscopy. *Analytical Chemistry*, **46**, 1110A–1112A.
- Fowler H, Forti L, Pereira-da-Silva V, Saes N (1986) Economics of grass-cutting ants. In: Fire ants and leaf-cutting ants (eds Lofgren CS, Vander Meer RK), pp. 18–35. Westview Pr, Boulder, CO.
- Gershenzon J, Dudareva N (2007) The function of terpene natural products in the natural world. *Nature chemical biology*, **3**, 408–414.
- Grell MN, Linde T, Nygaard S et al. (2013) The fungal symbiont of Acromyrmex leaf-cutting ants expresses the full spectrum of genes to degrade cellulose and other plant cell wall polysaccharides. BMC Genomics, 14, 928.
- Gutiérrez JL, Jones CG (2006) Physical ecosystem engineers as agents of biogeochemical heterogeneity. *BioScience*, **56**, 227–236.
- Hammer TJ, Bowers MD (2015) Gut microbes may facilitate insect herbivory of chemically defended plants. *Oecologia*, 1–14.
- Hansen AK, Moran NA (2011) Aphid genome expression reveals host-symbiont cooperation in the production of amino acids. *PNAS*, **108**, 2849–2854.
- Hansen AK, Moran NA (2013) The impact of microbial symbionts on host plant utilization by herbivorous insects. *Molecular Ecology*, **23**, 1473–1496.
- Hart AG, Ratnieks FLW (2000) Leaf caching in Atta leafcutting ants: discrete cache formation through positive feedback. *Animal Behaviour*, **59**, 587–591.
- Herz H, Beyschlag W, Hölldobler B (2007) Assessing herbivory rates of leaf-cutting ant (*Atta colombica*) colonies through short-term refuse deposition counts. *Biotropica*, **39**, 476–481.
- Hess M, Sczyrba A, Egan R et al. (2011) Metagenomic Discovery of Biomass-Degrading Genes and Genomes from Cow Rumen. Science, 331, 463–467.
- Hollingsworth PM, Graham SW, Little DP (2011) Choosing and Using a Plant DNA Barcode (D Steinke, Ed,). *PLoS ONE*, **6**, e19254–13.
- Howard JJ (1988) Leafcutting and diet selection: relative influence of leaf chemistry and physical features. *Ecology*, **69**, 250–260.
- Howard JJ, Cazin J Jr., Wiemer DF (1988) Toxicity of terpenoid deterrents to the leafcutting ant *Atta cephalotes* and its mutualistic fungus. *Journal of Chemical Ecology*, **14**, 59–69.
- Hubbell SP, Howard JJ, Wiemer DF (1984) Chemical leaf repellency to an attine ant: seasonal distribution among potential host plant species. *Ecology*, 1067–1076.
- Huntemann M, Ivanova NN, Mavromatis K et al. (2016) The standard operating procedure of the DOE-JGI Metagenome Annotation Pipeline (MAP v.4). Standards in Genomic

- Sciences, 1–5.
- Kanehisa M, Goto S, Sato Y et al. (2014) Data, information, knowledge and principle: back to metabolism in KEGG. Nucleic Acids Research, 42, D199–205.
- Khadempour L, Burnum KE, Baker ES *et al.* (2016) The fungal cultivar of leaf-cutter ants produces specific enzymes in response to different plant substrates. *Molecular Ecology*, **25**, 5795–5805.
- King BC, Waxman KD, Nenni NV et al. (2011) Arsenal of plant cell wall degrading enzymes reflects host preference among plant pathogenic fungi. Biotechnology for biofuels, 4, 4–18.
- Kohl KD, Connelly JW, Dearing MD, Forbey JS (2016) Microbial detoxification in the gut of a specialist avian herbivore, the Greater Sage-Grouse (A Endo, Ed,). FEMS Microbiology Letters, 363, fnw144–6.
- Kohl KD, Weiss RB, Cox J, Dale C, Denise Dearing M (2014) Gut microbes of mammalian herbivores facilitate intake of plant toxins (N van Dam, Ed,). *Ecology Letters*, **17**, 1238–1246.
- Kooij PW, Schiott M, Boomsma JJ, De Fine Licht HH (2011) Rapid shifts in *Atta cephalotes* fungus-garden enzyme activity after a change in fungal substrate (Attini, Formicidae). *Insectes sociaux*, **58**, 145–151.
- Kudo T (2009) Termite-microbe symbiotic system and its efficient degradation of lignocellulose. *Bioscience, Biotechnology, and Biochemistry*, **73**, 2561–2567.
- Langenheim JH (1994) Higher plant terpenoids: A phytocentric overview of their ecological roles. *Journal of Chemical Ecology*, **20**, 1223–1280.
- LeBlanc JG, Milani C, de Giori GS et al. (2013) Bacteria as vitamin suppliers to their host: a gut microbiota perspective. Current opinion in biotechnology, 24, 160–168.
- Li Z, Wright A-DG, Liu H et al. (2014) Bacterial community composition and fermentation patterns in the rumen of sika deer (*Cervus nippon*) fed three different diets. *Microbial Ecology*, **69**, 307–318.
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, **15**, 31–21.
- Mariaca RG, Berger TFH, Gauch R et al. (1997) Occurrence of Volatile Mono- and Sesquiterpenoids in Highland and Lowland Plant Species as Possible Precursors for Flavor Compounds in Milk and Dairy Products. *Journal of Agricultural and Food Chemistry*, **45**, 4423–4434.
- Massey FP, Hartley SE (2009) Physical defences wear you down: progressive and irreversible impacts of silica on insect herbivores. *Journal of Animal Ecology*, **78**, 281–291.
- Mattson WJ (1980) Herbivory in relation to plant nitrogen content. *Annual Review Of Ecology And Systematics*, **11**, 119–161.
- Mayhé-Nunes A, Jaffe K (1998) On the biogeography of Attini (Hymenoptera: Formicidae). *Ecotropicos*, **11**, 45–54.
- Moreira-Soto RD, Sanchez E, Currie CR, Pinto-Tomás AA (2017) Ultrastructural and microbial analyses of cellulose degradation in leaf-cutter ant colonies. *Journal of General*

- Microbiology, 163, 1578-1589.
- Moutinho P, Nepstad DC, Davidson EA (2003) Influence of leaf-cutting ant nests on secondary forest growth and soil properties in Amazonia. *Ecology*, **84**, 1265–1276.
- Muegge BD, Kuczynski J, Knights D *et al.* (2011) Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science*, **332**, 970–974.
- Nagamoto NS, Carlos AA, Moreira SM et al. (2009) Differentiation in Selection of Dicots and Grasses by the Leaf-Cutter Ants Atta capiguara, Atta laevigata and Atta sexdens rubropilosa. Sociobiology, **54**, 127–138.
- Nagamoto NS, Garcia MG, Forti LC *et al.* (2011) Microscopic evidence supports the hypothesis of high cellulose degradation capacity by the symbiotic fungus of leaf-cutting ants. *Journal of Biological Research-Thessaloniki*, **16**, 308–312.
- North RD, Jackson CW, Howse PE (1997) Evolutionary aspects of ant-fungus interactions in leaf-cutting ants. *Trends in Ecology & Evolution*, **12**, 386–389.
- Oksanen J, Blanchet FG, Kindt R et al. (2013) vegan: Community ecology package. CRAN.R-project.org.
- Pinto-Tomás AA, Anderson MA, Suen G et al. (2009) Symbiotic nitrogen fixation in the fungus gardens of leaf-cutter ants. Science, 326, 1120–1123.
- Popper ZA, Tuohy MG (2010) Beyond the Green: Understanding the Evolutionary Puzzle of Plant and Algal Cell Walls. *Plant physiology*, **153**, 373–383.
- R Core Team (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Raffa KF (2013) Terpenes tell different tales at different scales: glimpses into the chemical ecology of conifer bark beetle microbial interactions. *Journal of Chemical Ecology*, **40**, 1–20.
- Rane RV, Walsh TK, Pearce SL *et al.* (2016) Are feeding preferences and insecticide resistance associated with the size of detoxifying enzyme families in insect herbivores? *Current Opinion in Insect Science*, **13**, 70–76.
- Rockwood LL, Hubbell SP (1987) Host-plant selection, diet diversity, and optimal foraging in a tropical leafcutting ant. *Oecologia*, **74**, 55–61.
- Roschard J, Roces F (2003) Cutters, carriers and transport chains: Distance-dependent foraging strategies in the grass-cutting ant *Atta vollenweideri. Insectes sociaux*, **50**, 237–244.
- Shipley LA, Forbey JS, Moore BD (2009) Revisiting the dietary niche: When is a mammalian herbivore a specialist? *Integrative and Comparative Biology*, **49**, 274–290.
- Silva LC, Camargo RS, Lopes JFS, Forti LC (2016) Mandibles of leaf-cutting ants: morphology related to food preference. *Sociobiology*, **63**, 881–8.
- Solomon SE (2007) Biogeography and evolution of widespread leafcutting ants, *Atta* spp. (Formicidae, Attini). University of Texas at Austin.
- Suen G, Scott JJ, Aylward FO, Currie CR (2011a) The microbiome of leaf-cutter ant fungus gardens. In: Handbook of Molecular Microbial Ecology, Volume III: Metagenomics in Different

- Habitats Metagenomics in Different Habitats. (ed de Bruijn FJ), pp. 367–379. John Wiley & Sons, Inc., Hoboken, NJ, USA.
- Suen G, Scott JJ, Aylward FO et al. (2010) An insect herbivore microbiome with high plant biomass-degrading capacity. PLoS genetics, 6, e1001129.
- Suen G, Teiling C, Li L et al. (2011b) The genome sequence of the leaf-cutter ant Atta cephalotes reveals insights into its obligate symbiotic lifestyle. *PLoS genetics*, **7**, e1002007.
- Talbot PHB (1977) The Sirex-Amylostereum-Pinus association. Annual Review of Phytopathology, 15, 41–54.
- Wang Y, DiGuistini S, Wang T-CT, Bohlmann J, Breuil C (2010) *Agrobacterium*-meditated gene disruption using split-marker in *Grosmannia clavigera*, a mountain pine beetle associated pathogen. *Current genetics*, **56**, 297–307.
- Wang Y, Lim L, DiGuistini S et al. (2012) A specialized ABC efflux transporter GcABC-G1 confers monoterpene resistance to *Grosmannia clavigera*, a bark beetle-associated fungal pathogen of pine trees. New Phytologist, 197, 886–898.
- Warnecke F, hl PLU, Ivanova N et al. (2007) Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. Nature Biotechnology, **450**, 560–565.
- Wetterer JK (1994) Nourishment and evolution in fungus-growing ants and their fungi. In: *Nourishment and Evolution in Insect Societies* (eds Hunt JH, Nalepa CA), pp. 309–328. Boulder.
- White TCR (1993) The Inadequate Environment. Springer-Verlag, Berlin.
- Winkelmann G (2002) Microbial siderophore-mediated transport. *Biochemical Society transactions*, **30**, 691–696.
- Winkler JB, Herbst M (2004) Do plants of a semi-natural grassland community benefit from long-term CO2 enrichment? *Basic and Applied Ecology*, **5**, 131–143.
- Wirth R, Beyschlag W, Ryel RJ, Holldobler B (1997) Annual foraging of the leaf-cutting ant *Atta colombica* in a semideciduous rain forest in Panama. *Journal of Tropical Ecology*, **13**, 1–741–75717.
- Wolfe BE, Kuo M, Pringle A (2012) *Amanita thiersii* is a saprotrophic fungus expanding its range in the United States. *Mycologia*, **104**, 22–33.



Grass-cutters

Dicot-cutters







