1 TITLE

Challenging the point neuron dogma: FS basket cells as 2-stage nonlinear integrators

AUTHORS

Alexandra Tzilivaki[1,2] and Panayiota Poirazi[1*]

[1] Institute of Molecular Biology and Biotechnology (IMBB), Foundation for Research
and Technology Hellas (FORTH), Heraklion, Greece

[2] Department of Biology, University of Crete, Heraklion, Greece.

[*] Correspondence: Panayiota Poirazi (poirazi@imbb.forth.gr)

ABSTRACT

Interneurons are critical for the proper functioning of neural circuits and are typically
considered to act as linear point neurons. However, exciting new findings reveal
complex, sub- and/or supralinear computations in the dendrites of various interneuron
types. These findings challenge the point neuron dogma and call for a new theory of
interneuron arithmetic. Using detailed, biophysically constrained models, we predict
that dendrites of FS basket cells in both the hippocampus and mPFC come in two
flavors: supralinear, supporting local sodium spikes within large-volume branches and
sublinear, in small-volume branches. Synaptic activation of varying sets of these
dendrites leads to somatic firing variability that cannot be explained by the point
neuron reduction. Instead, a 2-stage Artificial Neural Network (ANN), with both sub-
and supralinear hidden nodes, captures the variance. We propose that FS basket cells
have substantially expanded computational capabilities sub-served by their non-linear
dendrites and act as a 2-layer ANN.

GABAergic interneurons play a key role in modulating neuronal activity and
transmission in multiple brain regions[1–5]. Among others, they are responsible for
controlling the excitability of both excitatory and inhibitory cells, modulating synaptic

plasticity and coordinating synchrony during neuronal oscillations[2,6–8,9]. GABAergic interneurons come in a variety of molecular profiles, anatomical features and electrophysiological properties[1,3,5,10]. Despite this variability, many interneuron types exhibit similar computations, the most common being a precise EPSP-spike coupling[2,11,12]. As they innervate a large number of cells near the site of action potential initiation[3,13], they are believed to generate a powerful widespread inhibition, also referred to as an inhibitory blanket[13].

Fast Spiking (FS) basket cells constitute one of the main types of hippocampal and neocortical interneurons.[6,13,14] They are distinguished from other subtypes by molecular markers –e.g. the expression of the Parvalbumin (PV) protein-, their anatomical features[15], synaptic connectivity patterns[13,16] and membrane mechanisms such as the presence of calcium permeable AMPA receptors[17,6,18] and the high density of potassium channels in their aspiny dendritic trees[5,14,19,20,6].

A growing body of literature recognizes the importance of FS basket cells in controlling executive functions such as working memory and attention as well as their role in neurodegenerative disorders[4,21,22]. However, little is known about the mechanistic underpinnings of FS basket cell contributions to these functions. Most studies have focused on the molecular and anatomical features of FS basket cells[7,12,15,19,23,24] and supported the dogma that these cells integrate inputs like linear – or at best sublinear- point neurons[25,26].

This dogma is based on the assumption that FS basket cells integrate synaptic inputs in a linear manner, completely ignoring potential dendritic infuences[6]. Active dendritic mechanisms however, are known to transform incoming information in non-trivial ways, thus greatly influencing output signals[27,28,29,30]. Despite its fundamental role in neuronal computations, dendritic integration has been studied mainly in excitatory pyramidal cells[27,31–38]. The current knowledge about FS basket cell dendrites entails a passive backpropagation of APs, low density of sodium channels[6] and high density of fast, high-threshold potassium channels in distal dendrites[6,39,40]. All of the above contribute to sublinear dendritic integration, coupled to fast and temporally precise AP initiation in response to synaptic input[12,25], largely in line with the simplified point-neuron view of interneurons.

Exciting new findings however, reveal that the dendrites of certain interneuron types

are much more powerful than originally assumed. For example, sublinear dendritic

EPSP integration along with supralinear calcium accumulations has been reported in

cerebellar Stellate Cells (SCs)[11,41].Moreover, RAD dendrite-targeting interneurons in

the CA1 area exhibit calcium supralinearities[42] while in the CA3, both calcium

nonlinearities and sodium spikes in FS basket cell dendrites during sharp wave

ripples, have been reported[2]. The exact nature of dendritic computations in FS basket

cells, however, is unknown. As a result, whether a linear point neuron or a more

sophisticated abstraction -like the two-stage[32] or multi-stage integration[43] proposed

for pyramidal neurons- can successfully capture their synaptic integration profile,

remains an open question.

To address this question, we developed detailed, biologically constrained biophysical

models of FS basket cells using anatomical reconstructions of both hippocampal [44]

and cortical (medial Prefrontal Cortex) neurons[45] (shown in figure 1). Synaptic

stimulation within the dendrites of model cells predicts the co-existence of two

distinct integration modes; some dendrites exhibit supralinear synaptic integration

while others operate in a sublinear mode (figure 2 and supplementary figures 4,5).

Morphological features such as dendritic length and/or diameter influence the

integration mode and these features differ between hippocampal and cortical neurons.

Interestingly, dendritic volume appears to be a consistent discriminating feature

among sub- and supralinear dendrites of both areas (figure 3). By generating a large

number of different spatial patterns of synaptic activation we find that spatially

dispersed inputs lead to higher firing rates than inputs clustered within a few dendrites

in both Hippocampus and PFC models (figure 4), opposite to respective simulations

in pyramidal neurons[46]. Moreover, a 2-layer Artificial Neural Network (ANN) with

both sub- and supralinear hidden nodes can predict the firing rate of the

aforementioned models much better than a linear ANN (figures 5, 6, Table 1,

Supplementary figure 7).

This is the first study that provides a systematic, cross-area analysis of dendritic

integration in FS basket cells. Our findings challenge the current dogma, whereby

interneurons are treated as linear summing devices, essentially void of dendrites. We

predict that the dendrites of FS basket cells in both Hippocampal and Neocortical

regions can operate in distinct non-linear modes. As a result, FS basket cells, similar

to pyramidal neurons[32], are better represented by a 2-stage integrator abstraction
rather than a point neuron.

Figure 1: Fast spiking basket cell reconstructions from 5 Hippocampal and 3 PFC
interneurons. Dendrites are shown in red. Firing patterns in response to 200 pA current
injection at the cell body are also shown for each morphology.

RESULTS

The models

A total of 8 biophysical model neurons were built within the NEURON simulation

environment[47]. We used realistic reconstructions of FS basket cells from the rat

hippocampus (5 cells)[44] and from the mPFC (3 cells)[45] of mice (figure 1). All

morphologies were downloaded from the Neuromorpho.org database and checked for

reconstruction accuracy (diameter) (See Online Methods). To ensure biological

relevance, ionic and synaptic conductances as well as basic membrane properties of

model cells were heavily validated against experimental data[6,12,15,19,40,48] (see

Supplementary Table 1-4 and Supplementary figures 1-3). Moreover, for consistency

reasons, the same set of biophysical mechanisms (type and distribution) was used in

all model cells. This is to our knowledge the first set of detailed, biologically realistic

models of FS basket cells from two brain areas.

Bi-modal dendritic integration in Fast Spiking Basket cells

The first step for deducing a realistic abstraction of FS basket cells is the systematic

characterization of dendritic/neuronal integration properties across a significant

number of neurons and dendrites. Towards this goal, we simulated gradually

increasing excitatory synaptic input to the dendrites of all neuronal models (total of

637 simulated dendrites) and recorded the voltage response both locally (figure 2) and

at the soma (Supplementary figure 5)[11,31]. Increasing numbers of synapses (1:1:20)

were uniformly distributed in each stimulated dendrite and activated synchronously

with a single pulse. Sodium conductances in somatic and axonal compartments were

set to zero, to avoid AP backpropagation contamination effects. We compared

measured EPSPs to their linearly expected values, given by the number of activated

inputs multiplied by the unitary EPSP. We found that within the same dendritic tree,

branches summate inputs either in a supralinear or a sublinear mode (figure 2,

supplementary figures 4,5). While there were differences in the number of dendrites

and proportions of sub- vs. supralinear dendrites, all of the morphologies tested

expressed both integration modes (Supplementary Table 5). Importantly, while both

modes have been suggested in distinct interneuron types[11,42], this is the first study that

predicts their co-existence within a single cell.

Figure 2: Bimodal dendritic integration in FS basket cell models. Representative recordings from supralinear (A,C) and sublinear (B,D) dendritic branches in Hippocampal (top) and PFC (bottom) biophysical model interneurons, in response to synaptic stimulation. Increasing numbers of synapses (1:1:20) were uniformly distributed within each activated branch and activated with a single pulse. The y-axis shows the actual peak depolarization caused by synaptic activation while the x-axis shows the expected peak depolarization that would result from the linear summation of unitary EPSPs. The dashed line indicates linearity.

To assess the robustness of this finding, we performed a sensitivity analysis whereby the cp-AMPA, NMDA, VGCCs, sodium and IA potassium conductances were varied by ±20% of their control value. We found no changes in the integration mode of dendrites (data not shown) and only insignificant alterations in the spike threshold of supralinear dendrites (Supplementary figure 6). These simulations suggest that under physiological conditions, FS basket cells are highly likely to express two types of dendritic integration modes.

Morphological determinants of dendritic integration modes

Morphological features of dendrites were previously shown to influence synaptic integration profiles[49]. We thus investigated whether specific anatomical features correlate with the expression of each integration mode. We found that the mean dendritic diameter was highly statistically different (p-value=4.1966e-54) among sub- (thinner) and supra-linear (thicker) dendrites in the hippocampus (figure 3C) while in

the PFC the dendritic length was a better determinant of sub- (shorter) vs. supra-
linearity (longer) (p-value=7.6543e-05) (figure 3B). Length was less significant in the
hippocampus (p-value=0.0064) (figure 3A) while diameter was not different among
sub- and supralinear dendrites in the PFC (p-value=0.2454) (figure 3D). Dendritic
volume considers both of the above features and serves as a robust morphological
determinant for all dendrites in both areas (p-value=1.8433e-09), (figure 3E). These
findings predict that morphology plays a crucial role in the spiking abilities of FS
basket cell dendrites (figure 3F).

Figure 3: Morphological determinants of dendritic integration modes. A-B: Total length
(μm) distributions of supralinear vs. sublinear dendrites in the hippocampus (A) and the PFC
(B). Statistically significant differences are observed for both sub- and supra-linear models, in
both areas. Significance is higher in PFC cells (p-value=7.6543e-05, 130 dendrites analyzed)
compared to Hippocampal (p-value=0.0064, 507 dendrites analyzed) FS basket cells. C-D:
Same as in A-B, for mean dendritic diameter (μm). Statistically significant differences are
observed in Hippocampal (p-value=4.1966e-54, 507 dendrites analyzed) but not in PFC FS
basket cells (p-value=0.2454, 130 dendrites analyzed). E. Dendritic Volume (μm^3) is a
common discriminating characteristic among supralinear (larger) and sublinear dendrites, for

both areas (p-value= p-value=1.8433e-09, 637 dendrites analyzed). F. Schematic illustration
of distinctive morphological features for supralinear and sublinear dendrites in Hippocampus
(left) and PFC (right).

Effect of bimodal dendritic integration on neuronal firing

In light of a bimodal dendritic integration in the modeled FS basket cells, the natural
question that arises concerns the functional implications of such a property. To
answer this question, we simulated a large variety of different spatial patterns of
synaptic activation and assessed their effect on neuronal output. Specifically, we
generated over 10,000 synaptic stimulus patterns, which comprised of 0 to 60
excitatory synapses (activated with random Poisson spike trains at 50 Hz) distributed
within a few, strongly activated branches (clustered) or randomly distributed within
the entire dendritic tree (dispersed).

Several stimulation protocols were devised, in which different numbers of synapses
were activated in various locations within the dendritic tree (see Online Methods).
Dendrites were selected at random and inputs were distributed uniformly within
selected dendrites. For the dispersed case, we allocated 5 or 10 synapses in randomly
selected dendrites, one at a time, while for the clustered case we allocated 10 or 20
synapses within an increasing number of branches. In all cases, the number of
activated synapses increased gradually up to a maximum of 60, as this number was
sufficient to induce spiking at gamma frequencies (30-100 Hz). This process was
repeated k times (k = number of dendrites in each cell) to ensure full coverage of the
entire tree. As expected given the two modes of dendritic integration, the localization
of activated inputs affected neuronal firing. For a given number of activated synapses,
dispersed activation led to higher somatic firing rates than clustered activation, both in
Hippocampal (p-value=2.0914e-21, figure 4A) as well as in PFC FS basket cells (p-
value=0.0051, figure 4B). Interestingly, this finding is opposite to what has been
reported for pyramidal neurons, in which synapse clustering increases firing rates[46].

Figure 4: Effect of bimodal dendritic integration on neuronal firing. Firing rate responses (in Hz) from one Hippocampal (A) and one PFC (B) model cell, in response to stimulation of increasing numbers of synapses (10 to 60) that are either randomly distributed throughout the entire dendritic tree (blue) or clustered within a few dendritic branches (pink).) Synapses are stimulated with a 50 Hz Poisson spike train. Indicative somatic traces in response to stimulation of 30 synapses are shown for the four cases. Red dots represent the synaptic allocation motif.

FS basket cells as 2-layer artificial neural networks

The non-linear synaptic integration taking place within the dendrites of cortical[33] and CA1[31,34] pyramidal neurons was previously described as a sigmoidal transfer function[50]. Based on this reduction, a single pyramidal neuron was proposed to integrate its synaptic inputs like a 2-layer artificial neural network, where dendrites provide the hidden layer and the soma/axon the output layer[32]. To assess whether a similar mathematical formalization could be ascribed to our FS basket cell models, we constructed linear and non-linear artificial neural networks (as graphically illustrated in figure 5) and asked which of them can better capture the spike variance of the biophysical models.

Figure 5: Reducing detailed compartmental models into mathematical abstractions. Two
types of abstractions were used: a) a Linear ANN, in which the input (number of synapses)
was linearly combined at the cell body and b) a 2-layer modular ANN, in which the input was
fed into two parallel, separated hidden layers. The supralinear-layer was fed with the number
of inputs landing onto supralinear branches while the sublinear layer was fed with the number
of inputs landing onto sub-linear dendrites. Neurons in both hidden layers were equipped with
nonlinear transfer functions, a step-sigmoid function in the supralinear layer and a saturating
linear function in the sublinear layer. The somatic transfer functions of both ANNs were
linear.

Specifically, four types of feedforward, backpropagation Artificial Neural Networks

(ANNs) were constructed (see Online Methods). In the *2-layer modular ANN*,

supralinear and sublinear dendrites were simulated as 2 parallel hidden layers

consisting of a step-sigmoidal and a saturating linear activation function,

respectively[51] (figure 5). The total number of activated synapses allocated to

supralinear and/or sublinear dendrites in the biophysical models was used as input to

the respective hidden layers. The output layer represented the soma/axon of the

biophysical model and consisted of a linear activation function. In the *linear ANN*,

there was only a single hidden layer consisting of linear activation functions (figure 5). We also constructed two ANNs with the exact same architecture as the linear one, but with either a) a step-sigmoidal (*2-layer supralinear ANN*) or b) a saturating linear (*2-layer sublinear ANN*) activation function in the hidden layer neurons (Supplementary figure 7). These ANNs represent FS basket cells with just one type of non-linear dendrites. The free parameters in all networks were identical (Supplementary Table 6).

For a given hippocampal and a given mPFC biophysical model cell, the *linear* and *2-layer modular ANNs* were trained using the number of synapses to supra-/sublinear dendrites as inputs to the respective hidden layers and the mean firing rate of the soma as target output. A randomly selected 70% of our synaptic activation data set (See Online Methods) was used to train the model and the rest to assess its generalization performance (15% Validation, 15% testing). Performance accuracy was estimated based on regression analysis between the ANN-generated firing rates and those produced by the biophysical models. The *2-layer modular ANN* reached an average performance accuracy of 96% and 95% (figure 6A, C) in predicting the spike rate variance in hippocampal and PFC models, respectively, while the *linear ANN* captured 85% and 75% of the spike rate variance, respectively (figure 6B, D). As expected, the *supralinear* and *sublinear ANNs* achieved intermediate accuracies for both hippocampal: 91%, 92% and PFC 90.8%, 92% models, indicating that both types of non-linear transfer functions are needed to capture the biophysical model variability (see Supplementary figure 7).

The relatively high performance of the *linear ANN* can be attributed to the wide range of activated synapses (2 to 60) which resulted in large differences in the somatic firing, irrespectively of synapse location, and can thus be captured by any linear model (also see[32]). To perform a fairer comparison, we also assessed the performance accuracy of *linear* and *2-layer modular ANNs* to the more challenging task of discriminating between input distributions corresponding to the exact same number of synapses. To do so, we subdivided the data into input categories corresponding to 20, 40 and 60 synapses, respectively. In this case, the *2-layer modular ANN* clearly outperformed the *linear ANN*, which failed to explain the variance produced by differences in input location (Table 1).

Taken together, this analysis suggests that a 2-layer artificial neural network that considers both types of dendritic non-linearities is a much better mathematical abstraction for FS basket cells than the currently assumed linear point neuron.

Figure 6: Challenging the point neuron dogma: FS basket cells as 2-stage nonlinear integrators. Linear regression analysis for 2-layer modular (A,C) and linear (B,D) ANNs for one indicative Hippocampal (top) and one indicative PFC (bottom) model cell. Actual Mean Firing Rates (Hz) correspond to the responses of the compartmental model when stimulating - with 50Hz Poisson spike trains- varying numbers of synapses (1 to 60), distributed in several ways (clustered or dispersed) within both sub- and supra-linear dendrites. Expected Mean Firing Rates (Hz) are those produced by the respective ANN abstraction when receiving the same input (number of stimulated synapses) in its respective sub-/supra- or linear input layer nodes.

Table 1. ANN regression performance (R) for individual sets of synapses			
ANN type	*20 synapses*	*40 synapses*	*60 synapses*
2-layer modular ANN	**0.8994(HPC)/ 0.8750 (PFC)**	**0.9508 (HPC) / 0.8869 (PFC)**	**0.9406(HPC) /0.8399(PFC)**
Linear ANN	0.4862 (HPC) / 0.4867 (PFC)	0.6296 (HPC) / 0.5527 (PFC)	0.6801 (HPC) / 0.5136 (PFC)
2-layer supralinear ANN	0.6486 (HPC) / 0.7468 (PFC)	0.8172 (HPC) / 0.7995 (PFC)	0.8474 (HPC) /0.6633 (PFC)
2-layer sublinear ANN	0.7645 (HPC) / 0.8130 (PFC)	0.8816 (HPC) / 0.8487 (PFC)	0.8617 (HPC) /0.7674 (PFC)

Table 1: Comparison of ANN prediction accuracy (measured as the correlation coefficient, R) for all four ANN reductions, when tested on three sets of synaptic inputs consisting of 20, 40 or 60 activated synapses, respectively. Synapses were randomly distributed in various ways/locations in the biophysical model cells and resulting firing rates were used as target vectors for the ANNs. The 2-layer modular ANN is clearly superior to the Linear ANN when it comes to capturing location-induced firing-rate variability.

DISCUSSION

The role of dendrites in interneuron computations is a rapidly emerging and debatable subject[39]. Several recent reports present exciting findings according to which dendrites may serve as key players[2,11,41,42,52]. For example, sodium spikes and supralinear calcium accumulation have recently been reported in the dendrites of FS basket cells[2,39,53], yet the consensus still favors the linear point neuron dogma[6,39,54]. The present study provides new insight into this ongoing debate by systematically analyzing the dendritic integration mode of FS basket cells in two widely studied areas: the Hippocampus and the PFC. We predict that dendrites of both cortical and hippocampal FS basket cells operate in one of two modes of synaptic integration: supralinear or sublinear (figure 2). Supralinearity is due to the generation of dendritic sodium spikes, and can be facilitated –or prohibited as in sublinear dendrites- by the morphology (diameter, length, volume, (figure 3)) of dendrites. Moreover, we find that somatic output is influenced by the spatial distribution of activated synapses, with dispersed stimulation inducing higher firing rates than clustered stimulation (figure 4). Due to these properties, a 2-layer Artificial Neural Network abstraction with both sub- and supra-linear hidden neurons captures the spiking profile of biophysical neurons with much higher accuracy compared to a linear ANN, analogous to a point neuron (figure 5, 6). These findings suggest that the dendrites of FS basket cells in both the

313   hippocampus and the cortex can support two types of non-linear computations and are
314   the first to explicitly challenge the point neuron dogma.

Mediators of supralinear and sublinear dendritic integration in FS basket cells

A bimodal dendritic integration is predicted for all hippocampal and PFC
morphologies analyzed. In all cases, supralinearity is due to the occurrence of
dendritic sodium spikes. Several mechanisms can influence the generation of such
dendritic spikes: ionic conductances (primarily of sodium currents but also potassium
currents) and morphological features. In our models, biophysical mechanisms are
constrained by existing experimental data and dendritic sodium conductances are kept
to a minimum (10 times smaller than the soma[6]), so as to minimize the probability of
non-physiological dendritic spiking. Sensitivity analysis further demonstrates that
results are robust to physiological variations in a wide range of dendritic
conductances. These findings strongly suggest that dendritic spiking in certain
dendrites of FS basket cells are highly likely to occur under physiological conditions,
in line with recent experimental reports[2].

Apart from sodium currents as a universal enabling mechanism, we find a key role of
morphology in gating local dendritic spikes. A combination of dendritic length and
mean diameter, or otherwise the dendritic volume, is statistically different between
sub- (smaller) and supralinear (larger) dendrites across all morphologies tested. These
results are in line with other studies reporting a similar effect of morphology on the
ability of dendrites to generate local spikes[55].

Functional coexistence of sub- and supra-linear dendrites within FS basket cells

Our simulations predict the co-existence of both sublinear and supralinear dendrites in
all simulated FS basket cells (figure 2, Supplementary figures 4,5). Similar bimodal
dendritic integration has been reported in hippocampal CA1 pyramidal neurons[31,34]
and predicted in PFC pyramidal neurons[33]. However, the functional consequences of
this coexistence in interneurons requires further investigation.

The existence of sublinear dendritic branches supports the idea of inhibitory neurons
acting as coincidence detectors by aggregating spatially disperse and nearly
synchronous synaptic inputs[6]. Moreover, sub-linear dendrites can compute complex
non-linear functions similar to those computed by sigmoidal dendrites[50], thus

substantially extending the processing capacity of these neurons compared to a linear

integrator. Why have two types of nonlinearity then?

One possibility is to enable the detection of few but highly correlated inputs: via

spatial clustering onto supralinear dendrites these inputs would reliably induce

dendritic spikes capable of overcoming the dampening effects of inhibitory

conductances, thus generating strong somatic responses. Another possibility entails

increases in flexibility through the ability to (a) engage intrinsic plasticity

mechanisms (e.g. regulation of potassium channels) and/or (b) to dynamically tune

the neuronal operation mode from generic (sublinear domination) to specific

(supralinear domination), depending on the behavioral state. As dendrites of FS

basket cells often cross layers and receive input from different afferent pathways[56],

another possibility is that feedback vs. feedforward pathways target dendrites with

distinct modes of integration. These scenarios can be tested in future studies engaging

network models and/or experimental probing.

Not that Simple: FS basket cells as 2-layer modular ANNs

Artificial Neural Network analysis demonstrates that a FS basket cell is better

described by a 2-stage abstraction, which takes into account both modes of dendritic

integration. This work, along the lines of the 2-stage model proposed for pyramidal

neurons[32], strongly challenges the prevailing point neuron dogma. The 2-stage

abstraction is supported by experimental reports of dendritic sodium spikes and

supralinear calcium accumulations[2] while it also explains sublinear dendritic

integration[6,11,26,57], providing a unifying framework for interneuron processing.

Possible limitations of our work include the imprecise modeling of ionic and synaptic

mechanisms given the shortage of sufficient information for FS basket cells. This

limitation is counteracted by the sensitivity analysis of the mechanisms that mostly

influence our findings and their consistency across several cortical and hippocampal

morphologies. Another limitation is the lack of inhibitory inputs (except from the

autaptic GABAa current that is incorporated in all models) and gap junctions on our

model cells. Inhibitory inputs consist of just 6% of all incoming contacts in Fast

Spiking interneurons[6,44,58]. Thus, our results are unlikely to be affected by inhibitory

inputs. FS basket cells in the hippocampus and the neocortex are highly

interconnected by gap junctions[6], that can speed the EPSP time course, boost the

efficacy of distal inputs and increase the average action potential frequency after

repetitive synaptic activation.[6] All of these effects would contribute to stronger

responses but unless gap junctions are spatially specific to certain branches and not

others, they are unlikely to influence the non-linear integration modes of dendrites.

Conclusion

This work provides a novel view of dendritic integration in FS basket cells, that

extends in hippocampal and cortical areas[59]. To our knowledge, we are the first to

suggest a new reductionist model for interneuron processing, in which dendrites play

a crucial role. Experimental validation of this new model is likely to change the way

we think about interneuron processing, attribute new and exciting roles to FS basket

cells and open new avenues for understanding interneuron contributions to brain

function.

Online Methods

Simulations were performed on a High-Performance Computing Cluster equipped

with 312 CPU cores and 1.150 Gigabytes of RAM, under 64-bit CentOS 6.7 Linux.

The source code will be publicly available in ModelDB upon acceptance for

publication.

Compartmental modeling

All 8 model neurons were implemented within the NEURON simulation environment

(version 7.3)[47]. Detailed morphological reconstructions of the 5 Fast spiking Basket

cells of the rat Hippocampus were adopted from Tukker et al. 2013[44], via the

NeuroMorpho.org database (figure 1). Due to the lack of axonal reconstructions, we

used the axon from the B13a.CNG.swc reconstruction for all 5 hippocampal neuron

models. The 3 PFC morphologies were adopted from Rotaru et al. 2011[45], via the

NeuroMorpho.org database (figure 1) and included their respective axons.

Dendritic branches with mean diameter values larger than 1.2 μm were excluded from

all simulations and data analysis procedures, based on Emri et al. 2001[60]. The NLM

Morphology Viewer Software was used to transform morphological reconstructions

into .hoc files.

Biophysical properties

All model neurons were calibrated with respect to their biophysical properties so as to

conform to experimental data. The same active and passive properties were used in all

model cells, with the exception of very small modifications in the conductances of

somatic/axonal Sodium and Kdr mechanisms (Supplementary Tables 1 and 2). The

latter were necessary to account for the influence of morphological variability on

neuronal responses.

Conductances of all active ionic mechanisms were adapted from Konstantoudaki et al.

2014[61]. Both hippocampal and PFC models include fast voltage-dependent sodium

channels (g_{nafin}), delay rectifier potassium channels (g_{kdrin}), slow inactivation

potassium channels (g_{slowin}), slow calcium dependent potassium channels (g_{kcain}), A-

type potassium channels for proximal and distal dendritic regions (g_{kadin}, g_{kapin}), h

currents (g_{hin}), and L-, N- and T- type voltage-activated calcium channels (g_{cal}, g_{can}

and g_{cat}, respectively).   Sodium current conductances were substantially larger in

axonal compared to somatic compartments, which were in turn an order of magnitude

larger than dendritic sodium conductances[6]. Moreover, dendritic branches located

beyond 100 microns from the soma (distal dendrites) had smaller sodium

conductances than proximal branches (located less than 100 microns from the soma)

as per[6,40]. A calcium buffering mechanism was included in all compartments. Details

about all biophysical mechanisms are listed in Supplementary Table 2.

Synaptic conductances

Calcium permeable (GluR2-lacking) AMPA, NMDA and autaptic GABAa synaptic

currents were incorporated in all model cells. Synaptic weights were validated so as to

reproduce the current waveforms depicted in Wang Gao et al 2009[62] and Bacci et al

2003[19] and are shown in Supplementary Table 4 and Supplementary figure 2.

Electrophysiological validation

All model neurons were heavily validated against experimental data in order to ensure

biological plausibility. Averaged electrophysiological values for the model cells and

respective experimental values are shown in Supplementary Table 3.

Bi-modal dendritic integration in Fast Spiking Basket cells

To map the dendritic integration profiles of our model neurons, we activated

increasing numbers of synapses (1 to 20, with step 1) in each dendrite and recorded

the voltage responses both locally and at the cell body for 100 ms. Synaptic input was

simulated as a single depolarizing pulse, as per Poirazi et al 2003a[31]. Sodium

conductances in somatic and axonal compartments were set to zero in order to avoid

backpropagation effects. 12 autaptic events were also activated in somatic

compartments[63].

Integration modes were deduced by comparing the measured dendritic/somatic

responses (Actual maximum EPSP) against what would be expected if synaptic

depolarizations summed linearly (Expected maximum EPSP). A dendrite was termed supralinear if Actual responses were larger than Expected, even if this was true only for a short range of synaptic inputs. A dendrite was considered sublinear if the Actual EPSPs were smaller than the Expected values for all synaptic inputs tested.

Sensitivity analysis was performed by modifying the conductances of NMDA, calcium-permeable AMPA receptors, Voltage gated Calcium Channels (VGCCs), Sodium, and A-type (proximal and distal) mechanisms by ± 20%. Increasing numbers of synapses (for 1 to 40) were used to assess possible changes in single branch integration. Results for all manipulations are shown in Supplementary figure 4.

Morphological determinants of dendritic integration mode

Mean dendritic diameter and total dendritic length for supralinear and sublinear dendrites were measured for all reconstructed neurons. Dendritic volume was calculated according to the following formula:

$$V = \left(\pi * \left(\frac{diam}{2} \right)^2 \right) * length \quad (\mu m^3)$$

Statistical analysis for all morphological features was performed using Student's t-test with equal sample sizes and assuming unequal variances (Welch's t-test).

Synaptic Stimulation Protocols

In all stimulation protocols, inputs were activated using a 50 Hz Poisson spike train. The maximum number of activated synapses was 60, as it was sufficient to induce firing at gamma related frequencies (30-100 Hz). The spatial arrangement of activated synapses was defined according to each of the following stimulation protocols:

Randomly dispersed, whole tree stimulation

Different numbers of synapses (N_{sysn} = 5, 10 up to 60) were randomly placed on randomly selected dendrites. For a given number of synapses N_{syn}, at each allocation step, one dendrite was chosen at random and one synapse was placed at a random location within this dendrite. For the selected dendrite, synaptic location was randomly changed 5 times. This process was repeated N times, where N was the number of dendrites for each model cell. This process ensured a realistic distribution of activated synapses within the entire dendritic tree of each modelled neuron.

Clustered, whole tree stimulation

The only difference of this protocol from the one described above is that each selected dendrite received a cluster (of size S_{clu} = 10 or 20) of synapses as opposed to a single synapse. For example, for N_{syn}=60 and S_{clu} = 10, a total of 6 dendrites were randomly selected to receive 10 synapses each. We followed the same experimental design as in the disperse, whole tree protocol. Thus, for a given number of synapses N_{syn}, at each allocation step, one dendrite was chosen at random and one synapse was placed at a random location within this dendrite. For the selected dendrite, synaptic location was randomly changed 5 times. This process was repeated N times, where N was the number of dendrites for each model cell.

Artificial Neural Network Models

We constructed four feed-forward, backpropagation Artificial Neural Networks with customized code written in MATLAB, version 2009: a) a 2-layer modular ANN whereby hidden neurons were divided in two parallel layers (1 & 2). In hidden layer

1, neurons consisted of supralinear (step-sigmoid) transfer functions while neurons in

hidden layer 2 consisted of sublinear (saturating linear) transfer functions. The two

types of transfer functions corresponded to respective supra- and sublinear dendrites

of the biophysical model cells[50]. b) a 2-layer ANN with one hidden layer, whereby all

hidden neurons had supralinear transfer functions, c) a 2-layer ANN with one hidden

layer, whereby all hidden neurons had sub-linear transfer functions and d) a linear

ANN whereby the hidden and output neurons had a linear transfer function (classical

linear point neuron). In all ANNs, the output neuron had a linear transfer function. All

ANNs were trained with the same input/output data sets and performance accuracy

was estimated according to the correlation among predicted (by the ANN) and actual

mean firing rates generated by the biophysical model, for a wide range of stimulation

protocols. The parameters of each ANN are listed in **Supplementary Table 6**. Firing

rate corresponding to this particular configuration.

ANN training/testing datasets:

Input to the four ANNs consisted of the number of synapses located in the modelled

dendritic branches and the target output consisted of the mean firing rate generated by

the biophysical model in response to synaptic stimulation. In the biophysical model,

these synapses were activated with the Dispersed and Clustered protocols described

above as well as five new protocols using the same pattern of repetition trials as

described above: 1) Randomly dispersed activation of synapses (Nsyn=2:2:60) in the

entire dendritic trees. 2,3) Clustered (Sclu=3, Nsyn=20) or 4,5) Dispersed (Nsyn 10)

synaptic allocation on supralinear dendrites and Clustered or Dispersed synaptic

allocation on sublinear dendrites, respectively. Data shown in Figure 5 and

Supplementary Figure 6 correspond to PFC3 and HIPP2 model neurons and are

representative of all model cells.

Linear Regression and Statistical Analysis.

We calculated the correlation coefficient (R) between Actual Mean Firing Rates

(Target rates, in Hz) generated from the biophysical models and Predicted Mean

Firing Rates (Predicted rates, in Hz) generated by the trained ANNs, respectively.

Statistical analysis between Target and Predicted firing rates was performed using

Student's t-test with equal sample sizes and assuming unequal variances (Welch's t-

test).

ACKNOWLEDGMENTS

This work was supported by the ERC Starting Grant dEMORY to Panayiota Poirazi,

by the Google Europe Scholarship to Alexandra Tzilivaki and by the Alexander C.

Onassis Benefit Foundation Scholarship to Alexandra Tzilivaki.

The authors would like to thank Stefanos Stamatiadis for providing valuable help on

simulation procedures and all the members of the Computational Biology Lab for

helpful discussions.

AUTHOR CONTRIBUTIONS

PP and AT designed the experiments. AT performed the simulations and analyzed the

data. AT and PP wrote the manuscript. PP supervised the project.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Supplementary Information

Table S1. Passive properties of biophysical models				
	Soma	*Axon*	*Proximal dendrites*	*Distal Dendrites*
Leak conductance (*g_pas*)[12]	1.315e-4 S/cm^2	3.55e-6 S/cm^2	1.315e-4 S/cm^2	1.34e-5 S/cm^2
Resting Membrane Potential (e_pas)[12,19]	-68 mV	-68 mV	-68 mV	-68 mV
Membrane capacitance (*cm*) [12]	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}
Axial Resistance (*Ra*) [12]	172 ohm/ cm	172 ohm/ cm	142 ohm/ cm	142 ohm/ cm

554 Supplementary Table S1: Passive properties common to the 8 biophysical models

Table S2. Active properties of biophysical models				
Ion channel (S/cm^2)	*Soma*	*Axon*	*Proximal dendrites*	*Distal Dendrites*
Na_v [6,26,64]	0.145(PFC1-3) /0.396 (Hipp1-3) /0.828(Hipp4,5)	0.675(PFC1-3) /1.296(Hipp1-3) /1.512(PFC4,5)	0.018	0.014
H_v	0.00001	X	x	X
Kdr_v	0.036 (PFC)/ 0.0432(Hipp)	0.108 (PFC)/ 0.144 (Hipp)	0.0009	0.009
$Kslow_v$	0.000725	X	x	X
Kct_v	0.0001	X	x	X
Kca_v	0.02	X	x	X
Ka_v (proximal type) [64]	0.0032	X	0.001	0.0009
Ka_v (distal type) [48]	x	X	x	0.00216
Cal_v	x	X	0.00003	0.00003
Can_v	x	X	0.00003	0.00003
Cat_v	x	X	0.0002	0.0002
Calcium buffering dynamics	Yes	No	Yes	Yes

556 Supplementary Table S2: Active membrane conductances across somatic, axonal, proximal
557 (<=100 microns from the soma) and distal (>100 microns from the soma) dendritic
558 compartments. Sodium current conductances are larger in axonal than somatic compartments.
559 Dendritic sodium conductances are ~10 times smaller than axo-somatic conductances.

Table S3. Synaptic mechanisms of biophysical models	
Synaptic Current	Conductance Weight
Autaptic GABAa	71.1e-4
Ca permeable AMPA	7,5e-4
NMDA	16e-4

562 Supplementary Table S3: Validated Synaptic conductance weight values of Autaptic GABAa

563 Calcium permeable AMPA and NMDA currents, used in all simulations .

Table S4. Electrophysiological properties of biophysical models		
Value	*Model*	*Experimental data*
rHeobase *(pA)*[15]	150.0 ± 30.0	123 ± 58
Input Resistance (Ohm)[15]	97.7 ± 30.0	182 ± 83
Spike threshold (mV)[15]	-37.0 ± 3.0	-34 ± 2
Spike amplitude (mV)[15]	52.0 ± 2.0	53.0 ± 8.0
Spike half width (msec)[15],[19]	0.5 ± 0.1	0.2 ± 0.01/0.38 ± 0.008
f-i slope (Hz/pA)[19]	0.26 ± 0.03	0.2 ± 0.002
AHP (mV)[19]	24.6 ± 2.6	24.3 ± 0.7/23 ± 5

566 Supplementary Table S4: Validation of electrophysiological properties.

Table S5. Nonlinearity distributions		
Cell ID	Number of supralinear dendrites	Number of Sublinear dendrites
Hipp 1	95	67
Hipp 2	13	38
Hipp 3	10	41
Hipp 4	89	98
Hipp 5	27	32
PFC 1	36	7
PFC 2	48	5
PFC 3	43	14

567 Supplementary Table S5: Number of supralinear and sublinear dendrites per cell.

Table S6. ANN properties	
Train function	Levenberg-Marquardt backpropagation
Error Function	Mean squared normalized error performance
Number of epochs	1000
Type of network	feedforward

569 Supplementary Table S6: Network parameters of all ANNs.

  Supplementary Figures

Supplementary Figure 1. Model cell firing profiles. Somatic Current-clamp traces of
Hippocampal (A) and PFC (B) model cells, after a depolarizing current injection in somata
(500 pA; 1000 ms) evoked a high-frequency firing pattern. A hyperpolarizing current
injection in somata (-300pA, 1000ms) induced a realistic hyperpolarizing response.

Supplementary Figure 2: Mean firing frequencies in response to injected currents of
different amplitudes (600 ms duration) in Hippocampal (A) and PFC (B) model cells.

Supplementary Figure 3: Validation of synaptic currents in Fast Spiking basket cells.
A,C: A three-step voltage clamp of voltage changes from −70 mV to 10 mV (duration 1 ms)
and back to −70 mV was used to produce a self-inhibitory (autaptic) current. During the
validation of this current, the reversal potential of Cl− was adjusted from −80 to −16 mV, in
order to reproduce the experimental set up of Bacci et al., 2003[19]. However, a physiological
reverse potential (−80 mV) was used for all other simulations. B,D: Model reproduction of
cp-AMPA (−70 mV) and NMDA (+60 mV) currents in response to stimulation of 2 synapses
as per Wang et al., 2009[62]. * Each trace represents the mean of all Hippocampal and PFC cells
respectively.

603 **Supplementary Figure 4:** Related to figure 2. Bimodal non-linear integration in Fast Spiking
604 basket cells. Supralinear (blue) and sublinear (magenta) dendrites shown in each model cell.

Hippocampus

PFC

Supplementary Figure 5: Related to figure 2. Bimodal non-linear integration in Fast Spiking basket cells. Representative Somatic EPSPs after stimulation (single pulse) of an increasing number of synapses (1:1:20), uniformly distributed within dendrites.

Supplementary Figure 6: Related to figure 2. Sensitivity analysis of biophysical dendritic

621   mechanisms reveals minor changes in the synaptic threshold for spike generation in

622   supralinear dendrites across Hippocampus (A) and PFC (B).

Supplementary Figure 7. Related to figure 6. Challenging the point neuron dogma: FS basket cells as 2-stage nonlinear integrators. Linear regression analysis for one hidden layer supralinear (A,C) and one hidden layer sublinear (B,D) ANNs for one indicative Hippocampal (top) and one indicative PFC (bottom) model cell. Actual Mean Firing Rates (Hz) correspond to the responses of the compartmental model when stimulating -with 50Hz Poisson spike trains- varying numbers of synapses (1 to 60), distributed in several ways (clustered or dispersed) within both sub- and supra-linear dendrites. Expected Mean Firing Rates (Hz) are those produced by the respective ANN abstraction when receiving the same input (number of stimulated synapses) in its respective sub-/supra- or linear input layer nodes.

References:

1. Buzsáki, G., Geisler, C., Henze, D. A. & Wang, X.-J. Interneuron Diversity
series: Circuit complexity and axon wiring economy of cortical interneurons.
Trends Neurosci. **27,** 186–193 (2004).

2. Chiovini, B. *et al.* Marti. *Neuron* **82,** 908–924 (2014).

3. Freund, T. F. Interneuron Diversity series: Rhythm and mood in perisomatic
inhibition. *Trends in Neurosciences* (2003). doi:10.1016/S0166-
2236(03)00227-3

4. Kann, O. The interneuron energy hypothesis: Implications for brain disease.
Neurobiol. Dis. **90,** 75–85 (2016).

5. Tremblay, R., Lee, S. & Rudy, B. GABAergic Interneurons in the Neocortex:
From Cellular Properties to Circuits. *Neuron* **91,** 260–292 (2016).

6. Hu, H., Gan, J. & Jonas, P. Interneurons. Fast-spiking, parvalbumin[+]
GABAergic interneurons: from cellular design to microcircuit function.
Science **345,** 1255263 (2014).

7. Klausberger, T. & Somogyi, P. Neuronal diversity and temporal dynamics: the
unity of hippocampal circuit operations. *Science* **321,** 53–7 (2008).

8. Van Aerde, K. I. *et al.* Flexible spike timing of layer 5 neurons during dynamic
beta oscillation shifts in rat prefrontal cortex. *J. Physiol.* **587,** 5177–5196
(2009).

9. Averkin, R. G., Szemenyei, V., Bordé, S. & Tamás, G. Identified Cellular
Correlates of Neocortical Ripple and High-Gamma Oscillations during
Spindles of Natural Sleep. *Neuron* (2016). doi:10.1016/j.neuron.2016.09.032

10. Somogyi, P. & Klausberger, T. Defined types of cortical interneurone structure
space and spike timing in the hippocampus. *J. Physiol.* **562,** 9–26 (2005).

11. Abrahamsson, T., Cathala, L., Matsui, K., Shigemoto, R. & DiGregorio, D. A.
Thin Dendrites of Cerebellar Interneurons Confer Sublinear Synaptic
Integration and a Gradient of Short-Term Plasticity. *Neuron* **73,** 1159–1172
(2012).

12. Nörenberg, A., Hu, H., Vida, I., Bartos, M. & Jonas, P. Distinct nonuniform
cable properties optimize rapid and efficient activation of fast-spiking
GABAergic interneurons. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 894–9 (2010).

13. Freund, T. F. & Katona, I. Perisomatic Inhibition. *Neuron* (2007).
doi:10.1016/j.neuron.2007.09.012

14. Lee, S.-H. *et al.* Parvalbumin-Positive Basket Cells Differentiate among
Hippocampal Pyramidal Cells. *Neuron* **82,** 1129–1144 (2014).

15. Povysheva, N. V *et al.* Parvalbumin-positive basket interneurons in monkey
and rat prefrontal cortex. *J. Neurophysiol.* **100,** 2348–60 (2008).

16.   Hioki, H. *et al.* Cell type-specific inhibitory inputs to dendritic and somatic
compartments of parvalbumin-expressing neocortical interneuron. *J. Neurosci.*
33, 544–55 (2013).

17.   Hainmüller, T., Krieglstein, K., Kulik, A. & Bartos, M. Joint CP-AMPA and
group I mGlu receptor activation is required for synaptic plasticity in dentate
gyrus fast-spiking interneurons. *Proc. Natl. Acad. Sci. U. S. A.* **111,** 13211–6
(2014).

18.   Geiger, J. R. P. *et al.* Relative abundance of subunit mRNAs determines gating
and Ca2+ permeability of AMPA receptors in principal neurons and
interneurons in rat CNS. *Neuron* **15,** 193–204 (1995).

19.   Bacci, A., Rudolph, U., Huguenard, J. R. & Prince, D. A. Cellular/Molecular
Major Differences in Inhibitory Synaptic Transmission onto Two Neocortical
Interneuron Subclasses.

20.   Goldberg, E. M. *et al.* K+ channels at the axon initial segment dampen near-
threshold excitability of neocortical fast-spiking GABAergic interneurons.
Neuron **58,** 387–400 (2008).

21.   Kim, D. *et al.* Distinct Roles of Parvalbumin- and Somatostatin-Expressing
Interneurons in Working Memory. *Neuron* (2016).
doi:10.1016/j.neuron.2016.09.023

22.   Kim, H., Ährlund-Richter, S., Wang, X., Deisseroth, K. & Carlén, M.
Prefrontal Parvalbumin Neurons in Control of Attention. *Cell* **164,** 208–218
(2016).

23.   Povysheva, N. V, Zaitsev, A. V, Gonzalez-Burgos, G. & Lewis, D. A.
Electrophysiological heterogeneity of fast-spiking interneurons: chandelier
versus basket cells. *PLoS One* **8,** e70553 (2013).

24.   Rudy, B. & McBain, C. J. Kv3 channels: voltage-gated K+ channels designed
for high-frequency repetitive firing. *Trends Neurosci.* **24,** 517–526 (2001).

25.   Martina, M. & Jonas, P. Functional differences in Na + channel gating between
fast-spiking interneurones and principal neurones of rat hippocampus. *J.*
Physiol. **505,** 593–603 (1997).

26.   Hu, H., Martina, M. & Jonas, P. Dendritic Mechanisms Underlying Rapid
Synaptic Activation of Fast-Spiking Hippocampal Interneurons. *Science (80-.).*
327, 52–58 (2010).

27.   Silver, R. A. Neuronal arithmetic. *Nat. Rev. Neurosci.* **11,** 474–489 (2010).

28.   Polsky, A., Mel, B. W. & Schiller, J. Computational subunits in thin dendrites
of pyramidal cells. *Nat. Neurosci.* **7,** 621–7 (2004).

29.   Papoutsi, A., Kastellakis, G., Psarrou, M., Anastasakis, S. & Poirazi, P. Coding
and decoding with dendrites. *J. Physiol. Paris* **108,** 18–27 (2014).

30.   Häusser, M. & Mel, B. Dendrites: bug or feature? *Curr. Opin. Neurobiol.* **13,**
372–383 (2003).

31. Poirazi, P., Brannon, T. & Mel, B. W. Arithmetic of Subthreshold Synaptic
Summation in a Model CA1 Pyramidal Cell. *Neuron* **37,** 977–987 (2003).

32. Poirazi, P., Brannon, T. & Mel, B. W. Pyramidal Neuron as Two-Layered
Neural Network. *Neuron* **37,** 989–999 (2003).

33. Larkum, M. E., Nevian, T., Sandler, M., Polsky, A. & Schiller, J. Synaptic
Integration in Tuft Dendrites of Layer 5 Pyramidal Neurons: A New Unifying
Principle. *Science (80-.).* **325,** (2009).

34. Losonczy, A. & Magee, J. C. Integrative properties of radial oblique dendrites
in hippocampal CA1 pyramidal neurons. *Neuron* **50,** 291–307 (2006).

35. Branco, T. & Häusser, M. *Synaptic Integration Gradients in Single Cortical
Pyramidal Cell Dendrites*. *Neuron* **69,** (2011).

36. Spruston, N. Pyramidal neurons: dendritic structure and synaptic integration.
Nat. Rev. Neurosci. **9,** 206–221 (2008).

37. Cazé, R. D., Jarvis, S., Foust, A. J. & Schultz, S. R. *Dendrites enable a robust
mechanism for neuronal stimulus selectivity*. *bioRxiv* (Cold Spring Harbor Labs
Journals, 2015). doi:10.1101/023200

38. Stuart, G. J. & Spruston, N. Dendritic integration: 60 years of progress. *Nat.
Neurosci.* **18,** 1713–1721 (2015).

39. Hu, H. & Vervaeke, K. Synaptic Integration in Cortical Inhibitory Neuron
Dendrites. *Neuroscience* **368,** 115–131 (2018).

40. Goldberg, J. H., Tamas, G. & Yuste, R. Ca2+ imaging of mouse neocortical
interneurone dendrites: Ia-type K+ channels control action potential
backpropagation. *J. Physiol.* **551,** 49–65 (2003).

41. Tran-Van-Minh, A., Abrahamsson, T., Cathala, L. & DiGregorio, D. A.
Differential Dendritic Integration of Synaptic Potentials and Calcium in
Cerebellar Interneurons. *Neuron* **91,** 837–850 (2016).

42. Katona, G. *et al.* Roller Coaster Scanning reveals spontaneous triggering of
dendritic spikes in CA1 interneurons. *Proc. Natl. Acad. Sci.* **108,** 2148–2153
(2011).

43. Branco and Hausser, M, T. The single dendritic branch as a fundamental
functional unit in the nervous system. *Curr. Opin. Neurobiol.* **20,** 494–502
(2010).

44. Tukker, J. J. *et al.* Distinct Dendritic Arborization and In Vivo Firing Patterns
of Parvalbumin-Expressing Basket Cells in the Hippocampal Area CA3. *J.
Neurosci.* **33,** (2013).

45. Rotaru, D. C., Yoshino, H., Lewis, D. A., Ermentrout, G. B. & Gonzalez-
Burgos, G. Glutamate Receptor Subtypes Mediating Synaptic Activation of
Prefrontal Cortex Neurons: Relevance for Schizophrenia. *J. Neurosci.* **31,** 142–
156 (2011).

46. Kastellakis, G., Cai, D. J., Mednick, S. C., Silva, A. J. & Poirazi, P. Synaptic
clustering within dendrites: An emerging theory of memory formation. *Prog.*
Neurobiol. **126,** 19–35 (2015).

47. Hines, M. L. & Carnevale, N. T. The NEURON simulation environment.
Neural Comp. **9,** 1179–1209 (1997).

48. Goldberg, J. H. & Yuste, R. Space matters: Local and global dendritic Ca2+
compartmentalization in cortical interneurons. *Trends Neurosci.* **28,** 158–167
(2005).

49. Tran-Van-Minh, A. *et al.* Contribution of sublinear and supralinear dendritic
integration to neuronal computations. *Front. Cell. Neurosci.* **9,** 67 (2015).

50. Cazé, R. D., Humphries, M., Gutkin, B., Tank, D. & DiGregorio, D. Passive
Dendrites Enable Single Neurons to Compute Linearly Non-separable
Functions. *PLoS Comput. Biol.* **9,** e1002867 (2013).

51. Cazé, R. D., Humphries, M. & Gutkin, B. Passive dendrites enable single
neurons to compute linearly non-separable functions. *PLoS Comput. Biol.* **9,**
e1002867 (2013).

52. Topolnik, L. Dendritic calcium mechanisms and long-term potentiation in
cortical inhibitory interneurons. *Eur. J. Neurosci.* **35,** 496–506 (2012).

53. Camire, O. & Topolnik, L. Dendritic Calcium Nonlinearities Switch the
Direction of Synaptic Plasticity in Fast-Spiking Interneurons. *J. Neurosci.* **34,**
3864–3877 (2014).

54. Hu, H., Martina, M. & Jonas, P. Dendritic Mechanisms Underlying Rapid
Synaptic Activation of Fast-Spiking Hippocampal Interneurons. *Science (80-.).*
327, 52–58 (2010).

55. Tran-Van-Minh, A. *et al.* Contribution of sublinear and supralinear dendritic
integration to neuronal computations. *Front. Cell. Neurosci.* **9,** 67 (2015).

56. Kohus, Z. *et al.* Properties and dynamics of inhibitory synaptic communication
within the CA3 microcircuits of pyramidal cells and interneurons expressing
parvalbumin or cholecystokinin. *J. Physiol.* **594,** 3745–74 (2016).

57. Martina, M. Distal Initiation and Active Propagation of Action Potentials in
Interneuron Dendrites. *Science (80-.).* **287,** 295–300 (2000).

58. Gulyás, A. I., Megías, M., Emri, Z. & Freund, T. F. Total Number and Ratio of
Excitatory and Inhibitory Synapses Converging onto Single Interneurons of
Different Types in the CA1 Area of the Rat Hippocampus. *J. Neurosci.* **19,**
(1999).

59. Kastellakis, G., Silva, A. J. & Poirazi, P. Linking Memories across Time via
Neuronal and Dendritic Overlaps in Model Neurons with Active Dendrites.
Cell Rep. **17,** 1491–1504 (2016).

60. Emri, Z., Antal, K., Gulyás, A. I., Megías, M. & Freund, T. F. Electrotonic
profile and passive propagation of synaptic potentials in three subpopulations

of hippocampal CA1 interneurons. *Neuroscience* **104,** 1013–1026 (2001).

61.   Konstantoudaki, X., Papoutsi, A., Chalkiadaki, K., Poirazi, P. & Sidiropoulou,
K. Modulatory effects of inhibition on persistent activity in a cortical
microcircuit model. *Front. Neural Circuits* **8,** 7 (2014).

62.   Wang, H.-X. & Gao, W.-J. Cell type-specific development of NMDA receptors
in the interneurons of rat prefrontal cortex. *Neuropsychopharmacology* **34,**
2028–40 (2009).

63.   Connelly, W. M. & Lees, G. Modulation and function of the autaptic
connections of layer V fast spiking interneurons in the rat neocortex. *J. Physiol.*
588, 2047–63 (2010).

64.   Goldberg, J. H., Yuste, R. & Tamas, G. Ca2+ imaging of mouse neocortical
interneurone dendrites: contribution of Ca2+-permeable AMPA and NMDA
receptors to subthreshold Ca2+dynamics. *J. Physiol.* **551,** 67–78 (2003).