Staphylococcus aureus **toxin LukSF dissociates from its membrane receptor target to enable**

renewed ligand sequestration

Authors:

Karita Haapasalo[1, 2, 4], Adam J. M. Wollman[3, 4], Carla de Haas[1], Kok van Kessel[1], Jos van Strijp[1], Mark

C. Leake[3, 5, *]

[1] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht University,

Utrecht, The Netherlands

[2] Department of Bacteriology and Immunology, and Research Programs Unit, Immunobiology,

University of Helsinki, Helsinki, 00014, Finland

[3] Biological Physical Sciences Institute, Departments of Physics and Biology, University of York,

York, YO10 5DD, United Kingdom

[4] These authors contributed equally

[5] Lead contact: Prof Mark Leake, Biological Physical Sciences Institute, Departments of Physics and

Biology, University of York, York YO10 5DD, UK. Tel: +44 (0)1904322697. Email:

mark.leake@york.ac.uk. Orcid ID**:** http://orcid.org/0000-0002-1715-1249.

[*] Correspondence: Email: mark.leake@york.ac.uk.

Keywords: bacterial toxin; pore formation; single-molecule; super-resolution; immune response.

ABSTRACT

Background: *Staphylococcus aureus* Panton Valentine Leukocidin (PVL) is a pore-forming toxin targeting the human C5a receptor (hC5aR), enabling this pathogen to battle the immune response by destroying phagocytes through targeted lysis. The mechanisms that contribute to rapid cell lysis are largely unexplored.

Results: Here we show that cell lysis may be enabled by a process of toxins targeting receptor clusters and receptor 'recycling' which allows multiple toxin pores to be formed close together. Using live cell single-molecule super-resolution imaging, Förster resonance energy transfer (FRET) and nanoscale total internal reflection fluorescence (TIRF) colocalization microscopy we visualized toxin pore formation in the presence of its natural docking ligand.

Conclusions: We demonstrate disassociation of hC5aR from toxin complexes and simultaneous binding of new ligands. This effect may free mobile receptors to amplify hyper inflammatory reactions in early stages of microbial infections and have implications for several other similar bi-component toxins and the design of new antibiotics.

BACKGROUND

S. aureus causes diseases ranging from superficial skin and soft tissue infections (SSTI) to severe invasive diseases like osteomyelitis and necrotizing pneumonia [1]. During the 1960s, methicillin-resistant *Staphylococcus aureus* (MRSA) was identified as a nosocomial pathogen [2]. In the 1990s, infection of previously healthy community-dwelling individuals with MRSA was reported [3]. Since then, these community-associated (CA-) MRSA have rapidly emerged worldwide [4]. Variants have also recently been identified which have reduced susceptibility to the antibiotic vancomycin [5] as well as complete resistance, so-called VRSA [6], and these forms of *S. aureus* pose a significant threat to human health. S. *aureus* and resistant variants have also evolved adaptations to evade attack from cells of the human immune system. However, the molecular processes which underlie these strategies are underexplored in living cells. There are compelling scientific and societal motivations to understand the mechanisms involved in immunogenic evasion strategies of *S. aureus*.

In the early 1930s, Panton and Valentine described a powerful leukocidal toxin produced by multiple *S. aureus* isolates, now denoted Panton-Valentine Leukocidin (PVL), years later shown to be cytotoxic to neutrophils, monocytes and macrophages but not to lymphocytes [7, 8]. The majority of CA-MRSA isolates carry the genes encoding PVL, partially due to the successful spread of the PVL carrying clone USA300 in the USA [3, 4, 9, 10], rarely present in hospital-acquired antimicrobial resistant MRSA and methicillin susceptible *S. aureus* isolates. Based on epidemiological studies, PVL is associated with primary skin infections in humans, osteomyelitis and, in particular, severe necrotizing pneumonia [11, 12]. Necrotizing pneumonia is a severe complication caused by bacterial lung infection. It is characterized by massive recruitment of neutrophils in the site of infection, diffuse pulmonary inflammation, septic shock, and respiratory failure. Both host factors and microbial virulence factors are thought to play an important role in the inflammation, however, it is unknown

how the interplay between these two factors affects the severity of the disease [13]. The specificity to

cell surface receptors makes it difficult to study PVL's role in *S. aureus* pathogenesis in a whole animal

model. It is possible that lysis of neutrophils by PVL is responsible for a reduced host defense response

allowing the pathogen to spread and cause eventual tissue damage. However, a previous study using a

rabbit animal model on necrotizing pneumonia suggests that PVL itself directly or indirectly causes

tissue injury and by this way induces local inflammation [14].

PVL is a pro-phage encoded bi-component, β-barrel pore-forming toxin (β-PFT) comprising

protein subunits LukS and LukF. LukS binding to the surface of target cells induces secondary LukF

binding; chemical and genetic analysis suggests that the resulting complex consists of a lytic pore-

forming hetero-octamer [15, 16]. Stoichiometric analysis *in vitro* of this complex suggests it is an

octamer of 4-plus-4 subunits [17]. In this complex only LukS is known to interact with the human C5a

receptor (hC5aR, CD88), a G-protein coupled seven-transmembrane receptor (GPCR). LukS targets at

least the extracellular N-terminus of hC5aR [18, 19], similar to the chemotaxis inhibitory protein of *S.*

aureus (CHIPS), but may also interact with the transmembrane receptor region [20]. C5aR is the ligand

for C5a, a powerful anaphylatoxin released during complement activation. Complement is a powerful

first line defense mechanism against invading pathogens which can be initiated through three

pathways: the classical, lectin, or alternative pathways. Activation of any of the three pathways on the

target leads to a rapid opsonization with C3b [21]. Further activation of complement leads to initiation

of the terminal pathway with release of C5a and formation of membrane attack complexes that are lytic

for Gram-negative but not Gram-positive bacteria [22, 23]. Therefore, in defense against Gram-positive

bacteria C3b-opsonization together with attraction and activation of neutrophils via C5a-C5aR

interaction are essential [24, 25]. In severe cases, formation of C5a can potentially lead to hyper

activation of the inflammatory response, an inability to regulate this potentially fatal reaction and

eventually harm the human host tissues. Because of this strong pro-inflammatory activity, therapeutic

interventions have recently focused on neutralizing antibodies against C5a and C5aR as potential

candidates for the treatment of severe inflammatory conditions such as bacterial induced sepsis [26,

27].

LukS binding to hC5aR inhibits C5a receptor binding which efficiently blocks neutrophil

activation [18]. LukS receptor binding alone is not sufficient for cell lysis but requires simultaneous

interaction between the leukocidin subunits and hC5aR. However, multiple possible subunit and

receptor combinations are theoretically possible and the spatiotemporal dynamics in functional

complexes in live cells between LukS, LukF and hC5aR is not yet known. In addition to PVL *S. aureus*

can produce a number of other β-PFTs with varying receptor and cell type specificities. From these

LukED, LukAB (or LukGH) and γ-hemolysin (composed of two compound pairs, HlgA/HlgB or

HlgC/HlgB) are classified as bi-component toxins like PVL while α-hemolysin is the prototypical β-

PFT that assembles into a pore through the oligomerization of seven monomeric polypeptides [28].

Next to bacterial toxins, an entire group of other pore forming proteins have been identified

which are involved in human innate immunity, indicating that pore-forming proteins are employed in

survival strategies for several types of organisms [29]. Development of methods to study dynamic

processes of pore formation by these toxins at a molecular level may improve our understanding of the

evolution of bacterial virulence and human immunity. There are several studies that have attempted to

explain the function of bacterial PFTs, including structural and subunit stoichiometry data from high

resolution X-ray crystallography and single-molecule fluorescence microscopy [17, 30, 31]. However,

these studies focused on pathogen instead of host factors and were thereby limited in excluding the

specific interaction between host cell receptor and bacterial toxin component, the first step required for

toxin oligomerization on the host cell membrane and the presence of the most potent factor mediating

the inflammatory response via C5a recognition in the site of infection [18].

Here, we used standard and single-molecule fluorescence detection with super-resolution

localization microscopy [32] to determine protein complex assembly on receptors in live and fixed cell

membranes. We studied human embryonic kidney (HEK) cells modified to express monomeric Green

Fluorescent Protein (mGFP) labeled hC5aR, exposed to Alexa dye-labeled *S. aureus* toxin components

LukS and LukF and imaged using standard total internal reflection fluorescence (TIRF) real time

microscopy (Additional file 1: Figure S1a) allowing us to monitor the spatiotemporal dynamics of

receptor and toxin molecules in the cell membrane. Our findings indicate that LukS binds on clusters of

membrane-integrated hC5aRs. The receptor-bound LukS then binds LukF leading to the formation of a

pore that is consistent with previous stoichiometric studies. However, when LukF is bound to the

complex, we observe fewer colocalized hC5aRs with toxin in fixed cells, more immobilized toxin

complexes in live cells and a significantly reduced Förster resonance energy transfer (FRET) signal,

indicating, unexpectedly, that pore formation leads to simultaneous dissociation of the receptors from

the complex. In addition, our biochemical data suggests that the dissociated receptor can then be

available for additional LukS molecules or the C5a generated during complement activation as a

response to LukSF-mediated cell lysis. This new finding suggests that a limited number of receptors

can be 'recycled' as docking for further toxin. This ensures that a sufficient number of pores will

damage nearby phagocytic cells, particularly important when high numbers of C5a anaphylatoxin are

blocking LukS, and potentially also enables simultaneous C5a mediated inflammatory response on

adjacent cells.

RESULTS

Maleimide-labeled LukSF mediates toxicity on human polymorphonuclear (PMN) and HEK

cells. To study LukSF pore formation on live cells using single-molecule fluorescence microscopy,

single cysteine substitutions on the exposed surface of the cap domain of the individual toxins

(Additional file 1: Figure S1b), K288C on LukF and K281C on LukS, were engineered to facilitate

maleimide labeling. These were denoted as the modified protein mLukF or mLukS. A second

substitution Y113H on LukS was chosen on the stem domain to facilitate pore formation of the LukS

mutant (mLukS), based on previous studies [17]. We compared the lytic activity of these mutants to

their unmodified wild type equivalents by measuring PMN membrane permeabilization after 30 min

toxin exposure using the DNA-binding fluorescent dye DAPI by flow cytometry. DAPI does not

penetrate intact cell membranes, and is therefore a good measure for cell permeability and cell death. In

this assay each of the wild type toxins was replaced with the modified protein either unlabeled (mLukF

or mLukS) or with a single Alexa647 dye molecule label (mLukF* or mLukS*) (Figure 1). All

modified toxins induced PMN permeabilisation reaching 100% at ~3 nM (Figure 1a), interchangeable

with the wild type equivalents. Only maleimide-labeled mLukF (mLukF*) lost activity and required

~30 nM to reach 100% permeabilization. Since the LukS component mediates the toxin recognition on

the target C5aR, we evaluated the binding potency of mLukS and mLukS* on PMNs. In this assay,

mLukS was able to inhibit the interaction of FITC-labeled wild type LukS on PMNs equally well as the

maleimide-labeled mLukS* (Figure 1b).

The LukSF toxin is known to be specific towards human cells expressing human C5aR (hC5aR) such as neutrophils, monocytes and macrophages but does not lyse cells that do not express the receptor [18]. To report on the spatiotemporal localization of the receptor and for determining the subunit stoichiometry in any observed receptor clusters we prepared HEK cells expressing hC5aR with a monomeric variant (bearing the obligate-monomer mutation A206K) of green fluorescent protein (mGFP) cloned in the C-terminal end of the receptor. This cell line also forms a monolayer on the coverslip and can be used for introducing a single dye on the cloned receptor, requirements for TIRF and single-molecule imaging. We verified the specificity and activity of the mutated and labeled toxins on the HEK-hC5aR cells. As expected, the toxins lysed only cells expressing hC5aR while control hCCR2 expressing cells, that do not bind LukS [33], remained intact (Figure 1d). We did not observe any binding of mLukF* on the same cells (Figure 1c) which is consistent with previous observations that LukF in the absence of LukS does not interact with PMN [16]. The unlabeled mLukS inhibited binding of mLukS* in a dose-dependent fashion (Figure 1e). HEK cells transfected with hC5aR required higher toxin concentrations for optimal binding and lysis by mLukSF as compared with PMN which is in agreement with previous data for the wild type variants [33]. We did not compare the hC5aR expression levels between these cells as PMNs also express another ligand for LukS, C5LR (or C5aR2) and because C5aR expression levels are not stable in neutrophils but can easily change in natural settings for example as a response to increased C5a levels [34].

To be able to analyze the dynamics of receptor and toxin interactions, we verified the conditions required for HEK cell lysis in time in the presence of mLukF and mLukS. Since the maleimide-labeled mLukF* required higher concentrations for efficient lysis of HEK cells, and because of the loss of molecules during washing cycles, the assay was optimized to have 20-fold excess of mLukF* (600 nM). Following pre-incubation of hC5aR-mGFP expressing HEK cells with LukS(wt) or mLukS,

LukF(wt) or mLukF* was added and the cellular uptake of DAPI was measured by flow cytometry

every 5 min. The wild type toxins LukF(wt) and LukS(wt) caused >80% cell toxicity within 10 min,

while closer to 20 min was required for significant lysis by the mLukF* and mLukS toxin combination

(Figure 1f).

Standard TIRF microscopy of live cells shows LukS colocalizes to hC5aR, and causes cell lysis

upon addition of LukF. For life cell imaging we first set the conditions to facilitate data acquisition of

dynamic events involved in the formation of LukSF nanopores in hC5aR-mGFP HEK cell membranes.

We sampled every 2.5 s at 50 ms exposure time per frame using standard (non single-molecule) total

internal reflection fluorescence (TIRF) microscopy, at very low excitation intensity to prevent

photobleaching. Cells were first imaged in the absence of toxin. In the green channel we observed

mGFP localization consistent with the cell membrane, manifest as relatively high apparent brightness

towards the cell boundaries consistent with the cell membrane curving away from the microscope

coverslip perpendicular to the TIRF excitation field. Controlled addition of mLukS* (labeled with

Alexa647) to the sample Petri dish followed by washing, while imaging simultaneously throughout,

resulted in colocalization of the hC5aR and mLukS (Figure 2, Additional file 6: Movie 1), with an

image structural similarity index of ~0.75. Further addition of mLukF resulted in complete lysis of the

cell, as defined by the observation of explosive release of membrane vesicles, after ~15 min (Figure 2,

Additional file 7: Movie 2). Colocalization of hC5aR-GFP, mLukS* (Alexa647) and mLukF*

(Alexa594) was also confirmed by three color experiments, imaging cells after addition of toxins and

washes until the start of lysis (Figure 2). In the given mLukF concentrations and time frame, where

imaging was possible without immediate cell lysis, also free hC5aR could be detected without mLukSF

colocalization.

Single-molecule TIRF microscopy of live cells shows LukS forms tetramers, and clusters hC5aR

before binding LukF. Using higher laser intensity TIRF excitation enabled rapid millisecond single

color channel sampling of single fluorophores faster than their molecular mobility in the cell membrane

[35], confirmed by imaging antibody-immobilized mGFP and Alexa dyes (Additional file 2: Figure

S2). Imaging live hC5aR-mGFP cells in these conditions saturated the camera CCD but after 1-2 min

of exposure, photobleaching was sufficient to reduce intensity and allow us to observe several distinct,

mobile, circular fluorescent foci at a mean surface density of ~1 per μm^2 in the planar membrane

regions which lie parallel to the TIRF field away from the cell boundaries (Figure 3a, Additional file 8:

Movie 3). We monitored the spatiotemporal dynamics of foci in the planar membrane regions using

automated tracking software [36] which allowed foci to be tracked for several seconds to a spatial

precision of ~40 nm [37], below the diffraction limit, thus enabling super-resolution localization data to

be obtained. The measured focus width (defined as the half width at half maximum determined from

their pixel intensity profile) was in the range 200-300 nm, consistent with the point spread function

(PSF) width of our microscope. By using step-wise photobleaching analysis we estimated

stoichiometry values for all detected fluorescent foci by employing a method which quantifies the

initial unbleached foci brightness and divides this by the measured brightness for the relevant single

dye reporter molecule (Additional file 2: Figure S2) [38]. These foci contained large numbers of

receptors with a mean stoichiometry of ~180 (Figure 3b, Table 1). Addition of mLukS and mLukF

increased the mean stoichiometry by >50% consistent with the toxin causing receptor clustering.

Imaging mLukS* incubated with hC5aR-mGFP cells revealed distinct foci (Figure 3a,

Additional file 9: Movie 4). The probability distribution of mLukS* stoichiometry values in live cells

in the absence of mLukF is shown in Figure 3c, rendered using a kernel density estimation which

generates an objective distribution that does not depend upon the size and location of subjective

histogram bins [39]. We measured a broad range of stoichiometry values, spanning a range from only a few LukS molecules per foci to several tens of molecules, with a mean of ~30 molecules per foci. Closer inspection of the stoichiometry values indicated an underlying periodicity to their probability distribution, which we investigated using Fourier spectral analysis [40]. The resulting power spectrum (Figure 3d) indicated a fundamental peak equivalent to a stoichiometry of 3.9 ± 0.2 molecules, suggesting that foci are composed of multiples of tetrameric mLukS* complexes.

Fluorescent foci, if separated by less than the diffraction-limited PSF width of our microscope, are detected as a single particle but with higher apparent stoichiometry. We therefore tested the hypothesis that the observed mLukS* foci stoichiometry distribution could be explained by the random overlap of isolated mLukS* tetramer foci. To do so we modeled the nearest-neighbor separations of individual mLukS* tetramers in the cell membrane as a random Poisson distribution [41] and used sensible ranges of tetramer surface density based on our single particle tracking results (Additional file 3: Figure S3). However, all random tetramer overlap models we explored showed poor agreement to the observed experimental stoichiometry distribution, but we found that random overlap of multimers of tetramers could account for the stoichiometry distribution well (Figure 3e). Optimized fits indicated that the random overlap of mLukS* foci with a stoichiometry in the range 4-20 molecules were able to best account for the experimental data. As hC5aR is clustered, this likely accounts for the clustering of mLukS* but not its tetrameric periodicity. These results are consistent with mLukS* binding to clusters of hC5aR as tetramers or forming tetrameric sub-structures.

We tested if there was a dependence of foci stoichiometry on incubation time with leukocidin. Acquiring a time course for mLukF* accumulation following pre-incubation of cells with mLukS was not feasible since unbound mLukF* had to be washed from the sample to prevent a prohibitively high fluorescent background. However, we were able to acquire time courses in which mLukF was added to

cells that had been pre-incubated with mLukS*. For these, the mLukS* foci stoichiometry distribution

was measured as a function of time after mLukF addition for several different fields of view, each

containing typically ~5 cells. We found that the mean hC5aR foci stoichiometry indicated no obvious

correlation to mLukF incubation time (Figure 3f), however the mean mLukS* foci stoichiometry

increased with time ($p < 0.05$).

By calculating the mean square displacement (MSD) as a function of time interval (τ) for each

tracked foci we could determine its apparent microscopic diffusion coefficient (D). The distribution of

D for hC5aR and mLukS*/mLukF (Additional file 4: Figure S4) had similar low value peaks at

~0.05 μm^2/s, consistent with immobile foci tracked with our localization precision spatial precision of

40 nm. Several mobile foci were also seen, which diffused at rates up to ~5 μm^2/s. Based on the

measured width of the immobile peak width on these distributions we set a threshold of 0.12 μm^2/s to

categorize foci as either immobile, which indicated a mean $D = 0.025 \pm 0.030$ μm^2/s ($\pm$SD), or mobile,

which indicated a mean $D = 0.47 \pm 0.40$ μm^2/s (Table 1). Plots of the measured MSD *vs.* τ relations for

mobile foci indicated a linear dependence indicative of free Brownian (i.e. 'normal') diffusion.

However, similar plots for immobile foci indicated an asymptotic dependence consistent with confined

diffusion [42], whose plateau was equivalent to a confinement diameter of ~400 nm (Additional file 4:

Figure S4). The relative proportion of mobile foci was ~35% of tracked foci for hC5aR, regardless of

toxin and similar for mLukS in the absence of mLukF. Addition of mLukF, caused a drop in the mobile

proportion by a factor of ~3 (Figure 3g) suggesting that LukF causes insertion of the complex and

possible disassociation of the LukSF complex from the hC5aR.

Single-molecule TIRF microscopy combined with colocalization analysis of fixed cells suggests

LukSF dissociates from the receptor. Due to the high image frame rate of single-molecule TIRF

microscopy, we were not able to simultaneously image two color channels on our microscope, rather

each channel was imaged separately in the same cells. Therefore to determine whether the toxin

remains bound to the receptor, and to quantify the relative stoichiometry of components, we imaged

fixed cells, halting cell lysis, using the same two spectrally distinct green/red dyes of mGFP and

Alexa647 to label receptor and toxin components, respectively, as for the live cell experiments. We

imaged cells incubated with mLukS*, followed by incubation with mLukF (Figure 4a upper panels) as

well as simultaneously with mLukS+mLukF* (Figure 4a lower panels) and observed foci with similar

stoichiometries (Table 1) to live cells but colocalized with hC5aR. Around 32% of the hC5aR foci were

found colocalized in the presence of mLukS* dropping to <10% in the presence of mLukF (Figure 4b).

This low percentage was within our estimate of the degree of random colocalization between the green

and red fluorophores, entirely down to chance, of ~10%. This suggests that in the presence of mLukF,

the toxin is not colocalized with the receptor and that mLukF causes disassociation from hC5aR. The

stoichiometry values for detected green hC5aR-mGFP foci were calculated and plotted against the

equivalent stoichiometry estimates for colocalized red foci of mLukS* and mLukF* respectively (Figue

4c and Additional file 5: Figure S5). In the presence of mLukS* but in the absence of mLukF, the

hC5aR-mGFP foci stoichiometry showed an approximately linear dependence on number of associated

mLukS* molecules, suggesting that each colocalized mLukS* molecule was associated on average

with ~4-5 hC5aR molecules. In the presence of labeled or unlabeled mLukF no dependence was

observed (Additional file 5: Figure S5, R^2<0) consistent with random association between toxin and

receptor. These results are unlikely to be due to fluorescence quenching, as it would need to be near

100% quenching to detect no Alexa fluorescence in the hC5aR-mGFP foci and the drop in

colocalization is observed independent of the labeled toxin used, either mLukS* with mLukF or

mLukF* with mLukS.

Live whole cell FRET and biochemical measurements also support LukSF disassociation: We

performed FRET experiments on FITC sortase-labeled hC5aR and Cy3-labeled mLukS or mLukF, as

donor and acceptor respectively, in live cells to further probe the association between toxin and

receptor. A FRET signal from whole cells of 75% efficiency was observed, with a statistically

siginfcant drop (p=0.008, Student's *t*-test) to 56% when incubated with unlabeled mLukF (Figure 4d

and e), as would be expected if the complex formation leads to dissociation of the toxin from the

receptor. In order to examine possible FRET between hC5aR and Cy3-labeled mLukF we performed

similar experiments on fixed cells. In these experiments a FRET efficiency of 60% was observed

between hC5aR and labeled mLukS dropping below 40% (p=0.023, Student's *t*-test) between hC5aR

and labeled mLukF. As expected, no FRET signal was observed in the negative control where only

Cy3-labeled mLukF was present. These results are also consistent with the finding that hC5aR

dissociates from the LukSF pore, although conformational or local environment changes cannot be

ruled out with FRET alone since the relatively high remaining signal might in principle also indicate

remaining association or other inter or intra hC5aR-Luk interaction. The greater drop in FRET when

measured with mLukF compared to mLukS might be caused by the 3-4 nm further distance of LukF

from hC5aR.

     To further confirm that the LukSF complex dissociates from the target receptor we used a

monoclonal PE-labeled anti-CD88 antibody to detect the liberation of free hC5aR receptors on the cell

membrane upon LukSF formation. We first confirmed the ability of both C5a and wild type LukS to

compete for binding of the anti-CD88 antibody to the hC5aR expressing HEK cells. Both ligands

showed clear inhibition of anti-CD88 binding at 100 nM concentrations while LukF was ineffective

(Figure 5a). However, when the hC5aR expressing cells were incubated with 100 nM of LukS followed

by incubation with increasing concentrations of wild type LukF to form an active toxin, a statistically

significant increase in anti-CD88 binding was detected at a LukF concentration of 1nM when compared

to no LukF (0 nM) . Addition of a control protein Ecb did not change anti-CD88 binding. Cell

permeabilization was measured in parallel and proved to be "sublytic" enabling proper detection of

liberated anti-CD88 without significant cell lysis (% of lysed cells < 10%) (Figure 5b). At a 3 nM LukF

concentration the proportion of dead cells increased above 10% which determined the maximum

concentration and increase in anti-CD88 binding that could be measured. The changes in C5aR

mobility, colocalization, and FRET with addition of LukF, combined with the biochemical evidence of

anti-CD88 rebinding on hC5aR upon LukF(wt) addition on LukS(wt) coated cells are strongly

indicative of disassociation of the LukSF complex.

Dissociation of LukSF pores from hC5aR allows rebinding of C5a or intact LukS on the receptor.

Since C5a is the natural ligand for hC5aR and can outcompete binding of LukS on the receptor we next

analyzed whether LukSF formation and disengagement of hC5aR would allow rebinding of C5a on the

receptor. We specifically chose to analyze binding of labeled C5a and not LukS in the presence of

increasing concentrations of LukF. This is because adding labeled LukS to cells coated with LukS (and

then washing) together with LukF will give several possibilities for association: for example, new pores

for free or unoccupied C5aR; intercalation with present bound LukS and lukF in pores; binding to free

or unoccupied C5aR but without pore formation. To detect C5a rebinding at higher LukF

concentrations we used a G130D LukF mutant that interacts with LukS but does not cause cell lysis.

Non lytic activity of this mutant in this assay was confirmed by DAPI staining that showed minimal

cell lysis even at higher concentrations (9% at a 300 nM LukFG130D concentration). At a 300 nM

concentration a significant increase in C5a binding was detected indicating that LukSF dissociates from

hC5aR enabling simultaneous rebinding of an hC5aR interacting ligand (Figure 5c). On the contrary,

addition of the control molecule, Ecb, did not cause increase in C5a binding suggesting that this was

due to rebinding of C5a to disengaged hC5aR and not for example because of increase in receptor

expression. This assay showed that C5a can potentially interact with these cells that are attacked by

LukSF. Because C5a is a potent anaphylatoxin that is generated during complement activation and

potentially plays a crucial role in *S. aureus* infections [43] we next analyzed whether LukSF could lead

to complement activation and C5a formation in an *ex vivo* full blood assay. We used soluble C5b-9 as a

marker for terminal complement activation and C5a formation, not C5a, because LukS is known to

compete with C5a for binding to hC5aR on neutrophils [18]. The presence of 200 nM of LukSF clearly

increased formation of soluble C5b-9 compared to full blood without any toxin or only LukS (Figure

5d), indicating that LukSF mediated cell lysis increases C5a formation and potentially also

inflammation in the site of infection.

DISCUSSION

To determine the stoichiometry of the toxin components without immobilizing the protein on a

surface or within a crystal we implemented our fluorescence imaging method which allows us to

monitor the actual pore formation mechanism within a living cell, including the target receptor crucial

for the complex formation. This kind of study on protein complex formation has not been done before

due primarily to the difficulty of labeling the components and the high native fluorescence background

in mammalian cells. Our covalent labeling strategy and high excitation intensity TIRF microscopy,

combined with advanced image analysis tools, opens the way for further studies into many other pore

forming toxins and processes involving membrane bound protein complex formation.

The finding that the toxin complexes are found in receptor clusters indicates that lysis of cells

depends on the local density of close proximity hC5a receptors that will initiate the pore formation

process by docking LukS close to the cell membrane such that four hC5aR-LukS dimers (assuming that

one LukS binds only one hC5aR, although hC5aR are randomly clustered on the membrane) can

interact with the free non-bound LukF that will eventually form an octamer (i.e. 4 by 4) and a

functional pore with LukS. These data also suggest that when proper assembly in an octamer is

ongoing/complete, hC5aR will dissociate from the complex and, at the same time, can interact with the

C5a that is formed during complement activation amplified by LukSF-mediated cell lysis itself. This is

logical because, in addition to invading microbes, apoptotic and necrotic cells are known to activate

complement [44]. The triggering of local complement activation by toxin damaged cells and the release

of locally generated C5a (Figure 5) and its interaction with adjacent cells such as endothelial or lung

epithelial cells [45] could explain the mechanism behind the exacerbated inflammation characteristics

exhibited in necrotizing pneumonia. This is an important finding, suggesting that the cause of infection

can dramatically affect the magnitude of the inflammatory response and is highly dependent on the

dynamics of microbial molecules interacting with human receptors. In addition, the disengaged hC5aR

is possibly also available for new toxins to bind, thus allowing the receptor to be recycled and reused

by additional LukS molecules. Our finding that C5a can rebind, doesn't only suggest a mechanism for

exaggerated immune response by LukSF, but also indicates that the dissociated free hC5aR does not

change in conformation due to previous contact with LukS and therefore would also be available for

binding with LukS. We find that roughly half of LukS complexes are immobile prior to LukF binding,

but that LukF binding then results in mostly immobile LukSF complexes. Speculatively, this result may

suggest that LukS binds hC5aR initially and then inserts itself transiently in the membrane

phospholipid bilayer via the exposed hydrophobic residues, following binding of LukF molecules to

LukS. This stable insertion of the LukSF complex into the cell membrane then leads to pore formation

across the whole cell membrane.

To characterize the hC5aR interaction with LukSF at a molecular level, we used malemide-

labeled toxins and HEK cells that expressed only hC5aR and not the second docking target for LukSF,

which are both present on human PMNs [46]. We verified that the interaction between maleimide-

labeled toxin component LukS and the cell surface receptor is required for the target recognition and

cell lysis similarly as shown before for wild type LukS [18] both for human PMNs and hC5aR

expressing HEK cells that were chosen for TIRF imaging because of their stability and ability to form

monolayers on the microscopy cover slip.

We observe clusters of pre-established hC5aR in the absence of LukS or LukF, but the addition

of LukS or LukF significantly increased the mean cluster stoichiometry. Fourier spectral analysis

combined with foci overlap modeling suggests that LukS complexes comprise a multimer of 4-5

hC5aR subunits each of which contain 4 LukS molecules, even before addition of LukF. Our findings

are consistent with the hetero-octamer model of 4-plus-4 LukS/LukF subunits [17, 30, 31], however,

our colocalization analysis also indicates the presence of stable complexes of LukS with hC5aR

independent of LukF.

By characterizing the mobility of hC5aR and LukS in live cells we find that roughly half of

hC5aR and LukS foci diffuse relatively freely in the cell membrane while the remainder are confined to

zones in the membrane of ~400 nm effective diameter. However, when LukF is present > 90% of LukS

foci become immobile (confined). If LukS were to undergo a conformational change following LukF

binding then this may potentially expose hydrophobic residues that could fascilitate insertion of the

toxin into the hydrophobic interior of the phospholipid bilayer. This hypothesis is strongly supported

by the β-barrel prepore-pore formation putative mechanism of γ-hemolysin. Here the residues

responsible for binding with the phospholipid head group are located at the bottom of the rim domain

whereas the stem domain forms an antiparallel β-barrel of which the bottom half comprises the

transmembrane portion of the pore [31]. This change from receptor associated LukS to cell membrane

associated LukSF complex can be seen as a change in the proportion of mobile (receptor associated

LukS) and immobile (toxin complexes inserted into cell membrane) foci detected in live cells. GPCRs

similarly are known to have heterogeneous mobility and lateral distribution properties in living cells at

different states for example before and after activation [47].

Crystallographic evidence from the monomeric LukF and LukS components and the intact γ-

hemolysin pore suggests that the pore is octameric formed from 4-plus-4 LukF/LukS subunits [30, 48,

49]. Our findings support this octamer model but unlike previous studies also indicate that LukS pre-

forms into a tetramer without LukF and that formation of this tetramer is facilitated by close proximity

C5aR clusters. The presence of LukS tetramers in the absence of LukF cannot be further explained by

our data. It is, however, possible that if LukS molecules would be associated as a tetramer when bound

on the receptor the conformational changes on LukS caused by interactions with LukF should enable

association of the LukF subunits to the complex. According the data presented here this is possible

because in these assays we first enabled LukS to bind on the receptors and eliminated the effect of

freshly formed complexes by free unbound LukS by a washing step before addition of LukF. Each

octamer component consists of cap, rim and stem domains. Here, the cap domain contains the site for

LukS/LukF interaction while the stem domain unfolds and forms the transmembrane β-barrel upon

pore formation. Within crystallization the 2-methyl-2,4-pentanediol (MPD) molecules are bound at the

base of the rim domain, and recognized by Trp177 and Arg198 residues, that may participate in

recognition of the phospholipid bilayer as suggested in a crystal structure of the LukF monomer [50].

In contrast, the structure of the γ-hemolysin suggests a membrane interaction site within residues

Tyr117, Phe119 and Phe139 on the same toxin component [30]. The crystal structure of LukED

determined recently reveals important details of the residues on LukE required for receptor

identification [51]. This component corresponds to the receptor binding component LukS on the LukSF

complex, scanning mutagenesis indicting that LukS residues Arg73, Tyr184, Thr244, His245 and

Tyr250, and to a lesser extent Tyr181, Arg242 and Tyr246, are involved in binding to the neutrophil

surface [52].

These results suggest that further binding sites for hC5aR on LukS could be possible in addition

to those identified in the LukS rim domain [52]. However, since the binding of LukS to neutrophils is

inhibited by the C5a it is likely that LukS has only one binding site on the receptor [20]. This is also

supported by the similar inhibition profiles of LukS and C5a towards anti-CD88 binding on hC5aR

shown in this study. Therefore, the association of LukS with approximately 4-5 hC5aR molecules could

be explained by the previous suggestion that C5aR forms homo-oligomers in living cells [53]. Our

findings imply that LukSF assembly is dependent on hC5aR cell membrane area density as opposed to

the effective hC5aR concentration when calculated over the whole of a target cell volume, such that

even when hC5aR cellular expression levels are low, for example when inflammatory mediators are

formed to limit the inflammation [34], a cell lysis response may still be achieved through the efficient

targeting of receptor clusters and recycling of the receptor molecules in the cell membrane to be re-

used by free non-bound LukS to get engaged in octamer pore formation. It is possible that

overexpression of hC5aR on HEK cells could lead to an increased ability to form hC5aR foci with

many receptors especially because of intracellular GFP tag although we mitigated this possibility by

cloning the receptor with a monomeric variant of GFP that does not cause GFP dimerization. Other

studies, however, have shown that hC5aR forms clusters of homodimers or heterodimers with the

second C5a receptor C5LR (C5aR2) or other GPCRs like CCR5 especially under high concentrations

of C5a [53-55].

Previous *in vitro* studies on LukSF pores formed on human leukocytes and rabbit erythrocytes

have found evidence for both octamers and hexamers, but importantly both suggest a LukS/LukF ratio

of 1:1 [17, 56, 57]. Interestingly we did not observe any correlation to the number of hC5aR present

with LukF incubation time once LukF was already bound to LukS. Moreover, when LukS was

incubated with LukF using sortase-labeled hC5aR cells a significant reduction was observed in the

FRET efficiency signal between LukS and C5aR. It is unlikely that the reduction that was observed in

FRET efficiency would be due to a conformational change because the cysteine mutation used for

maleimide labeling was designed to be exposed on the cap domain of LukS and LukF (Additional file

1: Figure S1) that in light of the structural data undergoes minimal or no conformational changes

during complex assembly [31]. Since our biochemical assays indicate that LukF does not bind directly

to hC5aR expressing cells and, that binding of LukF to LukS results in an increased distance between

the receptor and the complex, this suggests that LukF binding to LukS results in LukS dissociating

from the receptor, released as a newly formed LukSF complex.

We cannot directly determine the cause of this behavior in our present study, however one

explanation may lie in the conformational change during the prepore-to-pore transition that has been

shown to occur on γ-hemolysin complexes subsequently after binding of LukF to LukS [30, 31].

Interestingly, this same study shows that during the pre-pore state the space for the transmembrane

region is occupied by the rim domain of the adjacent octamer in a LukSF crystal. One explanation for

these observations, that remains to be explored, is that in addition to the stem domain the residues

within the rim domain that interact with the receptor might also have different orientations in the pre-

pore state when compared to the pore state. In addition to using the maleimide labeled mLukF and

mLukS and fluorescence microscopy the putative dissociation of the hC5aR from the LukSF complex

was further verified by using the wild type LukS and LukF proteins in an assay where LukS-coated

hC5aR cells were incubated with increasing concentrations of LukF. Here, increase in anti-CD88

binding also clearly indicates LukSF dissociation. In all of the assays where we could observe 20-30%

receptor dissociation we used sublytic concentrations of LukF to be able to measure healthy cells with

normal membrane fluidity and natural behavior rather than dead cells. This kind of receptor

disengagement has been shown before by at least the cytotoxin intermedilysin which interacts with a

GPI-anchored complement regulatory molecule on the cell membrane [58]. Moreover, dissociation of

LukSF complex is also supported by electron microscopy of LukSF on human leukocyte membrane

fragments. Here, the ring-shaped oligomers with outer and inner diameters of 9 nm and 3 nm were

shown without a receptor [57].

In this study we also show, for the first time to our knowledge, that the dissociated receptor can

be reused by free unbound C5a. This indicates that LukS binding on the receptor does not change

receptor conformation and thereby can putatively also be reused by additional LukS. In our full blood

model we observed that LukSF mediated cell lysis clearly increased complement activation and C5a

formation. The increase in C5a concentration in the site of infection could potentially limit the

availability of hC5aR for LukS molecules on neutrophils and thereby reduce lytic activity of the toxin

as C5a has previously been shown to reduce LukSF mediated lysis *in vitro* [18]. Rebinding of C5a on

the receptor may therefore indicate that in natural settings where all components (i.e. LukS, LukF and

C5a) are present C5a can outcompete binding of LukSF on the target cells. Therefore, recycling of the

receptor could be one strategy for the toxin to ensure that a sufficient number of pores will damage the

cells especially when limited number of receptors are available.

There are several steps on the leukocidin complex assembly that may be critical for the function

of the toxin. Based on our observations we provide new information on leukocidin-receptor interactions

and propose two additional stages to the processes of pore formation and the mechanism by which

LukSF potentially induces inflammation (Figure 6a). Stage 1 is the binding of LukS to hC5aR clusters.

The first step in this process is the target recognition of LukS binding to the membrane receptor. We

detected LukS-hC5aR complexes on clusters of receptors indicating that pore formation takes place in

these clusters. Stage 2 is the binding of 4 LukF molecules to to 4 LukS molecules resulting in a hetero-

octamer LukSF nanopore in the neutrophil cell membrane. Stage 3 is then the dissociation of the

receptors from the LukSF complex enabling the repcetor to be resued for subsequent binding free

unbound ligand to generate more nanopores in the cell membrane and enhance the damage to the

neutrophil. We measured a correlation between the number of LukS and hC5aR molecules present in

LukS-hC5aR complexes, but with no obvious correlation between the number of LukF and hC5aR

molecules when LukF was added to the LukS-hC5aR complex. In addition to previous studies [14] we

suggest that LukSF-mediated cell lysis and dissociation from hC5aR can potentially amplify *S. aureus*

mediated inflammation in the site of infection (Figure 6). The direct lysis of neutrophils is enhanced by

newly formed LukSF complexes that are formed on the cell membrane hC5aR via reattachment of new

LukS. Neutrophil lysis activates the complement system and the newly generated C5a induces

cytokine/chemokine production and neutrophil chemotaxis via the C5a/C5aR signaling pathway on

adjacent cells. Furthermore, the increased vasodilation and vascular permeability (Figure 6b) leads to

massive neutrophil accumulation and tissue injury at the site of bacterial infection [59].

CONCLUSIONS

In summary, our findings that the receptors of targeted host cells dissociate rapidly from the

leukocidin complex upon formation of a harmful toxin pore, freeing up mobile receptor 'seeds' that can

diffuse to other parts of the cell membrane, suggest a hitherto undiscovered strategy used by microbes

to kill human immune cells. This enables a limited number of receptors to be recycled as docking for

the leukocidin or potentially the anaphylatoxin C5a to ensure that enough pores will form to damage

the host cell and simultaneously maintain or possibly amplify the inflammation in the site of infection.

This discovery may generalize to other bi-component toxins which employ a similar docking receptor

like the C5aR receptor, including the family of *Staphylococcal* bi-component leukocidins of

HlgC/HlgB, HlgA/HlgB, LukE/LukD (CXCR1, CXCR2, CCR5), and LukM/LukF' for bovine CCR2.

These results highlight the importance of leukocidin-receptor interactions in pore formation and may

facilitate further understanding in the role of pore-forming toxins in *S. aureus* infections. This new

mechanistic insight may prove valuable to the development of future antibacterial and anti-

inflammatory therapies, especially important in light of the growing menace of global antimicrobial

resistance.

MATERIALS AND METHODS

Experimental model and subject details

PMN isolation, Cell Lines, and Transfections. Human blood was obtained from healthy volunteers

and the polymorphonuclear (PMN) cells were isolated by Ficoll/Histopaque centrifugation [60].

Informed consent was obtained from all subjects, in accordance with the Declaration of Helsinki and

the Medical Ethics Committee of the University Medical Center Utrecht (METC-protocol 07-125/C

approved 1 March 2010). To ensure a truly monomeric state and prevent GFP mediated clustering of

the receptor, a fusion construct of hC5aR with the monomeric GFP variant mGFP with A206K

mutation (also denoted GFPmut3) [61, 62] was made at the C-terminus (primers used listed in Table 2)

or a sortase A LPXTGG sequence was made in the N-terminus and cloned into pIRESpuro vectors

(Table 2) by PCR. The amplification reaction was performed in three separate amplification steps using

overlap extension PCR on hC5aR and mGFP templates. hC5aR (Accession number of human C5aR =

NM_00173) was used as the template using enzymes and purification kits as described above. The

clones were ligated into the vectors and transferred into TOP10 *E. coli* competent cells, then amplified

and sequenced similarly to the toxin clones described previously. The pIRESpuro/hC5aR-mGFP vector

was transfected into Human Embryonic Kidney (HEK) 293T cells (a HEK cell line, Invitrogen), stably

expressing G protein Gα16, using Lipofectamine-2000 Reagent according to manufacturer's

instructions (Thermo Fisher Scientific). After 24–48 hr, transfected cells were harvested with 0.05%

trypsin. To obtain a uniform, stable culture, cells were sub-cloned in a concentration of 0.5 cells/well in

a 96-well plate in Dulbecco's modified Eagle's medium (DMEM, Lonza) supplemented with 10% fetal

calf serum (Invitrogen) 100 U/ml penicillin/ 100 µg/ml streptomycin (PS; Invitrogen), 1 µg/ml

Hygromycin and 250 µg/ml Puromycin. For N-terminal labelling of the sortase A recognition sequence

containing HEK cells with FITC were successfully performed in two steps as described previously

[63]. The expression of hC5aR was analyzed by incubating the cells in 50 µl RPMI (Invitrogen)

supplemented with 0.05% human serum albumin (Sanquin), RPMI-HSA, at 5×10^6 cell/ml

concentration for 45 min with PE-conjugated anti-CD88 and detected by flow cytometry. The presence

of mGFP or FITC-LPXTG was detected directly by flow cytometry.

Recombinant Protein Production and Purification. Polyhistidine-tagged LukS and LukF were cloned

and expressed using an *E. coli* expression system. For maleimide-based labeling a single-cysteine

mutation was designed to the LukS and LukF components based on previous data and the crystal

structure of the octameric pore [30]. An additional mutation Y113H was included in LukS to facilitate

oligomerization of the maleimide-labeled protein [17]. The target genes were amplified by PCR (Table

2) from the wild type sequences using Phusion High-Fidelity DNA polymerase (Thermo Scientific)

[18]. The PCR product was cloned into a slightly modified pRSET expression vector (Invitrogen),

resulting in expression of proteins with an N-terminal 6xHIS-tag. For LukF mutant G130D we useda

gBlock (custom dsDNA sequence via Integrated DNA Technologies) to incorporate the LukF in the

pRSET vector. Clones were sequenced to verify the correct sequence. The recombinant proteins were

expressed in Rosetta Gami (DE3) pLysS *E. coli* using 1mM IPTG induction and isolated by a native

isolation method. The expressed proteins were purified according to the manufacturer's instructions

(Invitrogen) using 1 ml Nickel HisTrap and Superdex 75 HiLoad columns (GE Health Care Life

Sciences). Toxin components were labeled with either Cy3 (GE Healthcare), Alexa Fluor® 594 or

Alexa Fluor® 647 C_2 Maleimide reagent according to the manufacturer's instructions (Thermo

Scientific) resulting in negligible unlabeled content. The labeling efficiency was 100% as determined

by protein concentrations using absorption at A280 and dye concentrations using absorption at A650 by

a Nanodrop ND-1000 Spectrophotometer.

Method details

Binding Assays. Binding of the maleimide-labeled proteins to PMN and HEK cells was confirmed by

flow cytometry. LukS-K281C-Y113H (mLukS) or wild type, LukS (wt) was labeled with FITC or

Alexa Fluor maleimide 647 or 594. For competition assays 3 µg/ml of the labeled protein and

increasing concentration of non-labeled mLukS or LukS(wt) was incubated with isolated PMNs or

HEK hC5aR-mGFP cells (5 $\times10^6$ cell/ml) in a total volume of 50 µl RPMI-HSA on ice. For binding

assays without competition the cells were incubated with increasing concentration of mLukF*. After

30 min incubation on ice, cells were washed, fixed with 1% paraformaldehyde and analyzed by flow

cytometry. HEK cells transfected with CCR2 receptor were used as negative control for mLukS

binding. To see inhibition of PE anti-CD88 (BD biosciences) binding by LukS(wt) or C5a hC5aR

expressing HEK cells were first incubated with increasing concentrations of LukS or C5a for 45 min at

4°C. Then 2 µl of anti-CD88/200,000 cells was added and incubated as previously. Cells were washed

once with RPMI-HSA, fixed with 1% paraformaldehyde and analyzed by flow cytometry. To detect

hC5aR dissociation using sublytic concentrations of LukSF hC5aR expressing HEK cells were

incubated with 100 nM of wild type LukS for 45 min at 4°C. After washing the unbound LukS sublytic

concentrations of wild type LukF was added to the cells and incubated for 20 min at 37°C and 5% CO_2

atmosphere. Percentages of lysed vs. non lysed cells were measured by using 1μg/ml of DAPI in the

reaction. For C5a rebinding assay 1 μM of C5a (Sigma) was labeled with NT-647 according to

manufacturer's instructions (Monolith NT™). Free label from the sample was removed by three times

centrifugation trough Amicon Ultra 0.5 mL centrifugal filters (Sigma). The hC5aR expressing HEK

cells were incubated with 1 μM of wild type LukS for 45 min at 4°C. After washing 20 nM of NT647-

C5a and increasing concentrations of wild type LukF was added to the cells and incubated for 20 min at

37°C and 5% CO_2 atmosphere. Cells were washed once with RPMI-HSA, fixed with 1%

paraformaldehyde and analyzed by flow cytometry. Percentages of lysed vs. non lysed cells were

measured by using 1μg/ml of DAPI in the reaction. *S. aureus* Ecb (extracellular complement binding

protein) was used as negative control as it interacts with another cell surface receptor, CR1 [64]. Flow

cytometry data were analyzed using FlowJo v10 software package.

Cell Permeability Assays. Isolated PMNs or HEK hC5aR-mGFP cells (5×10^6 cell/ml) were exposed to

labeled and unlabeled mixtures as appropriate of mLukF/mLukS recombinant proteins at equimolar

concentrations in a volume of 50 μl RPMI-HSA with 1 μg/ml of DAPI. Cells were incubated for

30 min at 37°C with 5% CO_2 and subsequently analyzed by flow cytometry. To calculate the lysis time

cells were first incubated with 150 nM of mLukS for 15 min. Then 600 nM of mLukF was added and

immediately subjected to flow cytometry analysis where the permeability was measured at several time

points. Cell lysis was defined as intracellular staining by DAPI. HEK cells transfected with human

CCR2 receptor was used as negative control for toxin-mediated lysis. Statistical differences between

means of repeated experiments were calculated using two-tailed Student *t*-tests.

Ex vivo complement activation assay. To maintain complement activity the blood samples were

anticoagulated with lepirudin (Refludan, Schering, Berlin, Germany). Increasing concentrations of

LukSF or LukS (0 to 2000 nM) was incubated in full blood for 30 min at 37°C under continuous

rotation (300 rpm). Complement activity was stopped by adding 10 mM EDTA in the suspension and

the plasma was separated from the blood cells by centrifugation at $5000 \times$ g. A 1:30 dilution of each

plasma sample was analyzed by SC5b 9 Enzyme Immunoassay according to manufacturer's

instructions (MicroVue SC5b 9 Plus Enzyme Immunoassay, Quidel). One *S. aureus* colony (1 x 10^8

cells) was used as a positive control for SC5b-9 formation. The bacteria were grown over night on

blood agar plate at 37°C 5% CO_2 atmosphere.

Fluorescence microscopy. Cells were imaged using a Nikon A1R/STORM microscope utilizing a x100

NA oil immersion Nikon TIRF objective lens. We used a total internal reflection fluorescence (TIRF)

microscopy module. We used laser excitation at wavelengths 488 nm (for mGFP), 561 nm (for

Alexa594) and 647 nm (for Alexa647) from a commercial multi laser unit fiber-coupled into the

microscope, capable of delivering maximum power outputs up to ~200 mW, with a depth of

penetration in the range ~100-130 nm for the TIRF excitation evanescent field. Fluorescent images

acquired on an iXon+ 512 EMCCD camera detector (Andor) at a magnification of 150nm/pixel. Green

and red channel images were obtained by imaging through separate GFP or Alexa647 filter sets. For

high laser excitation intensity single-molecule millisecond imaging, green channel images to determine

mGFP localization were acquired continuously using 488 nm wavelength laser excitation over a period

of ~5 min through a GFP filter set, then the filter set was manually switched to Alexa647 as for red

channel images acquisition continuously using 647 nm wavelength laser excitation until complete

photobleaching of the sample after 1-2 min. For photobleaching laser powers ranged between 15 mW

(Alexa 647) to 100 mW (mGFP). For fixed cell analysis cells were either incubated first with mLukS or

mLukS*, washed and incubated with mLukF or mLukF*, or incubated first just with mLukS* with

mLukF* absent, then washed and fixed with 1% paraformaldehyde.

For fluorescence imaging the HEK cells were grown on 0.1% poly-L-lysine which coated

8-well chambered cover glass slides (Ibidi) in standard growth conditions described above. To analyze

the deposition of mLukS* on live cells the cells were first imaged in PBS buffer in the absence of

toxin. Here, a 256×256 pixel area covered approximately one cell per field of view. Then the cells

were incubated for 2 min with 5 µg/ml of Alexa594 maleimide-labeled LukS in RPMI-HSA and the

cells were carefully washed with PBS keeping the imaging area and focus constant. Because of the fast

bleaching of the Alexa647 label a more stable Alexa594 label was used for the LukS deposition

imaging. The deposition of mLukS was detected for 10 min and the lysis of the cell was recorded for

15 min after addition of 600 nM of unlabeled mLukF. Cells were imaged in TIRF at 50 ms per frame

with the laser automatically switched between 488 nm/0.22 mW, 647 nm/3 mW and 561 nm/3 mW or

488 nm/0.22 mW and 561 nm/3 mW (Figure 2).

FRET experiments. The Cy3-labeled LukS or non-labeled LukS and Cy3-labeled LukF or non-labeled

LukF or Cy-labeled LukF and controls (only FITC-labeled cells or unlabeled cells with only Cy3-

labeled LukS) were incubated at 4°C for 30 min and then 10 min at 37°C, washed 2 x with RPMI-HSA

and fixed as before. The cells were in PBS during imaging. FITC experiments were performed using

Leica TCS SP5 microscope, using a 62× oil immersion objective lens, and FRET Sensitized Emission

Wizard in Leica Application Suite Advanced Fluorescence (LAS AF). Images were acquired using

488 nm and 543 nm wavelength lasers and a laser power of 27% 12.0 (A.U.) and a scan size of

512×512, 800 ms, 50 ms per frame, beam splitter TD 488/543/633.

FRET efficiency ε was calculated using the donor, directly excited acceptor and donor excited

acceptor intensity from n=5-10 manual regions of interest inside cells for each experiment, using the

following formula [65]:

$$\varepsilon \; = \frac{\beta(\mathrm{B} - \mathrm{D}) \; - \gamma\mathrm{A}}{\mathrm{A}}$$

B = Intensity signal, donor excited acceptor

D = Intensity signal, donor

A = Intensity Signal, directly excited acceptor

β = calibration for ratio of measured intensities of $B_{\text{donor channel}}/A_{\text{donor channel}}$

γ = calibration for ratio of measured intensities of $B_{\text{acceptor channel}}/A_{\text{acceptor channel}}$

Single-molecule imaging of live and fixed cells. GFP and Alexa647 fluorescence micrograph time

series of fixed and live cells were sampled taken at. 20 ms per frame. Green channel images were

acquired continuously using 488 nm wavelength laser excitation over a period of *ca.* 5 min via the GFP

filter set. Then the filter set was manually switched to that for Alexa647 and red channel images were

acquired continuously using 647 nm wavelength laser excitation was until complete photobleaching of

the sample after 1-2 min. The step-wise single-molecule fluorescence photobleaching was analyzed

both for live and fixed cells. For live cell photobleaching analysis the cells were incubated with 150 nM

of labeled or unlabeled mLukS as required for *ca.* 15 min. After washing with PBS, 600 nM of labeled

or unlabeled mLukF was added and the imaging was done immediately within 10-15 min. If labeled

LukF was added the wells were washed with PBS before analysis. Also samples with only LukS and

without toxins were analyzed. For fixed cell analysis the cells were incubated first with mLukS or

mLukS* for 30 min at +4°C in RPMI-HSA, washed with same buffer and incubated for 10 min at 37°C

with mLukF or mLukF*, or the same protocol was followed but using mLukS* alone with mLukF*

absent. Then the cells were washed and fixed with 1% paraformaldehyde. 1M mercaptoethylamine

(MEA) buffer was used for fixed cell analysis. Photobleaching of recombinant mGFP and mLukS-

Alexa647 were also separately analyzed in a tunnel slide comprising two pieces of double-sided tape

forming a channel sandwiched between a standard glass microscope slide and a plasma cleaned

coverslip. Proteins solutions (1μg/ml) were immobilized onto the coverslip coated by anti-GFP or anti-

His antibodies respectively with phosphate buffered saline washes in between.

Quantification and statistical analysis

Binding and permeability assays. Statistical significance between repeated (n>1) experiments was

analyzed using two-tailed Student's *t*-tests where using a standard p-value threshold of <0.05 as

indicating statistical significance. Means and standard deviations of repeated experiments are shown in

error bars, unless indicated otherwise.

Image analysis. Basic image extraction, cropping and quantification was done using NIS-Elements

microscope imaging software and Image J. More advanced foci tracking was done using bespoke

software written in MATLAB (Mathworks) [36] which enabled automatic detection and localization of

individual fluorescent foci to within 40 nm lateral precision (Additional file 2: Figure S2a). The

software identifies candidate foci by a combination of pixel intensity thresholding and image

transformation. The intensity centroid and characteristic intensity, defined as the sum of the pixel

intensities inside a 5-pixel radius circular region of interest around the foci intensity centroid minus the

local background and corrected for non-uniformity in the excitation field are determined by repeated

Gaussian masking. If the signal-to-noise ratio of a foci (the intensity per pixel/background standard

deviation per pixel) is greater than a pre-set threshold, nominally here set at 0.4 based on prior

simulations, it is accepted and fitted with a 2D radial Gaussian function to determine its width. Foci in

consecutive frames within a single PSF width, and not different in intensity or width by greater than a

factor of two, are linked into the same track.

Foci intensity was used to quantify stoichiometry information. As foci photobleach over time

during continuous laser excitation their intensity falls in a stepwise manner due to photobleaching of an

integer number of fluorophore tags in each sampling time window. By quantifying the size of a single

step, the characteristic intensity of a single fluorophore can be obtained and thus the stoichiometry of

the foci from its initial intensity. The step size is found from the periodicity in the distribution of foci

intensities corroborated by the pairwise distance (PwD) distribution of these intensities and the Fourier

spectrum of the PwD which contains peaks at the characteristic intensity and harmonics at multiples of

this value (Additional file: Figure S2d-e).

Here, the copy number of hC5aR-mGFP was comparatively high such that the TIRF images

were initially saturated in regards to pixel intensity output. After ~20 s of photobleaching the non-

saturated foci intensity values were fitted by an exponential function which characterized the rate of

intensity decay, equivalent to an exponential photobleach time of ~20 s, and extrapolated back to zero

time to determine the initial foci intensity (Additional file: Figure S2f). The Alexa647 dye also

bleached during 647 nm wavelength laser excitation but images were not initially saturated, but also in

some images which were exposed to the 488nm laser and then the 647nm laser, also bleached by the

488 nm wavelength laser. In these images a fixed correction factor of 6x, determined by comparing to

images exposed to the 647 nm laser first, was used. The stoichiometry of each foci was then

determined as the initial intensity divided by the intensity of the appropriate single fluorescent dye tag

(i.e. either mGFP or Alexa647 in this case).

We characterized the mobility of tracked foci by calculating their MSD as a function of time

interval (τ). For each detected foci the MSD was calculated from the measured intensity centroid

($x(t),y(t)$) at time t assuming a foci track of N consecutive image frames at a time interval $\tau=n\Delta t$ where

n is a positive integer and Δt is the frame integration time (here 20 ms):

$$MSD(\tau) = MSD(n\Delta t) = \frac{1}{N-1-n} \sum_{i=1}^{N-1-n} \left(\left(x(i\Delta t + n\Delta t) - x(i\Delta t)\right)^2 + \left(y(i\Delta t + n\Delta t) - y(i\Delta t)\right)^2 \right)$$

$$= 4D\tau + 4\sigma^2$$

The lateral (xy) localization precision is given by σ which we determine to be 40 nm. We fitted a

straight line to each separate MSD relation. Assuming a line fit has an optimized gradient g to the first

4 points (defined as the first 3 measured MSD data points for $n=1$, 2 and 3, in addition to a 4th data

corresponding to $n=0$ obtained from constraining the intercept be $4\sigma^2$ to within measurement error of

the localization precision) then estimated the microscopic diffusion coefficient D as $g/4.\Delta t$. For

immobile foci, tracks were collated and compiled to generate a mean MSD vs. τ relation which was

fitted to an asymptotic rising exponential function as an analytical model for confined diffusion of

MSD plateau equal to $L^2/6$ where L is the effective confinement diameter [42], enabling us to estimate

the confinement diameter.

Colocalization analysis. The extent of colocalization between red and green detected foci was

determined using a method which calculated the overlap integral between each green and red foci pair,

whose centroids were within ~1 PSF width (~3 pixels). Assuming two normalized, two-dimensional

Gaussian intensity distributions g_1 and g_2, for green and red foci respectively, centred around (x_1, y_1)

with sigma width σ_1, and around (x_2, y_2) with width σ_2, the overlap integral ν is analytically determined

as:

$$\nu = e^{\left(-\Delta r^2 / 2\left(\sigma_1^2 + \sigma_2^2\right)\right)}$$

736 Where:

$$\Delta r^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$$

We use a criterion of an overlap integral of 0.75 or above to indicate putative colocalization

[41] since this corresponds to a foci centroid separation equivalent to the localization precision in this

case. By quantifying the standard deviation on the number of detected foci in each channel we estimate

that the standard error of colocalisation proportion under our typical imaging conditions is

approximately 9%.

Random foci overlap models. We calculated the probability of foci overlap in a single color channel by

first estimating a sensible range of foci surface density n. For the lower limit we used the number of

foci tracks detected in a 20 image frame time window, for the upper limit we used the average

measured value of the background-corrected pixel intensity value divided by the intensity of a single

fluorophore (equivalent to ~1 mLukS* molecule per pixel). We implemented these probability

estimates into a surface density model which assumed a random Poisson distribution for nearest-

neighbor separation [40, 41, 66-69]. This model indicates that the probability that a nearest-neighbor

separation is greater than w is given by $\exp(-\pi w^2 n)$. The probability of overlap for each density

estimate (Additional file 4: Figure S4) was convolved with a real molecular stoichiometry distribution

and a Gaussian function $p(x)$ of stoichiometry (x):

$$p(x) = 2\pi\sigma^2 \exp(-\frac{(x-n)^2}{2\sigma^2\sqrt{n}})$$

where σ is the width of single fluorophore intensity distribution (~0.7 molecules), and n is the real

molecular stoichiometry. The tetramer model assumes $n=4$, then all higher order stoichiometries are

due to overlapping PSFs. The tetramer oligomer molecule assumed an equal number of multimerized

tetramers up to 5, which gave the best fit to the data.

The same strategy was used to model the random overlap probability for green and red color

channel fluorescent foci in dual color imaging experiments to assess the extent of apparent

colocalization due to random overlap between hC5aR and mLukS*/F*. The probability that a nearest-

neighbor separation is greater than w for foci of two different types is the same as a single type

multiplied by 2/3. (38)

Software

All our bespoke software developed is freely and openly accessible

via https://sourceforge.net/projects/york-biophysics/ (65).

Statistical tests and replicates

All statistical tests used are two-tailed unless stated otherwise. For single-molecule TIRF imaging each

cell can be defined as a biological replicate sampled from the cell population. We chose sample sizes of

5-7 cells yielding thousands of foci, generating reasonable estimates for stoichiometry and diffusion

coefficient distributions. Technical replicates are not possible with the irreversible photobleaching

assay.

Ethics statement

Human polymorphonuclear (PMN) cells, obtained from healthy volunteers were isolated by

Ficoll/Histopaque centrifugation [60]. Informed written consent was obtained from all subjects, in

accordance with the Declaration of Helsinki and the Medical Ethics Committee of the University

Medical Center Utrecht (METC-protocol 07-125/C approved 1 March 2010).

Acknowledgments

We thank Piet Aerts and Angelino Tromp for assistance in sample preparation and labeling and Esther

van 't Veld and Richard Wubbolts (Utrecht) for assistance with light microscopy and Dr Karin Strijbis

(Veterinary School, UU) providing the sortase A enzyme. This work was supported by The Finnish

Cultural Foundation (grants 00131060 and 00142390), the Biological Physical Sciences Institute,

Royal Society, MRC (grant MR/K01580X/1), BBSRC (grant BB/N006453/1), the EPSRC Physics of

Life UK network and the Wellcome Trust [ref: 204829] through the Centre for Future Health (CFH) at

the University of York, UK.

Conflict of interest

All the authors declare that they have no conflict of interests.

References

1.   DeLeo FR, Otto M, Kreiswirth BN, Chambers HF. Community-associated meticillin-resistant

Staphylococcus aureus. Lancet. 2010;375(9725):1557-68.

2.   Barrett FF, McGehee RF, Jr., Finland M. Methicillin-resistant Staphylococcus aureus at Boston

City Hospital. Bacteriologic and epidemiologic observations. N Engl J Med. 1968;279(9):441-8.

3.   Udo EE, Pearman JW, Grubb WB. Genetic analysis of community isolates of methicillin-

resistant Staphylococcus aureus in Western Australia. J Hosp Infect. 1993;25(2):97-108.

4.   Vandenesch F, Naimi T, Enright MC, Lina G, Nimmo GR, Heffernan H, et al. Community-

acquired methicillin-resistant Staphylococcus aureus carrying Panton-Valentine leukocidin genes:

worldwide emergence. Emerg Infect Dis. 2003;9(8):978-84.

5.   Hiramatsu K, Aritaka N, Hanaki H, Kawasaki S, Hosoda Y, Hori S, et al. Dissemination in

Japanese hospitals of strains of Staphylococcus aureus heterogeneously resistant to vancomycin.

Lancet. 1997;350(9092):1670-3.

6.   Koch G, Yepes A, Forstner KU, Wermser C, Stengel ST, Modamio J, et al. Evolution of

resistance to a last-resort antibiotic in Staphylococcus aureus via bacterial competition. Cell.

2014;158(5):1060-71.

7.   Meyer F, Girardot R, Piemont Y, Prevost G, Colin DA. Analysis of the specificity of Panton-

Valentine leucocidin and gamma-hemolysin F component binding. Infect Immun. 2009;77(1):266-73.

8.   Gauduchon V, Werner S, Prevost G, Monteil H, Colin DA. Flow cytometric determination of

Panton-Valentine leucocidin S component binding. Infect Immun. 2001;69(4):2390-5.

9.   Otter JA, French GL. Molecular epidemiology of community-associated meticillin-resistant

Staphylococcus aureus in Europe. Lancet Infect Dis. 2010;10(4):227-39.

10.   Naimi TS, LeDell KH, Como-Sabetti K, Borchardt SM, Boxrud DJ, Etienne J, et al.

Comparison of community- and health care-associated methicillin-resistant Staphylococcus aureus

infection. JAMA. 2003;290(22):2976-84.

11.   Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, et al. Involvement of

Panton-Valentine leukocidin-producing Staphylococcus aureus in primary skin infections and

pneumonia. Clin Infect Dis. 1999;29(5):1128-32.

12.   Gillet Y, Issartel B, Vanhems P, Fournet JC, Lina G, Bes M, et al. Association between

Staphylococcus aureus strains carrying gene for Panton-Valentine leukocidin and highly lethal

necrotising pneumonia in young immunocompetent patients. Lancet. 2002;359(9308):753-9.

13.   Perret M, Badiou C, Lina G, Burbaud S, Benito Y, Bes M, et al. Cross-talk between

Staphylococcus aureus leukocidins-intoxicated macrophages and lung epithelial cells triggers

chemokine secretion in an inflammasome-dependent manner. Cellular microbiology. 2012;14(7):1019-

36.

14.   Diep BA, Chan L, Tattevin P, Kajikawa O, Martin TR, Basuino L, et al. Polymorphonuclear

leukocytes mediate Staphylococcus aureus Panton-Valentine leukocidin-induced lung inflammation

and injury. Proc Natl Acad Sci U S A. 2010;107(12):5587-92.

15.   Jayasinghe L, Bayley H. The leukocidin pore: evidence for an octamer with four LukF subunits

and four LukS subunits alternating around a central axis. Protein Sci. 2005;14(10):2550-61.

16.   Colin DA, Mazurier I, Sire S, Finck-Barbancon V. Interaction of the two components of

leukocidin from Staphylococcus aureus with human polymorphonuclear leukocyte membranes:

sequential binding and subsequent activation. Infect Immun. 1994;62(8):3184-8.

17.   Das SK, Darshi M, Cheley S, Wallace MI, Bayley H. Membrane protein stoichiometry

determined from the step-wise photobleaching of dye-labelled subunits. Chembiochem. 2007;8(9):994-

9.

18.   Spaan AN, Henry T, van Rooijen WJ, Perret M, Badiou C, Aerts PC, et al. The staphylococcal

toxin Panton-Valentine Leukocidin targets human C5a receptors. Cell host & microbe. 2013;13(5):584-

94.

19.   Postma B, Kleibeuker W, Poppelier MJ, Boonstra M, Van Kessel KP, Van Strijp JA, et al.

Residues 10-18 within the C5a receptor N terminus compose a binding domain for chemotaxis

inhibitory protein of Staphylococcus aureus. J Biol Chem. 2005;280(3):2020-7.

20.   Spaan AN, Schiepers A, de Haas CJ, van Hooijdonk DD, Badiou C, Contamin H, et al.

Differential Interaction of the Staphylococcal Toxins Panton-Valentine Leukocidin and gamma-

Hemolysin CB with Human C5a Receptors. J Immunol. 2015;195(3):1034-43.

21.   Walport MJ. Complement. First of two parts. N Engl J Med. 2001;344(14):1058-66.

22.   Fearon DT, Austen KF. Activation of the alternative complement pathway with rabbit

erythrocytes by circumvention of the regulatory action of endogenous control proteins. The Journal of

experimental medicine. 1977;146(1):22-33.

23.   Berends ET, Mohan S, Miellet WR, Ruyken M, Rooijakkers SH. Contribution of the

complement Membrane Attack Complex to the bactericidal activity of human serum. Molecular

immunology. 2015;65(2):328-35.

24.   Easmon CS, Glynn AA. Comparison of subcutaneous and intraperitoneal staphylococcal

infections in normal and complement-deficient mice. Infection and immunity. 1976;13(2):399-406.

25.   Mullaly SC, Kubes P. The role of TLR2 in vivo following challenge with Staphylococcus

aureus and prototypic ligands. J Immunol. 2006;177(11):8154-63.

26.   Woodruff TM, Nandakumar KS, Tedesco F. Inhibiting the C5-C5a receptor axis. Mol Immunol.

2011;48(14):1631-42.

27.   Ricklin D, Lambris JD. Progress and Trends in Complement Therapeutics. Adv Exp Med Biol.

2013;735:1-22.

28.   DuMont AL, Torres VJ. Cell targeting by the Staphylococcus aureus pore-forming toxins: it's

not just about lipids. Trends Microbiol. 2014;22(1):21-7.

29.   Dal Peraro M, van der Goot FG. Pore-forming toxins: ancient, but never really out of fashion.

Nat Rev Microbiol. 2016;14(2):77-92.

30.   Yamashita K, Kawai Y, Tanaka Y, Hirano N, Kaneko J, Tomita N, et al. Crystal structure of the

octameric pore of staphylococcal gamma-hemolysin reveals the beta-barrel pore formation mechanism

by two components. Proc Natl Acad Sci U S A. 2011;108(42):17314-9.

31.   Yamashita D, Sugawara T, Takeshita M, Kaneko J, Kamio Y, Tanaka I, et al. Molecular basis

of transmembrane beta-barrel formation of staphylococcal pore-forming toxins. Nat Commun.

2014;5:4897.

32.   Chiu SW, Leake MC. Functioning nanomachines seen in real-time in living bacteria using

single-molecule and super-resolution fluorescence imaging. Int J Mol Sci. 2011;12(4):2518-42.

33.   Spaan AN, van Strijp JAG, Torres VJ. Leukocidins: staphylococcal bi-component pore-forming

toxins find their receptors. Nat Rev Microbiol. 2017;15(7):435-47.

34.   Unnewehr H, Rittirsch D, Sarma JV, Zetoune F, Flierl MA, Perl M, et al. Changes and

regulation of the C5a receptor on neutrophils during septic shock in humans. J Immunol.

2013;190(8):4215-25.

35.   Plank M, Wadhams GH, Leake MC. Millisecond timescale slimfield imaging and automated

quantification of single fluorescent protein molecules for use in probing complex biological processes.

Integr Biol (Camb). 2009;1(10):602-12.

36.   Miller H, Zhou Z, Wollman AJ, Leake MC. Superresolution imaging of single DNA molecules

using stochastic photoblinking of minor groove and intercalating dyes. Methods. 2015;88:81-8.

37.   Wollman AJ, Leake MC. Millisecond single-molecule localization microscopy combined with

convolution analysis and automated image segmentation to determine protein concentrations in

complexly structured, functional cells, one cell at a time. Faraday Discuss. 2015;184:401-24.

38.   Leake MC, Chandler JH, Wadhams GH, Bai F, Berry RM, Armitage JP. Stoichiometry and

turnover in single, functioning membrane protein complexes. Nature. 2006;443(7109):355-8.

39.   Leake MC. Analytical tools for single-molecule fluorescence imaging in cellulo. Phys Chem

Chem Phys. 2014;16(25):12635-47.

40.   Leake MC, Greene NP, Godun RM, Granjon T, Buchanan G, Chen S, et al. Variable

stoichiometry of the TatA component of the twin-arginine protein transport system observed by in vivo

single-molecule imaging. Proc Natl Acad Sci U S A. 2008;105(40):15376-81.

41.   Llorente-Garcia I, Lenn T, Erhardt H, Harriman OL, Liu LN, Robson A, et al. Single-molecule

in vivo imaging of bacterial respiratory complexes indicates delocalized oxidative phosphorylation.

Biochim Biophys Acta. 2014;1837(6):811-24.

42.   Robson A, Burrage K, Leake MC. Inferring diffusion in single live cells at the single-molecule

level. Philos Trans R Soc Lond B Biol Sci. 2013;368(1611):20120029.

43.   von Kockritz-Blickwede M, Konrad S, Foster S, Gessner JE, Medina E. Protective role of

complement C5a in an experimental model of Staphylococcus aureus bacteremia. J Innate Immun.

2010;2(1):87-92.

44.   Walport MJ. Complement. Second of two parts. N Engl J Med. 2001;344(15):1140-4.

45.   Pandya PH, Wilkes DS. Complement system in lung disease. Am J Respir Cell Mol Biol.

2014;51(4):467-73.

46.   Li R, Coulthard LG, Wu MC, Taylor SM, Woodruff TM. C5L2: a controversial receptor of

complement anaphylatoxin, C5a. FASEB J. 2013;27(3):855-64.

47.   Veya L, Piguet J, Vogel H. Single Molecule Imaging Deciphers the Relation between Mobility

and Signaling of a Prototypical G Protein-coupled Receptor in Living Cells. J Biol Chem.

2015;290(46):27723-35.

48.   Pedelacq JD, Maveyraud L, Prevost G, Baba-Moussa L, Gonzalez A, Courcelle E, et al. The

structure of a Staphylococcus aureus leucocidin component (LukF-PV) reveals the fold of the water-

soluble species of a family of transmembrane pore-forming toxins. Structure. 1999;7(3):277-87.

49.   Guillet V, Roblin P, Werner S, Coraiola M, Menestrina G, Monteil H, et al. Crystal structure of

leucotoxin S component: new insight into the Staphylococcal beta-barrel pore-forming toxins. J Biol

Chem. 2004;279(39):41028-37.

50.   Olson R, Nariya H, Yokota K, Kamio Y, Gouaux E. Crystal structure of staphylococcal LukF

delineates conformational changes accompanying formation of a transmembrane channel. Nat Struct

Biol. 1999;6(2):134-40.

51.   Nocadello S, Minasov G, Shuvalova L, Dubrovska I, Sabini E, Bagnoli F, et al. Crystal

structures of the components of the Staphylococcus aureus leukotoxin ED. Acta Crystallogr D Struct

Biol. 2016;72(Pt 1):113-20.

52.   Laventie BJ, Guerin F, Mourey L, Tawk MY, Jover E, Maveyraud L, et al. Residues essential

for Panton-Valentine leukocidin S component binding to its cell receptor suggest both plasticity and

adaptability in its interaction surface. PLoS One. 2014;9(3):e92094.

53.   Rabiet MJ, Huet E, Boulay F. Complement component 5a receptor oligomerization and

homologous receptor down-regulation. J Biol Chem. 2008;283(45):31038-46.

54.   Moreno-Fernandez ME, Aliberti J, Groeneweg S, Kohl J, Chougnet CA. A Novel Role for the

Receptor of the Complement Cleavage Fragment C5a, C5aR1, in CCR5-Mediated Entry of HIV into

Macrophages. AIDS Res Hum Retroviruses. 2016;32(4):399-408.

55.   Croker DE, Halai R, Fairlie DP, Cooper MA. C5a, but not C5a-des Arg, induces upregulation of

heteromer formation between complement C5a receptors C5aR and C5L2. Immunol Cell Biol.

2013;91(10):625-33.

56.   Miles G, Movileanu L, Bayley H. Subunit composition of a bicomponent toxin: staphylococcal

leukocidin forms an octameric transmembrane pore. Protein Sci. 2002;11(4):894-902.

57.   Sugawara N, Tomita T, Sato T, Kamio Y. Assembly of Staphylococcus aureus leukocidin into a

pore-forming ring-shaped oligomer on human polymorphonuclear leukocytes and rabbit erythrocytes.

Biosci Biotechnol Biochem. 1999;63(5):884-91.

58.   LaChapelle S, Tweten RK, Hotze EM. Intermedilysin-receptor interactions during assembly of

the pore complex: assembly intermediates increase host cell susceptibility to complement-mediated

lysis. J Biol Chem. 2009;284(19):12719-26.

59.   Guo RF, Ward PA. Role of C5a in inflammatory responses. Annu Rev Immunol. 2005;23:821-

52.

60.   Veldkamp KE, Heezius HC, Verhoef J, van Strijp JA, van Kessel KP. Modulation of neutrophil

chemokine receptors by Staphylococcus aureus supernate. Infect Immun. 2000;68(10):5908-13.

61.   Zacharias DA, Violin JD, Newton AC, Tsien RY. Partitioning of lipid-modified monomeric

GFPs into membrane microdomains of live cells. Science. 2002;296(5569):913-6.

62.   von Stetten D, Noirclerc-Savoye M, Goedhart J, Gadella TW, Jr., Royant A. Structure of a

fluorescent protein from Aequorea victoria bearing the obligate-monomer mutation A206K. Acta

Crystallogr Sect F Struct Biol Cryst Commun. 2012;68(Pt 8):878-82.

63.   Hirota N, Yasuda D, Hashidate T, Yamamoto T, Yamaguchi S, Nagamune T, et al. Amino acid

residues critical for endoplasmic reticulum export and trafficking of platelet-activating factor receptor.

The Journal of biological chemistry. 2010;285(8):5931-40.

64.   Amdahl H, Haapasalo K, Tan L, Meri T, Kuusela PI, van Strijp JA, et al. Staphylococcal

protein Ecb impairs complement receptor-1 mediated recognition of opsonized bacteria. PLoS One.

2017;12(3):e0172675.

65.   Wouters FS, Verveer PJ, Bastiaens PI. Imaging biochemistry inside cells. Trends Cell Biol.

2001;11(5):203-11.

66.   Badrinarayanan A, Reyes-Lamothe R, Uphoff S, Leake MC, Sherratt DJ. In vivo architecture

and action of bacterial structural maintenance of chromosome proteins. Science. 2012;338(6106):528-

31.

67.   Delalez NJ, Wadhams GH, Rosser G, Xue Q, Brown MT, Dobbie IM, et al. Signal-dependent

turnover of the bacterial flagellar switch protein FliM. Proc Natl Acad Sci U S A. 2010;107(25):11347-

51.

68.   Reyes-Lamothe R, Sherratt DJ, Leake MC. Stoichiometry and architecture of active DNA

replication machinery in Escherichia coli. Science. 2010;328(5977):498-501.

69.   Lenn T, Leake MC, Mullineaux CW. Clustering and dynamics of cytochrome bd-I complexes

in the Escherichia coli plasma membrane in vivo. Mol Microbiol. 2008;70(6):1397-407.

Legends

Figure 1. Toxin functionality on PMN and HEK cells. (A) PMN cell permeability in the presence of

unlabeled LukSK281CY113H and LukFK288C (mS+mF, number of biological replicates n=2), Alexa-

labeled mS* or mF* and wild type LukF (F(wt)) or LukS, (S(wt)) (n=1), compared to PMN cell

permeability of S(wt) and F(wt), (n=3); (B) Inhibition of 3 µg/ml of FITC-labeled S(wt) (n=3) and mS*

(n=1) binding to PMN cell by mS; Permeability dose dependencies for (A) and (B) are shown with a

polynomial spline fit, statistical significance indicated between low (0.3 and 0.001 nM) and high (300

nM) toxin concentrations using Student's *t*-test. Error bars indicate SD. (C) Column indicating binding

responses for mF* on hC5aR cells (n=2). *** Indicates a statistically significant difference (p<0.001)

between mS* binding on HEK-hC5aR cells compared to HEK-CCR2 and mF* binding on these cells.

(D) Permeability of hC5aR transfected HEK cells using unlabelled mS and mF, and Alexa-labeled mS*

and mF* compared to wild type S(wt) and F(wt) (n=2). (E) Inhibition of 3 µg/ml of mS* binding by

mS on HEK-hC5aR cells, (n=3). CCR2-transfected HEK cells used as negative controls for toxin

binding and lysis in (C, D) n=2 or (C) one representative experiment. Dose dependency shown with

polynomial spline fit. Statistical significance calculated between low (0.3 and 0.001 nM) and high

(300 nM) toxin concentrations using Student's *t*-test. (F) Permeability response of hC5aR-transfected

HEK cells following incubation with unlabeled mS and Alexa maleimide-labeled mF* or wild type

toxins F(wt) and S(wt) (n=3). Statistical significance calculated between 15 min and 0 min time points

using Student's *t*-test. Error bars indicate SD. Percentages of mean fluorescence intensities is shown as

relative to the maximum intensity in each individual experiment (B, C and E). Permeability of the cells

were analyzed after 30 minutes incubation at +37°C while the inhibition assays were analyzed after 45

minutes incubation at +4°C.

Figure 2. Colocalization of LukS with hC5aR on HEK cells. (A) (left panel) TIRF image of hC5aR-mGFP on the surface of a HEK cell before addition of toxin; (right panels) zoom-in of yellow dashed square of left panel immediately following 2 min incubation with Alexa647-labeled LukSK281CY113H (mS*-Alexa647). (B) Equivalent images of same cell of (B) after > 15 min incubation with LukFK288C (mF). (C) (upper panel) TIRF image of colocalization of Alexa594- and Alexa647-labeled mF and mS (mF*Alexa594 and mS*Alexa647) with hC5aR-mGFP on HEK cells; (lower panel) zoom-in of yellow-dashed square of upper panel with colocalized foci indicated (white arrow).

Additional file 1: Figure S1. Construction of recombinant leukocidin proteins. (A) Schematic of TIRF imaging assay. (B) (left panel) Crystal structure of octameric pore complex of γ-hemolysin (PDB ID:3B07). K273/Y111 and K289 on S and F components of γ-hemolysin corresponds to our engineered mutations, K281C/Y113H and K288C, on LukSF marked in their equivalent places on γ-hemolysin; (right panel) SDS-PAGE of the unlabeled LukSK281CY113H and LukFK288C (mLukS and mLukF) and Alexa-labeled mLukS* and mLukF* toxin components, bands visible at locations consistent with molecular weight of 33 kDa and 34 kDa for LukS and LukF respectively.

Figure 3. Spatiotemporal dynamics of hC5aR, LukS and LukF in live cells. (A) Images of HEK cells treated with LukSK281CY113H (mS) and Alexa-labeled LukFK288C (mF*) showing brightfield (left), hC5aR-mGFP (middle) and mF* (right). (B) Probability distribution for stoichiometry of hC5aR in absence and presence of Alexa-labeled mS (mS*) and mF*, and (C) of mS* foci, indicating (D) tetramer periodicity from Fourier spectral analysis. (E) A random tetramer overlap model cannot account for mS* experimental stoichiometry data ($R^2 < 0$), but a tetramer-multimer model results in

excellent agreement (R^2=0.85). (F) hC5aR and mS* stoichiometry as a function of incubation time. Proportion of immobile and mobile colocalized foci in the (G) presence and absence of mS and mF. Error bars show standard error of the mean from n=5-15 image subregions.

Additional file 2: Figure S2. Fluorescent protein characterization. A. Fluorescence micrograph of immobilized mGFP and an intensity *vs.* time trace for one foci showing a single photobleach step. Raw data in light blue and edge-preserving Chung-Kennedy filtered data in dark blue. B. As A for LukS-Alexa647, C. LukS-Alexa647 micrograph (red) with found foci indicated as white circles. D. Intensity distribution of Alexa 647 foci intensities from whole photobleach experiment showing periodicity at ~3,500 counts on our camera detector. E. Pairwise distance distribution of intensity in D with Fourier spectrum (inset) showing peak at ~4,000 counts. F. GFP foci intensity (natural log) time traces (green) with linear fits (red).

Additional file 3: Figure S3. Density of LukS spots. A. micrograph of mLukS* (white) with found foci (orange circles) B. Probability distribution of overlap frequency using spots in A to calculate density. C. Zoom in of mLukS micrograph. D. Probability distribution of overlap frequency using intensity in C to calculate maximum density estimate.

Additional file 4: Figure S4. Mobility analysis. A. The probability distribution of microscopic diffusion coefficient showing the threshold for immobility as black dotted line and B. the mean squared

1040 displacement against time interval for mobile (upper) and immobile (lower) of hC5aR. C. and D.

1041 similar for mLukS/F*. An insert showing a zoomed in portion of the plot is shown in C. to better

1042 illustrate the division between mobile and immobile.

1043

Figure 4. Relative stoichiometry of hC5aR, LukS and LukF in fixed cells. (A) Micrographs of fixed

hC5aR-mGFP HEK cells treated with LukSK281CY113H (mS) and LukFK288C (mF) showing

hC5aR-mGFP (left) and Alexa647 (middle) and merge (right) on Alexa-labeled mS (mS*) above and

mF (mF*) below (B) Proportion of foci colocalized and not colocalized, treated with mS, mS*+mF and

mS+mF* for hC5aR. Error bars show standard error of the mean from n=4 image subregions. (C)

Heatmap of correlation between hC5aR and mS stoichiometry (red dash line indicates 4 mS per hC5aR

molecule), R^2 ~0.15 (D) and (E) FRET images and efficiencies. The FRET experiment was performed

in live and fixed sortase-tagged FITC-hC5aR expressing cells. Live cells (number of biological

replicates n=2) were incubated in the presence of Cy3-labeled mS* for 1 h at +4°C and washed, after

which unlabeled mF was added. FRET was analyzed before (mS*) or after addition of mF (mS*+mF).

FRET from fixed cells (n=3) was analyzed in the presence of mS* or unlabeled mS and Cy3-labeled

mF* respectively (n=2). Statistical significance between cells with only mS and both of the toxin

components, mS and mF, was analyzed using Student's *t*-test. Error bars indicate SD.

1057

1058 **Additional file 5: Figure S5. Colocalization analysis.** The probability distribution of linked (A) and

1059 unlinked (B) hC5aR and similar for mLukS* (C. and D.). E. and F. False-color heatmap scatter plots

1060 indicating that h5CaR stoichiometry is uncorrelated to mLukS or mLukF stoichiometry in the presence

1061 of mLukF.

1062

1063

Figure 5. LukSF dissociation and rebinding of C5a on hC5aR expressing cells. (A) Inhibition of

anti-CD88 binding on hC5aR expressing HEK cells using increasing concentrations (horizontal axis) of

wild type LukS, S(wt) and C5a. Wild type LukF, F(wt), is used as a negative control for inhibition of

anti-CD88 binding (number of biological repeats n=2). The values are normalized against the

maximum binding observed with only anti-CD88. (B) Disengagement of hC5aR from LukSF was

observed as increase in PE conjugated anti-CD88 binding (right vertical axis, indicated with bars) on

S(wt) pre-coated cells using increasing but sublytic concentrations (horizontal axis, indicated with dots)

of F(wt). The values are normalized against the maximum binding observed with LukS incubated cells

with only anti-CD88. Minimal cell lysis (% of lysed cells, left vertical axis) detected in F(wt)

concentrations below 3 nM (n=3). (C) Rebinding of constant amount of NT647 labeled C5a (647-C5a)

on hC5aR upon LukSF formation analyzed by incubating S(wt) pre-coated cells with increasing

concentrations (horizontal axis) of LukF mutant G130D that associates with LukS but does not lead to

cell lysis (n=2). The values are normalized against the maximum binding observed with only 657-C5a.

(D) Effect of LukSF mediated cell lysis on complement activation and C5a formation on full blood

measured by using C5b-9 as a marker for complement activation in plasma (n=3). Maximal C5a

formation is observed by incubating full blood with live *S. aureus* bacteria. Ecb (B and C), F(wt) (A) or

S(wt) (D) are used as negative controls in the assays. Percentages of mean fluorescence intensities is

shown as relative to the maximal intensity in each individual experiment (A-C). Statistical

significances are calculated using Student's *t*-test. Error bars indicate SD.

Figure 6. Model for LukSF-receptor binding and the mechanism of LukSF-induced

inflammation. (A). LukS (PDB ID: 1T5R) binds on hC5aR (structure based on angiotensin receptor data PDB ID: 4YAY, cyan dashed box) as a soluble monomer on the cell membrane. Each LukS monomer binds one hC5aR molecule via the receptor interacting residues R73, Y184, Y250, T244 (marked with blue dots) within a cluster of approximately 4-5 hC5aR homo-oligomers Upon binding to hC5aR LukS exposes residues for LukF (PDB ID: 1LKF) binding (interface indiacted by dashed ellipse). In these tight clusters each LukF can bind to two LukS monomers via two interfaces. Binding of LukF on LukS and formation of the octameric pore (PDB ID: 3B07) causes dissociation of the receptors from the complex because of leakage of the cell membrane and possibly also since the receptor binding region (marked with a circle) is buried between the monomers in the complex. The detached hC5aR molecule can be reused by its ligands LukS or C5a anaphylatoxin (PDB ID: 1KJS). (B) Zoom out of (A), illustrating the putative mechanism of LukSF induced inflammation.

1083

1084

1085 **Additional file 6: Movie S1 (.avi).** Deposition of mLukS on hC5aR-mGFP HEK cells. The movie is

1086 shown in two clips before (no toxin) and after addition of Alexa 595 labeled mLukS (add mLukS*).

1087 This movie was recorded for *ca.* 13 min and displayed here at 100x speed.

1088

1089 **Additional file 7: Movie S2 (.avi).** Lysis of hC5aR-mGFP HEK cells incubated with Alexa594 labeled

1090 mLukS* and mLukF. The cells were preincubated with mLukS* and the lysis of the cells were

1091 monitored for *ca.* 13 min after addition of mLukF (add mLukF). The red arrow points to the vesicles

1092 released during cell lysis. This movie was recorded for *ca.* 13 min and displayed here at 100x speed.

1093

1094 **Additional file 8: Movie S3 (.avi).** Imaging live hC5aR-mGFP cells. After 1-2 min of exposure,

1095 several distinct, mobile, circular fluorescent foci in the planer membrane regions were observed. Movie

1096 is displayed in real time.

1097

1098 **Additional file 9: Movie S4 (.avi).** Imaging mLukS* incubated with hC5aR-mGFP cells. Several

1099 distinct, mobile, circular fluorescent foci were observed. Movie is displayed in real time.

1100

1101

1102

1103

1104

1105

1106

1107

1108

Figures

Figure 1. Toxin functionality on PMN and HEK cells.

Figure 2. Colocalization of LukS with hC5aR on HEK cells.

Additional file 1: Figure S1. Construction of recombinant leukocidin proteins.

Figure 3. Spatiotemporal dynamics of hC5aR, LukS and LukF in live cells.

Additional file 2: Figure S2. Fluorescent protein characterization.

Additional file 3: Figure S3. Density of LukS spots.

Additional file 4: Figure S4. Mobility analysis.

Figure 4. Relative stoichiometry of hC5aR, LukS and LukF in fixed cells.

Additional file 5: Figure S5. Colocalization analysis.

1109

Figure 5. LukSF dissociation and rebinding of C5a on hC5aR expressing cells.

Figure 6. Model for LukSF-receptor binding and the mechanism of LukSF-induced inflammation.

1110   Table 1. Mean and standard deviation stoichiometry and diffusion constant in live and fixed cells.

1111   Live cells:

	Mean [SD] stoichiometry (molecules)	Number of foci	Mobile foci mean [SD] D (μ^2m/s)	Immobile foci mean [SD] D (μ^2m/s)
C5a	185[224]	5346	0.48[0.43]	0.03[0.03]
C5a+mLukS*	281[213]	8272	0.49[0.42	0.02[0.03]
C5a+mLukS*+mLukF	291[248]	6605	0.46[0.42]	0.03[0.03]
C5a+mLukS+mLukF*	223[202	4981	0.47[0.40]	0.02[0.03]
mLukS*	29[22]	999	0.44[0.45]	0.03[0.03]
mLukS*+mLukF	84[89]	841	0.34[0.35]	0.01[0.02]
mLukS+mLukF*	6[4]	557	0.40[0.44]	0.02[0.03]

1112

1113   Fixed cells:

	Number and (%) of colocalized C5a foci	Number and (%) of C5a foci not colocalized	Mean [SD] stoichiometry of colicalized C5a foci (molecules)	Mean [SD] stoichiometry of C5a foci which are not colocalized (molecules)
C5a+mLukS*	89(32)	193(68)	36[26]	35[24]
C5a+mLukS*+mLukF	84(7)	1079(93)	37[25]	32[24]
C5a+mLukS+mLukF*	59(83)	283(17)	26[20]	26[20]
	Number and (%) of colocalized Luk foci	Number and (%) of Luk foci not colocalized	Mean [SD] stoichiometry of colocalized Luk foci (molecules)	Mean [SD] stoichiometry of Luk foci which are not colocalized (molecules)
mLukS*	88(7)	1198(93)	72[47]	61[44]
mLukS*+mLukF	86(88)	658(12)	121[95]	136[94]
mLukS+mLukF*	60(5)	1186(95)	20[16]	18[15]

1114

1115

1116

1117

1118

1119

1120

1121

1122

Table 2. Cloning sequence details

Clone (restriction site)	Primers, gBlock (5'-3')
3'mGFP(NotI)stop	ATATGCGGCCGCTTATTTGTATAGTTCATCCATG
5'KOZ-hC5aR(BamHI)	ATATGGATCCGCCGCCACCATGAACTCCTTCAATTATAC
5'hC5aR-mGFP	AGACCCAGGCAGTGAGTAAAGGAGAAGAACTTTTC
3'hC5aR-mGFP	TCTCCTTTACTCACTGCCTGGGTCTTCTGGGCCATAG
5'N-sor-hC5aR	CGGGATCCGCCGCCACCATGCTACCCGAGACTGGAGGCGGAGGTGGCAACTCCTTCAATTATACCAC
3'hC5aR-not	ATATGCGGCCGCCTACACTGCCTGGGTCTTCTG
5'LukF-	CGGGATCCGCTCAACATATCACACCTGTAAG
3'LukF-K288C(NotI)	ATATGCGGCCGCTTAGCTCATAGGATTTTTTTCCTTAGATTGAGTATCTATTAAGCAAACTGTATGATTTTCCCAATC
3'LukS-K281C(NotI)	ATATGCGGCCGCTCAATTATGTCCTTTCACGCAAATTTCATGAGTTTTCC
5'LukS-Y113H	GTCAAACATTAGGTCATAACATAGGTGGTAATTTTAATAG
3'LukS-Y113H	TTACCACCTATGTTATGACCTAATGTTTGACTAAC
5'LukS(BamHI)	CGGGATCCAAAGCTGATAACAATATTGAG
LukFG130D gBlock BamHI/ NotI pRSET B C-his overlap	CTTTAAGAAGGAGATATACATATGGGATCCCAACATATCACACCTGTAAGTGAGAAAAAGGTTGATGATAAAATTACTTTGTACAAAACAACTGCAACATCAGATTCCGATAAGTTAAAAATTTCTCAGATTTTAACTTTTAATTTTATTAAAGATAAAAGTTATGATAAAGATACATTAATACTCAAAGCTGCTGGAAACATTTATTCTGGCTATACAAAGCCAAATCCAAAAGACACTATTAGTTCTCAATTTTATTGGGGTTCTAAGTACAACATTTCAATTAATTCAGATTCTAATGACTCAGTAAACGTTGTAGATTATGCACCTAAAAATCAAAATGAAGAATTTCAAGTACAACAAACGGTAGGTTATTCTTATGGTGGAGATATTAATATCTCTAACGGCTTGTCAGGTGATGGTAATGGTTCAAAATCTTTTTCAGAGACAATTAACTATAAACAAGAAAGCTATAGAACTAGCTTAGATAAAAGAACTAATTTCAAAAAAATTGGTTGGG

| | ATGTTGAAGCACATAAAATTATGAATAATGGTTGGGGACCATATGGCAGAGATAGTTATCATTCA ACTTATGGTAATGAAATGTTTTTAGGCTCAAGACAAAGCAACTTAAATGCTGGACAAAACTTCTT GGAATATCACAAAATGCCAGTGTTATCCAGAGGTAACTTCAATCCAGAATTTATTGGTGTCCTAT CTCGAAAACAAAACGCTGCAAAAAAATCAAAAATTACTGTTACTTATCAAAGAGAAATGGATAGA TATACAAACTTTTGGAATCAACTTCACTGGATAGGTAATAATTATAAAGATGAAAATAGAGCAAC TCATACATCAATTTATGAAGTTGATTGGGAAAATCATACAGTTAAATTAATAGATACTCAATCTA AGGAAAAAAATCCTATGAGCGCGGCCGCACACCATCACCATCACCATTAA |

1123

1124