Staphylococcus aureus **toxin LukSF dissociates from its membrane receptor target to enable**

renewed ligand sequestration

Authors:

Karita Haapasalo[1, 2, 4], Adam J. M. Wollman[3, 4], Carla de Haas[1], Kok van Kessel[1], Jos van Strijp[1], Mark

C. Leake[3, 5, *]

[1] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht University,

Utrecht, The Netherlands

[2] Department of Bacteriology and Immunology, and Research Programs Unit, Immunobiology,

University of Helsinki, Helsinki, 00014, Finland

[3] Biological Physical Sciences Institute, Departments of Physics and Biology, University of York,

York, YO10 5DD, United Kingdom

[4] These authors contributed equally

[5] Lead contact: Prof Mark Leake, Biological Physical Sciences Institute, Departments of Physics and

Biology, University of York, York YO10 5DD, UK. Tel: +44 (0)1904322697. Email:

mark.leake@york.ac.uk. Orcid ID**:** http://orcid.org/0000-0002-1715-1249.

[*] Correspondence: Email: mark.leake@york.ac.uk.

Keywords: bacterial toxin; pore formation; single-molecule; super-resolution; immune response.

ABSTRACT

Staphylococcus aureus Panton Valentine Leukocidin (PVL) is a pore-forming toxin targeting the human C5a receptor (hC5aR), enabling this pathogen to battle the immune response by destroying phagocytes through targeted lysis. The mechanisms that contribute to rapid cell lysis are largely unexplored. Here we show that cell lysis may be enabled by a process of toxins targeting receptor clusters and receptor 'recycling' which allows multiple toxin pores to be formed close together. Using live cell single-molecule super-resolution imaging, Förster resonance energy transfer (FRET) and nanoscale total internal reflection fluorescence (TIRF) colocalization microscopy we visualized toxin pore formation in the presence of its natural docking ligand. We demonstrate disassociation of hC5aR from toxin complexes and simultaneous binding of new ligands. This effect may free mobile receptors to amplify hyper inflammatory reactions in early stages of microbial infections and have implications for several other similar bi-component toxins and the design of new antibiotics.

INTRODUCTION

S. aureus causes diseases ranging from superficial skin and soft tissue infections (SSTI) to severe invasive diseases like osteomyelitis and necrotizing pneumonia (1). During the 1960s, methicillin-resistant *Staphylococcus aureus* (MRSA) was identified as a nosocomial pathogen (2). In the 1990s, infection of previously healthy community-dwelling individuals with MRSA was reported (3). Since then, these community-associated (CA-) MRSA have rapidly emerged worldwide (4). Variants have also recently been identified which have reduced susceptibility to the antibiotic vancomycin (5) as well as complete resistance, so-called VRSA (6), and these forms of *S. aureus* pose a significant threat to human health. S. *aureus* and resistant variants have also evolved adaptations to evade attack from cells of the human immune system. However, the molecular processes which underlie these strategies are underexplored in living cells. There are compelling scientific and societal motivations to understand the mechanisms involved in immunogenic evasion strategies of *S. aureus*.

In the early 1930s, Panton and Valentine described a powerful leukocidal toxin produced by multiple *S. aureus* isolates, now denoted Panton-Valentine Leukocidin (PVL), years later shown to be cytotoxic to neutrophils, monocytes and macrophages but not to lymphocytes (7, 8). The majority of CA-MRSA isolates carry the genes encoding PVL, partially due to the successful spread of the PVL carrying clone USA300 in the USA (3, 4, 9, 10), rarely present in hospital-acquired antimicrobial resistant MRSA and methicillin susceptible *S. aureus* isolates. Based on epidemiological studies, PVL is associated with primary skin infections in humans, osteomyelitis and, in particular, severe necrotizing pneumonia (11, 12). Necrotizing pneumonia is a severe complication caused by bacterial lung infection. It is characterized by massive recruitment of neutrophils in the site of infection, diffuse pulmonary inflammation, septic shock, and respiratory failure. Both host factors and microbial

virulence factors are thought to play an important role in the inflammation, however, it is unknown

how the interplay between these two factors affects the severity of the disease (13).

PVL is a pro-phage encoded bi-component, β-barrel pore-forming toxin (β-PFT) comprising protein

subunits LukS and LukF. LukS binding to the surface of target cells induces secondary LukF binding;

chemical and genetic analysis suggests that the resulting complex consists of a lytic pore-forming

hetero-octamer (14, 15). Stoichiometric analysis *in vitro* of this complex suggests it is an octamer of 4-

plus-4 subunits (16). In this complex only LukS is known to interact with the human C5a receptor

(hC5aR, CD88), a G-protein coupled seven-transmembrane receptor (GPCR). LukS targets at least the

extracellular N-terminus of hC5aR (17, 18), similar to the chemotaxis inhibitory protein of *S. aureus*

(CHIPS), but may also interact with the transmembrane receptor region (19). C5aR is the ligand for

C5a, a powerful anaphylatoxin released during complement activation. Complement is a powerful first

line defense mechanism against invading pathogens which can be initiated through three pathways: the

classical, lectin, or alternative pathway. Activation of any of the three pathways on the target leads to a

rapid opsonization with C3b (20). Further activation of complement leads to initiation of the terminal

pathway with release of C5a and formation of membrane attack complexes that are lytic for Gram-

negative but not Gram-positive bacteria (21, 22). Therefore, in defense against Gram-positive bacteria

C3b-opsonization together with attraction and activation of neutrophils via C5a-C5aR interaction are

essential (23, 24). In severe cases, formation of C5a can potentially lead to hyper activation of the

inflammatory response, an inability to regulate this potentially fatal reaction and eventually harm the

human host tissues. Because of this strong pro-inflammatory activity, therapeutic interventions have

recently focused on neutralizing antibodies against C5a and C5aR as potential candidates for the

treatment of severe inflammatory conditions such as bacterial induced sepsis (25, 26).

LukS binding to hC5aR inhibits C5a receptor binding which efficiently blocks neutrophil

activation (17). LukS receptor binding alone is not sufficient for cell lysis but requires simultaneous

interaction between the leukocidin subunits and hC5aR. However, multiple possible subunit and

receptor combinations are theoretically possible and the spatiotemporal dynamics in functional

complexes in live cells between LukS, LukF and hC5aR is not yet known. In addition to PVL *S. aureus*

can produce a number of other β-PFTs with varying receptor and cell type specificities. From these

LukED, LukAB (or LukGH) and γ-hemolysin (composed of two compound pairs, HlgA/HlgB or

HlgC/HlgB) are classified as bi-component toxins like PVL while α-hemolysin is the prototypical β-

PFT that assembles into a pore through the oligomerization of seven monomeric polypeptides (27).

Next to bacterial toxins, an entire group of other pore forming proteins have been identified

which are involved in human innate immunity, indicating that pore-forming proteins are employed in

survival strategies for several types of organisms (28). Development of methods to study dynamic

processes of pore formation by these toxins at a molecular level may improve our understanding of the

evolution of bacterial virulence and human immunity. There are several studies that have attempted to

explain the function of bacterial PFTs, including structural and subunit stoichiometry data from high

resolution X-ray crystallography and single-molecule fluorescence microscopy (16, 29, 30). However,

these studies focused on pathogen instead of host factors and were thereby limited in excluding the

specific interaction between host cell receptor and bacterial toxin component, the first step required for

toxin oligomerization on the host cell membrane and the presence of the most potent factor mediating

the inflammatory response via C5a recognition in the site of infection (17).

Here, we used single-molecule fluorescence detection with super-resolution localization

microscopy (31) to determine protein complex assembly on receptors in live and fixed cell membranes.

We studied human embryonic kidney (HEK) cells modified to express monomeric Green Fluorescent

Protein (mGFP) labeled hC5aR, exposed to Alexa dye-labeled *S. aureus* toxin components LukS and

LukF and imaged using total internal reflection fluorescence (TIRF) real time microscopy (Figure 2-

figure supplement 1a) allowing us to monitor the spatiotemporal dynamics of receptor and toxin

molecules in the cell membrane. Our findings indicate that LukS binds on clusters of membrane-

integrated hC5aRs. The receptor-bound LukS then binds LukF leading to the formation of a pore that is

consistent with previous stoichiometric studies. However, when LukF is bound to the complex, we

observe fewer colocalized hC5aRs with toxin in fixed cells, more immobilized toxin complexes in live

cells and a significantly reduced Förster resonance energy transfer (FRET) signal, indicating,

unexpectedly, that pore formation leads to simultaneous dissociation of the receptors from the complex.

In addition, our biochemical data suggests that the dissociated receptor can then be available for

additional LukS molecules or the C5a generated during complement activation as a response to LukSF

mediated cell lysis. This new finding suggests that a limited number of receptors can be 'recycled' as

docking for further toxin. This ensures that a sufficient number of pores will damage nearby

phagocytic cells, particularly important when high numbers of C5a anaphylatoxin are blocking LukS,

and potentially also enables simultaneous C5a mediated inflammatory response on adjacent cells.

RESULTS

Maleimide-labeled LukSF mediates toxicity on human polymorphonuclear (PMN) and HEK

cells. To study LukSF pore formation on live cells using single-molecule fluorescence microscopy,

single cysteine substitutions on the exposed surface of the cap domain of the toxin complex (Figure 2 –

figure supplement 1b), K281C on LukS and K288C on LukF, were engineered to facilitate maleimide

labeling of LukF and LukS. These were denoted as the modified protein mLukF or mLukS. A second

substitution Y113H on LukS was chosen on the stem domain to facilitate pore formation of the LukS

mutant (mLukS), based on previous studies (16). We compared the lytic activity of these mutants to

their unmodified wild type equivalents by measuring PMN membrane permeabilization after 30 min

toxin exposure using a DAPI fluorescent dye to measure the DNA of lysed cells by flow cytometry.

DAPI does not penetrate intact cell membranes, and is therefore a good measure for cell permeability

and cell death. In this assay each of the wild type toxins was replaced with the modified protein either

unlabeled (mLukF or mLukS) or with a single Alexa647 dye molecule label (mLukF* or mLukS*)

(Figure 1). All modified proteins displayed toxicity phenotypes towards PMN cells with 100% cell

lysis detected at concentrations of 1-10 nM (Figure 1a), compared to wild type equivalents at ~3 nM.

Similarly, both mLukF and mLukF* displayed toxicity towards PMN cells, with the labeled toxin

mLukF* requiring closer to 30 nM concentration for complete lysis of the cells within 30 mincompared

to the wild type LukF(wt). Since the LukS component mediates the toxin recognition on the target cells

we next studied the binding of mLukS or mLukS* on PMNs. In this assay, mLukS was able to inhibit

the interaction of FITC-labeled wild type LukS on PMN cells equally well as the maleimide-labeled

mLukS* (Figure 1b).

The LukS/LukF toxin is known to be specific towards human cells expressing human C5aR

(hC5aR) such as neutrophils, monocytes and macrophages but does not lyse cells that do not express

the receptor (17). To further investigate the specificity of the mutated and labeled toxins on human

cells we prepared HEK cells overexpressing hC5aR, which unlike neutrophils can be easily cultured for

genetic manipulation and optical imaging experiments. We used a monomeric variant of green

fluorescent protein (mGFP) cloned in the C-terminal end of the receptor to report on the spatiotemporal

localization of the receptor and for determining the subunit stoichiometry in observed receptor clusters.

As expected, the toxins lysed only cells expressing hC5aR while the CCR2 expressing cells remained

intact (Figure 1d). We did not observe any binding of mLukF* on the same cells (Figure 1c) which is

consistent with previous observations that LukF in the absence of LukS does not interact with PMN

cells (15). The toxin inhibited binding of mLukS* in a dose-dependent fashion (Figure 1e).

After analyzing the concentration required for efficient LukS binding and cell lysis we wanted

to quantify the dynamics of cell lysis in the presence of mLukF and mLukS. Since the maleimide-

labeled mLukF required up to 10-fold higher concentrations for complete lysis of HEK cells within 30

min compared to PMNs, and because of the loss of molecules during washing cycles, the assay was

optimized to have 20-fold excess of mLukF* (600 nM for mLukF*). Following pre-incubation of

hC5aR-mGFP expressing cells with LukS(wt) or mLukS the second toxin component was added and

the cellular uptake of DAPI was measured as before using flow cytometry. The wild type toxins

LukF(wt) and LukS(wt) caused significant cell lysis (defined as >80% of all sampled cells in the

population) within 10 min, while closer to 20 min was required for significant lysis by the mLukF* and

mLukS toxin combination (Figure 1f).

LukS colocalizes to hC5aR then addition of LukF triggers cell lysis. A similar analysis was

performed by monitoring the lysis of live cells, sampling every 2.5 s at 50 ms exposure time per frame

using total internal reflection fluorescence (TIRF) microscopy, at very low excitation intensity to

prevent photobleaching and facilitate data acquisition of dynamic events involved in the formation of

LukSF nanopores in cell membranes. Here, hC5aR-mGFP cells were first imaged in the absence of

toxin. In the green channel we observed mGFP localization consistent with the cell membrane,

manifest as relatively high apparent brightness towards the cell boundaries consistent with the cell

membrane curving away from the microscope coverslip perpendicular to the TIRF excitation field.

Controlled addition of mLukS* (labeled with Alexa647) to the sample petri dish followed by washing,

while imaging simultaneously throughout, resulted in colocalization of the hC5aR and mLukS (Figure

2, Movie 1). Further addition of mLukF resulted in complete lysis of the cell, as defined by the

observation of explosive release of membrane vesicles, after ~15 min (Figure 2, Movie 2).

Colocalization of hC5aR, mLukS* and mLukF* (labeled with Alexa594) was also confirmed by three

color experiments, imaging cells after addition of toxins and washes until the start of lysis (Figure 2).

LukS forms tetramers and clusters hC5aR before binding LukF. Using higher laser intensity TIRF

excitation enabled rapid millisecond single channel sampling of single fluorophores faster than their

molecular mobility (32), confirmed by imaging antibody-immobilized GFP and Alexa dyes (Figure 3 –

figure supplement 1). Imaging live hC5aR-GFP cells in these conditions saturated the camera CCD but

after 1-2 min of exposure, photobleaching was sufficient to reduce intensity and allow us to observe

several distinct, mobile, circular fluorescent foci at a mean surface density of ~1 per μm^2 in the planer

membrane regions which lie parallel to the TIRF field away from the cell boundaries (Figure 3a, Movie

3). We monitored the spatiotemporal dynamics of foci in the planar membrane regions using automated

tracking software (33) which allowed foci to be tracked for several seconds to a spatial precision of

~40 nm (34). The measured foci width (half width at half maximum determined from their pixel

intensity profile) was in the range 200-300 nm, consistent with the point spread function (PSF) width of

our microscope. By using step-wise photobleaching analysis we estimated stoichiometry values for all

detected fluorescent foci by employing a method which quantifies the initial unbleached foci brightness

and divides this by the measured brightness for the relevant single dye reporter molecule (Figure 3 –

figure supplement 1) (35). These foci contained large numbers of receptors of up to ~2,500 hC5aR

molecules with a mean stoichiometry of ~180 (Figure 3b, Table 1 – source data 1). Addition of mLukS

and mLukF increased the mean stoichiometry by >50% consistent with the toxin causing receptor

clustering.

Imaging mLukS* incubated with hC5aR-GFP cells revealed distinct foci (Figure 3a, Movie 4).

The probability distribution of mLukS* stoichiometry values in live cells in the absence of mLukF is

shown in Figure 3c, rendered using a kernel density estimation which generates an objective

distribution that does not depend upon the size and location of subjective histogram bins (36). We

measured a broad range of stoichiometry values, spanning a range from only a few LukS molecules per

foci to several tens of molecules, with a mean of ~30 molecules per foci. Closer inspection of the

stoichiometry values indicated an underlying periodicity to their probability distribution, which we

investigated using Fourier spectral analysis (37). The resulting power spectrum (Figure 3d) indicated a

fundamental peak equivalent to a stoichiometry of 3.9 ± 0.2 molecules, suggesting that foci are

composed of multiples of tetrameric mLukS* complexes.

Fluorescent foci, if separated by less than the diffraction-limited PSF width of our microscope,

are detected as a single particle but with higher apparent stoichiometry. We therefore tested the

hypothesis that the observed mLukS* foci stoichiometry distribution could be explained by the random

overlap of isolated mLukS* tetramer foci. To do so we modeled the nearest-neighbor separations of

individual mLukS* tetramers in the cell membrane as a random Poisson distribution (38) and used

sensible ranges of tetramer surface density based on our single particle tracking results (Figure 3 –

Figure supplement 2). However, all random tetramer overlap models we explored showed poor

agreement to the observed experimental stoichiometry distribution, but we found that random overlap

of multimers of tetramers could account for the stoichiometry distribution well (Figure 3e). Optimized

fits indicated that the random overlap of mLukS* foci with a stoichiometry in the range ~4-20

molecules were able to best account for the experimental data.

We tested if there was a dependence of foci stoichiometry on incubation time with leukocidin.

Acquiring a time course for mLukF* accumulation following pre-incubation of cells with mLukS was

not feasible since unbound mLukF* had to be washed from the sample to prevent a prohibitively high

fluorescent background. However, we were able to acquire time courses in which mLukF was added to

cells that had been pre-incubated with mLukS*. For these, the mLukS* foci stoichiometry distribution

was measured as a function of time after mLukF addition for several different fields of view, each

containing typically ~5 cells. We found that the mean hC5aR foci stoichiometry indicated no obvious

correlation to mLukF incubation time (Figure 3f), however the mean mLukS* foci stoichiometry

increased with time (p<0.05).

By calculating the mean squared displacement (MSD) as a function of time interval (τ) for each

tracked foci we could determine its microscopic diffusion coefficient (D). The distribution of D for

hC5aR and mLukS/F* (Figure 3 – figure supplement 2) had similar low value peaks at ~0.05 μm^2/s,

consistent with immobile foci tracked with our localization precision spatial precision of 40 nm.

Several mobile foci were also seen, which diffused at rates up to ~5 μm^2/s. Based on the measured

width of the immobile peak width on these distributions we set a threshold of 0.12 μm^2/s to categorize

foci as either immobile, which indicated a mean $D=0.025 \pm 0.030$ μm^2/s ($\pm$SD), or mobile, which

indicated a mean $D=0.47 \pm 0.40$ μm^2/s (Table 1 - source data 1). Plots of the measured MSD *vs.* τ

relations for mobile foci indicated a linear dependence indicative of free Brownian (i.e. 'normal')

diffusion. However, similar plots for immobile foci indicated an asymptotic dependence consistent with

confined diffusion (39), whose plateau was equivalent to a confinement diameter of ~400 nm (Figure 3

– figure supplement 3). The relative proportion of mobile foci was ~35% of tracked foci for hC5aR,

regardless of toxin and similar for mLukS in the absence of mLukF. Addition of mLukF, caused a drop

in the mobile proportion by a factor of ~3 (Figure 3g) suggesting that LukF causes insertion of the

complex and possible disassociation from the hC5aR.

LukSF pores disassociate, allowing further toxin to bind hC5aR. To determine whether the toxin

remains bound to the receptor, and to quantify the relative stoichiometry of components, we imaged

fixed cells, halting cell lysis, using the same two spectrally distinct green/red dyes of mGFP and

Alexa647 to label receptor and toxin components, respectively, as for the live cell experiments. We

imaged cells incubated with mLukS*, followed by incubation with mLukF and mLukS+mLukF* and

observed foci (Figure 4a) with similar stoichiometries (Figure 4 – figure supplement 1, Table 1 - source

data 1) to live cells but colocalized with hC5aR. Only <10% of the toxin was detected colocalized with

hC5aR but ~32% of the hC5aR was found colocalized in the presence of mLukS* dropping to <10% in

the presence of mLukF (Figure 4b). This low percentage was within our estimate of the degree of

random colocalisation between the green and red fluorophores, entirely down to chance, of ~10%. This

suggests that in the presence of mLukF, the toxin is not colocalised with the receptor such that mLukF

causes at disassociation from hC5aR. The stoichiometry values for detected green hC5aR-mGFP foci

were calculated and plotted against the equivalent stoichiometry estimates for colocalized red foci of

mLukS* and mLukF* respectively (Figue 4c and Figure 4 – figure supplement 1). In the presence of

mLukS* but in the absence of mLukF, the hC5aR-mGFP foci stoichiometry showed an approximately

linear dependence on number of associated mLukS* molecules, suggesting that each colocalized

mLukS* molecule was associated on average with ~4-5 hC5aR molecules. In the presence of labeled or

unlabeled mLukF no dependence was observed (Figure 4 – figure supplement 1, $R^2<0$) consistent with

random association between toxin and receptor

We performed FRET experiments on FITC sortase-labeled hC5aR and Cy3-labeled mLukS or

mLukF, as donor and acceptor respectively, in live cells to further probe the association between toxin

and receptor. A FRET signal from whole cells of ~75% efficiency was observed, dropping to ~56%

when incubated with unlabeled mLukF (Figure 4d and e), as would be expected if the complex

formation leads to dissociation of the toxin from the receptor. In order to examine possible FRET

between hC5aR and Cy3-labeled mLukF we performed similar experiments on fixed cells. In these

experiments a FRET efficiency of ~60% was observed between hC5aR and labeled mLukS dropping

below 40% between hC5aR and labeled mLukF. These results are also consistent with disassociation of

hC5aR from the pore, although conformational or local environment changes cannot be ruled out with

FRET alone since the relatively high remaining signal might in principle also indicate remaining

association or other inter or intra hC5aR-Luk interaction. The greater drop in FRET when measured

with mLukF compared to mLukS might be caused by the 3-4 nm further distance of LukF from hC5aR.

The collated findings taken altogether, however, from the changes in mobility, colocalization and

FRET with addition of mLukF are strongly indicative of disassociation of the LukSF complex.

To further confirm that the LukSF complex dissociates from the target receptor we used a

monoclonal PE-labeled anti-CD88 antibody to detect the increase of free hC5aR receptors on the cell

membrane upon LukSF formation. When C5a or wild type LukS was incubated with the hC5aR

expressing HEK cells without LukSF both ligands showed clear inhibition of anti-CD88 binding at 100

nM concentrations (Figure 5a). However, when the hC5aR expressing cells were incubated with 100

nM of LukS followed by incubation with increasing concentrations of wild type LukF a statistically

significant increase in anti-CD88 binding was detected at a LukF concentration of 1 nM when

compared to a 0 nM LukF concentration. This indicates that this was a "sublytic" concentration at

which rebinding of anti-CD88 could be detected without significant cell lysis (% of lysed cells < 10%)

(Figure 5b). At a 3 nM LukF concentration the proportion of dead cells increased above 10% of all

measured cells.

Since C5a is the natural ligand for hC5aR and can outcompete binding of LukS on the receptor

we next analyzed whether LukSF formation and disengagement of hC5aR would allow rebinding of

C5a on the receptor. To detect C5a rebinding at higher LukF concentrations we used a G130D LukF

mutant that interacts with LukS but does not cause cell lysis. Non lytic activity of this mutant in this

assay was confirmed by DAPI staining that showed minimal cell lysis even at higher concentrations

(9% at a 300 nM LukFG130D concentration). At a 300 nM concentration a significant increase in C5a

binding was detected indicating that LukSF dissociates from hC5aR enabling simultaneous rebinding

of an hC5aR interacting ligand (Figure 5c). This assay showed that C5a can potentially interact with

these cells that are attacked by LukSF. Because C5a is a potent anaphylatoxin that is generated during

complement activation and potentially plays a crucial role in *S. aureus* infections (41) we next analyzed

whether LukSF could lead to complement activation and C5a formation in an *ex vivo* full blood assay.

We used soluble C5b-9 as a marker for terminal complement activation and C5a formation, not C5a,

because LukS is known to compete with C5a for binding to hC5aR on neutrophils (17). The presence

of 200 nM of LukSF clearly increased formation of soluble C5b-9 compared to full blood without any

toxin or only LukS (Figure 5d), indicating that LukSF mediated cell lysis increases C5a formation and

potentially also inflammation in the site of infection.

DISCUSSION

S. aureus toxin components LukF and LukS are encoded by two co-transcribed genes of a prophage

integrated in the *S. aureus* chromosome (42). Most of the *S. aureus* clinical isolates investigated have

the genes encoding α-hemolysin, γ-hemolysin, LukAB and LukED but only 5% of those have LukSF

(PVL). This ratio for PVL, however, is different among CA-MRSA isolates where 85% of the strains

carry the *pvl* gene linking the toxin epidemiologically to these strains (10). The specificity to cell

surface receptors makes it difficult to study PVL's role in *S. aureus* pathogenesis in a whole animal

model. It is possible that lysis of neutrophils by PVL is responsible for a reduced host defense

mechanisms allowing the pathogen to spread and cause eventual tissue damage. However, a previous

study using a rabbit animal model on necrotizing pneumonia suggests that PVL itself directly or

indirectly causes tissue injury and by this way induces local inflammation (43). Our finding that the

toxin complexes are found in receptor clusters indicates that lysis of cells depends on the local density

of close proximity hC5a receptors that will initiate the pore formation process by docking LukS close

to the cell membrane such that four hC5aR-LukS dimers (assuming that one LukS binds one hC5aR)

can interact with the free non-bound LukF that will eventually form an octamer (i.e. 4 by 4) and a

functional pore with LukS. These data also suggest that when proper assembly in an octamer is

ongoing/complete, hC5aR will dissociate from the complex and, at the same time, interact with the C5a

that is formed during complement activation amplified by LukSF-mediated cell lysis. This is logical

because, in addition to invading microbes, apoptotic and necrotic cells are known to activate

complement (44). The triggering of local complement activation by toxin damaged cells and the release

of locally generated C5a (Figure 5) and its interaction with adjacent cells such as endothelial or lung

epithelial cells (45) could explain the mechanism behind the exacerbated inflammation characteristics

exhibited in necrotizing pneumonia. This is an important finding, suggesting that the cause of infection

can dramatically affect the magnitude of the inflammatory response and is highly dependent on the

dynamics of microbial molecules interacting with human receptors. In addition, the disengaged hC5aR

is possibly available for new toxins to bind, as C5a has been shown to here, thus allowing the receptor

to be recycled and reused by additional LukS molecules. Our finding that C5a can rebind indicates that

the dissociated free hC5aR does not change in conformation due to previous contact with LukS and

therefore would also be available for binding with LukS. We find that roughly half of LukS complexes

are immobile prior to LukF binding, but that LukF binding then results in mostly immobile LukSF

complexes. Speculatively, this result may suggest that LukS binds hC5aR initially and then inserts

itself transiently in the membrane phospholipid bilayer via the exposed hydrophobic residues,

following binding of LukF molecules to LukS. This conformation would in principle then lead to pore

formation across the whole cell membrane, thus resulting in a more stable insertion of the LukSF

complex into the cell membrane.

Human C5aR expressing cells such as human PMNs are susceptible to LukSF-mediated lysis

and we show that the interaction between maleimide-labeled toxin component LukS and the cell

surface receptor is required for the target recognition and cell lysis similarly as shown before for wild

type LukS (17). By characterizing the mobility of hC5aR and LukS in live cells we find that roughly

half of hC5aR and LukS foci diffuse relatively freely in the cell membrane while the remainder, rising

to > 90% of LukS when LukF is present, are confined to zones in the membrane of ~400 nm effective

diameter. We cannot directly determine the cause of this confinement in our present study. However, if

LukS were to undergo a conformational change following LukF binding then this may potentially

expose hydrophobic residues that could then insert the toxin into the hydrophobic interior of the

phospholipid bilayer; in other words, hydrophobobicity-mediated anchoring. This hypothesis is

strongly supported by the β-barrel prepore-pore formation putative mechanism of γ-hemolysin. Here

the residues responsible for binding with the phospholipid head group are located at the bottom of the

rim domain whereas the stem domain forms an antiparallel β-barrel of which the bottom half comprises

the transmembrane portion of the pore (30). This change from receptor associated LukS to cell

membrane associated LukSF complex can be seen as a change in the proportion of mobile (receptor

associated LukS) and immobile (toxin complexes inserted into cell membrane) foci detected in live

cells. GPCRs similarly are known to have heterogeneous mobility and lateral distribution properties in

living cells at different states for example before and after activation (46).

Crystallographic evidence from the monomeric LukF and LukS components and the intact γ-

hemolysin pore suggests that the pore is octameric formed from 4-plus-4 LukF/LukS subunits (29, 47,

48). Our findings support this octamer model but unlike previous studies also indicate that higher

stoichiometry clusters are formed on live cell membranes which have a measured periodicity of

stoichiometry equivalent to 4 LukS molecules, which cannot be accounted for by random overlap of the

diffraction-limited fluorescence images of individual tetrameric LukS or LukF complexes, but are

explained by a model which assumes the presence of octamer clusters. Each octamer component

consists of cap, rim and stem domains. Here, the cap domain contains the site for LukS/LukF

interaction while the stem domain unfolds and forms the transmembrane β-barrel upon pore formation.

Within crystallization the 2-methyl-2,4-pentanediol (MPD) molecules are bound at the base of the rim

domain, and recognized by Trp177 and Arg198 residues, that may participate in recognition of the

phospholipid bilayer as suggested in a crystal structure of the LukF monomer (49). In contrast, the

structure of the γ-hemolysin suggests a membrane interaction site within residues Tyr117, Phe119 and

Phe139 on the same toxin component (29). The crystal structure of LukED determined recently reveals

important details of the residues on LukE required for receptor identification (50). This component

corresponds to the receptor binding component LukS on the LukSF complex, scanning mutagenesis

indicting that LukS residues Arg73, Tyr184, Thr244, His245 and Tyr250, and to a lesser extent Tyr181,

Arg242 and Tyr246, are involved in binding to the neutrophil surface (51).

To determine the stoichiometry of the toxin components without immobilizing the protein on a

surface or within a crystal we implemented our fluorescence imaging method which allows us to

monitor the actual pore formation mechanism within a living cell, including the target receptor crucial

for the complex formation. This kind of study on protein complex formation has not been done before

due primarily to the difficulty of labeling the components and the high native fluorescence background

in mammalian cells. Our covalent labeling strategy and high excitation intensity TIRF microscopy,

combined with advanced image analysis tools, opens the way for further studies into many other pore

forming toxins and processes involving membrane bound protein complex formation.

Fourier spectral analysis combined with foci overlap modeling suggests that LukS complexes

comprise a multimer of ~4-5 hC5aR subunits each of which contain 4 LukS molecules. Our findings

are consistent with the hetero-octamer model of 4-plus-4 LukS/LukF subunits (16, 29, 30), but with the

refinement that 4-5 hetero-octamers associate as higher order multimers in the functional cell

membrane. However, our colocalization analysis also indicates the presence of stable complexes of

LukS with hC5aR independent of LukF. Similarly, we observe clusters of pre-established hC5aR in

the absence of LukS or LukF, but the addition of LukS or LukF significantly increased the mean cluster

stoichiometry. These results suggest that further binding sites for hC5aR on LukS could be possible in

addition to those identified in the LukS rim domain (51). However, since the binding of LukS to

neutrophils is inhibited by the C5a it is likely that LukS has only one binding site on the receptor (19).

This is also supported by the similar inhibition profiles of LukS and C5a towards anti-CD88 binding on

hC5aR shown in this study. Therefore, the association of LukS with approximately 4-5 hC5aR

molecules could be explained by the previous suggestion that C5aR forms homo-oligomers in living

cells (52). Our findings imply that LukSF assembly is dependent on hC5aR cell membrane area density

as opposed to the effective hC5aR concentration when calculated over in the whole of a target cell

volume, such that even when hC5aR cellular expression levels are low a cell lysis response may still be

achieved through the efficient targeting of receptor clusters and recycling of the receptor molecules in

the cell membrane to be re-used by free non-bound LukS to get engaged in octamer pore formation.

Previous *in vitro* studies on LukSF pores formed on human leukocytes and rabbit erythrocytes

have found evidence for both octamers and hexamers, but importantly both suggest a LukF/LukS ratio

of 1:1 (16, 53, 54). Interestingly we did not observe any correlation to the number of hC5aR present

with LukF incubation time once LukF was already bound to LukS. Moreover, when LukS was

incubated with LukF using sortase-labeled hC5aR cells a significant reduction was observed in the

FRET efficiency signal. It is unlikely that the reduction that was observed in FRET efficiency would be

due to a conformational change because the cysteine mutation used for maleimide labeling was

designed to be exposed on the cap domain of LukS and LukF (Figure 2 – figure supplement 1) that in

light of the structural data undergoes minimal or no conformational changes during complex assembly

(30). Since our biochemical assays indicate that LukF does not bind directly to hC5aR expressing cells

and, that binding of LukF to LukS results in an increased distance between the receptor and the

complex, this suggests that LukF binding to LukS results in LukS dissociating from the receptor,

released as a newly formed LukSF complex.

We cannot directly determine the cause of this behavior in our present study, however one

explanation may lie in the conformational change during the prepore-to-pore transition that has been

shown to occur on γ-hemolysin complexes subsequently after binding of LukF to LukS (29, 30).

Interestingly, this same study shows that during the pre-pore state the space for the transmembrane

region is occupied by the rim domain of the adjacent octamer in a LukSF crystal. One explanation for

these observations, that remains to be explored, is that in addition to the stem domain the residues

within the rim domain that interact with the receptor might also have different orientations in the pre-

pore state when compared to the pore state. The putative dissociation of the hC5aR from the LukSF

complex was further verified by incubating LukS-coated hC5aR cells with increasing concentrations of

LukF. Here, increase in anti-CD88 binding also clearly indicates LukSF dissociation. This kind of

receptor disengagement has been shown before by at least the cytotoxin intermedilysin which interacts

with a GPI-anchored complement regulatory molecule on the cell membrane (55). Moreover,

dissociation of LukSF complex is also supported by electron microscopy of LukSF on human leukocyte

membrane fragments. Here, the ring-shaped oligomers with outer and inner diameters of 9 nm and

3 nm were shown without a receptor (54).

In this study we also show, for the first time to our knowledge, that the dissociated receptor can

be reused by free unbound C5a. This indicates that LukS binding on the receptor does not change

receptor conformation and thereby can putatively also be reused by additional LukS. In our full blood

model we observed that LukSF mediated cell lysis clearly increased complement activation and C5a

formation. The increase in C5a concentration in the site of infection could potentially limit the

availability of hC5aR for LukS molecules on neutrophils and thereby reduce lytic activity of the toxin

as C5a has previously been shown to reduce LukSF mediated lysis *in vitro* (17). Rebinding of C5a on

the receptor may therefore indicate that in natural settings where all components (i.e. LukS, LukF and

C5a) are present C5a can outcompete binding of LukSF on the target cells. Therefore, recycling of the

receptor could be one strategy for the toxin to ensure that a sufficient number of pores will damage the

cells especially when limited number of receptors are available.

There are several steps on the leukocidin complex assembly that may be critical for the function

of the toxin. Based on our observations we provide new information on leukocidin-receptor interactions

and propose two additional stages to the processes of pore formation and the mechanism by which

LukSF potentially induces inflammation (Figure 6). Stage 1 is the binding of LukS to hC5aR clusters.

The first step in this process is the target recognition of LukS binding to the membrane receptor. The

LukS-hC5aR complexes were detected on clusters of receptors indicating that pore formation takes

place in these clusters. Stage 2 is the dissociation of the LukSF complex from the receptors and

rebinding of free unbound ligand back to the receptor. We measured a correlation between the number

of LukS and hC5aR molecules present in LukS-hC5aR complexes, but with no obvious correlation

between the number of LukF and hC5aR molecules when LukF was added to the LukS-hC5aR

complex. In addition to previous studies (43) we suggest that LukSF-mediated cell lysis and

dissociation from hC5aR can potentially amplify *S. aureus* mediated inflammation in the site of

infection (Figure 6). The direct lysis of neutrophils is enhanced by newly formed LukSF complexes

that are formed on the cell membrane hC5aR via reattachment of new LukS. Neutrophil lysis activates

the complement system and the newly generated C5a induces cytokine/chemokine production and

neutrophil chemotaxis via the C5a/C5aR signaling pathway on adjacent cells. Furthermore, the

increased vasodilation and vascular permeability leads to massive neutrophil accumulation and tissue

injury at the site of bacterial infection (56).

In summary, our findings that the receptors of targeted host cells dissociate rapidly from the

leukocidin complex upon formation of a harmful toxin pore, freeing up mobile receptor 'seeds' that can

diffuse to other parts of the cell membrane, suggest a hitherto undiscovered strategy used by microbes

to kill human immune cells that enables a limited number of receptors to be recycled as docking for the

leukocidin or potentially the anaphylatoxin C5a to ensure that enough pores will form to damage the

host cell and simultaneously maintain or possibly amplify the inflammation in the site of infection. This

discovery may generalize to other bi-component toxins which employ a similar docking receptor like

the C5aR receptor, including the family of *Staphylococcal* bi-component leukocidins of HlgC/HlgB,

HlgA/HlgB, LukE/LukD (CXCR1, CXCR2, CCR5), and LukM/LukF' for bovine CCR2. These results

highlight the importance of leukocidin-receptor interactions in pore formation and may facilitate further

understanding in the role of pore-forming toxins in *S. aureus* infections. This new mechanistic insight

may prove valuable to the development of future antibacterial and anti-inflammatory therapies,

especially important in light of the growing menace of global antimicrobial resistance.

MATERIALS AND METHODS

 Experimental model and subject details

PMN isolation, Cell Lines, and Transfections. Human blood was obtained from healthy volunteers

and the polymorphonuclear (PMN) cells were isolated by Ficoll/Histopaque centrifugation (57).

Informed consent was obtained from all subjects, in accordance with the Declaration of Helsinki and

the Medical Ethics Committee of the University Medical Center Utrecht (METC-protocol 07-125/C

approved 1 March 2010). A fusion construct of hC5aR with the monomeric GFP variant mGFP with

A206K mutation (also denoted GFPmut3) (58) was made at the C-terminus (primers used listed in

Table 2 - source data 2) or a sortase A LPXTGG sequence was made in the N-terminus and cloned into

pIRESpuro vectors (Table 2 - source data 2) by PCR. The amplification reaction was performed in

three separate amplification steps using overlap extension PCR on hC5aR and mGFP templates. hC5aR

(Accession number of human C5aR = NM_00173) was used as the template using enzymes and

purification kits as described above. The clones were ligated into the vectors and transferred into

TOP10 *E. coli* competent cells, then amplified and sequenced similarly to the toxin clones described

previously. The pIRESpuro/hC5aR-mGFP vector was transfected into Human Embryonic Kidney

(HEK) 293T cells (a HEK cell line, Invitrogen), stably expressing G protein Gα16, using

Lipofectamine-2000 Reagent according to manufacturer's instructions (Thermo Fisher Scientific).

After 24–48 hr, transfected cells were harvested with 0.05% trypsin. To obtain a uniform, stable

culture, cells were sub-cloned in a concentration of 0.5 cells/well in a 96-well plate in Dulbecco's

modified Eagle's medium (DMEM, Lonza) supplemented with 10% fetal calf serum (Invitrogen) 100

U/ml penicillin/ 100 µg/ml streptomycin (PS; Invitrogen), 1 µg/ml Hygromycin and 250 µg/ml

Puromycin. For N-terminal labelling of the sortase A recognition sequence containing HEK cells with

FITC were successfully performed in two steps as described previously (59). The expression of hC5aR

was analyzed by incubating the cells in 50 µl RPMI (Invitrogen) supplemented with 0.05% human

serum albumin (Sanquin), RPMI-HSA, at 5×10^6 cell/ml concentration for 45 min with PE-conjugated

anti-CD88 and detected by flow cytometry. The presence of mGFP or FITC-LPXTG was detected

directly by flow cytometry.

Recombinant Protein Production and Purification. Polyhistidine-tagged LukS and LukF were cloned

and expressed using an *E. coli* expression system. For maleimide-based labeling a single-cysteine

mutation was designed to the LukS and LukF components based on previous data and the crystal

structure of the octameric pore (29). An additional mutation Y113H was included in LukS to facilitate

oligomerization of the maleimide-labeled protein (16). The target genes were amplified by PCR (Table

2 - source data 2) from the wild type sequences using Phusion High-Fidelity DNA polymerase (Thermo

Scientific) (17). The PCR product was cloned into a slightly modified pRSET expression vector

(Invitrogen), resulting in expression of proteins with an N-terminal 6xHIS-tag. For LukF mutant

G130D we useda gBlock (custom dsDNA sequence via Integrated DNA Technologies) to incorporate

the LukF in the pRSET vector. Clones were sequenced to verify the correct sequence. The recombinant

proteins were expressed in Rosetta Gami (DE3) pLysS *E. coli* using 1mM IPTG induction and isolated

by a native isolation method. The expressed proteins were purified according to the manufacturer's

instructions (Invitrogen) using 1 ml Nickel HisTrap and Superdex 75 HiLoad columns (GE Health Care

Life Sciences). Toxin components were labeled with either Cy3 (GE Healthcare), Alexa Fluor® 594 or

Alexa Fluor® 647 C_2 Maleimide reagent according to the manufacturer's instructions (Thermo

Scientific) resulting in negligible unlabeled content. The labeling efficiency was 100% as determined

by protein concentrations using absorption at A280 and dye concentrations using absorption at A650 by

a Nanodrop ND-1000 Spectrophotometer.

Method details

Binding Assays. Binding of the maleimide-labeled proteins to PMN and HEK cells was confirmed by

flow cytometry. LukS-K281C-Y113H (mLukS) or wild type, LukS (wt) was labeled with FITC or

Alexa Fluor maleimide 647 or 594. For competition assays 3 µg/ml of the labeled protein and

increasing concentration of non-labeled mLukS or LukS(wt) was incubated with isolated PMNs or

HEK hC5aR-mGFP cells (5×10^6 cell/ml) in a total volume of 50 µl RPMI-HSA on ice. For binding

assays without competition the cells were incubated with increasing concentration of mLukF*. After

30 min incubation on ice, cells were washed, fixed with 1% paraformaldehyde and analyzed by flow

cytometry. HEK cells transfected with CCR2 receptor were used as negative control for mLukS

binding. To see inhibition of PE anti-CD88 (BD biosciences) binding by LukS(wt) or C5a hC5aR

expressing HEK cells were first incubated with increasing concentrations of LukS or C5a for 45 min at

4°C. Then 2 µl of anti-CD88/200,000 cells was added and incubated as previously. Cells were washed

once with RPMI-HSA, fixed with 1% paraformaldehyde and analyzed by flow cytometry. To detect

hC5aR dissociation using sublytic concentrations of LukSF hC5aR expressing HEK cells were

incubated with 100 nM of wild type LukS for 45 min at 4°C. After washing the unbound LukS sublytic

concentrations of wild type LukF was added to the cells and incubated for 20 min at 37°C and 5% CO_2

atmosphere. Percentages of lysed vs. non lysed cells were measured by using 1µg/ml of DAPI in the

reaction. For C5a rebinding assay 1 µM of C5a (Sigma) was labeled with NT-647 according to

manufacturer's instructions (Monolith NT™). Free label from the sample was removed by three times

centrifugation trough Amicon Ultra 0.5 mL centrifugal filters (Sigma). The hC5aR expressing HEK

cells were incubated with 1 µM of wild type LukS for 45 min at 4°C. After washing 20 nM of NT647-

C5a and increasing concentrations of wild type LukF was added to the cells and incubated for 20 min at

37°C and 5% CO_2 atmosphere. Cells were washed once with RPMI-HSA, fixed with 1%

paraformaldehyde and analyzed by flow cytometry. Percentages of lysed vs. non lysed cells were

measured by using 1µg/ml of DAPI in the reaction. *S. aureus* Ecb (extracellular complement binding

protein) was used as negative control as it interacts with another cell surface receptor, CR1 (60). Flow

cytometry data were analyzed using FlowJo v10 software package.

Cell Permeability Assays. Isolated PMNs or HEK hC5aR-mGFP cells (5×10^6 cell/ml) were exposed to

labeled and unlabeled mixtures as appropriate of mLukF/mLukS recombinant proteins at equimolar

concentrations in a volume of 50 µl RPMI-HSA with 1 µg/ml of DAPI. Cells were incubated for

30 min at 37°C with 5% CO_2 and subsequently analyzed by flow cytometry. To calculate the lysis time

cells were first incubated with 150 nM of mLukS for 15 min. Then 600 nM of mLukF was added and

immediately subjected to flow cytometry analysis where the permeability was measured at several time

points. Cell lysis was defined as intracellular staining by DAPI. HEK cells transfected with human

CCR2 receptor was used as negative control for toxin-mediated lysis. Statistical differences between

means of repeated experiments were calculated using two-tailed Student *t*-tests.

Ex vivo complement activation assay. To maintain complement activity the blood samples were

anticoagulated with lepirudin (Refludan, Schering, Berlin, Germany). Increasing concentrations of

LukSF or LukS (0 to 2000 nM) was incubated in full blood for 30 min at 37°C under continuous

rotation (300 rpm). Complement activity was stopped by adding 10 mM EDTA in the suspension and

the plasma was separated from the blood cells by centrifugation at $5000 \times g$. A 1:30 dilution of each

plasma sample was analyzed by SC5b 9 Enzyme Immunoassay according to manufacturer's

instructions (MicroVue SC5b 9 Plus Enzyme Immunoassay, Quidel). One *S. aureus* colony (1×10^8

cells) was used as a positive control for SC5b-9 formation. The bacteria were grown over night on

blood agar plate at 37°C 5% CO_2 atmosphere.

Fluorescence microscopy. Cells were imaged using a Nikon A1R/STORM microscope utilizing a x100

NA oil immersion Nikon TIRF objective lens. We used a total internal reflection fluorescence (TIRF)

microscopy module. We used laser excitation at wavelengths 488 nm (for mGFP), 561 nm (for

Alexa594) and 647 nm (for Alexa647) from a commercial multi laser unit fiber-coupled into the

microscope, capable of delivering maximum power outputs up to ~200 mW, with a depth of

penetration in the range ~100-130 nm for the TIRF excitation evanescent field. Fluorescent images

acquired on an iXon+ 512 EMCCD camera detector (Andor) at a magnification of 150nm/pixel. Green

and red channel images were obtained by imaging through separate GFP or Alexa647 filter sets. For

high laser excitation intensity single-molecule millisecond imaging, green channel images to determine

mGFP localization were acquired continuously using 488 nm wavelength laser excitation over a period

of ~5 min through a GFP filter set, then the filter set was manually switched to Alexa647 as for red

channel images acquisition continuously using 647 nm wavelength laser excitation until complete

photobleaching of the sample after 1-2 min. For photobleaching laser powers ranged between 15 mW

(Alexa 647) to 100 mW (mGFP). For fixed cell analysis cells were either incubated first with mLukS or

mLukS*, washed and incubated with mLukF or mLukF*, or incubated first just with mLukS* with

mLukF* absent, then washed and fixed with 1% paraformaldehyde.

For fluorescence imaging the HEK cells were grown on 0.1% poly-L-lysine which coated

8-well chambered cover glass slides (Ibidi) in standard growth conditions described above. To analyze

the deposition of mLukS* on live cells the cells were first imaged in PBS buffer in the absence of

toxin. Here, a 256×256 pixel area covered approximately one cell per field of view. Then the cells

were incubated for 2 min with 5 µg/ml of Alexa594 maleimide-labeled LukS in RPMI-HSA and the

cells were carefully washed with PBS keeping the imaging area and focus constant. Because of the fast

bleaching of the Alexa647 label a more stable Alexa594 label was used for the LukS deposition

imaging. The deposition of mLukS was detected for 10 min and the lysis of the cell was recorded for

15 min after addition of 600 nM of unlabeled mLukF. Cells were imaged in TIRF at 50 ms per frame

with the laser automatically switched between 488 nm/0.22 mW, 647 nm/3 mW and 561 nm/3 mW or

488 nm/0.22 mW and 561 nm/3 mW (Figure 2).

FRET experiments. The Cy3-labeled LukS or non-labeled LukS and Cy3-labeled LukF or non-labeled

LukF and controls (only FITC-labeled cells or unlabeled cells with only Cy3-labeled LukS) were

incubated at 4°C for 30 min and then 10 min at 37°C, washed 2 x with RPMI-HSA and fixed as before.

The cells were in PBS during imaging. FITC experiments were performed using Leica TCS SP5

microscope, using a 62× oil immersion objective lens, and FRET Sensitized Emission Wizard in Leica

Application Suite Advanced Fluorescence (LAS AF). Images were acquired using 488 nm and 543 nm

wavelength lasers and a laser power of 27% 12.0 (A.U.) and a scan size of 512×512, 800 ms, 50 ms

per frame, beam splitter TD 488/543/633.

FRET efficiency ε was calculated using the donor, directly excited acceptor and donor excited

acceptor intensity from n=5-10 manual regions of interest inside cells for each experiment, using the

following formula (61):

$$\varepsilon = \frac{\beta(B - D) - \gamma A}{A}$$

604   B = Intensity signal, donor excited acceptor
605   D = Intensity signal, donor
606   A = Intensity Signal, directly excited acceptor
607   β = calibration for ratio of measured intensities of $B_{donor\ channel}/A_{donor\ channel}$
608   γ = calibration for ratio of measured intensities of $B_{acceptor\ channel}/A_{acceptor\ channel}$
609

Single-molecule imaging of live and fixed cells. GFP and Alexa647 fluorescence micrograph time

series of fixed and live cells were sampled taken at. 20 ms per frame. Green channel images were

acquired continuously using 488 nm wavelength laser excitation over a period of *ca.* 5 min via the GFP

filter set. Then the filter set was manually switched to that for Alexa647 and red channel images were

acquired continuously using 647 nm wavelength laser excitation was until complete photobleaching of

the sample after 1-2 min. The step-wise single-molecule fluorescence photobleaching was analyzed

both for live and fixed cells. For live cell photobleaching analysis the cells were incubated with 150 nM

of labeled or unlabeled mLukS as required for *ca.* 15 min. After washing with PBS, 600 nM of labeled

or unlabeled mLukF was added and the imaging was done immediately within 10-15 min. If labeled

LukF was added the wells were washed with PBS before analysis. Also samples with only LukS and

without toxins were analyzed. For fixed cell analysis the cells were incubated first with mLukS or

mLukS* for 30 min at +4°C in RPMI-HSA, washed with same buffer and incubated for 10 min at 37°C

with mLukF or mLukF*, or the same protocol was followed but using mLukS* alone with mLukF*

absent. Then the cells were washed and fixed with 1% paraformaldehyde. 1M mercaptoethylamine

(MEA) buffer was used for fixed cell analysis. Photobleaching of recombinant mGFP and mLukS-

Alexa647 were also separately analyzed in a tunnel slide comprising two pieces of double-sided tape

forming a channel sandwiched between a standard glass microscope slide and a plasma cleaned

coverslip. Proteins solutions (1μg/ml) were immobilized onto the coverslip coated by anti-GFP or anti-

His antibodies respectively with phosphate buffered saline washes in between.

Quantification and statistical analysis

Binding and permeability assays. Statistical significance between repeated (n>1) experiments was analyzed using two-tailed Student's *t*-tests where using a standard p-value threshold of <0.05 as indicating statistical significance. Means and standard deviations of repeated experiments are shown in error bars, unless indicated otherwise.

Image analysis. Basic image extraction, cropping and quantification was done using NIS-Elements microscope imaging software and Image J. More advanced foci tracking was done using bespoke software written in MATLAB (Mathworks) (33) which enabled automatic detection and localization of individual fluorescent foci to within 40 nm lateral precision (Figure 3 – figure supplement 1a). The software identifies candidate foci by a combination of pixel intensity thresholding and image transformation. The intensity centroid and characteristic intensity, defined as the sum of the pixel intensities inside a 5-pixel radius circular region of interest around the foci intensity centroid minus the local background and corrected for non-uniformity in the excitation field are determined by repeated Gaussian masking. If the signal-to-noise ratio of a foci (the intensity per pixel/background standard deviation per pixel) is greater than a pre-set threshold, nominally here set at 0.4 based on prior simulations, it is accepted and fitted with a 2D radial Gaussian function to determine its width. Foci in consecutive frames within a single PSF width, and not different in intensity or width by greater than a factor of two, are linked into the same track.

Foci intensity was used to quantify stoichiometry information. As foci photobleach over time during continuous laser excitation their intensity falls in a stepwise manner due to photobleaching of an integer number of fluorophore tags in each sampling time window. By quantifying the size of a single step, the characteristic intensity of a single fluorophore can be obtained and thus the stoichiometry of

the foci from its initial intensity. The step size is found from the periodicity in the distribution of foci

intensities corroborated by the pairwise distance (PwD) distribution of these intensities and the Fourier

spectrum of the PwD which contains peaks at the characteristic intensity and harmonics at multiples of

this value (Figure 3- figure supplement 1d-e).

Here, the copy number of hC5aR-mGFP was comparatively high such that the TIRF images

were initially saturated in regards to pixel intensity output. After ~20 s of photobleaching the non-

saturated foci intensity values were fitted by an exponential function which characterized the rate of

intensity decay, equivalent to an exponential photobleach time of ~20 s, and extrapolated back to zero

time to determine the initial foci intensity (Figure 3 – figure supplement 1f). The Alexa647 dye also

bleached during 647 nm wavelength laser excitation but images were not initially saturated, but also in

some images which were exposed to the 488nm laser and then the 647nm laser, also bleached by the

488 nm wavelength laser. In these images a fixed correction factor of 6x, determined by comparing to

images exposed to the 647 nm laser first, was used. The stoichiometry of each foci was then

determined as the initial intensity divided by the intensity of the appropriate single fluorescent dye tag

(i.e. either mGFP or Alexa647 in this case).

We characterized the mobility of tracked foci by calculating their MSD as a function of time

interval (τ). For each detected foci the MSD was calculated from the measured intensity centroid

($x(t)$,y(t)) at time t assuming a foci track of N consecutive image frames at a time interval $\tau=n\Delta t$ where

n is a positive integer and Δt is the frame integration time (here 20 ms):

$$MSD(\tau) = MSD(n\Delta t) = \frac{1}{N-1-n} \sum_{i=1}^{N-1-n} \left(\left(x(i\Delta t + n\Delta t) - x(i\Delta t)\right)^2 + \left(y(i\Delta t + n\Delta t) - y(i\Delta t)\right)^2 \right)$$

$$= 4D\tau + 4\sigma^2$$

The lateral (*xy*) localization precision is given by σ which we determine to be 40 nm. We fitted a

straight line to each separate MSD relation. Assuming a line fit has an optimized gradient *g* to the first

4 points (defined as the first we 3 measured MSD data points for *n*=1, 2 and 3, in addition to a 4th data

corresponding to *n*=0 obtained from constraining the intercept be $4\sigma^2$ to within measurement error of

the localization precision) then estimated the microscopic diffusion coefficient *D* as *g*/4.Δt. For

immobile foci, tracks were collated and compiled to generate a mean MSD vs. τ relation which was

fitted to an asymptotic rising exponential function as an analytical model for confined diffusion of

MSD plateau equal to $L^2/6$ where *L* is the effective confinement diameter (39), enabling us to estimate

the confinement diameter.

Colocalization analysis. The extent of colocalization between red and green detected foci was

determined using a method which calculated the overlap integral between each green and red foci pair,

whose centroids were within ~1 PSF width (~3 pixels). Assuming two normalized, two-dimensional

Gaussian intensity distributions g_1 and g_2, for green and red foci respectively, centred around (x_1, y_1)

with sigma width σ_1, and around (x_2, y_2) with width σ_2, the overlap integral ν is analytically determined

as:

$$\nu = e^{\left(-\Delta r^2/2\left(\sigma_1^2+\sigma_2^2\right)\right)}$$

Where:

$$\Delta r^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$$

We use a criterion of an overlap integral of 0.75 or above to indicate putative colocalization

(38) since this corresponds to a foci centroid separation equivalent to the localization precision in this

case. By quantifying the standard deviation on the number of detected foci in each channel we estimate

that the standard error of colocalisation proportion under our typical imaging conditions is

approximately 9%.

Random foci overlap models. We calculated the probability of foci overlap in a single color channel by

first estimating a sensible range of foci surface density *n*. For the lower limit we used the number of

foci tracks detected in a 20 image frame time window, for the upper limit we used the average

measured value of the background-corrected pixel intensity value divided by the intensity of a single

fluorophore (equivalent to ~1 mLukS* molecule per pixel). We implemented these probability

estimates into a surface density model which assumed a random Poisson distribution for nearest-

neighbor separation (37, 38, 40, 62-64). This model indicates that the probability that a nearest-

neighbor separation is greater than *w* is given by $\exp(-\pi w^2 n)$. The probability of overlap for each

density estimate (Figure 3 – figure supplement 3) was convolved with a real molecular stoichiometry

distribution and a Gaussian function *p(x)* of stoichiometry (*x*):

$$p(x) = 2\pi\sigma^2 \exp(-\frac{(x - n)^2}{2\sigma^2\sqrt{n}})$$

where σ is the width of single fluorophore intensity distribution (~0.7 molecules), and *n* is the real

molecular stoichiometry. The tetramer model assumes *n=4*, then all higher order stoichiometries are

due to overlapping PSFs. The tetramer oligomer molecule assumed an equal number of multimerized

tetramers up to 5, which gave the best fit to the data.

The same strategy was used to model the random overlap probability for green and red color

channel fluorescent foci in dual color imaging experiments to assess the extent of apparent

colocalization due to random overlap between hC5aR and mLukS*/F*. The probability that a nearest-

neighbor separation is greater than w for foci of two different types is the same as a single type

multiplied by 2/3. (38)

Statistical tests and replicates

All statistical tests used are two-tailed unless stated otherwise. For single-molecule TIRF imaging each

cell can be defined as a biological replicate sampled from the cell population. We chose sample sizes of

5-7 cells yielding thousands of foci, generating reasonable estimates for stoichiometry and diffusion

coefficient distributions. Technical replicates are not possible with the irreversible photobleaching

assay.

Ethics statement

Human polymorphonuclear (PMN) cells, obtained from healthy volunteers were isolated by

Ficoll/Histopaque centrifugation (57). Informed written consent was obtained from all subjects, in

accordance with the Declaration of Helsinki and the Medical Ethics Committee of the University

Medical Center Utrecht (METC-protocol 07-125/C approved 1 March 2010).

Acknowledgments

We thank Piet Aerts and Angelino Tromp for assistance in sample preparation and labeling and Esther van 't Veld and Richard Wubbolts (Utrecht) for assistance with light microscopy and Dr Karin Strijbis (Veterinary School, UU) providing the sortase A enzyme. This work was supported by The Finnish Cultural Foundation (grants 00131060 and 00142390), the Biological Physical Sciences Institute, Royal Society, MRC (grant MR/K01580X/1), BBSRC (grant BB/N006453/1), the EPSRC Physics of Life UK network and the Wellcome Trust [ref: 204829] through the Centre for Future Health (CFH) at the University of York, UK.

Conflict of interest

All the authors declare that they have no conflict of interests.

References

1.   DeLeo FR, Otto M, Kreiswirth BN, Chambers HF. Community-associated meticillin-resistant Staphylococcus aureus. Lancet. 2010;375(9725):1557-68.

2.   Barrett FF, McGehee RF, Jr., Finland M. Methicillin-resistant Staphylococcus aureus at Boston City Hospital. Bacteriologic and epidemiologic observations. N Engl J Med. 1968;279(9):441-8.

3.   Udo EE, Pearman JW, Grubb WB. Genetic analysis of community isolates of methicillin-resistant Staphylococcus aureus in Western Australia. J Hosp Infect. 1993;25(2):97-108.

4.   Vandenesch F, Naimi T, Enright MC, Lina G, Nimmo GR, Heffernan H, et al. Community-acquired methicillin-resistant Staphylococcus aureus carrying Panton-Valentine leukocidin genes: worldwide emergence. Emerg Infect Dis. 2003;9(8):978-84.

5.   Hiramatsu K, Aritaka N, Hanaki H, Kawasaki S, Hosoda Y, Hori S, et al. Dissemination in

Japanese hospitals of strains of Staphylococcus aureus heterogeneously resistant to vancomycin.

Lancet. 1997;350(9092):1670-3.

6.   Koch G, Yepes A, Forstner KU, Wermser C, Stengel ST, Modamio J, et al. Evolution of

resistance to a last-resort antibiotic in Staphylococcus aureus via bacterial competition. Cell.

2014;158(5):1060-71.

7.   Meyer F, Girardot R, Piemont Y, Prevost G, Colin DA. Analysis of the specificity of Panton-

Valentine leucocidin and gamma-hemolysin F component binding. Infect Immun. 2009;77(1):266-73.

8.   Gauduchon V, Werner S, Prevost G, Monteil H, Colin DA. Flow cytometric determination of

Panton-Valentine leucocidin S component binding. Infect Immun. 2001;69(4):2390-5.

9.   Otter JA, French GL. Molecular epidemiology of community-associated meticillin-resistant

Staphylococcus aureus in Europe. Lancet Infect Dis. 2010;10(4):227-39.

10.   Naimi TS, LeDell KH, Como-Sabetti K, Borchardt SM, Boxrud DJ, Etienne J, et al.

Comparison of community- and health care-associated methicillin-resistant Staphylococcus aureus

infection. JAMA. 2003;290(22):2976-84.

11.   Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, et al. Involvement of

Panton-Valentine leukocidin-producing Staphylococcus aureus in primary skin infections and

pneumonia. Clin Infect Dis. 1999;29(5):1128-32.

12.   Gillet Y, Issartel B, Vanhems P, Fournet JC, Lina G, Bes M, et al. Association between

Staphylococcus aureus strains carrying gene for Panton-Valentine leukocidin and highly lethal

necrotising pneumonia in young immunocompetent patients. Lancet. 2002;359(9308):753-9.

13.   Perret M, Badiou C, Lina G, Burbaud S, Benito Y, Bes M, et al. Cross-talk between

Staphylococcus aureus leukocidins-intoxicated macrophages and lung epithelial cells triggers

chemokine secretion in an inflammasome-dependent manner. Cellular microbiology. 2012;14(7):1019-

36.

14.   Jayasinghe L, Bayley H. The leukocidin pore: evidence for an octamer with four LukF subunits

and four LukS subunits alternating around a central axis. Protein Sci. 2005;14(10):2550-61.

15.   Colin DA, Mazurier I, Sire S, Finck-Barbancon V. Interaction of the two components of

leukocidin from Staphylococcus aureus with human polymorphonuclear leukocyte membranes:

sequential binding and subsequent activation. Infect Immun. 1994;62(8):3184-8.

16.   Das SK, Darshi M, Cheley S, Wallace MI, Bayley H. Membrane protein stoichiometry

determined from the step-wise photobleaching of dye-labelled subunits. Chembiochem. 2007;8(9):994-

9.

17.   Spaan AN, Henry T, van Rooijen WJ, Perret M, Badiou C, Aerts PC, et al. The staphylococcal

toxin Panton-Valentine Leukocidin targets human C5a receptors. Cell host & microbe. 2013;13(5):584-

94.

18.   Postma B, Kleibeuker W, Poppelier MJ, Boonstra M, Van Kessel KP, Van Strijp JA, et al.

Residues 10-18 within the C5a receptor N terminus compose a binding domain for chemotaxis

inhibitory protein of Staphylococcus aureus. J Biol Chem. 2005;280(3):2020-7.

19.   Spaan AN, Schiepers A, de Haas CJ, van Hooijdonk DD, Badiou C, Contamin H, et al.

Differential Interaction of the Staphylococcal Toxins Panton-Valentine Leukocidin and gamma-

Hemolysin CB with Human C5a Receptors. J Immunol. 2015;195(3):1034-43.

20.   Walport MJ. Complement. First of two parts. N Engl J Med. 2001;344(14):1058-66.

21.   Fearon DT, Austen KF. Activation of the alternative complement pathway with rabbit

erythrocytes by circumvention of the regulatory action of endogenous control proteins. The Journal of

experimental medicine. 1977;146(1):22-33.

22.   Berends ET, Mohan S, Miellet WR, Ruyken M, Rooijakkers SH. Contribution of the

complement Membrane Attack Complex to the bactericidal activity of human serum. Molecular

immunology. 2015;65(2):328-35.

23.   Easmon CS, Glynn AA. Comparison of subcutaneous and intraperitoneal staphylococcal

infections in normal and complement-deficient mice. Infection and immunity. 1976;13(2):399-406.

24.   Mullaly SC, Kubes P. The role of TLR2 in vivo following challenge with Staphylococcus

aureus and prototypic ligands. J Immunol. 2006;177(11):8154-63.

25.   Woodruff TM, Nandakumar KS, Tedesco F. Inhibiting the C5-C5a receptor axis. Mol Immunol.

2011;48(14):1631-42.

26.   Ricklin D, Lambris JD. Progress and trends in complement therapeutics. Advances in

experimental medicine and biology. 2013;735:1-22.

27.   DuMont AL, Torres VJ. Cell targeting by the Staphylococcus aureus pore-forming toxins: it's

not just about lipids. Trends Microbiol. 2014;22(1):21-7.

28.   Dal Peraro M, van der Goot FG. Pore-forming toxins: ancient, but never really out of fashion.

Nat Rev Microbiol. 2016;14(2):77-92.

29.   Yamashita K, Kawai Y, Tanaka Y, Hirano N, Kaneko J, Tomita N, et al. Crystal structure of the

octameric pore of staphylococcal gamma-hemolysin reveals the beta-barrel pore formation mechanism

by two components. Proc Natl Acad Sci U S A. 2011;108(42):17314-9.

30.   Yamashita D, Sugawara T, Takeshita M, Kaneko J, Kamio Y, Tanaka I, et al. Molecular basis

of transmembrane beta-barrel formation of staphylococcal pore-forming toxins. Nat Commun.

2014;5:4897.

31.   Chiu SW, Leake MC. Functioning nanomachines seen in real-time in living bacteria using

single-molecule and super-resolution fluorescence imaging. Int J Mol Sci. 2011;12(4):2518-42.

32.   Plank M, Wadhams GH, Leake MC. Millisecond timescale slimfield imaging and automated

quantification of single fluorescent protein molecules for use in probing complex biological processes.

Integr Biol (Camb). 2009;1(10):602-12.

33.   Miller H, Zhou Z, Wollman AJ, Leake MC. Superresolution imaging of single DNA molecules

using stochastic photoblinking of minor groove and intercalating dyes. Methods. 2015;88:81-8.

34.   Wollman AJ, Leake MC. Millisecond single-molecule localization microscopy combined with

convolution analysis and automated image segmentation to determine protein concentrations in

complexly structured, functional cells, one cell at a time. Faraday Discuss. 2015;184:401-24.

35.   Leake MC, Chandler JH, Wadhams GH, Bai F, Berry RM, Armitage JP. Stoichiometry and

turnover in single, functioning membrane protein complexes. Nature. 2006;443(7109):355-8.

36.   Leake MC. Analytical tools for single-molecule fluorescence imaging in cellulo. Phys Chem

Chem Phys. 2014;16(25):12635-47.

37.   Leake MC, Greene NP, Godun RM, Granjon T, Buchanan G, Chen S, et al. Variable

stoichiometry of the TatA component of the twin-arginine protein transport system observed by in vivo

single-molecule imaging. Proc Natl Acad Sci U S A. 2008;105(40):15376-81.

38.   Llorente-Garcia I, Lenn T, Erhardt H, Harriman OL, Liu LN, Robson A, et al. Single-molecule

in vivo imaging of bacterial respiratory complexes indicates delocalized oxidative phosphorylation.

Biochim Biophys Acta. 2014;1837(6):811-24.

39.   Robson A, Burrage K, Leake MC. Inferring diffusion in single live cells at the single-molecule

level. Philos Trans R Soc Lond B Biol Sci. 2013;368(1611):20120029.

40.   Badrinarayanan A, Reyes-Lamothe R, Uphoff S, Leake MC, Sherratt DJ. In vivo architecture

and action of bacterial structural maintenance of chromosome proteins. Science. 2012;338(6106):528-

31.

41.   von Kockritz-Blickwede M, Konrad S, Foster S, Gessner JE, Medina E. Protective role of

complement C5a in an experimental model of Staphylococcus aureus bacteremia. J Innate Immun.

2010;2(1):87-92.

42.   Prevost G, Cribier B, Couppie P, Petiau P, Supersac G, Finck-Barbancon V, et al. Panton-

Valentine leucocidin and gamma-hemolysin from Staphylococcus aureus ATCC 49775 are encoded by

distinct genetic loci and have different biological activities. Infect Immun. 1995;63(10):4121-9.

43.   Diep BA, Chan L, Tattevin P, Kajikawa O, Martin TR, Basuino L, et al. Polymorphonuclear

leukocytes mediate Staphylococcus aureus Panton-Valentine leukocidin-induced lung inflammation

and injury. Proc Natl Acad Sci U S A. 2010;107(12):5587-92.

44.   Walport MJ. Complement. Second of two parts. N Engl J Med. 2001;344(15):1140-4.

45.   Pandya PH, Wilkes DS. Complement system in lung disease. Am J Respir Cell Mol Biol.

2014;51(4):467-73.

46.   Veya L, Piguet J, Vogel H. Single Molecule Imaging Deciphers the Relation between Mobility

and Signaling of a Prototypical G Protein-coupled Receptor in Living Cells. J Biol Chem.

2015;290(46):27723-35.

47.   Pedelacq JD, Maveyraud L, Prevost G, Baba-Moussa L, Gonzalez A, Courcelle E, et al. The

structure of a Staphylococcus aureus leucocidin component (LukF-PV) reveals the fold of the water-

soluble species of a family of transmembrane pore-forming toxins. Structure. 1999;7(3):277-87.

48.   Guillet V, Roblin P, Werner S, Coraiola M, Menestrina G, Monteil H, et al. Crystal structure of

leucotoxin S component: new insight into the Staphylococcal beta-barrel pore-forming toxins. J Biol

Chem. 2004;279(39):41028-37.

49.   Olson R, Nariya H, Yokota K, Kamio Y, Gouaux E. Crystal structure of staphylococcal LukF

delineates conformational changes accompanying formation of a transmembrane channel. Nat Struct

Biol. 1999;6(2):134-40.

50.   Nocadello S, Minasov G, Shuvalova L, Dubrovska I, Sabini E, Bagnoli F, et al. Crystal

structures of the components of the Staphylococcus aureus leukotoxin ED. Acta Crystallogr D Struct

Biol. 2016;72(Pt 1):113-20.

51.   Laventie BJ, Guerin F, Mourey L, Tawk MY, Jover E, Maveyraud L, et al. Residues essential

for Panton-Valentine leukocidin S component binding to its cell receptor suggest both plasticity and

adaptability in its interaction surface. PLoS One. 2014;9(3):e92094.

52.   Rabiet MJ, Huet E, Boulay F. Complement component 5a receptor oligomerization and

homologous receptor down-regulation. J Biol Chem. 2008;283(45):31038-46.

53.   Miles G, Movileanu L, Bayley H. Subunit composition of a bicomponent toxin: staphylococcal

leukocidin forms an octameric transmembrane pore. Protein Sci. 2002;11(4):894-902.

54.   Sugawara N, Tomita T, Sato T, Kamio Y. Assembly of Staphylococcus aureus leukocidin into a

pore-forming ring-shaped oligomer on human polymorphonuclear leukocytes and rabbit erythrocytes.

Biosci Biotechnol Biochem. 1999;63(5):884-91.

55.   LaChapelle S, Tweten RK, Hotze EM. Intermedilysin-receptor interactions during assembly of

the pore complex: assembly intermediates increase host cell susceptibility to complement-mediated

lysis. J Biol Chem. 2009;284(19):12719-26.

56.   Guo RF, Ward PA. Role of C5a in inflammatory responses. Annu Rev Immunol. 2005;23:821-

52.

57.   Veldkamp KE, Heezius HC, Verhoef J, van Strijp JA, van Kessel KP. Modulation of neutrophil

chemokine receptors by Staphylococcus aureus supernate. Infect Immun. 2000;68(10):5908-13.

58.   Zacharias DA, Violin JD, Newton AC, Tsien RY. Partitioning of lipid-modified monomeric

GFPs into membrane microdomains of live cells. Science. 2002;296(5569):913-6.

59. Hirota N, Yasuda D, Hashidate T, Yamamoto T, Yamaguchi S, Nagamune T, et al. Amino acid

residues critical for endoplasmic reticulum export and trafficking of platelet-activating factor receptor.

The Journal of biological chemistry. 2010;285(8):5931-40.

60. Amdahl H, Haapasalo K, Tan L, Meri T, Kuusela PI, van Strijp JA, et al. Staphylococcal

protein Ecb impairs complement receptor-1 mediated recognition of opsonized bacteria. PLoS One.

2017;12(3):e0172675.

61. Wouters FS, Verveer PJ, Bastiaens PI. Imaging biochemistry inside cells. Trends Cell Biol.

2001;11(5):203-11.

62. Delalez NJ, Wadhams GH, Rosser G, Xue Q, Brown MT, Dobbie IM, et al. Signal-dependent

turnover of the bacterial flagellar switch protein FliM. Proc Natl Acad Sci U S A. 2010;107(25):11347-

51.

63. Reyes-Lamothe R, Sherratt DJ, Leake MC. Stoichiometry and architecture of active DNA

replication machinery in Escherichia coli. Science. 2010;328(5977):498-501.

64. Lenn T, Leake MC, Mullineaux CW. Clustering and dynamics of cytochrome bd-I complexes

in the Escherichia coli plasma membrane in vivo. Mol Microbiol. 2008;70(6):1397-407.

Legends

Figure 1. Toxin functionality on PMN and HEK cells. (A) PMN cell permeability in the presence of

unlabeled LukSK281CY113H and LukFK288C (mS+mF, number of biological replicates n=2), Alexa-

labeled mS* or mF* and wild type LukF (F(wt)) or LukS, (S(wt)) (n=1), compared to PMN cell

permeability of S(wt) and F(wt), (n=3); (B) Inhibition of 3 µg/ml of FITC-labeled S(wt) (n=3) and mS*

(n=1) binding to PMN cell by mS; Permeability dose dependencies for (A) and (B) are shown with a

polynomial spline fit, statistical significance indicated between low (0.3 and 0.001 nM) and high (300

nM) toxin concentrations using Student's *t*-test. Error bars indicate SD. (C) Column indicating binding

responses for mF* on hC5aR cells (n=2). *** Indicates a statistically significant difference (p<0.001)

between mS* binding on HEK-hC5aR cells compared to HEK-CCR2 and mF* binding on these cells.

(D) Permeability of hC5aR transfected HEK cells using unlabelled mS and mF, and Alexa-labeled mS*

and mF* compared to wild type S(wt) and F(wt) (n=2). (E) Inhibition of 3 µg/ml of mS* binding by

mS on HEK-hC5aR cells, (n=3). CCR2-transfected HEK cells used as negative controls for toxin

binding and lysis in (C, D) n=2 or (C) one representative experiment. Dose dependency shown with

polynomial spline fit. Statistical significance calculated between low (0.3 and 0.001 nM) and high

(300 nM) toxin concentrations using Student's *t*-test. (F) Permeability response of hC5aR-transfected

HEK cells following incubation with unlabeled mS and Alexa maleimide-labeled mF* or wild type

toxins F(wt) and S(wt) (n=3). Statistical significance calculated between 15 min and 0 min time points

using Student's *t*-test. Error bars indicate SD. Percentages of mean fluorescence intensities is shown as

relative to the maximum intensity in each individual experiment (B, C and E).

Figure 2. Colocalization of LukS with hC5aR on HEK cells. (A) (left panel) TIRF image of hC5aR-

mGFP on the surface of a HEK cell before addition of toxin; (right panels) zoom-in of yellow dashed

square of left panel immediately following 2 min incubation with Alexa647-labeled

LukSK281CY113H (mS*-Alexa647). (B) Equivalent images of same cell of (B) after > 15 min

incubation with LukFK288C (mF). (C) (upper panel) TIRF image of colocalization of Alexa594- and

Alexa647-labeled mF and mS (mF*Alexa594 and mS*Alexa647) with hC5aR-mGFP on HEK cells;

(lower panel) zoom-in of yellow-dashed square of upper panel with colocalized foci indicated (white

arrow).

Figure 2- figure supplement 1. Construction of recombinant leukocidin proteins. (A) Schematic of

TIRF imaging assay. (B) (left panel) Crystal structure of octameric pore complex of γ-hemolysin (PDB

ID:3B07). K273/Y111 and K289 on S and F components of γ-hemolysin corresponds to our engineered

mutations, K281C/Y113H and K288C, on LukSF marked in their equivalent places on γ-hemolysin;

(right panel) SDS-PAGE of the unlabeled LukSK281CY113H and LukFK288C (mLukS and mLukF)

and Alexa-labeled mLukS* and mLukF* toxin components, bands visible at locations consistent with

molecular weight of 33 kDa and 34 kDa for LukS and LukF respectively.

Figure 3. Spatiotemporal dynamics of hC5aR, LukS and LukF in live cells. (A) Images of HEK

cells treated with LukSK281CY113H (mS) and Alexa-labeled LukFK288C (mF*) showing brightfield

(left), hC5aR-GFP (middle) and mF* (right). (B) Probability distribution for stoichiometry of hC5aR in

absence and presence of Alexa-labeled mS (mS*) and mF*, and (C) of mS* foci, indicating (D)

tetramer periodicity from Fourier spectral analysis. (E) A random tetramer overlap model cannot

account for mS* experimental stoichiometry data ($R^2<0$), but a tetramer-multimer model results in

excellent agreement ($R^2=0.85$). (F) hC5aR and mS* stoichiometry as a function of incubation time.

Proportion of immobile and mobile colocalized foci in the (G) presence and absence of mS and mF.

Error bars show standard error of the mean from n=5-15 image subregions.

Figure 3 – figure supplement 1. Fluorescent protein characterization. A. Fluorescence micrograph

of immobilized GFP and an intensity *vs.* time trace for one foci showing a single photobleach step.

Raw data in light blue and edge-preserving Chung-Kennedy filtered data in dark blue. B. As A for

LukS-Alexa647, C. LukS-Alexa647 micrograph (red) with found foci indicated as white circles. D.

Intensity distribution of Alexa 647 foci intensities from whole photobleach experiment showing

periodicity at ~3,500 counts on our camera detector. E. Pairwise distance distribution of intensity in D

with Fourier spectrum (inset) showing peak at ~4,000 counts. F. GFP foci intensity (natural log) time

traces (green) with linear fits (red).

Figure 3 – figure supplement 2. Density of LukS spots. A. micrograph of mLukS* (white) with

found foci (orange circles) B. Probability distribution of overlap frequency using spots in A to calculate

density. C. Zoom in of mLukS micrograph. D. Probability distribution of overlap frequency using

intensity in C to calculate maximum density estimate.

Figure 3 – figure supplement 3. Mobility analysis. A**.** The probability distribution of microscopic

diffusion coefficient showing the threshold for immobility as black dotted line and B. the mean squared

displacement against time interval for mobile (upper) and immobile (lower) of hC5aR. C. and D.

similar for mLukS/F*.

Figure 4. Relative stoichiometry of hC5aR, LukS and LukF in fixed cells. (A) Micrographs of fixed

hC5aR-GFP HEK cells treated with LukSK281CY113H (mS) and LukFK288C (mF) showing hC5aR-

GFP (left) and Alexa647 (middle) and merge (right) on Alexa-labeled mS (mS*) above and mF (mF*)

below (B) Proportion of foci colocalized and not colocalized, treated with mS, mS*+mF and mS+mF*

for hC5aR (green) and mLuk* (orange). Error bars show standard error of the mean from n=4 image

subregions. (C) Heatmap of correlation between hC5aR and mS stoichiometry (red dash line indicates

4 mS per hC5aR molecule), R^2 ~0.15 (D) and (E) FRET images and efficiencies. The FRET

experiment was performed in live and fixed sortase-tagged FITC-hC5aR expressing cells. Live cells

(number of biological replicates n=2) were incubated in the presence of Cy3-labeled mS* for 1 h at

+4°C and washed, after which unlabeled mF was added. FRET was analyzed before (mS*) or after

addition of mF (mS*+mF). FRET from fixed cells (n=3) was analyzed in the presence of mS* or

unlabeled mS and Cy3-labeled mF* respectively (n=2). Statistical significance between cells with only

mS and both of the toxin components, mS and mF, was analyzed using Student's *t*-test. Error bars

indicate SD.

Figure 4 – figure supplement 1. Colocalization analysis. The probability distribution of linked (A)

and unlinked (B) hC5aR and similar for mLukS* (C. and D.). E. and F. False-color heatmap scatter

plots indicating that h5CaR stoichiometry is uncorrelated to mLukS or mLukF stoichiometry in the

presence of mLukF.

Figure 5. LukSF dissociation and rebinding of C5a on hC5aR expressing cells. (A) Inhibition of

anti-CD88 binding on hC5aR expressing HEK cells using increasing concentrations (horizontal axis) of

wild type LukS, S(wt) and C5a. Wild type LukF, F(wt), is used as a negative control for inhibition of

anti-CD88 binding (number of biological repeats n=2). (B) Disengagement of hC5aR from LukSF was

observed as increase in PE conjugated anti-CD88 binding (right vertical axis) on S(wt) pre-coated cells

using increasing but sublytic concentrations (horizontal axis) of F(wt). Minimal cell lysis (% of lysed

cells, left vertical axis) detected in F(wt) concentrations below 3 nM (n=3). (C) Rebinding of constant

amount of NT647 labeled C5a (647-C5a) on hC5aR upon LukSF formation analyzed by incubating

S(wt) pre-coated cells with increasing concentrations (horizontal axis) of LukF mutant G130D that

associates with LukS but does not lead to cell lysis (n=2). (D) Effect of LukSF mediated cell lysis on

complement activation and C5a formation on full blood measured by using C5b-9 as a marker for

complement activation in plasma (n=3). Maximal C5a formation is observed by incubating full blood

with live *S. aureus* bacteria. Ecb (B and C), F(wt) (A) or S(wt) (D) are used as negative controls in the

assays. Percentages of mean fluorescence intensities is shown as relative to the maximal intensity in

each individual experiment (A-C). Statistical significances are calculated using Student's *t*-test. Error

bars indicate SD.

Figure 6. Model for LukSF-receptor binding and the mechanism of LukSF-induced

inflammation. (A). LukS (PDB ID: 1T5R) binds on hC5aR (structure based on angiotensin receptor

data PDB ID: 4YAY) as a soluble monomer on the cell membrane. Each LukS monomer binds one

hC5aR molecule via the receptor interacting residues R73, Y184, Y250, T244 (marked with blue dots)

within a cluster of approximately 4-5 hC5aR homo-oligomers Upon binding to hC5aR LukS exposes

residues for LukF (PDB ID: 1LKF) binding. In these tight clusters each LukF can bind to two LukS

monomers via two interfaces. (B). Binding of LukF on LukS and formation of the octameric pore (PDB

ID: 3B07) causes dissociation of the receptors from the complex because of leakage of the cell membrane and possibly also since the receptor binding region (marked with a circle) is buried between the monomers in the complex. (C) The detached hC5aR molecule can be reused by its ligands LukS or C5a anaphylatoxin (PDB ID: 1KJS). (D) Schematic illustrating the putative mechanism of LukSF induced inflammation.

1016

1017

1018 **Movie 1.** Deposition of mLukS on hC5aR-mGFP HEK cells. The movie is shown in two clips before

1019 (no toxin) and after addition of Alexa 595 labeled mLukS (add mLukS*). This movie was recorded for

1020 *ca.* 13 min and displayed here at 100x speed.

1021

1022 **Movie 2.** Lysis of hC5aR-mGFP HEK cells incubated with Alexa594 labeled mLukS* and mLukF.

1023 The cells were preincubated with mLukS* and the lysis of the cells were monitored for *ca.* 13 min after

1024 addition of mLukF (add mLukF). The red arrow points to the vesicles released during cell lysis. This

1025 movie was recorded for *ca.* 13 min and displayed here at 100x speed.

1026

1027 **Movie 3.** Imaging live hC5aR-mGFP cells. After 1-2 min of exposure, several distinct, mobile,

1028 circular fluorescent foci in the planer membrane regions were observed. Movie is displayed in real

1029 time.

1030

1031 **Movie 4.** Imaging mLukS* incubated with hC5aR-mGFP cells. Several distinct, mobile, circular

1032 fluorescent foci were observed. Movie is displayed in real time.

1033

1034

1035

1036

1037

1038

1039

1040

1041

Figures

Figure 1. Toxin functionality on PMN and HEK cells.

Figure 2. Colocalization of LukS with hC5aR on HEK cells.

Figure 2- figure supplement 1. Construction of recombinant leukocidin proteins.

Figure 3. Spatiotemporal dynamics of hC5aR, LukS and LukF in live cells.

Figure 3 – figure supplement 1. Fluorescent protein characterization.

Figure 3 – figure supplement 2. Density of LukS spots.

Figure 3 – figure supplement 3. Mobility analysis.

Figure 4. Relative stoichiometry of hC5aR, LukS and LukF in fixed cells.

Figure 4 – figure supplement 1. Colocalization analysis.

1042

Figure 5. LukSF dissociation and rebinding of C5a on hC5aR expressing cells.

Figure 6. Model for LukSF-receptor binding and the mechanism of LukSF-induced inflammation.