Running title: Comprehensive Distribution Modeling

Modeling Avian Full Annual Cycle Distribution and Population Trends

with Citizen Science Data

Daniel Fink*, Tom Auer, Viviana Ruiz-Gutierrez, Wesley M. Hochachka,

Alison Johnston, Frank A. La Sorte, and Steve Kelling

Cornell Lab of Ornithology, Cornell University, Ithaca, New York, USA.

*Corresponding author: Email: daniel.fink@cornell.edu

Authors' contributions. *DF, WMH, and STK conceived and designed this*

study. DF designed the statistical methodology. TA and DF designed the

computational methodology, processed data, and distribution models. TA,

VRG, WMH, AJ, and FAL designed the analysis of the model products. DF

wrote the first draft of the manuscript, and all authors contributed

substantially to revisions. All the authors have approved the final version of

this manuscript and agree to be accountable for all aspects of the work.

Data Accessibility: *Should the manuscript be accepted, the data supporting*

the results will be archived in an appropriate public repository such as

Dryad or Figshare and the data DOI will be included at the end of the

article.

Abstract

Information on species' distributions and abundances, environmental associations, and

how these change over time are central to the study and conservation of wildlife

populations. This information is challenging to obtain at relevant scales across range-

wide extents for two main reasons. First, local and regional processes that affect

populations vary throughout the year and across species' ranges, requiring fine-scale,

year-round information across broad — sometimes hemispheric — spatial extents.

Second, while citizen science projects can collect data at these scales, using these data

requires additional steps to address known sources of bias. Here we present an analytical

framework to address these challenges and generate year-round, range-wide distributional

information using citizen science data. To illustrate this approach, we apply the

framework to Wood Thrush (*Hylocichla mustelina*), a long distance Neotropical migrant

and species of conservation concern, using data from the citizen science project eBird.

We estimate relative occupancy and abundance with enough spatiotemporal resolution to

support inference across a range of spatial scales throughout the annual cycle. This

includes intra-annual estimates of the range (quantified as the area of occupancy), intra-

annual estimates of the associations between species and features of their local

environment, and inter-annual season-specific trends in relative abundance. This is the

first example of an analysis to capture intra- and inter-annual distributional dynamics

across the entire range of a broadly distributed, highly mobile species.

Keywords: *conservation, biogeography, macroecology, eBird, bird*

migration, biodiversity monitoring, Wood Thrush

MAIN TEXT:

(a) Introduction

Information about species' distribution and abundance are essential to the fields of

applied ecology and conservation biology, for they are critical in the study of the

processes that limit and regulate populations. Because the biotic and abiotic processes

affecting species' populations vary seasonally and regionally, it is also important to

generate this information both at the spatiotemporal *scales* at which these processes

operate and across the full spatiotemporal *extents* over which these processes vary

(Heffernan *et al.* 2014). Finally, for broadly distributed species, information that supports

inference *across spatial and temporal scales* is necessary to understand how local,

regional, and seasonal-scale processes interact to affect entire populations at continental

extents.

For most well studies species groups, we still lack basic information on species

distributions, especially at relevant spatiotemporal resolutions and extents. A limiting

factor has been the ability to collect sufficient quantities of observational data both at fine

scales and across broad spatiotemporal extents. Current information on species

distributions often suffer from strong regional biases (Boakes *et al.* 2010) in coverage.

Although range maps are available for a growing number of taxa, these often provide

coarse spatial (Hurlbert & Jetz 2007) and temporal resolutions (Ridgeley *et al.* 2012).

Further, most of this information depicts only the most basic information on species'

range, often as expert drawn polygons, lacking more informative and ecologically

valuable measures of occupancy rates, relative abundance, and habitat associations within

a species' range.

Most information on species distributions is also static and does not capture intra- or

inter-annual dynamics. The majority of studies on amphibians, reptiles, birds, and

mammals have been conducted only during the breeding season (Marra *et al.* 2015). Even

for birds, one of the best-surveyed classes of organisms, continental-scale trends in

abundance are routinely estimated only for the breeding season and only in North

America (Sauer & Link 2011) and Europe (European Bird Census Council 2016). The

inability to estimate changes in distribution and abundance is a serious deficiency in our

ecological knowledgebase, and underscores the need for scalable approaches to collect

biodiversity data and model intra- and inter-annual dynamics.

Citizen science projects that use crowdsourcing techniques to engage the public have

been very successful at collecting observational data across large areas throughout the

year (Dickinson *et al.* 2010). However, using these data to generate robust distributional

information is fraught with analytical challenges (Hochachka *et al.* 2012; Bird *et al.*

2014). These challenges have led to the development and application of a number of new

analytical tools. For example, bias related to heterogeneity in species' detection is a

significant challenge when analyzing citizen science data. Several approaches have been

used to deal with heterogeneous and imperfect observation processes, from the inclusion

of relevant fixed and random effects (Sauer & Link, 2011) to the development of

detection process models (Kéry & Royle, 2015). When citizen scientist participants

choose where and when to conduct their surveys, site selection biases lead to repeated

surveys in popular search locations and missing surveys in areas that are hard to access.

Data-sampling methods have proven useful to mitigate the effects of these site-selection

biases (Robinson *et al.* 2017). The ability to accurately capture the complex species-

environment relationships underlying regional patterns of species' distributions and

abundance is another challenge. Machine and statistical learning models have proven to

be efficient tools for learning these species-environment relationships from large sets of

environmental covariates (Elith & Leathwich 2009). Using citizen science data to

estimate abundance often presents the statistical challenge of "zero inflation" where too

many zero counts can degrade model performance. A wide variety of new abundance

models have been proposed to deal with zero-inflation (Denes *et. al.* 2015).

Most of the methodological developments discussed above have been used to study

regional-scale patterns of species' distribution and abundance during a single season of

the year. To generate distributional information across larger spatiotemporal extents with

citizen science data, requires tacking two additional challenges. First, most broad-scale

citizen science projects exhibit strong variation in data density across large regions which

can degrade model performance. Adaptive knot designs (Gelfand *et al.* 2012) and

partitioning methods (Fink *et al.* 2013) have been proposed to deal with this challenge by

adding multi-scale structure to broad extent analyses. Second, the spatial and temporal

variation in a species' response to the same environmental conditions, statistical non-

stationarity, can also degrade model performance. Non-stationary regression techniques

function to add multi-scale structure to analyses and have proven to be a useful solution

to this challenge (Fink *et al.* 2010; Finley 2011).

While previous studies have dealt with one or more of the analytical challenges outlined

above (e.g. Johnston *et al.* 2015), to date, none have dealt with all of these challenges

simultaneously at the relevant scales necessary to study broadly distributed species across

the annual cycle. Here, we describe a simple analytical framework capable of estimating

species' relative occupancy and abundance, year-round and range-wide with enough

spatial and temporal resolution to support inference across a range scales. This includes

intra-annual estimates of species ranges' (quantified as the area of occupancy), intra-

annual estimates of the associations between species and features of their local

environment, and inter-annual trends in relative abundance. For convenience, we will

refer to this suite of parameter estimates as Cross-Scale Full-Annual Cycle (CS-FAC)

distributional information.

As a case study, we analyzed data from the global citizen science project eBird (Sullivan

et al. 2014) for the long-distance migratory songbird, Wood Thrush (*Hylocichla*

mustelina) that breeds in eastern North America and winters largely in Mesoamerica. The

Wood Thrush is a species of conservation concern, having suffered steep population

declines over the past several decades. Despite numerous regional studies (e.g. Rushing

et al. 2017), comprehensive information on abundance and distribution is still lacking.

We used the CS-FAC analysis to fill important information gaps for this species,

illustrating a method that is widely applicable to other species and taxonomical groups.

(b) Materials and methods

Observational Data

The bird observation data were obtained from the global bird monitoring project, eBird

(Sullivan *et al.* 2014) using the eBird Reference Dataset (ERD2016, Fink *et al.* 2017).

We used a subset of data in which the time, date, and location of the survey period were

reported and observers recorded the number of individuals of all bird species detected

and identified during the survey period, resulting in a complete 'checklist' of species on

the survey (Sullivan *et al.* 2009). The checklists used here were restricted to those using

the 'stationary', 'traveling', or 'areal search' protocols from January 1, 2004 to December

31, 2016 within the spatial extent between 180° to 30° W Longitude. Areal surveys were

restricted to those covering less than 56 sq. km. and traveling surveys were restricted to

those ≤ 15km. This resulted in a dataset consisting of 14 million checklists, of which 10%

were withheld for model validation.

Predictor Data

We incorporated three classes of predictors in the models: (1) Four observation-effort

predictors to account for variation in detection rates, (2) Three temporal predictors to

account trends at different scales, and (3) 79 site-specific predictors from remote sensing

data to capture associations of birds with elevation and a variety of habitats across the

hemisphere. The effort predictors included the duration spent searching for birds, whether

the observer was stationary or not, the distance traveled during the search, and the

number of people in the search party. The observation time of the day was used to model

variation in availability for detection, e.g. variation in behavior such as participation in

the dawn chorus (Diefenbach *et al.* 2007). The day of the year (1-366) on which the

search was conducted was used to capture intra-annual variation and the year of the

observation was included to account for inter-annual variation.

Elevation defines basic abiotic conditions that influence species distributions. To account

for the effects of elevation and topography, each checklist location was associated with

elevation, eastness, and northness at $1km^2$ resolution (Amatulli *et al.* 2017). To account

for species' local habitat-selectivity each checklist was linked to a series of covariates

derived from the NASA MODIS land cover data (Friedl *et al.* 2010). We selected this

data product for its moderately high spatial resolution, annual temporal resolution, and

global coverage. We used the University of Maryland (UMD) land cover classifications

(Hansen *et al.* 2000) and derived water cover classes from the MODIS Land Cover Type

QA Science Dataset resulting in a class label for each 500m pixel into one of 19 classes

(Table 1).

Checklists were linked to the MODIS data by-year from 2004-2013 (checklist data after

2013 are matched to the 2013 data). All cover classes were summarized within a 2.8km ×

2.8km (784 hectare) neighborhood centered on the checklist location. In each

neighborhood, we computed the proportion of each class in the neighborhood (PLAND).

To describe the spatial configuration of each class within each neighborhood we

computed three statistics using FRAGSTATS (McGarigal *et al.* 2012; VanDerWal *et al.*

2014): LPI an index of the largest contiguous patch, PD an index of the patch density,

and ED an index of the edge density.

Analysis Overview

To meet the analytical challenges of CS-FAC modeling with eBird data, we adopted an

ensemble modeling strategy combining pattern discovery and inference based on the

Spatio-Temporal Exploratory Model (STEM; Fink *et al.* 2010). STEM is an ensemble of

regional-seasonal regression models generated by repeatedly subsampling and

partitioning the study extent into 100 randomly located grids of overlapping

spatiotemporal blocks, called *stixels*, and then fitting an independent regression model,

called a *base model*, within each stixel. Together, the base models are used to form an

ensemble of local parameter estimates distributed uniformly across the study extent.

Using the fact that stixels overlap in space and time, parameters at a given location and

time are estimated by averaging across overlapping base models. Combining estimates

across the ensemble controls for inter-model variability (Efron 2014), providing a simple

way to control for overfitting while naturally adapting to non-stationary species-

environment relationships (Fink *et al.* 2010). Utilizing the fact that data are subsampled

for each base model, resampling techniques are employed to generate uncertainty

estimates for the ensemble parameter estimates. All analysis was conducted in R, version

3.4.2 (R Development Core Team 2017) and deployed using Apache Spark 2.1 (Zaharia,

et. al. 2016).

In the following sections, we describe the STEM base models, ensemble design, and the

spatial case-control sampling procedure. Then we describe how we used the ensemble to

estimate four population parameters: (1) landscape-scale relative occupancy and

abundance, (2) landscape-scale range boundaries quantified as the area of occupancy, (3)

regional-scale habitat use and avoidance, and (4) regional-scale trends in occupancy and

abundance.

STEM Base Models

Within each stixel, predictor-response (i.e. species-environment) relationships were

assumed to be stationary. To estimate relative occupancy and abundance from the large

predictor set while accounting for high numbers of zero counts, we used a two-step Zero-

Inflated Boosted Regression Tree (ZI-BRT) model (Johnston *et. al.* 2015; Ridgeway *et al.*

2017). Effort and time covariates were included in both steps of the ZI-BRT to account

for variation in detectability and variation in availability for detection. To generate

estimates of the binary un/occupied state, the occupancy threshold value that maximized

the Kappa statistic (Cohen 1960) was recorded for each ZI-BRT base model.

While the ZI-BRT base model can produce good estimates of the spatial patterns of

occupancy and abundance from large sets of predictors, it is not well suited to making

inference about inter-annual trends. Instead, we used the Zero-Inflated Generalized

Additive Model (ZI-GAM) fit with the mcv package (Wood *et al.* 2016) to estimate

trends, employing the additive structure to focus inference. The model equations defining

the occupancy and abundance sub-models had the forms

$logit(\psi) = \beta_0 + s(Year) + s(\psi spat) + s(lat, lon) + s(duration) + s(distance)$

$+ s(number) + stationary$

$log(N) = \alpha_0 + s(Year) + s(Nspat) + s(lat, lon),$

where ψ is the occupancy and N is the abundance. Inter-annual variation in occupancy

and abundance were both modeled using smooth functions, *s,* of the *Year*. To account for

variation in the observation process, smooth functions of search *duration*, search *distance*,

and the *number* of observers were included. The *stationary* indicator, was used to identify

stationary counts, accounting for differences in search protocols. Spatial variation for

both occupancy and abundance was modeled at two scales. Coarse-scale spatial patterns

were captured by smooth two dimensional smooth functions of latitude and longitude

with a limited number of knots. Fine-scale variation was modeled as smooth functions of

ZI-BRT derived covariates describing landscape-scale variation in occupancy, *ψspat*, and

abundance, *Nspat*.

The ZI-BRT derived covariates were constructed to encapsulate landscape-scale spatial

variation as a pair of covariates that could be used within the ZI-GAM. To do this we

leveraged the strength of the ZI-BRT model as a high-dimensional regression model. The

idea was to use the ZI-BRT model to adaptively select and estimate intra-seasonal

landscape-scale effects from the large set of predictors describing land and water cover

class and elevation. This was done by fitting the ZI-BRT model as described above, then

predicting the expected relative occupancy and abundance for each checklist in the

training data to generate the two covariates. Because we want to describe inter-annual

variation in the ZI-GAM, an important part of generating these predictors was removing

any inter-annual variation from the covariates. This was done simply by holding the

checklist year predictor constant when predicting the covariate values. Similarly, to

remove variation in detection associated with the effort predictors, all effort predictors

were also held constant.

By including the derived covariates in the ZI-GAM as smooth additive functions, the Zi-

GAM was able to adaptively calibrate the covariate information. Note, that by allowing

the day of the year predictor to vary when computing the derived covariates, the

covariates captured strong intra-seasonal changes in the spatial patterns of occupancy and

abundance. This is especially useful for modeling data from non-stationary periods.

Finally, we note that both ZI-BRT and ZI-GAM were fit independently within each stixel

in the ensemble. Thus, averaging across the ensemble naturally controls for the joint

variation of the ZI-BRT, the ZI-BRT derived covariates, and the ZI-GAM.

Ensemble Design

Stixel size controls a bias-variance tradeoff (Fink *et al.* 2013) and must strike a balance

between stixels that are large enough to achieve sufficient sample sizes, controlling

variance, and small enough to assume stationarity, controlling bias. For estimating

occupancy and abundance with ZI-BRT base model, $10°$ longitude $\times$ $10°$ latitude $\times$ 30-

contiguous day stixels were small enough to meet these requirements across much of the

northern study extent. To account for the relatively low data density south of $12°$ north

latitude we doubled the length of stixels to $20°$ longitude $\times$ $20°$ latitude $\times$ 30-days. For

estimating trends with ZI-GAM base model larger sample sizes were required to insure

representative sampling across years. To estimate breeding season trends, the same $10°$

longitude $\times$ $10°$ latitude $\times$ 30-contiguous day stixels were used. To estimate trends during

the non-breeding season, when data density is lower, we increased stixel size to cover the

spatiotemporal region from southern Mexico to Panama from December 1 to February 28.

See Appendix S1 in Supporting Information for additional information about the

specification of the stixel ensemble design.

Spatial Case-Control Sampling

Within each stixel, a spatial case-control sampling strategy was used to address the

challenges of highly imbalanced data and site selection bias. Imbalanced data arise when

there are a very small number of species detections and a very large number of non-

detections. This is a modeling concern because binary regression methods, like the first

component of ZI-BRT model, become overwhelmed by the non-detections and perform

poorly (Robinson *et al.* 2017). The low detection rates of many species, especially along

seasonal range boundaries, can generate highly imbalanced training data and make data

imbalance a defining challenge for broad-scale, year-round modeling. By sampling

non/detection cases separately, case-control sampling (e.g. Fithian & Hastie 2014)

improves data balance and model performance. Additionally, to alleviate spatial biases

caused by the eBird site selection process, spatially balanced samples were drawn as part

of the case-control sampling.

To generate spatially balanced samples for the case-control sampling, training data were

drawn from a randomly located regular grid, with one checklist randomly selected per

grid cell. North of 12° latitude, the grid cells were 10km × 10km and south of the cutoff

they were 20km × 20km. As part of the case-control sampling, detection data were over-

sampled, using the same spatially balanced procedure, when they represented less than

25% of the spatially balanced data. Because boosting, used in the ZI-BRT base models, is

driven more by the set of distinct data points than the number of tied data points

generated when oversampling, the spatial predictors of tied checklists were jittered to

break the ties (Mease et al. 2007). Finally, because case control sampling changes the

training data prevalence we back-transformed occupancy estimates to match the original

stixel prevalence rate.

For the ZI-GAM models, we also stratified the spatial case control samples by year and

restricted the total number of samples per year to control for inter-annual increases in

sample sizes resulting from increases in eBird participation rates. eBird participation rates

have been increasing at 20-30% per year since 2005. We set 2007 as our *reference year*,

fixing the sample size of spatially balanced training data for each year to equal that of

2007. We selected 2007 as the reference year as a balance between the amount of per-

year information available to estimate trends and the length of the trend.

Local Relative Occupancy and Abundance

We estimated relative occupancy and abundance once per week for all 52 weeks of the

calendar year. Estimates were made at 614,575 locations across the terrestrial Western

Hemisphere from a regular spatial grid with a density of one location per 8.4km × 8.4km

grid cell. Estimates at each location and date were made based on predictor values at that

location from all base models that contained the location and date. Then we averaged

across the model estimates using the upper 5% trimmed mean, a robust estimator

designed to guard against bias due to large outlying values. Uncertainty of the estimates

was estimated using the subsampling approach of Politis *et al.* (2009) following the

computational strategy of Geyer (2013). See the Appendix S2 for more information about

the subsampling procedures.

For all estimates we controlled for variation in detection rates associated with search

effort by holding predictors for search duration, protocol, search length, and number of

observers constant. Thus, the quantity we used to estimate relative occupancy was

defined as the probability that an average eBird participant would detect the species on a

search from 7–8AM while traveling 1 km on the given day at the given location. Relative

abundance was estimated as the expected count of individuals of the species on the same

standardized checklist. Although this approach accounts for variation in detection rates, it

can not directly estimate the absolute detection probability. For this reason, we refer to

the quantities estimated by the model as *relative* measures of occupancy and abundance.

Local Area of Occupancy

To estimate the Area of Occupancy (AOO) we predicted the binary un/occupied state for

all weeks using the same 8.4km × 8.4km spatial grid of locations used for relative

occupancy and abundance estimates. At each location and week, the AOO was estimated

as the proportion of base model occupancy estimates that were larger than the thresholds

for the corresponding base models. We call this the Proportion Above local Threshold

(PAT) estimator. We say the site was occupied, and, therefore within the species' range,

if the PAT estimator exceeded a specified value. One benefit of this ensemble estimator

is that it naturally adapts to regional and seasonal variation in species prevalence and

detectability. By averaging across the ensemble, PAT controls for inter-model variation

in both occupancy and base model threshold estimates.

Regional Habitat Associations

For each base model, we quantified the strength and direction of association for each

cover class predictor. Predictor importance (PI) statistics measured the strength of the

over-all contribution of individual predictors as the change in predictive performance

between the model that includes all predictors and the same model with permuted values

of the given predictor (Breiman 2001). PI statistics capture both positive and negative

effects arising from both additive and interacting model components. Partial Dependence

(PD) statistics measured the functional form of the additive association for each

individual cover class predictor by averaging out the effects of all other predictors (Hastie

et al. 2009). To measure the direction of association, we estimated the slope of each PD

estimate using simple linear regression. Because the PI and PD statistics account for the

effects of all other predictors in the base models, they account for variation in

detectability associated with effort and the time of day.

To examine how species' habitat use and avoidance varied among regions and seasons,

we computed regional trajectories of the strength and direction of the cover class

associations. Given the region and the set of predictors to compare, the PI statistics were

standardized to sum to 1 across the predictor set for each base model within the specified

region. Then, a loess smoother (Cleveland *et al.* 1992) was used to estimate the

trajectories of relative predictor importance throughout the year for each predictor.

Similarly, a loess smoother was used to estimate the proportion of increasing PD

estimates throughout the year for each predictor. Predictors with proportions greater than

70% were considered to be increasing and predictors with proportions less than 30%

were considered to be decreasing. Predictors with inconsistent directions, those between

30 and 70%, were excluded from summaries.

Regional Trends

We estimated the inter-annual trends using ensembles of partial effect estimates for year

from the ZI-GAM base models. These partial effects are regional-scale estimates that

describe the systematic change in relative abundance averaged across the stixel, after

accounting for landscape-scale spatial variation associated with elevation and the cover

classes. The partial effects of abundance are quantified on the log-link scale where they

can be interpreted in units of percent-per-year change.

Three types of trend summaries were computed. First, to understand how seasonal trends

varied geographically, we used a GAM (Wood *et al.* 2016) to generate a spatially explicit

estimate of the average percent change per year based on the ensemble of base models

with stixel centroid dates falling within a specified season. Second, to identify regions

consistently estimated to be in decline, we use a binary response GAM to generate the

corresponding spatial estimate of the probability of decline. Third, we estimated the

expected trend within a specified region and season by averaging across the set of partial

effect estimates with stixel centroids lying within the specified extent. To generate

estimates of uncertainty for the expected trend, we computed 90% point-wise confidence

intervals from a Monte Carlo sample of expected trends. To insure that the uncertainty

estimates were conservative, each expected trend was based on an average of 5 partial

effect estimates, far fewer than the number available.

Model Validation

To assess model quality, we validated the model's ability to predict the observed patterns

of occupancy and abundance using independent validation data. The statistics were

evaluated using a Monte Carlo design of 25 spatially balanced samples to help control for

the uneven spatial distribution of the validation data (Fink *et al.* 2010; Roberts *et al.*

2017). To quantify the predictive performance for the AOO we used the Area Under the

Curve (AUC) and Kappa (Cohen 1960) statistics to describe the models' ability to

classify un/occupied sites (Freeman & Moisen 2008). AUC measures a model's ability to

discriminate between positive and negative observations (Fielding & Bell 1997) as the

probability that the model will rank a randomly chosen positive observation higher than a

randomly chosen negative one. Cohen's Kappa statistic (Cohen 1960) was designed to

measure classification performance taking into account the background prevalence. To

quantify the quality of the relative occupancy estimate as a rate within the AOO, we

evaluated AUC and Kappa. To quantify the quality of the abundance estimates we

computed Spearman's Rank Correlation (SRC) and the percent Poisson Deviance

Explained (P-DE). SRC measures how well the abundance estimates rank the observed

abundances and the P-DE measures the correspondence between the magnitude of the

estimated counts and observed counts.

(c) Results

Weekly AOO, Relative Occupancy and Abundance

Using the Wood Thrush as exemplar analysis, we generated CS-FAC estimates of AOO,

relative occupancy and abundance at a spatiotemporal resolution of 8.4km × 8.4km ×

1week (Fig 1). A site was considered occupied if the PAT estimator was at least 0.05,

meaning that at least one individual of the species is expected to be detected on at least 1

out of every 20 independent, standardized eBird surveys of the site on the given day.

Unoccupied grid cells were considered to have zero occupancy and abundance, thus the

AOO was depicted as the boundary between pixels with and without color.

To assess the accuracy of estimates, we calculated range-wide validation estimates based

on spatially balanced samples of independent eBird observations. AOO weekly median

AUC scores were between 0.72 and 0.98 with mean 0.81 (Fig 2a) and AOO weekly

median Kappa scores were between 0.16 and 0.38 with mean 0.24 (Fig 2b). Relative

occupancy weekly median AUC scores were between 0.66 and 0.86 with mean 0.75 (Fig

2c) and relative occupancy weekly median Kappa scores were between 0.18 and 0.51

with mean 0.30 (Fig 2d). Relative abundance weekly median P-DE scores were between

0.01 and 0.71 with mean 0.20 (Fig 2e) and relative abundance weekly median SRC

scores were between 0.24 and 0.52 with mean 0.31 (Fig 2f). Weeks with insufficient

validation data were shown as zero. These weeks occurred during the migrations, when

detection rates and counts are lowest. Variation in predictive performance was highest

during the non-breeding season for all metrics, reflecting lower data densities in

Mesoamerica.

The AOO shows the seasonal changes in the population range size and shape while the

abundance estimates capture regional and seasonal variation in population structure

within the range. The breeding season range fills in the eastern deciduous forests east of

the Great Plains with highest population concentrations in the Appalachian Mountains

(Fig 1a). During autumn migration, the population concentrates in the southern part of the

Appalachian Mountains (Fig 1b) before crossing the Gulf of Mexico into Central

America. The winter distribution (Fig 1c) is concentrated in the Yucatán Peninsula, with

lower concentrations extending north into Veracruz and south to Costa Rica and Panama.

During the spring migration (Fig 1d), Wood Thrush crosses the Gulf, concentrating on

the Gulf Coast and again in the southern part of the Appalachian Mountains.

Seasonal Habitat Use and Avoidance

To quantify changes in habitat use and avoidance throughout the annual cycle, we made

weekly estimates of the association between Wood Thrush occupancy and the amount of

each habitat class in the local landscape (Fig 3). For each week, the associations were

summarized across the population core area, the 5° longitude × 5° latitude area located at

the population center for that week. For each cover class, values were combined for both

PLAND and LPI predictors to describe the relative strength and direction of the

association. Larger absolute values indicate stronger associations and the sign of the

value indicates class use or avoidance. Classes with inconsistent direction of association,

were removed, resulting in total weekly relative importance that sums to less than 1.

The accuracy of the habitat associations follows from the strong validation results (Fig 2).

The Wood Thrush breeding season is characterized by the strong positive association

with deciduous broadleaf forest and the non-breeding season is characterized by the

strong positive association tropical broadleaf forest. During the migrations, the

population is associated with a wider variety of cover classes, and a more even

distribution of associations, both positive and negative. This includes a notable positive

association with the urban developed class.

Inter-annual Seasonal Trends

We estimated the 2004–16 inter-annual trends during the breeding (May 30–July 3) and

non-breeding (Dec 1–Feb 28) seasons. The Wood Thrush population declined across

most of its range during the breeding season (Fig 4a). The steepest declines reached 3 to

4% per year along the east coast, and the southeastern and northwest portions of the

breeding range (Fig 4a). The green contour indicates where at least 95% of base model

trend estimates were declining, a large region including areas in the northeast, the Mid-

Atlantic coast, the Piedmont, and the Appalachian Mountains (Fig 4a). The breeding

season trend in the Ohio/West Virginia area (Fig 4b), declined at an average of 1% per

year, though this trend did show slight increases from 2009 to 2011. In the southern

Appalachian Mountains area (Fig 4a) the breeding season trend declined at an average of

3% per year, with the strongest declines from 2004-10 (Fig 4b). The winter population

(Fig 5) has declined at an average rate of 5.6% per year, with the largest declines from

2004-2010.

(d) Discussion

In this paper, we address challenges related to obtaining cross-scale, full annual cycle

information on patterns of abundance and distribution of bird species using citizen

science data. The resolution, extent, and comprehensiveness of the information generated

with this methodology is unprecedented, and has the potential to increase our knowledge

of information-poor species, regions and seasons (Runge *et al.* 2015).

The approach we present generated robust inferences about species' ranges, occupancy

and abundance (Fig. 2), habitat associations, and seasonal trends, confirming the

accuracy and utility of the approach. More broadly, the analysis presented here

demonstrates how citizen science data can be used to generate accurate species-level

information for broad-scale biodiversity monitoring like those outlined by the Group on

Earth Observations Biodiversity Observation Network (Kissling *et al.* 2017). It is worth

noting that without a single, comprehensive source of information, making population-

wide assessments requires the additional steps to acquire, analyze, and calibrate disparate

sources of information. Similarly, without critical ancillary information describing

participant search effort and information to infer the absence of species (e.g., complete

checklists), we would have been unable to account for the bias of imperfect detection. For

this reason, we advocate for other citizen science projects to collect ancillary information

sufficient to untangle the complexities of heterogeneous observation and ecological

processes.

To demonstrate how this methodology can be used to estimate complex patterns of

species' movement, phenology, and population concentration across the full annual cycle

we analyzed eBird data for Wood Thrush. This analysis captured important, known

patterns of movement, phenology, and concentration during the breeding (Fig 1a) and

non-breeding seasons (Fig 1c). Additionally, theses results filled important knowledge

gaps, providing novel population-level information during the less well-studied stages of

the annual cycle, such as migration and the overwintering period (Fig 1b and Fig 1d).

The estimated patterns of habitat use and avoidance (Fig 3) were consistent with

documented seasonal (Zuckerberg *et al.* 2016) and regional (Evans *et al.* 2011) patterns.

Notably, this is the first comprehensive population-level analysis of habitat associations

for not just Wood Thrush, but for any Neotropical migrant. In general, comprehensive

information on species' habitat associations will be useful for conservation planning and

prioritization, especially outside of the breeding season.

The population trend estimate for the non-breeding season is, to the best of our

knowledge, the first population-wide trend estimate for Wood Thrush outside the

breeding season. This trend estimate fills an important gap in understanding the role of

the autumn migration and non-breeding season on overall Wood Thrush population

health. The population-wide rate of decline during the non-breeding season (-5.6%) was

larger than the regional breeding season rates (1-3%), suggesting that limitation is

happening during the autumn migration and/or the non-breeding season. Although this is

consistent with the demographic models of Taylor & Stutchbury (2016), it contradicts the

results from Rushing *et al.* (2017), suggesting that there might be regional variation in

population declines in Central America.

Another novel aspect of the methodology is the ability to use citizen science data to

estimate trends in relative abundance - a task usually left to monitoring programs which

employ more stringent sampling protocols and are hard to deploy across broad extents.

The potential to use eBird data to accurately estimate population trends will grow with

the increasing volume and density of data. Increasing volumes and density of data will

further improve the precision and spatiotemporal resolution of trend estimates across a

wider geographic area than is currently possible. This can be seen when comparing the

Wood Thrush breeding and non-breeding season trends (Fig 4 & 5). The increased

volume and density of data in the breeding season, made it possible to estimate trends

with greater spatial resolution and higher precision than in the non-breeding season. The

increasing availability of population trends during the non-breeding season will help to

refine our understanding of where and when populations are limited or regulated,

complimenting migratory connectivity information derived from individual-level tracking

data.

Aside from filling knowledge gaps, the comprehensive nature of CS-FAC information

can be used for other novel and important applications. With CS-FAC distributional

information, it is straightforward to make population-wide comparisons and

prioritizations and to coordinate conservation activities across regions and seasons.

Moreover, once regions of interest have been identified, the resolution of CS-FAC can be

leveraged to seamlessly compare and prioritize landscapes within regions (e.g. Reynolds

et al. 2017). In addition, the impact of regional and seasonal scale processes can be

integrated across space throughout the year, making it possible to carry out accurate

multi-scale population-wide impact assessments. This is important for studying a variety

of broad-scale environmental and anthropogenic effects, many of which are themselves

multi-scale processes, from land-use change to ecosystem services (e.g., La Sorte *et al.*

2017).   The potential of our approach to to integrate effects also addresses an important

multi-scale challenge in climate change studies (Ådahl *et al.* 2006; Small-Lorenz *et al.*

2013) where nearly all facets of climate (e.g. temperature and precipitation) exhibit

strong regional-scale intra-annual variation.

(e) Acknowledgements

We thank the eBird participants for their contributions, the eBird team for their support,

and reviewers for their constructive suggestions. This work was funded by The Leon

Levy Foundation, The Wolf Creek Foundation, NASA (NNH12ZDA001N-ECOF),

Microsoft Azure Research Award (CRM: 0518680), and the National Science Foundation

(ABI sustaining: DBI-1356308; computing support from CNS-1059284 and CCF-

1522054).

(f) References

Ådahl, E., Lundberg, P., Jonzen, N. (2006) From climate change to population change: the need to consider annual life cycles. *Glob. Change Biol.* 12, 1627– 1633. (doi:10.1111/j.1365-2486.2006.01196.x)

Amatulli, G., Domisch S., Tuanmu M.-N., Parmentier B., Ranipeta A., Malczyk J., and Jetz W. (2017). A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. *Scientific Data*, in press (accepted).

Bird, T.J., Bates, A.E., Lefcheck, J.S., Hill, N.A., Thomson, R.J., Edgar, G.J., Stuart-Smith, R.D., Wotherspoon, S., Krkosek, M., Stuart-Smith, J.F. and Pecl, G.T., (2014). Statistical solutions for error and bias in global citizen science datasets. *Biological Conservation*, 173, 144-154.

Breiman, L., (2001). Random forests. *Machine learning*, 45(1), 5-32.

Boakes, E.H., McGowan, P.J., Fuller, R.A., Chang-qing, D., Clark, N.E., O'Connor, K. and Mace, G.M., (2010). Distorted views of biodiversity: spatial and temporal bias in species occurrence data. *PLoS biology*, 8(6), p.e1000385.

Bohrer, G., Beck, P.S., Ngene, S.M., Skidmore, A.K. and Douglas-Hamilton, I., (2014). Elephant movement closely tracks precipitation-driven vegetation dynamics in a Kenyan forest-savanna landscape. *Movement Ecology*, 2(1), 2.

Cleveland, W.S., Grosse, E. and Shyu, W.M., (1992). Local regression models. *Statistical models in S*, 2, 309-376.

Cohen, J., (1960). A coefficient of agreement for nominal scales. *Educational and psychological measurement*, 20(1), 37-46.

Denes, F.V., Silveira, L.F. and Beissinger, S.R., (2015). Estimating abundance of

    unmarked animal populations: accounting for imperfect detection and other sources

    of zero inflation. *Methods in Ecology and Evolution*, 6(5), 543-556.

Dickinson, J.L., Zuckerberg, B. and Bonter, D.N., (2010). Citizen science as an

    ecological research tool: challenges and benefits. *Annual review of ecology,*

    evolution, and systematics, 41, 149-172.

Diefenbach, D.R., Marshall, M.R., Mattice, J.A. and Brauning, D.W., (2007).

    Incorporating availability for detection in estimates of bird abundance. *The Auk*,

    124(1), 96-106.

Elith, J. and Leathwick, J.R., (2009). Species distribution models: ecological explanation

    and prediction across space and time. Annual review of ecology, evolution, and

    systematics, 40, 677-697.

Efron, B., (2014). Estimation and accuracy after model selection. *Journal of the*

    American Statistical Association, 109(507), 991-1007.

European Bird Census Council, (2016) *Trends of common birds in Europe, 2016 update.*

    Available at: http://www.ebcc.info/index.php?ID=612. Last accessed 06 December

    2017.

Evans, M., Gow, E., Roth, R. R. , Johnson, M. S. and Underwood, T. J. (2011). Wood

    Thrush (Hylocichla mustelina), version 2.0. In *The Birds of North America*

    (Rodewald, P. G. editor). Cornell Lab of Ornithology, Ithaca, New York, USA.

    https://doi.org/10.2173/bna.246

Faaborg, J., Holmes, R.T., Anders, A.D., Bildstein, K.L., Dugger, K.M., Gauthreaux,

    S.A., Heglund, P., Hobson, K.A., Jahn, A.E., Johnson, D.H. and Latta, S.C., (2010).

Recent advances in understanding migration systems of New World land birds. *Ecological monographs*, 80(1), 3-48.

Fielding, A.H. and Bell, J.F., (1997). A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environmental conservation*, 24(1), 38-49.

Fink, D., Hochachka, W.M., Zuckerberg, B., Winkler, D.W., Shaby, B., Munson, M.A., Hooker, G., Riedewald, M., Sheldon, D. and Kelling, S., (2010). Spatiotemporal exploratory models for broad_scale survey data. *Ecological Applications*, 20(8), 2131-2147.

Fink, D., Damoulas, T. and Dave, J., (2013), July. Adaptive Spatio-Temporal Exploratory Models: Hemisphere-wide species distributions from massively crowdsourced eBird data. In: Twenty-Seventh AAAI Conference on Artificial Intelligence (AAAI-13) July 14–18, 2013 in Bellevue, Washington, USA.

Fink, D., Damoulas, T., Bruns, N.E., La Sorte, F.A., Hochachka, W.M., Gomes, C.P. and Kelling, S., (2014). Crowdsourcing meets ecology: hemisphere-wide spatiotemporal species distribution models. *AI magazine*, 35(2), 19-30.

Fink, D., Auer,T., Obregon, F., Hochachka,W.M., Iliff,M., Sullivan,B., Wood,C., Davies,I., and Kelling, S. (2017). *The eBird Reference Dataset Version 2016 (ERD2016)*. Available at: http://ebird.org/ebird/data/download/erd. Last accessed 13 December 2017.

Finley, A.O., (2011). Comparing spatially-varying coefficients models for analysis of ecological data with non-stationary and anisotropic residual dependence. *Methods in Ecology and Evolution*, 2(2),143-154.

Fithian, W. and Hastie, T., (2014). Local case-control sampling: Efficient subsampling in imbalanced data sets. *Annals of statistics*, 42(5), 1693.

Freeman, E. A. and Moisen, G. (2008). PresenceAbsence: An R Package for Presence-Absence Model Analysis. *Journal of Statistical Software*, 23(11):1-31. http://www.jstatsoft.org/v23/i11

Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X. (2010). MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets. *Remote Sensing of Environment,* 114, 168–182.

Gelfand, A.E., Banerjee, S. and Finley, A.O., (2012). Spatial design for knot selection in knot-based dimension reduction models. In: *Spatio-Temporal Design: Advances in Efficient Data Acquisition*, eds. Mateu, J. and Müller, W.G., John Wiley & Sons., pp.142-169.

Geyer, C. J. (2013) *5601 Notes: The Subsampling Bootstrap*, Available at: http://www.stat.umn.edu/geyer/5601/notes/sub.pdf. Last accessed 06 December 2017.

Hansen, M. C., DeFries, R. S., Townshend, J. R. G., & Sohlberg, R. (2000). Global land cover classification at the 1km spatial resolution using a classification tree approach. *International Journal of Remote Sensing*, 21, 1331-1364.

Haste, T., Tibshirani, R., J. Friedman. (2009). *The elements of statistical learning (2nd ed)*. Springer series in statistics, New York.

Heffernan, J.B., Soranno, P.A., Angilletta, M.J., Buckley, L.B., Gruner, D.S., Keitt, T.H., Kellner, J.R., Kominoski, J.S., Rocha, A.V., Xiao, J. and Harms, T.K., (2014). Macrosystems ecology: understanding ecological patterns and processes at continental scales. *Frontiers in Ecology and the Environment*, 12(1), 5-14.

Hochachka, W.M., Fink, D., Hutchinson, R.A., Sheldon, D., Wong, W.K. and Kelling, S. (2012). Data-intensive science applied to broad-scale citizen science. *Trends in ecology & evolution*, 27(2), 130-137.

Hurlbert, A.H. and Jetz, W. (2007). Species richness, hotspots, and the scale dependence of range maps in ecology and conservation. *PNAS*, 104(33), 13384-13389.

Johnston, A., Fink, D., Reynolds, M.D., Hochachka, W.M., Sullivan, B.L., Bruns, N.E., Hallstein, E., Merrifield, M.S., Matsumoto, S. and Kelling, S. (2015). Abundance models improve spatial and temporal prioritization of conservation resources. *Ecological Applications*, 25(7), 1749-1756.

Kissling, W.D., Ahumada, J.A., Bowser, A., Fernandez, M., Fernández, N., García, E.A., Guralnick, R.P., Isaac, N.J., Kelling, S., Los, W. and McRae, L. (2017). Building essential biodiversity variables (EBVs) of species distribution and abundance at a global scale. *Biological reviews*. doi: 10.1111/brv.12359

Kearney, M. & Porter, W. (2009). Mechanistic niche modelling: combining physiological and spatial data to predict species' ranges. *Ecology Letters*, 12, 334–350.

Kéry, M. and Royle, J.A., (2015). *Applied Hierarchical Modeling in Ecology: Analysis of distribution, abundance and species richness in R and BUGS: Volume 1: Prelude and Static Models.* Academic Press.

La Sorte, F.A., Fink, D., Blancher, P.J., Rodewald, A.D., Ruiz-Gutierrez, V., Rosenberg, K.V., Hochachka, W.M., Verburg, P.H. and Kelling, S., (2017). Global change and the distributional dynamics of migratory bird populations wintering in Central America. *Global Change Biology*, 23(12), 5284-5296.

Marra P.P., Cohen, E.B., Loss, S.R., Rutter, J.E., Tonra, C.M. (2015). A call for full
annual cycle research in animal ecology. *Biol. Lett.* 11: 20150552.
http://dx.doi.org/10.1098/rsbl.2015.0552

Mease, D., Wyner, A. and Buja, A., (2007). Cost-weighted boosting with jittering and
over/under-sampling: JOUS-boost. *J. Machine Learning Research*, 8, 409-439.

McGarigal, K., Cushman, S.A., and Ene, E. (2012). *FRAGSTATS v4: Spatial Pattern*
Analysis Program for Categorical and Continuous Maps. Available at:
http://www.umass.edu/landeco/research/fragstats/fragstats.html.

Norris, D.R. and Marra, P.P. (2007). Seasonal interactions, habitat quality, and
population dynamics in migratory birds. *The Condor*, 109(3), 535-547.

Pereira, J. A., and A. J. Novaro. (2014). Habitat-specific demography and conservation of
Geoffroy's cats in a human-dominated landscape. *Journal of Mammalogy* 95, 1025–
1035.

Politis, D. N., Romano, J.P., and Wolf, M. (1999). *Subsampling, Springer Series in*
Statistics, Springer-Verlag New York, Inc.

R Development Core Team (2017). *R: A language and environment for statistical*
computing. R Foundation for Statistical Computing http://www.R–project.org,
Vienna, Austria.

Reynolds, M.D., Sullivan, B.L., Hallstein, E., Matsumoto, S., Kelling, S., Merrifield, M.,
Fink, D., Johnston, A., Hochachka, W.M., Bruns, N.E. and Reiter, M.E., 2017.
Dynamic conservation for migratory species. *Science Advances*, 3(8), p.e1700707.

Ridgeley, R.S., Allnutt, T.F., Brooks, T., NcNicol, D.K., Mehlman, D.W., Young, B.E. and Zook, J.R., BirdLife International. 2012. Digital distribution maps of the birds of the Western Hemisphere, version, 5.

Ridgeway, G. with contributions from others. (2017). gbm: Generalized Boosted Regression Models. R package version 2.1.3. https://CRAN.R-project.org/package=gbm

Roberts, D.R., Bahn, V., Ciuti, S., Boyce, M.S., Elith, J., Guillera-Arroita, G., Hauenstein, S., Lahoz-Monfort, J.J., Schröder, B., Thuiller, W. and Warton, D.I., (2017). Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography*, 40, 913–929.

Robinson, O.J., Ruiz-Gutierrez, V., Fink, D. (2017). Correcting for bias in distribution modeling for rare species using citizen science data. *Diversity and Distributions*, in press (accepted).

Runge, C.A., Watson, J.E., Butchart, S.H., Hanson, J.O., Possingham, H.P. and Fuller, R.A., (2015). Protected areas and global conservation of migratory birds. *Science*, 350(6265), pp.1255-1258.

Rushing, C.S., Hostetler, J.A., Sillett, T.S., Marra, P.P., Rotenberg, J.A. and Ryder, T.B., (2017). Spatial and temporal drivers of avian population dynamics across the annual cycle. *Ecology*, 98(11), 2837–2850.

Sauer, J.R. and Link, W.A., (2011). Analysis of the North American breeding bird survey using hierarchical models. *The Auk*, 128(1), 87-98.

Sherry, T.W. & Holmes, R.T. (1995) Summer versus winter limitation of populations: what are the issues and what is the evidence. In: *Ecology and Management of*

Neotropical Migratory Birds: a Synthesis and Review of Critical Issues (eds T.E.

Martin & D.M. Finch), Oxford University Press, New York, pp. 85–120.

Small-Lorenz, S.L., Culp, L.A., Ryder, T.B., Will, T.C., Marra, P.P. (2013) A blind spot

in climate change vulnerability assessments. *Nat. Clim. Change* 3, 91 – 93.

(doi:10.1038/nclimate1810)

Sullivan, B.L., Wood, C.L., Iliff, M.J., Bonney, R.E., Fink, D. and Kelling, S., (2009).

eBird: A citizen-based bird observation network in the biological sciences. *Biological*

Conservation, 142(10), 2282-2292.

Sullivan, B.L., Aycrigg, J.L., Barry, J.H., Bonney, R.E., Bruns, N., Cooper, C.B.,

Damoulas, T., Dhondt, A.A., Dieterich, T., Farnsworth, A. and Fink, D., (2014). The

eBird enterprise: an integrated approach to development and application of citizen

science. *Biological Conservation*, 169, 31-40.

Wood, S.N., Pya, N. and Säfken, B., (2016). Smoothing parameter and model selection

for general smooth models. *Journal of the American Statistical Association*,

111(516), pp.1548-1563.

VanDerWal, J., Falconi, L., Januchowski, S., Shoo, L. and Storlie, C., (2014).

SDMTools: Species Distribution Modelling Tools: Tools for processing data

associated with species distribution modelling exercises. *R package version*, 1-1.

Zaharia, M., Xin, R.S., Wendell, P., Das, T., Armbrust, M., Dave, A., Meng, X., Rosen,

J., Venkataraman, S., Franklin, M.J. and Ghodsi, A., (2016). Apache Spark: A unified

engine for big data processing. *Communications of the ACM*, 59(11), 56-65.

Zuckerberg, B., Fink, D., La Sorte, F.A., Hochachka, W.M. and Kelling, S., (2016).

Novel seasonal land cover associations for eastern North American forest birds

identified through dynamic species distribution modelling. *Diversity and Distributions*, 22(6), 717-730.

Table 1: Land and water cover classes used for distribution modeling. All cover classes were summarized within a 2.8km × 2.8km (784 hectares) landscape centered on each checklist location. Within each landscape, we computed the proportion of each class, and three descriptions of the spatial configuration of the class within the landscape.

Figure 1: Wood Thrush estimates of area of occupancy and relative abundance at 8.4km × 8.4km resolution during (a) breeding (June 20), (b) autumn migration (October 3), (c) non-breeding (December 12), and (d) spring migration (March 28) seasons. Positive abundance is only shown in areas estimated to be occupied and the area of occupancy is depicted as the boundary between pixels with and without color. Brighter colors indicate areas occupied with higher abundance. Relative abundance was measured as the expected count of the species on a standardized 1km survey conducted from 7-8AM.

Figure 2: Boxplots of range-wide weekly predictive performance for area of occupancy, relative occupancy and abundance estimates across 25 Monte Carlo samples of spatially balanced validation data. (a) AUC and (b) Kappa scores for area of occupancy estimates. (c) AUC and (d) Kappa scores for relative occupancy estimates. (e) Spearman's Rank Correlation and (f) Percent Deviance Explained scores for relative abundance estimates.

Figure 3: The weekly relative importance for the amount of each land and water cover class for the core Wood Thrush population. Positive importance indicates class use and negative importance indicates class avoidance. The strength of the association with each class is proportional to the width of the class color. Classes with inconsistent direction of

association were removed, resulting in total weekly relative importance that sums to less than 1.

Figure 4: Wood Thrush breeding trend map and regional estimates. The map shows the average percent change per year in relative abundance from 2004–16 during the breeding season (May 30– July 3). The green contour indicates the region where the probability of decline is at least 95%. Breeding season regional trends and 90% point-wise confidence intervals are shown for the (A) Ohio-West Virginia and (B) Southern Appalachian regions as the deviation from the mean on the log scale.

Figure 5: Wood Thrush non-breeding trend map and regional estimate. The map shows the average percent change per year in relative abundance from 2004–16 during the non-breeding season (Dec 1–Feb 28). The regional trend and 90% point-wise confidence intervals are shown as the deviation from the mean on the log scale.

MODIS Class Codes	Land Cover	Water Cover
0		Shallow Ocean
1	Evergreen Needleleaf Forest	
2	Evergreen Broadleaf Forest	Ocean coastlines and lake shores
3	Deciduous Needleleaf Forest	Shallow inland water
4	Deciduous Broadleaf Forest	
5	Mixed Forest	Deep Inland Water
6	Closed Shrublands	Moderate or continental ocean
7	Open Shrublands	Deep Ocean
8	Woody Savannas	
9	Savannas	
10	Grasslands	
12	Croplands	
13	Urban and built-up	
16	Barren	

Table 1

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supporting Information for

"Modeling Avian Full Annual Cycle Distribution and Population Trends with Citizen Science Data"

Daniel Fink, Tom Auer, Viviana Ruiz-Gutierrez, Wesley M. Hochachka, Alison Johnston, Frank A. La Sorte, and Steve Kelling

This document includes supplemental information on the

1. STEM ensemble design, and

2. Subsampling procedures to estimate uncertainty of the occupancy and abundance estimates.

These are described in the appendices below.

Appendix S1: Ensemble Design

The ensemble of stixels was designed as a Monte Carlo sample of 100 randomly located spatiotemporal partitions of the spatiotemporal study extent. This results in a sample of stixels uniformly distributed through out the spatiotemporal extent of study. Averaging across this sample helps control for biases associated with the arbitrary partitioning of data into stixels. We also use the Monte Carlo sample to generate estimates of uncertainty by incorporating subsampling into the selection of training data within stixels.

An important part of the STEM implementation was determining the spatial and temporal dimensions of the stixels. When averaging across the ensemble, stixel size controls an important bias-variance tradeoff (Fink *et al* 2010; Fink *et al.* 2013). Stixel size needs to

be chosen small enough to capture local predictor-response (i.e. species-environment)

relationships, controlling the bias of base model estimates. Stixel size also needs to be

chosen large enough to meet the minimum sample size requirements necessary for fitting

the base models: this controls the variance when averaging across the ensemble.

As stixel size gets smaller, the training data sample size within a stixel also decreases. When training sample sizes are too small to fit base models, the number of base model estimates available for ensemble averaging also decreases, increasing the variance of the ensemble estimator. The number of stixels used to compute a local estimate across the ensemble is called the *ensemble support*. Ensemble support is important because it determines effectiveness of ensemble averaging to control inter-model variability. In general, ensemble support follows patterns of data density, filtered through a combination of stixel geometry and the base model minimum sample size requirements.

Because of the irregular and often sparse distribution of eBird data, selecting the spatial and temporal dimensions of stixels necessary to maintain ensemble support is a non-trivial process. Our goal was to select the stixel size parameters to maximize the spatiotemporal coverage of the analysis while maintaining sufficient ensemble support to guarantee good model performance. To operationalize this, we required an ensemble support of at least 50 stixels, throughout at least 75% the Western Hemisphere for each week of the year. Estimates of occupancy and abundance were only produced when ensemble support was above 50.

We began by specifying the temporal dimension of the partitions to be 30 contiguous days. A 30 day window is small enough to capture a wide variety of migration patterns across a diverse set of terrestrial species using eBird data (Johnston *et al.* 2015; La Sorte *et al.* 2017). For simplicity, we decided to divide space into longitude-latitude squares

using un-projected latitude × longitude space. To account for the relatively low data density (Fink *et al.* 2013) in the southern part of the study area we doubled the length of stixel squares south of 12° latitude. Given the 30-day temporal dimension and minimum sample size requirements (see below for details), we found that a stixel length of 10° in the north and 20° in the south met our requirements, resulting in a minimum weekly coverage of 79% of the terrestrial Western Hemisphere in December and up to 90% coverage of the Western Hemisphere in July. Figure S1 shows realizations of 3 randomly located spatial partitions used to define STEM stixels for the analysis. This image shows how stixels overlap across the randomly located partitions and it shows how stixel size varies between North and South America.

The base model minimum sample size requirements affect base model bias and variance as well as ensemble support. To fit a base model, we required that the training data meet the following three criteria. First, we required a minimum sample size of 50 checklists. Second, to insure minimum spatial coverage within each base model, we required that at least 50 cells from the spatially balanced case-control sampling procedure (see below for details) contained checklists. Finally, to insure a minimum signal to predict positive relative occupancy and abundance, we required at least 10 species detections, i.e. non-zero counts, among the checklists. To guard against the effects of replicate surveys at popular birding locations, only one detection per day is considered from each location.

FIGURE S1: Realizations of three randomly located spatial partitions used to define STEM stixels. This image shows how stixels overlap across the randomly located partitions and it shows how stixel size varies between North and South America. One hundred randomized partitions were used for the analysis.

Appendix S2: Subsampling procedures to estimate uncertainty

The ensemble means are used to estimate relative occupancy and abundance. Consequently, the variation across the ensemble itself provides a conservative estimate of uncertainty for the ensemble mean (Efron 2014). A straightforward, brute force approach way to generate more accurate estimates of uncertainty for the ensemble mean can be computed by bootstrapping the ensemble mean, however, this is computationally prohibitive.

Instead, we employed a subsampling approach (Politis *et al.* 2009) following the

computational strategy of Geyer (2013). We faced two challenges to implement this

approach. First, the sample size, here, the ensemble support, was very small, 100 at most.

Second, the computational efficiency of the approach was very important because we

needed to compute uncertainty estimates for up to 86M quantities per species (= 614K

locations / week * 52 weeks * 2 estimates per location [occupancy & abundance] +

another 28M for the 14M training & testing checklists * 2 estimates per checklist). To

deal these challenges we selected a set of parameter settings that balanced the quality of

the interval estimates with the computational costs of generating them.

When the sample size was less than 10, subsampling was not performed and quantiles of

the original sample were to estimate uncertainty. For sample sizes greater than or equal to

ten, we computed the upper 90^{th} confidence limit and lower 10^{th} confidence limit. The

subsampling was performed with two different sizes to facilitate estimation of the rate

parameter used to correct the uncertainty estimates. Following Geyer (2013), we

subsampled the square root of the ensemble support value and the -1.5 power of the

ensemble support value.

To check these parameter settings, a small simulation test was run. We found that for

sample sizes of 25 or less, the rate parameter estimates tended to be too small, resulting in

intervals that were too small and had poor coverage. To mitigate this, we adjusted the rate

parameter estimate upwards by 0.5 of the rate parameter's standard error, producing more

conservative uncertainty estimates. In the cases where the rate parameter estimate was negative, subsampling was not performed and quantiles of the entire sample were used producing conservative uncertainty estimates. Note that ensemble support requirements for the occupancy and abundance estimates excludes most of the estimates suffering from these small sample size complications.

References

Efron, B., (2014). Estimation and accuracy after model selection. Journal of the American Statistical Association, 109(507), 991-1007.

Fink, D., Hochachka, W.M., Zuckerberg, B., Winkler, D.W., Shaby, B., Munson, M.A., Hooker, G., Riedewald, M., Sheldon, D. and Kelling, S., (2010). Spatiotemporal exploratory models for broad_scale survey data. *Ecological Applications*, 20(8), 2131-2147.

Fink, D., Damoulas, T. and Dave, J., (2013), July. Adaptive Spatio-Temporal Exploratory Models: Hemisphere-wide species distributions from massively crowdsourced eBird data. In: *Twenty-Seventh AAAI Conference on Artificial Intelligence* (AAAI-13) July 14–18, 2013 in Bellevue, Washington, USA.

Geyer, C. J. (2013) 5601 Notes: The Subsampling Bootstrap, Available at: http://www.stat.umn.edu/geyer/5601/notes/sub.pdf. Last accessed 06 December 2017.

Johnston, A., Fink, D., Reynolds, M.D., Hochachka, W.M., Sullivan, B.L., Bruns, N.E., Hallstein, E., Merrifield, M.S., Matsumoto, S. and Kelling, S. (2015). Abundance models improve spatial and temporal prioritization of conservation resources. *Ecological Applications*, 25(7), 1749-1756.

La Sorte, F.A., Fink, D., Blancher, P.J., Rodewald, A.D., Ruiz-Gutierrez, V., Rosenberg, K.V., Hochachka, W.M., Verburg, P.H. and Kelling, S., (2017). Global change and the distributional dynamics of migratory bird populations wintering in Central America. *Global Change Biology*, 23(12), 5284-5296.Mease *et al.* 2007

Politis, D. N., Romano, J.P., and Wolf, M. (1999). *Subsampling*, *Springer Series in Statistics*, Springer-Verlag New York, Inc.