Whole genome sequencing for predicting clarithromycin resistance in *Mycobacterium*

abscessus

Samuel Lipworth[1]*, Natasha Hough[1]*, Laura Leach[2], Markus Morgan[2], Katie Jeffery[2], Monique

Andersson[2], Esther Robinson[3], E. Grace Smith[3], Derrick Crook[1], Tim Peto[1] and Timothy

Walker[1]

*Contributed equally

[1] Nuffield Department of Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK,

OX3 9DU

[2] Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Oxford, UK, OX3

9DU

[3] Public Health England Regional Mycobacterial Reference Laboratory, Birmingham Heartlands

Hospital, Birmingham, UK, B9 5SS

Running title: WGS for M. abscessus clarithromycin susceptibility prediction

Corresponding author:

Samuel Lipworth

Email: samuel.lipworth@medsci.ox.ac.uk,

Tel: 07743448071

Abstract

Mycobacterium abscessus is emerging as an important pathogen in chronic lung diseases with

concern regarding patient to patient transmission. The recent introduction of routine whole

genome sequencing (WGS) as a replacement for existing reference techniques in England

provides an opportunity to characterise the genetic determinants of resistance. We conducted a

systematic review to catalogue all known resistance determining mutations. This knowledge

was used to construct a predictive algorithm which was tested on a collection of 203

sequentially acquired clinical isolates for which there was paired genotype/phenotype data. A

search for novel resistance determining mutations was conducted using an heuristic algorithm.

The sensitivity of existing knowledge for predicting resistance in clarithromycin was 76.19%

(95% CI 52.8 - 91.8%) and the specificity was 100%. Subspecies alone was a poor predictor of

resistance to clarithromycin. Eight potential new resistance conferring SNPs were identified.

WGS demonstrates probable resistance determining SNPs in regions the NTM-DR line probe

cannot detect. These mutations are potentially clinically important as they all occurred in

samples predicted to be inducibly resistant, and for which a macrolide would therefore currently

be indicated. We were unable to explain all resistance, raising the possibility of the involvement

of other as yet unidentified genes.

Introduction

Mycobacterium abscessus is a rapidly growing nontuberculous mycobacterium (NTM) of increasing clinical concern with a rising burden of associated pulmonary disease(1). It poses a significant problem, particularly in patients with cystic fibrosis (CF), where infection is associated with a more rapid decline in lung function and can be a barrier to transplantation(2). Of particular concern are the findings from recent work that have suggested person-to-person transmission of virulent clones amongst the CF population within a healthcare setting(3, 4), although not all studies have supported this(5, 6).

The taxonomy of *M. abscessus* is contentious. It is currently divided into three subspecies: *M. abscessus* subspecies *abscessus (Maba), M. abscessus* subspecies *massiliense (Mabm)*, and *M. abscessus* subspecies *bolletii (Mbol)(7)*. The organism has intrinsic resistance to multiple antibiotics including β-lactams, rifampicin and aminoglycosides due to the synergistic action of the cell envelope and genetic factors termed the 'intrinsic resistome'(8). Treatment requires prolonged courses of multiple antibiotics, but outcomes are thought to vary across the different subspecies. *Mabm* has been associated with clarithromycin susceptibility and favourable treatment outcomes, whereas *Maba* has been associated with inducible macrolide resistance and poorer treatment outcomes(9). However, the American Thoracic Society/Infectious Diseases Society of America and British Thoracic Society (ATS/IDSA and BTS) guidelines recommend including a macrolide in treatment regimens where samples are either susceptible, or demonstrate inducible resistance(10, 11).

Whole genome sequencing has been implemented in stages across England since December 2016, replacing existing reference techniques for mycobacterial identification. As a consequence, there is now the opportunity to explore the molecular determinants of drug

resistance for all clinical NTM isolates, as has been successfully achieved in a range of different

bacteria(12–14). As there is a particularly strong correlation between *in vitro* susceptibility and

clinical response to macrolide treatment of *M. abscessus*(15, 16), we have undertaken a study

to assess the feasibility of predicting clarithromycin susceptibility from whole genome

sequencing data for all three subspecies of *M. abscessus.*

Methods

Literature search

We first conducted a systematic review of the literature to search for known drug resistance

conferring mutations in *M. abscessus.* Pubmed was searched with the terms '*Mycobacterium*

abscessus' AND 'clarithromycin' OR 'macrolide' OR 'drug resistance' OR 'antibiotic resistance',

looking for English language articles published up to April 2018. To be included in the final list,

articles had to contain genotyping of coding regions relevant to clarithromycin resistance in *M.*

abscessus in addition to paired drug susceptibility data. Studies looking at both clinical and

non-clinical samples were included. 298 abstracts were screened for relevance and 81 full text

articles were obtained of which 26 met the inclusion criteria (Figure 1).

Sample selection and sequencing

We next sought all available clinical isolates (N = 180) which had undergone whole genome

sequencing by the Public Health laboratory in Birmingham (UK) as part of the routine diagnostic

workflow, and for which paired phenotypic data were also available. We supplemented this with

23 isolates for which the same data was available from a WGS archive at the University of

Oxford. Isolates were collected between May 2014 and January 2017 and no prior selection

according to site of isolation nor whether confirmed M. abscessus complex disease by

guidelines was made. Clinical samples were cultured in liquid mycobacterial growth indicator

tubes from which an aliquot was removed to be prepared for WGS as previously described(17).

Libraries for Illumina Miseq sequencing were prepared using the Illumina Nextera XT protocol

with manual library normalisation. Samples where batched 12 to 16 per flow cell and paired end

sequencing was performed with the MiSeq reagent kit V2. Bioinformatics was performed using

the PHE bioinformatics pipeline as previously described(12, 17). Briefly, reads were mapped to

the *Maba* reference genome (NC_010397.1) with Stampy v1.22 and variants called using

Samtools v0.1.18 (Only variants with >= 5 high-quality reads, mean quality per base >= 25 and

> 90% high-quality bases were retained as variants; heterozygous variants with >10% minor

variant were not retained). A self-self blast approach was used to mask repetitive regions. Sub-

species were identified by computing maximum likelihood (ML) phylogenetic trees incorporating

published representative isolates from each subspecies. A whole genome SNP alignment was

used as input to IQ-TREE OMP version 1.5.5 using a generalised time reversible model. The

erm(41) and rplV genes were manually inspected for insertions/deletions from aligned fasta files

using Seaview version 4.6.2. All newly sequenced data has been uploaded to NCBI under

project accession number PRJNA420644.

Drug susceptibility testing

Phenotypic drug susceptibility testing (DST) was performed at the Public Health England (PHE)

National Mycobacterial reference service laboratory in London. DST was performed using

Standard Broth Microdilution method using 96-well microtitre plates (Sensititre) and prepared as

previously described(18). Plates were read at day three post-inoculation, and if poor growth

again at day 5, according to Clinical and Laboratory Standards Institute (CLSI) guidelines (19).

Isolates are classed as either susceptible (S MIC ≦ 2 µg/ml), resistant (R ≧ 8 µg/ml) or

intermediate (I), according to CLSI breakpoints and if deemed Clarithromycin sensitive the

plates are re-incubated and read again at days 7, 14 and 21.

Genotypic prediction of clarithromycin susceptibility

We used BioPython software to extract base calls from whole genome sequence FASTA files,

comparing these to a list of genomic loci which our literature search indicated were associated

with clarithromycin resistance (table 2). We then predicted phenotypes algorithmically (Figure

2). A resistant phenotype was predicted where any mutations were present at *rrl* positions 2058

or 2059, or where the less well characterized *rrl*_A2057G or *rrl*_A2082C or *rrl*_G2069A

mutations were seen. In the absence of these mutations, susceptibility was predicted where an

isolate had a truncated *erm(41)* gene or a C nucleotide at position 28 in *erm(41)*. Inducible

resistance was predicted where a wild type call (T) was present at position 28 in *erm41*

(meaning that either susceptible or resistant could be correct calls depending on the temporal

relationship between antibiotic administration and sampling time). However, if an *erm41*_C19T

mutation was also present, susceptibility was predicted instead of inducible resistance. In cases

where there was a null call at *rrl* 2058/2059, we did not make a prediction but the number of

resistance and susceptible alleles at those gene positions were inspected and compared to the

phenotype (Figure 2). Statistics quoted were calculated using R Studio v1.1.383.

Search for novel resistance conferring mutations

We attempted to characterise new resistance mutations within genes linked to drug resistance

from the literature search. To maximise power for discovering new potential resistance

mutations, we included all genomes available to us . All mutations in these genes or their

promoter regions were extracted using Python software from variant call files. Phylogenetic

SNPs were identified by considering each subspecies in turn, assumed to be benign and

excluded from further analysis.

We considered variants at the level of SNPs in promoter regions or rRNA and amino acid

changes in coding regions.   A mutation was characterised as causing resistance if it occurred

as the only mutation in the relevant region in a resistant isolate or if it was always associated

with resistance when it was seen and did not co-occur with any other mutations known to cause

resistance. Mutations were characterised as consistent with susceptibility ('benign') if all isolates

were susceptible when the mutation occurred alone or if it occurred only in susceptible isolates.

This methodology is similar to one previously published by our group(12). We assumed no prior

knowledge in this section of the analysis and the identification of known resistance SNPs was

used as an internal validation of our approach.

Results

We studied 143 *Maba*, 20 *Mabb* and 40 *Mabm* genomes. Genotypic predictions were made on

the basis of mutations identified by the literature search. All relevant mutations identified were

contained in the genes *rrl* and *erm(41)* (figure 2 and table 2). The genes rplV, whiB7 and rpld

were also considered of potential interest and were additionally searched for variants.

Genotypic predictions

Inducible resistance was predicted in 91 isolates, meaning that these could be either

susceptible or resistant on phenotypic testing. Of these, 65/91 (71%) were reported as

phenotypically resistant. After excluding isolates for which no prediction could be made due to

missing data in key genes (n = 38) as well as those were we made a prediction of inducibly

resistant (n = 91), sensitivity was 76% (95% CI 53 - 92 %) and specificity was 100%. The very

major error rate (WGS prediction sensitive, phenotype resistant) was 3.0% (95% CI 0.4 - 5.6%)

and the major error rate (phenotype susceptible, WGS prediction resistant) was 0%. Positive

predictive value was 100% and negative predictive value was 90.7% (95% CI 82.0 - 95.5%) (Table

3).

Clarithromycin resistance in the subspecies

81/143 *Maba* were resistant, 58 sensitive and 4 intermediate. For *Mabb* 18/20 were resistant

and for *Mabm* 19/40 were resistant (table 1).   There was one *Mabm* isolate carrying a full

length *erm(41)* gene which was phenotypically resistant to clarithromycin. This was not

unexpected from a genotypic perspective as it harboured a wild type thymine nucleotide at

position 28 *erm(41)*, associated with inducible resistance.

Mechanisms of resistance

The negative predictive value of a truncated *erm(41)* gene for clarithromycin susceptibility was

53% (21/39 - the was one *Mabm* isolate with a full length erm(41). In 9/18 instances, resistance

in the presence of a truncated *erm(41)* could be explained by a mutation in position 2058 or

2059 in *rrl*. No coverage at all was seen at these positions for 4/18 isolates, whilst coverage was

inadequate to determine a consensus nucleotide call in 2/18. In the later two cases, minor

alleles were however present that could potentially explain the resistant phenotypes. No

genomic explanation could be identified for the remaining three discordant isolates (table 1).

All isolates which had any mutation of positions 2057, 2058 or 2059 in *rrl* were resistant to

clarithromycin (15/203 (7.4%)). Such a mutation was found in 2 *Mabb*, 8 *Mabm* and 5 *Maba*

isolates. We did not observe any isolates with an *rrl* mutation which also harboured a T28C

mutation in *erm(41)*. Where this occurred in isolates reported in the literature, they were always

resistant.

Of 36 isolates with a T28C mutation in *erm(41)* and no other relevant mutations, 83% (30) were

susceptible to clarithromycin, 11% (4) had intermediate susceptibility and 6% (2) were resistant.

This mutation was exclusively found in *Maba* isolates. We did not identify any drug resistance

associated mutations in any of these intermediate or resistant isolates. Across all three

subspecies, of 91 isolates with the T28_erm41 call associated with inducible resistance (and no

other relevant mutation), 71% (65) were resistant and 29% (26) susceptible.

De novo search for resistance determining mutations

The search for potential novel resistance determining mutations for clarithromycin revealed 13

SNPs of interest (table 4). Of these, five have previously been described in the literature.

There were additionally four SNPs (*rrl*_A2746T, *rrl*_G836A, *rrl*_T2674G and *rrl*_T636C) which

were only ever seen in resistant isolates but always co-occurred with known resistance

determining SNPs. There was one phenotypically resistant isolate which harboured 18 novel

SNPs. On performing a nucleotide BLAST of a 120 base region encompassing all of these

SNPs, there was a 99% (E 2 x10[-53]) match with *Streptococcal spp.* This therefore likely

represents sample contamination with flora from the nasopharynx. No new resistance

associated variants were discovered in rplV, rpld or whiB7.

Discussion

We conducted a systematic review of drug resistance determining mutations for clarithromycin

in *M. abscessus* and used the results to make genotypic predictions. The sensitivity of this

approach was 76% and the positive predictive value 100%. The prevalence of resistance

amongst our collection of isolates was high compared to that which has been reported

elsewhere(9, 20–22).

These results show that for clarithromycin, drug resistance can be predicted from WGS data as

it has been previously through targeted PCR and line probe assays such as the GenoType

NTM-DR. Assessment of the genotype of *erm(41)* with molecular diagnostics allows prediction

of its functional status which has been thought to correlate to treatment outcome(10). Similarly,

as the absence of a functional *erm(41)* gene has been associated with good therapeutic

outcomes its molecular detection ought to be beneficial to patients(9), although in our study this

alone was not an adequate predictor of in vitro resistance. Furthermore there exist samples

with a genotypic prediction of inducible resistance that show early time-point phenotypic

resistance(23). With DST alone, patients with such isolates would not be treated with a

macrolide however it is unclear whether this is the optimal strategy. Discriminating isolates with

such discordance is an important step towards answering this question.

In addition to mutations identified in the literature search, we also managed to identify mutations

that plausibly explain some of the drug resistance we failed to predict. However, these

mutations will require validation against an independent data set. Using routinely collected

diagnostic data to improve our understanding of the molecular determinants of drug resistance

is a key advantage WGS has over line-probe assays or PCR. The eight previously undescribed

mutations we report in this work could be of clinical importance because they all occur in

samples which the existing literature predicts to be inducibly resistant. As BTS guidelines

recommend that patients with such isolates should be given a macrolide, it is important to

determine further whether these SNPs are true resistance-determinants, and whether macrolide

therapy should be avoided in their presence.

Previous authors have suggested that it is clinically useful to discriminate between subspecies,

(9) as *Mabm* is typically associated with durable susceptibility to clarithromycin and *Mabb* and

Maba with inducible resistance (unless the T28C mutation is present). We found identifying

sub-species alone to be an inadequate predictor of *in vitro* clarithromycin phenotype. There

were three *Mabm* isolates in our collection that were resistant to clarithromycin and had no

mutations known to be relevant. Mougari and colleagues found that in 39/40 *Mabm* selected for

clarithromycin resistance, this could be explained by an rrl mutation at positions 2058/2059 with

a further sample containing an rplV insertion(24). All of our isolates contained this 'insertion'

(also present in the NC_010397.1 reference) which was associated with susceptibility to

clarithromycin except in the presence of a relevant rrl mutation.

In keeping with previous reports, we identified an isolate of *Mabm* with a full length *erm(41)* and

a thymine nucleotide at position 28(25). This likely represents recombination between the

subspecies. A recent study showed the Hain GenoType NTM-DR line probe assay incorrectly

predicted subspecies in 8% of samples, presumably because it lacks the whole genome

resolution provided by sequencing and is vulnerable to between species recombination(26).

Despite analysing all mutations occurring in *erm(41)* and *rrl* for the full collection of genomes,

we were unable to predict all clarithromycin resistance. This may be because there are other

genes implicated or due to unreliable DST results. Future work should aim to select discordant

genotypes and identify additional infrequently occurring genetic loci implicated in clarithromycin

resistance, for example by using genome wide association approaches. All of the new

clarithromycin resistance mutations we discovered occurred in isolates which we originally

predicted to be inducibly resistant. It could be that some or all of these SNPs are compensatory

mutations which act to reduce a fitness cost of the expression of *erm*, which has been

experimentally demonstrated in other bacteria(27). There were four SNPs which only occurred

in resistant samples but were always seen with a known drug resistance causing SNP, possibly

also representing compensatory mutations.

Key weaknesses of our study include that we were unable establish a temporal relationship

between antibiotic prescribing and inducible phenotypic resistance as we did not have the

relevant ethics approval to link to patient records. If for example, any SNPs on our list of novel

mutations were observed in isolates from patients who had never previously had macrolide

therapy, it would be much more likely that they were genuine resistance conferring rather than

compensatory mutations. In addition it is possible that some of the genomes were same patient

replicates over a number of months/years, although this may have also diversified the range of

mutations observed. We chose to include all available samples to maximise detection of low

frequency resistance determining SNPs meaning there was no validation set available. Our list

of novel resistance determining SNPs will therefore require validation on an independent

dataset before being applied to the clinical setting.

In summary, WGS can detect all of the same mutations as the Hain NTM-DR line probe assay,

allows more accurate identification of sub-species and demonstrates probable resistance

determining SNPs in regions the line probe cannot detect which if further validated may change

management. In addition, identification of subspecies alone inadequately predicts macrolide

resistance in *M. abscessus*. As more paired genome/DST data becomes available in the near

future, and we learn more about the molecular determinants of drug resistance, it is likely that

sensitivity of WGS resistance prediction will improve. Given that WGS data is already being

produced in the UK for the purposes of molecular epidemiology, it would now be possible to

phase out existing molecular tests and replicating their results *in silico*, from WGS data at no

additional cost.

Transparency declaration

The authors have no conflicts of interest to declare.

Funding

No specific funding was acquired for this work. All sequencing data was created as part of the

routine NHS clinical diagnostic service.

References

1. Prevots DR, Shaw PA, Strickland D, Jackson LA, Raebel MA, Blosky MA, Montes de Oca

R, Shea YR, Seitz AE, Holland SM, Olivier KN. 2010. Nontuberculous mycobacterial lung

disease prevalence at four integrated health care delivery systems. Am J Respir Crit Care

Med 182:970–976.

2. Esther CR Jr, Esserman DA, Gilligan P, Kerr A, Noone PG. 2010. Chronic Mycobacterium

abscessus infection and lung function decline in cystic fibrosis. J Cyst Fibros 9:117–123.

3. Bryant JM, Grogono DM, Greaves D, Foweraker J, Roddick I, Inns T, Reacher M, Haworth

CS, Curran MD, Harris SR, Peacock SJ, Parkhill J, Floto RA. 2013. Whole-genome

sequencing to identify transmission of Mycobacterium abscessus between patients with

cystic fibrosis: a retrospective cohort study. Lancet 381:1551–1560.

4.   Bryant JM, Grogono DM, Rodriguez-Rincon D, Everall I, Brown KP, Moreno P, Verma D,

Hill E, Drijkoningen J, Gilligan P, Esther CR, Noone PG, Giddings O, Bell SC, Thomson R,

Wainwright CE, Coulter C, Pandey S, Wood ME, Stockwell RE, Ramsay KA, Sherrard LJ,

Kidd TJ, Jabbour N, Johnson GR, Knibbs LD, Morawska L, Sly PD, Jones A, Bilton D,

Laurenson I, Ruddy M, Bourke S, Bowler IC, Chapman SJ, Clayton A, Cullen M, Daniels T,

Dempsey O, Denton M, Desai M, Drew RJ, Edenborough F, Evans J, Folb J, Humphrey H,

Isalska B, Jensen-Fangel S, Jönsson B, Jones AM, Katzenstein TL, Lillebaek T,

MacGregor G, Mayell S, Millar M, Modha D, Nash EF, O'Brien C, O'Brien D, Ohri C, Pao

CS, Peckham D, Perrin F, Perry A, Pressler T, Prtak L, Qvist T, Robb A, Rodgers H,

Schaffer K, Shafi N, van Ingen J, Walshaw M, Watson D, West N, Whitehouse J, Haworth

CS, Harris SR, Ordway D, Parkhill J, Floto RA. 2016. Emergence and spread of a human-

transmissible multidrug-resistant nontuberculous mycobacterium. Science 354:751–757.

5.   Harris KA, Underwood A, Kenna DTD, Brooks A, Kavaliunaite E, Kapatai G, Tewolde R,

Aurora P, Dixon G. 2014. Whole-genome sequencing and epidemiological analysis do not

provide evidence for cross-transmission of Mycobacterium abscessus in a cohort of

pediatric cystic fibrosis patients. Clin Infect Dis 60:1007–1016.

6.   Tortoli E, Kohl TA, Trovato A, Baldan R, Campana S, Cariani L, Colombo C, Costa D,

Cristadoro S, Di Serio MC, Manca A, Pizzamiglio G, Rancoita PMV, Rossolini GM, Taccetti

G, Teri A, Niemann S, Cirillo DM. 2017. Mycobacterium abscessus in patients with cystic

fibrosis: low impact of inter-human transmission in Italy. Eur Respir J 50.

7.   Magee JG, Ward AC. 2015. MycobacteriumBergey's Manual of Systematics of Archaea

and Bacteria. John Wiley & Sons, Ltd.

8.   Nessar R, Cambau E, Reyrat JM, Murray A, Gicquel B. 2012. Mycobacterium abscessus: a
new antibiotic nightmare. J Antimicrob Chemother 67:810–818.

9.   Koh W-J, Jeon K, Lee NY, Kim B-J, Kook Y-H, Lee S-H, Park YK, Kim CK, Shin SJ, Huitt
GA, Daley CL, Kwon OJ. 2011. Clinical significance of differentiation of Mycobacterium
massiliense from Mycobacterium abscessus. Am J Respir Crit Care Med 183:405–410.

10.   Haworth CS, Banks J, Capstick T, Fisher AJ, Gorsuch T, Laurenson IF, Leitch A, Loebinger
MR, Milburn HJ, Nightingale M, Ormerod P, Shingadia D, Smith D, Whitehead N, Wilson R,
Andres Floto R. 2017. British Thoracic Society Guideline for the management of non-
tuberculous mycobacterial pulmonary disease (NTM-PD). BMJ Open Respiratory Research
4:e000242.

11.   Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM,
Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S, von Reyn CF, Wallace
RJ Jr, Winthrop K, ATS Mycobacterial Diseases Subcommittee, American Thoracic
Society, Infectious Disease Society of America. 2007. An official ATS/IDSA statement:
diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. Am J
Respir Crit Care Med 175:367–416.

12.   Walker TM, Kohl TA, Omar SV, Hedge J, Del Ojo Elias C, Bradley P, Iqbal Z, Feuerriegel
S, Niehaus KE, Wilson DJ, Clifton DA, Kapatai G, Ip CLC, Bowden R, Drobniewski FA,
Allix-Béguec C, Gaudin C, Parkhill J, Diel R, Supply P, Crook DW, Smith EG, Walker AS,
Ismail N, Niemann S, Peto TEA, Modernizing Medical Microbiology (MMM) Informatics
Group. 2015. Whole-genome sequencing for prediction of Mycobacterium tuberculosis drug
susceptibility and resistance: a retrospective cohort study. Lancet Infect Dis 15:1193–1202.

13.   Eyre DW, De Silva D, Cole K, Peters J, Cole MJ, Grad YH, Demczuk W, Martin I, Mulvey

MR, Crook DW, Walker AS, Peto TEA, Paul J. 2017. WGS to predict antibiotic MICs for

Neisseria gonorrhoeae. J Antimicrob Chemother 72:1937–1947.

14. Gordon NC, Price JR, Cole K, Everitt R, Morgan M, Finney J, Kearns AM, Pichon B, Young

B, Wilson DJ, Llewelyn MJ, Paul J, Peto TEA, Crook DW, Walker AS, Golubchik T. 2014.

Prediction of Staphylococcus aureus antimicrobial resistance by whole-genome

sequencing. J Clin Microbiol 52:1182–1191.

15. Jeon K, Kwon OJ, Lee NY, Kim B-J, Kook Y-H, Lee S-H, Park YK, Kim CK, Koh W-J. 2009.

Antibiotic treatment of Mycobacterium abscessus lung disease: a retrospective analysis of

65 patients. Am J Respir Crit Care Med 180:896–902.

16. Choi H, Kim S-Y, Kim DH, Huh HJ, Ki C-S, Lee NY, Lee S-H, Shin S, Shin SJ, Daley CL,

Koh W-J. 2017. Clinical Characteristics and Treatment Outcomes of Patients with Acquired

Macrolide-Resistant Mycobacterium abscessus Lung Disease. Antimicrob Agents

Chemother 61.

17. Votintseva AA, Pankhurst LJ, Anson LW, Morgan MR, Gascoyne-Binzi D, Walker TM,

Quan TP, Wyllie DH, Del Ojo Elias C, Wilcox M, Walker AS, Peto TEA, Crook DW. 2015.

Mycobacterial DNA extraction for whole-genome sequencing from early positive liquid

(MGIT) cultures. J Clin Microbiol 53:1137–1143.

18. Broda A, Jebbari H, Beaton K, Mitchell S, Drobniewski F. 2013. Comparative drug

resistance of Mycobacterium abscessus and M. chelonae isolates from patients with and

without cystic fibrosis in the United Kingdom. J Clin Microbiol 51:217–223.

19. Committee for Clinical Laboratory Standards N. 2000. Susceptibility Testing Mycobacteria,

Nocardia, and Other Aerobic Actinomycetes: Tentative Standard.

20. Hatakeyama S, Ohama Y, Okazaki M, Nukui Y, Moriya K. 2017. Antimicrobial susceptibility

testing of rapidly growing mycobacteria isolated in Japan. BMC Infect Dis 17:197.

21. Li YM, Tong XL, Xu HT, Ju Y, Cai M, Wang C. 2016. Prevalence and Antimicrobial

Susceptibility of Mycobacterium abscessus in a General Hospital, China. Biomed Environ

Sci 29:85–90.

22. Cowman S, Burns K, Benson S, Wilson R, Loebinger MR. 2016. The antimicrobial

susceptibility of non-tuberculous mycobacteria. J Infect 72:324–331.

23. Bastian S, Veziris N, Roux A-L, Brossier F, Gaillard J-L, Jarlier V, Cambau E. 2011.

Assessment of clarithromycin susceptibility in strains belonging to the Mycobacterium

abscessus group by erm(41) and rrl sequencing. Antimicrob Agents Chemother 55:775–

781.

24. Mougari F, Bouziane F, Crockett F, Nessar R, Chau F, Veziris N, Sapriel G, Raskine L,

Cambau E. 2017. Selection of Resistance to Clarithromycin in Mycobacterium abscessus

Subspecies. Antimicrob Agents Chemother 61.

25. Shallom SJ, Moura NS, Olivier KN, Sampaio EP, Holland SM, Zelazny AM. 2015. New

Real-Time PCR Assays for Detection of Inducible and Acquired Clarithromycin Resistance

in the Mycobacterium abscessus Group. J Clin Microbiol 53:3430–3437.

26. Kehrmann J, Kurt N, Rueger K, Bange F-C, Buer J. 2016. GenoType NTM-DR for

Identifying Mycobacterium abscessus Subspecies and Determining Molecular Resistance.

J Clin Microbiol 54:1653–1655.

27. Gupta P, Sothiselvam S, Vázquez-Laslop N, Mankin AS. 2013. Deregulation of translation

due to post-transcriptional modification of rRNA explains why *erm* genes are inducible. Nat

Commun 4:ncomms2984.

28. Maurer FP, Castelberg C, Quiblier C, Böttger EC, Somoskövi A. 2014. Erm(41)-dependent

inducible resistance to azithromycin and clarithromycin in clinical isolates of Mycobacterium

abscessus. J Antimicrob Chemother 69:1559–1563.

29. Hanson KE, Slechta ES, Muir H, Barker AP. 2014. Rapid molecular detection of inducible

macrolide resistance in Mycobacterium chelonae and M. abscessus strains: a replacement

for 14-day susceptibility testing? J Clin Microbiol 52:1705–1707.

30. Brown-Elliott BA, Vasireddy S, Vasireddy R, Iakhiaeva E, Howard ST, Nash K, Parodi N,

Strong A, Gee M, Smith T, Wallace RJ Jr. 2015. Utility of sequencing the erm(41) gene in

isolates of Mycobacterium abscessus subsp. abscessus with low and intermediate

clarithromycin MICs. J Clin Microbiol 53:1211–1215.

31. Nie W, Duan H, Huang H, Lu Y, Chu N. 2015. Species Identification and Clarithromycin

Susceptibility Testing of 278 Clinical Nontuberculosis Mycobacteria Isolates. Biomed Res

Int 2015:506598.

32. Ramírez A, de Waard JH, Araque M. 2015. Molecular mechanisms of clarithromycin

resistance in Mycobacterium abscessus complex clinical isolates from Venezuela. J Glob

Antimicrob Resist 3:205–209.

33. Rubio M, March F, Garrigó M, Moreno C, Español M, Coll P. 2015. Inducible and Acquired

Clarithromycin Resistance in the Mycobacterium abscessus Complex. PLoS One

10:e0140166.

34. Chua KYL, Bustamante A, Jelfs P, Chen SC-A, Sintchenko V. 2015. Antibiotic susceptibility

of diverse Mycobacterium abscessus complex strains in New South Wales, Australia.

Pathology 47:678–682.

35. Kim J, Sung H, Park J-S, Choi S-H, Shim T-S, Kim M-N. 2016. Subspecies distribution and

446   macrolide and fluoroquinolone resistance genetics of Mycobacterium abscessus in Korea.

447   Int J Tuberc Lung Dis 20:109–114.

448  36. Jeong SH, Kim S-Y, Huh HJ, Ki C-S, Lee NY, Kang C-I, Chung DR, Peck KR, Shin SJ, Koh

449   W-J. 2017. Mycobacteriological characteristics and treatment outcomes in extrapulmonary

450   Mycobacterium abscessus complex infections. Int J Infect Dis 60:49–56.

451  37. Carvalho NFG de, Pavan F, Sato DN, Leite CQF, Arbeit RD, Chimara E. 2018. Genetic

452   correlates of clarithromycin susceptibility among isolates of the Mycobacterium abscessus

453   group and the potential clinical applicability of a PCR-based analysis of erm(41). J

454   Antimicrob Chemother 73:862–866.

455  38. Chew KL, Cheng JWS, Hudaa Osman N, Lin RTP, Teo JWP. 2017. Predominance of

456   clarithromycin-susceptible Mycobacterium massiliense subspecies: Characterization of the

457   Mycobacterium abscessus complex at a tertiary acute care hospital. J Med Microbiol

458   66:1443–1447.

459  39. Koh W-J, Jeong B-H, Kim S-Y, Jeon K, Park KU, Jhun BW, Lee H, Park HY, Kim DH, Huh

460   HJ, Ki C-S, Lee NY, Kim HK, Choi YS, Kim J, Lee S-H, Kim CK, Shin SJ, Daley CL, Kim H,

461   Kwon OJ. 2017. Mycobacterial Characteristics and Treatment Outcomes in Mycobacterium

462   abscessus Lung Disease. Clin Infect Dis 64:309–316.

463  40. Zhu YC, Mitchell KK, Nazarian EJ, Escuyer VE, Musser KA. 2015. Rapid prediction of

464   inducible clarithromycin resistance in Mycobacterium abscessus. Mol Cell Probes 29:514–

465   516.

466  41. Wallace RJ Jr, Meier A, Brown BA, Zhang Y, Sander P, Onyi GO, Böttger EC. 1996.

467   Genetic basis for clarithromycin resistance among isolates of Mycobacterium chelonae and

468   Mycobacterium abscessus. Antimicrob Agents Chemother 40:1676–1681.

469   42. Liu W, Li B, Chu H, Zhang Z, Luo L, Ma W, Yang S, Guo Q. 2017. Rapid detection of

470   mutations in erm(41) and rrl associated with clarithromycin resistance in Mycobacterium

471   abscessus complex by denaturing gradient gel electrophoresis. J Microbiol Methods

472   143:87–93.

473   43. Kim S-Y, Shin SJ, Jeong B-H, Koh W-J. 2016. Successful antibiotic treatment of pulmonary

474   disease caused by Mycobacterium abscessus subsp. abscessus with C-to-T mutation at

475   position 19 in erm(41) gene: case report. BMC Infect Dis 16:207.

483 **Figure 1:** Flow diagram showing stages of the systematic literature search

486   **Figure 2:** decision algorithm for predicting drug resistance in *M. abscessus* based on the

487   literature search with numbers of isolates meeting each predictive criterion shown. Bracketed

488   numbers represent (N resistant / N sensitive). * 4 isolates had intermediate susceptibility

	erm41 pos. 28	erm41 length	erm41 pos. 19	rrl. pos 2269	rrl pos. 2270	rrl pos. 2271	N	Phenotype	Prediction
Maba	T	full	C	A	C	A	4	1R	R
	T	full	C	A	T	A	1	1R	R
	C	full	C	A	A	A	36	4I, 2R, 30S	S
	T	full	C	A	A	A	77	53R, 24S	IR
	Excluded due to inadequate coverage over rrl 2270-2271						25		
Mbol	T	full	C	G	A	A	1	1R	R
	T	full	C	A	G	A	1	1R	R
	T	full	C	A	A	A	13	11R, 2S	IR
	Excluded due to inadequate coverage over rrl 2270-2271						5		
Mabm	T	truncated	C	A	C	A	2	2R	R
	T	truncated	C	A	G	A	2	2R	R
	T	truncated	C	A	A	G	5	5R	R
	T	truncated	C	A	A	A	22	19S, 3R	S
	T	full	C	A	A	A	1	1R	IR
	Excluded due to inadequate coverage over rrl 2270-2271						8		

Table 1 - summary of genotypes and corresponding clarithromycin phenotypes for the 203 isolates. Pos - M. abscessus numbering position in gene, prediction - genotypic prediction using the algorithm shown in Figure 2, Maba - *M abscessus absecssus*, Mbol - *M. abscessus bolletii*, Mabm - *M. abscessus massiliense*, N – total number of isolates with genotype, S – Sensitive, R – Resistant, IR – Inducible Resistance

erm(41) length	erm(41) position 28	rrl position 2058	rrl position 2059	Other	Phenotype
Full	T	A	A		Inducible resistance(3, 23–26, 28–38) Sensitive(34)
Truncated		A	A		Sensitive(9, 24–26, 29, 32, 34–39)
Full	C	A	A		Sensitive(24–26, 28–34, 36–38, 40)
Full or Truncated	C or T	G	A		Resistant(3, 21, 23–26, 28, 33, 37–42)
Full or truncated	C or T	C	A		Resistant(3, 24–26, 28, 39, 42)
Full or	T	T	A		Resistant(23,

truncated					24, 26)
Full		A	C		Resistant(23, 24, 41, 42)
Full or truncated	T or C	A	G		Resistant(21, 24–26, 28, 33, 38, 41)
Truncated		A	T		Resistant(23)
Full	T	A	A	C19T erm(41)	Sensitive(43)
Truncated		A	A	A2057G rrl	Resistant(33)
Full	Unknown	Unknown	Unknown	A2082C rrl + G2069C rrl	Resistant(42)

Table 2: Resistance determining mutations for clarithromycin identified in the literature search

	In vitro phenotype		
Genomic Prediction	Sensitive	Resistant	Intermediate
No Prediction*	6	32	0
Inducible resistance	26	65	0

Resistant	0	16	0
Sensitive	49	5	4
Sensitivity	76.19% (95% CI 52.8 - 91.8%)		
Specificity	100%		
Positive predictive value	100%		
Negative predictive value	90.7% (95% CI 82.0 - 95.5%)		

Table 3 - WGS Predictions vs DST phenotype for clarithromycin.

Sensitivity/Specificity/PPV/NPV are calculated excluding isolates predicted to be inducibly resistant and those where no prediction was made due to inadequate coverage at key positions.

Position	Nucleotide/Amino acid change	Rule met
rrl 2039	A > G	1
rrl 1401	T > C	2
rrl 371	T > C	2
rrl 795	G > A	1
rrl 2270*	A > C	1
rrl 2270*	A > G	2

rrl 2271*	A > G	2
rrl 2270 *	A > T	2
erm(41) 131	A > V	2
rrl 2279	G > A	2
rrl 2269*	A > G	2
erm (41) -31**	A > T	2
rrl 1932	A > G	2

Table 4: Mutations (both novel and previously described) detected during *de novo* search for resistance determining SNPs. Rule 1 = occurs as only SNP in relevant regions in resistant isolate, rule 2 = all samples resistant when SNP occurs, never seen in sensitive isolate. All numbering is relative to M. abscessus. * mutation already described in literature - *M. abscessus rrl* numbering 2270/2271 is *E. coli* numbering 2058/2059. ** mutation in erm(41) promoter region, 31 bases upstream of start of coding region.