Title: BMP Signaling Downstream of the Highwire E3 Ligase Sensitizes Nociceptors

Authors

Ken Honjo[3] and W. Daniel Tracey Jr[1,2,]

1 Department of Biology 2 Gill Center for Biomolecular Sciences, Indiana University, Bloomington

Indiana, 47405. 3 Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai,

Tsukuba, Ibaraki 305-8572, Japan.

Correspondence:

Ken Honjo (honjo.ken.ga@u.tsukuba.ac.jp), Dan Tracey (dtracey@indiana.edu)

Short Title: The roles of Hiw and BMP signaling in regulating nociception

Abstract (max 300 words)

A comprehensive understanding of the molecular machinery important for nociception is essential to

improving the treatment of pain. Here, we show that the BMP signaling pathway regulates

nociception downstream of the E3 ubiquitin ligase *highwire* (*hiw*). *Hiw* loss of function in nociceptors

caused antagonistic and pleiotropic phenotypes with simultaneous insensitivity to noxious heat but

sensitized responses to optogenetic activation of nociceptors. Thus, *hiw* functions to both positively

and negatively regulate nociceptors. We find that a sensory transduction-independent sensitization

pathway was associated with BMP signaling. BMP signaling in nociceptors was up-regulated in *hiw*

mutants, and nociceptor-specific expression of *hiw* rescued all nociception phenotypes including the

increased BMP signaling. Blocking the transcriptional output of the BMP pathway with dominant

negative Mad suppressed nociceptive hypersensitivity that was induced by interfering with *hiw*. The

up-regulated BMP signaling phenotype in *hiw* genetic mutants could not be suppressed by mutation

in *wallenda* suggesting that *hiw* regulates BMP in nociceptors via a *wallenda* independent pathway.

In a newly established Ca^{2+} imaging preparation, we observed that up-regulated BMP signaling

caused a significantly enhanced Ca^{2+} signal in the axon terminals of nociceptors that were stimulated

by noxious heat. This response likely accounts for the nociceptive hypersensitivity induced by

elevated BMP signaling in nociceptors. Finally, we showed that acute activation of BMP signaling in

nociceptors was sufficient to sensitize nociceptive responses to optogenetically-triggered nociceptor

activation without altering nociceptor morphology. Overall, this study demonstrates the previously

unrevealed roles of the Hiw-BMP pathway in the regulation of nociception and provides the first

direct evidence that up-regulated BMP signaling physiologically sensitizes responses of nociceptors

and nociception behaviors.

Author Summary (150-200 words)

Although pain is a universally experienced sensation that has a significant impact on human lives

and society, the molecular mechanisms of pain remain poorly understood. Elucidating these

mechanisms is particularly important to gaining insight into the clinical development of currently

incurable chronic pain diseases. Taking an advantage of the powerful genetic model organism

Drosophila melanogaster (fruit flies), we unveil the Highwire-BMP signaling pathway as a novel

molecular pathway that regulates the sensitivity of nociceptive sensory neurons. Highwire and the

molecular components of the BMP signaling pathway are known to be widely conserved among

animal phyla, from nematode worms to humans. Since abnormal sensitivity of nociceptive sensory

neurons can play a critical role in the development of chronic pain conditions, a deeper

understanding of the regulation of nociceptor sensitivity has the potential to advance effective

therapeutic strategies to treat difficult pain conditions.

Introduction

In spite of its clear medical importance, the molecular mechanisms of pain signaling remain poorly understood. Pain pathways in large part depend on sensory input from specialized sensory neurons called nociceptors (1). Since the activation of nociceptors leads to pain sensation and the sensitization of nociceptors is thought to be a major contributor of pain pathogenesis, understanding the molecular mechanisms controlling nociceptor function is essential for improving the treatment of pain (2).

Drosophila melanogaster is a powerful model system for neurogenetic studies of nociception. Larval *Drosophila* show stereotyped behavioral responses to potentially tissue-damaging stimuli, such as noxious heat or harsh mechanical stimulation (3). The most unambiguous larval nociception behavior involves a corkscrew like rolling around the long body axis (termed nocifensive escape locomotion (NEL) or simply "rolling"). Since rolling is specifically triggered by noxious stimuli and is clearly separable from normal larval locomotion, the analysis of NEL provides a robust behavioral paradigm to study nociception. Class IV multidendritic (md) neurons are polymodal nociceptors that are necessary for thermal and mechanical nociception in larvae (4). Optogenetic activation of the Class IV neurons is sufficient for triggering NEL(4, 5). Accumulating evidence in studies of fly nociception suggests that the molecular pathways of nociception are conserved between *Drosophila* and mammals (3, 6-15).

To identify genes important for nociceptor function, we recently performed thermal nociception

screens in which we targeted the RNAi knockdown of nociceptor-enriched genes in a

nociceptor-specific manner (16). In this screen, we found that two RNAi lines targeting *highwire (hiw)*

caused driver dependent hypersensitivity in thermal nociception assays (revealed as a rapid

response to a threshold heat stimulus) indicating a potential role for *hiw* as a negative regulator of

nociceptor activity (16). *hiw* is an evolutionally conserved gene encoding an E3 ubiquitin ligase,

whose function has been implicated in various aspects of neuronal development, synaptic function,

and neuronal degeneration (17). However, in contrast, very little is known about the roles of *hiw* in

sensory processing and in controlling behavior. Here, we present additional and more specific

evidence that *hiw* plays an important role in the regulation of behavioral nociception and nociceptor

sensitivity through the bone morphogenetic protein (BMP) pathway.

Results

Highwire regulates the sensitivity of nociceptors

To further investigate the potential function of *hiw* in nociception that was suggested by our

previous study, we tested mutants for a strong loss-of-function allele of *hiw* (hiw^{ND8}) in thermal

nociception assays (18). Unexpectedly, we found that genetic mutants of *hiw* showed insensitivity to

a noxious temperature probe of 42 or 46°C, which was, surprisingly, the opposite of the previously

described *hiw* RNAi phenotype (Fig 1A and data not shown). Similar thermal insensitivity was also

seen with other *hiw* alleles (Fig S1). Although *hiw* is widely expressed in the nervous system (18),

nociceptor-specific restoration of *hiw* expression rescued this insensitivity (Fig 1A), indicating that

hiw function in nociceptors is sufficient for restoration of normal thermal nociception and the relevant

site of action was in nociceptors.

Intrigued by the clear phenotypic distinction between genetic mutants and RNAi animals, we

further dissected the nociception phenotype of *hiw* mutants by employing an optogenetic strategy.

Optical activation of larval nociceptors via the blue light-gated cation channel Channelrhodopsin-2

(ChR2) is sufficient to induce larval NEL (4, 5). Since nociceptor activation by ChR2 circumvents

sensory transduction but still depends on the machinery essential for downstream signaling (Fig 1B),

this technique has been utilized to distinguish genes that are important for primary sensory function

from those that function in downstream aspects of signaling, such as action potential

generation/propagation and/or synaptic transmission (10, 19). Using low intensity blue light (3.8 klux),

which elicits NEL in about 20-30% of control animals expressing ChR2::YFP in nociceptors (Fig 1C),

we found that the hiw^{ND8} mutants had a significantly increased probability to show NEL, indicating

that the mutant for this allele is hypersensitive in response to optogenetic activation of nociceptors

(Fig 1C) even though it was insensitive in thermal nociception assays. Tissue specific rescue

experiments again showed that nociceptor specific expression of *hiw* was sufficient to rescue this

optogenetic hypersensitivity (Fig 1C). Taken together, these findings suggested that *hiw* has multiple,

but dissociable, effects in the regulation of nociceptors. On the one hand*, hiw* regulated a sensory

transduction-dependent function causing insensitivity, but it also regulated a function further

downstream of sensory transduction that caused hypersensitivity. Thus, the hypersensitivity seen in

our earlier RNAi experiments is likely reflective of effects on the latter process.

To further examine *hiw*'s role, we tested the effects of expressing *hiw△RING* in nociceptors.

The *hiw△RING* transcript encodes a mutated form of *hiw* lacking the RING domain that is

responsible for E3 ligase activity (20, 21). This mutated protein has been proposed to function as a

dominant-negative poison subunit in multimeric Hiw E3 ligase complexes. Similar to our original

observations with *hiw RNAi*, expression of *hiw△RING* in nociceptors resulted in significant

hypersensitivity in thermal nociception (Fig 1D and S2). This manipulation also caused

hypersensitive optogenetic nociception responses (Fig 1E). As *hiw* encodes a large protein with

many functional domains, and phenotypes of *hiw* mutants are known to show varied sensitivity to

gene dosage (Wu et al., 2005), the observed similarity between *hiw△RING* overexpression and *hiw*

RNAi is suggestive of dosage-dependent effects of *hiw* in nociceptors. For instance, the dominant

negative approach may lead to an incomplete loss of function for *hiw* that is similar to the effects of

RNAi.

Hiw attenuates BMP signaling in nociceptors

It has been very recently shown that the canonical BMP pathway in nociceptors is required for

nociceptive sensitization after tissue damage in *Drosophila* (22). Since the BMP signaling pathway

has also been proposed to be a downstream pathway regulated by Hiw in motoneurons (23), we

tested whether the BMP signaling pathway is regulated downstream of Hiw in nociceptors. We first

examined the level of phosphorylated Mad (pMad) in nociceptor nuclei by quantitative

immunohistochemistry, which is an established method for evaluating the activation level of

intracellular BMP signaling (24-30). In nociceptor nuclei, *hiw* genetic mutants showed significantly

elevated pMad levels (33%) in comparison to wild-type, even when processed together in the same

staining solution (see also Materials and Methods) (Fig 2 A, B and F). A similarly modest change in

pMad accumulation in motor neuron nuclei is associated with effects on presynaptic function and

morphology at the neuromuscular junction (NMJ) (31, 32). An increased accumulation of pMad in the

nucleus and the cytoplasm was observed in nociceptors expressing *hiw△RING* (Fig 2C and F).

Expression of wild-type *hiw* in nociceptors of *hiw* mutant animals rescued the elevated pMad level

(Fig 2D and F). We also confirmed that our immunohistochemistry successfully detected the

increase of nuclear pMad caused by expressing the constitutively active form of *thick veins* (tkv^{QD}),

which activates the intracellular BMP signaling cascade independently of BMP ligands (33) (Fig 2E

and F). These data together suggest that BMP signaling is negatively regulated downstream of *hiw* in

larval nociceptors. In the larval motoneurons, it is known that pMad signals can be locally detected at

synaptic boutons as well as nuclei (25, 34, 35). However, no detectable pMad signals were observed

at synaptic terminals in larval nociceptors (Fig 2G).

Next we tested whether up-regulated BMP signaling in nociceptors is responsible for the

hypersensitive nociceptive responses caused by *hiw* loss-of-function. mad^{1} encodes a

dominant-negative form of Mad with disrupted DNA-binding ability (36). When mad^1 was expressed

together with $hiw\triangle RING$ in nociceptors, the hypersensitive phenotype that was normally induced by

the expression of $hiw\triangle RING$ alone was not detected (Fig 2H). Since neither expressing mad^1

together with $hiw\triangle RING$ nor expressing mad^1 alone in nociceptors induced insensitivity to noxious

heat (Fig S3), these results indicate that hypersensitive nociception caused by weak *hiw* loss of

function requires an intact BMP signaling pathway that normally operates through mad. This result is

consistent with the elevated pMad observed with *hiw* loss of function as playing a causal role in the

hypersensitive phenotypes.

The MAP kinase kinase kinase (MAPKKK) *wallenda* (*wnd*) is a well-characterized target

substrate of Hiw ligase (17). Hiw negatively regulates the protein level of Wnd, and Hiw-Wnd

interaction is crucial for normal synaptic growth, but not for normal synaptic function in NMJ (30,

37-39). In addition, *hiw* interacts with *wnd* in Class IV neurons in the regulation of dendritic and

axonal morphology (40). In larval motoneurons, it has been suggested that *wnd* is not involved in the

regulation of BMP signaling (30). To test whether *wnd* is involved in the control of BMP signaling

downstream of *hiw* in nociceptors, we examined a genetic interaction between *hiw* and *wnd* in double

mutants. A *wnd* mutation in *hiw* mutant background did not suppress the elevated nuclear pMad level

in nociceptors that we observed in the *hiw* mutant (Fig 3A-D and F), nor did *wnd* single mutants show

altered nuclear pMad accumulation relative to controls (Fig 3E and F). Interestingly, significant

up-regulation of nuclear pMad signal was observed in nociceptors overexpressing *wnd*, but not with

a kinase-dead version of *wnd* (Fig S4). Taken together, these results suggest that elevated nuclear

pMad in *hiw* mutant nociceptors does not depend on the activity of Wnd, although overexpression of

wnd with GAL4/UAS can cause elevated BMP signaling in nociceptors.

To gain insight into which regions of Hiw protein are involved in attenuating BMP signaling in

nociceptors, we performed an expression study of a series of Hiw dominant negatives with various

deletions, as established by Tian et al. (38) (Fig 4A). Expressing HiwNT (N-terminal half of Hiw)

caused an over 200% increase in nuclear pMad signals compared to controls (Fig 4B, C and H).

HiwCT (C-terminal half of Hiw) and Hiw△RCC1 resulted in 99% and 68% increases in nuclear pMad

signals, respectively (Fig 4D, E and H). HiwCT and Hiw△RCC1 also caused marked accumulation of

pMad signals in the cytoplasm of nociceptors (Fig 4D and E), which was also observed with

Hiw△Ring expression (Fig 2C). This cytoplasmic accumulation of pMad signals is unlikely due to

technical variability of immunostaining since the control samples treated in the same staining

solutions with HiwCT or Hiw△RCC1 never developed such accumulations and cells nearby the

nociceptors showed the normal pMad signal. In contrast, Hiw△HindIII and HiwCT1000 (C-terminal

only region of Hiw) did not cause any changes in nuclear pMad signals in nociceptors (Fig 4C, F and

H). Thus, the attenuation of BMP signaling in nociceptors through Hiw appears dependent on

different regions of Hiw from those that have been proposed to be involved in the regulation of NMJ

morphology (Hiw△RCC1, and Hiw△HindIII function as dominant-negative in NMJ morphology while

HiwNT and HiwCT1000 do not (38)). Because both HiwNT and HiwCT, which are largely

non-overlapping N-terminal and C-terminal halves of Hiw, caused increased nuclear pMad signals, it

appears that multiple regions of the Hiw protein must be intact for normal suppression of BMP

signaling in nociceptors.

Elevated BMP signaling in nociceptors induces behavioral nociceptive hypersensitivity

Although a previous study by Follansbee et al. suggests that the canonical BMP signaling pathway in

larval nociceptors is a necessary component for nociceptive sensitization after tissue-damage,

whether up-regulation of BMP signaling in nociceptors is sufficient to sensitize nociception has not

been proven and potential mechanisms leading to sensitization are unknown. Because our data

support the notion that the up-regulation of BMP signaling in nociceptors plays a key role in inducing

sensitized nociception, we tested this hypothesis by examining whether artificial up-regulation of

intracellular BMP signaling in nociceptors is sufficient to induce nociceptive hypersensitivity. In

thermal nociception assays, animals with nociceptor-specific up-regulation of BMP signaling by

expressing tkv^{QD} did exhibit significant hypersensitivity (Fig 5A and S2), and tkv^{QD} expressors also

showed hypersensitive responses in optogenetic nociception. This suggests that elevated BMP

signaling in nociceptors was able to sensitize nociception through a mechanism that was

downstream of sensory transduction (Fig 5B). Although the dendritic structure of nociceptors in tkv^{QD}

overexpressors was not significantly altered (Fig 5C-E), overexpression of tkv^{QD} caused

overextension and overexpansion of nociceptor axon termini (Fig 5F-H). Combined, these data

demonstrate that elevated BMP signaling in nociceptors is sufficient to sensitize thermal and

optogenetic nociception behaviors in addition to axon terminal branching.

Elevated BMP signaling increases Ca^{2+} responses in nociceptor terminals

Since nociceptor-specific up-regulation of BMP signaling sensitizes thermal and optogenetic

nociception at the behavioral level, we next explored whether the up-regulation of intracellular BMP

signaling actually sensitizes physiological responses of nociceptors. To observe neuronal responses

of larval nociceptors to a range of thermal stimuli, we developed a preparation for optical recording

from axon terminals of the nociceptive neurons. We then observed these terminals while we applied

a thermal ramp stimulus was locally applied to the larval body wall (Fig 6A). To monitor Ca^{2+}, the

genetically encoded sensor GCaMP6m was expressed under the control of *ppk-GAL4* (41). In

control animals we observed a steep increase of the GCaMP6m signal in nociceptors when the

ramping temperature reached the 39-47°C temperature range (Fig 6B and D). We found that

nociceptors expressing *tkv^{QD}* showed a significantly greater increase of GCaMP6m signals through

36-50 °C in comparison to those in controls (Fig 6C and D), while basal fluorescence levels of

GCaMP6m (F$_0$) were comparable between the control and *tkv^{QD}*-expressing nociceptors (Fig 6E).

These results suggest that the significantly greater increase of GCaMP6m signals observed in

nociceptors expressing *tkv^{QD}* is due to the greater level of Ca^{2+} influx triggered by the heat ramp

stimulus, and not to unintended transcriptional upregulation of GCaMP6m or to the observed

increase in axonal branching. Thus, elevated BMP signaling in nociceptors results in exaggerated

Ca^{2+} signals at the terminals of nociceptors in response to heat in the noxious range. This conclusion

is consistent with the behavioral nociceptive sensitization induced by the same intracellular

up-regulation of BMP signaling in nociceptors.

Acute up-regulation of BMP signaling induces nociceptive hypersensitivity

Chronic up-regulation of BMP signaling in nociceptors caused sensitization of behavioral nociception

responses of larvae and an increased Ca^{2+} response of nociceptors to noxious heat, but also

expansion of nociceptor terminals. To further separate the physiological and developmental effects of

BMP up-regulation in nociceptors, we performed an acute up-regulation experiment of BMP signaling.

Using the temperature sensitive repressor of GAL4 activity (GAL80^{ts}) (42), we activated expression

of *tkv^{QD}* in larval nociceptors by shifting *ppk-GAL4 UAS-Chr2::YFP tub-GAL80^{ts}* animals to 30°C for

24 hours. We then tested the larvae for sensitized optogenetic nociception. The acute induction of

tkv^{QD} induced hypersensitivity in the optogenetic nocifensive responses and also significantly

increased nuclear pMad levels relative to controls (Fig 7A and B). However, no detectable axonal

overgrowth was induced by acute *tkv^{QD}* expression (Fig 7C and D). Unfortunately, we were not able

to investigate the effects of this manipulation on nociception responses with a 39°C thermal stimulus

because the prolonged incubation at 30°C interfered with 39°C NEL behavior in both controls and

experimental animals (Fig S5). This latter finding indicates that the sensitivity of thermal nociception

in *Drosophila* is modulated by the ambient temperature of the environment. Collectively, these data

demonstrate that acute activation of BMP signaling in nociceptors is sufficient to sensitize larval

nociceptive response in the absence of detectable changes to axonal morphology. Taken together

with our Ca^{2+} imaging results, these data suggest that a physiological, rather than a developmental,

role for BMP signaling in the regulation of nociceptor sensitivity.

Discussion

Identifying novel conserved molecular pathways that regulate nociception in model animals is a

promising strategy for understanding the molecular basis of pain signaling and pain pathogenesis

(43, 44). Using *Drosophila*, we found that both the E3 ligase Hiw and the downstream BMP signaling

pathway play crucial roles in regulating nociceptor sensitivity.

Hiw's complexed roles in regulating nociceptor functions

The data we present in this study suggest that *hiw* may play at least two distinct roles in the

regulation of nociceptor sensitivity. We found that strong loss-of-function mutants of *hiw* showed

insensitivity to noxious heat but hypersensitivity to optogenetic stimulation of nociceptors (Fig 1A and

C). Since expressing wild-type *hiw* in nociceptors of *hiw* mutants rescued both phenotypes, we argue

that *hiw* in nociceptors is responsible for these two ostensibly opposing phenotypes (Fig 1A and C).

We also found that nociceptor-specific expression of *hiw*RNAi or *hiw*△*RING* caused only

hypersensitivity (Fig 1D and E) (16), indicating that the process that governs hypersensitivity is

separable from the cause of insensitivity. As insensitivity was epistatic to hypersensitivity in thermal

nociception assays, we used optogenetics to show that hypersensitivity is present in *hiw* genetic

mutants as well as in previously described RNAi animals. The use of optogenetic stimulation of

neurons allowed us to bypass the endogenous sensory transduction step and to reveal this role. Our

data suggest that *hiw* is a) required for the negative regulation of a neural pathway that is

downstream sensory transduction   and b) required to confer normal sensitivity to noxious heat via

sensory transduction pathways. As strong *hiw* loss of function causes reduced dendritic arbors (40)

while *hiw* RNAi does not (16), it is possible that the reduced dendrite phenotype accounts for the

insensitivity of the strong *hiw* alleles. Consistent with this hypothesis, many mutants that cause

insensitive thermal nociception are associated with a reduction in the dendritic arbor (16). The

phenotypic difference between strong loss-of-function mutants and RNAi or Hiw dominant-negative

animals suggests that insensitive and hypersensitive phenotypes observed in *hiw* mutants have

different sensitivity to the dosage of *hiw*. This has also been seen in the larval motor neuron system

where it has been demonstrated that two different phenotypes of *hiw* in larval NMJ (overgrowth of

synaptic boutons and diminished synaptic function) are separable by their different sensitivity to the

dosage of *hiw* (20).

Our data also suggest that *hiw* may regulate distinct molecular pathways in motor neurons

and in nociceptors. In the larval NMJ, mutations of *hiw* or expression of *hiw△RING* cause a

diminished evoked excitatory junction potential (EJP) due to decreased quantal content in synaptic

vesicles (18, 20, 45). However, this diminished evoked EJP phenotype is apparently opposite to the

hypersensitive nociception phenotype observed in this study. Thus the downstream targets and/or

pathways of Hiw in nociceptors may be distinct from those in motor neurons.

We identified the BMP signaling pathway as an important signaling pathway in nociceptors

that is regulated downstream of *hiw*. In fly motor neurons, it has been proposed that BMP signaling is

a direct target of Hiw ligase (23). However, following a later study reporting that pMad up-regulation

was not detected in motor neuron nuclei in *hiw* mutants (30) and controversy has arisen over this

interaction. We found that nuclear pMad signals were up-regulated in *hiw* genetic mutants, and that

this molecular phenotype was rescued by wild-type *hiw* expression in nociceptors (Fig 2). In addition,

we also detected striking accumulations of pMad in both the nuclei and cytoplasm of nociceptors

expressing Hiw dominant negative proteins (Fig 2 and 4). Finally, using *UAS-mad^1*, we showed that

the Mad-dependent pathway is responsible for the hypersensitive thermal nociception caused by

hiw△RING expression (Fig 2H). Our data therefore supports the idea that the nociceptor BMP

signaling pathway is regulated downstream from *hiw*.

Although we demonstrated that BMP signaling is downstream of *hiw* in nociceptors, we have

yet to determine the precise mechanism for Hiw regulation of BMP signaling. Our genetic analysis

suggests that BMP signaling in nociceptors is regulated independently from the *wnd* pathway (Fig 3).

Wnd is the best characterized target substrate of Hiw in the regulation of NMJ morphology (30, 37-40,

46). Our expression analysis using various *hiw* deletion series showed that the set of *hiw* deletion

constructs that induced up-regulation of BMP signaling in nociceptors was not identical to the set that

induced abnormal synaptic morphology in motoneurons (38). This finding is somewhat consistent

with the existence of a Wnd-independent mechanism in the regulation of BMP signaling in

nociceptors, since the Hiw-Wnd pathway plays a pivotal role in regulating synaptic morphology in

larval NMJ.

Intriguingly, our expression study of the *hiw* deletion series showed that the expression of

HiwNT caused a prominent accumulation of nuclear pMad, while the expression of HiwCT or

Hiw△RCC1 caused accumulation of pMad signals in both the nuclei and cytoplasm in nociceptors

(Fig 4C-E). These data raise the possibility that Hiw is involved in two different mechanisms which

regulate pMad: one pathway affecting nuclear pMad and another for cytoplasmic pMad. Given that

hiw is a large protein with many functional domains for interacting with multiple molecules, the notion

that *hiw* is involved in multiple processes regulating various aspects of neuronal functions in both

motor neurons and nociceptive sensory neurons is reasonable. Further studies are necessary to

reveal the mechanisms of Hiw-dependent regulation of BMP signaling in nociceptors.

Physiological effects of BMP signaling in nociceptor axon terminals

We have presented a new physiological preparation for investigating the calcium levels in nociceptor

terminals with a physiologically relevant noxious heat stimulus. This allowed us to demonstrate that

up-regulation of BMP signaling in nociceptors sensitizes the physiological responses of nociceptors

in response to noxious heat in addition to its effects on behavior (Fig 5 and 6). We also demonstrated

that acute activation of intracellular BMP signaling in nociceptors is sufficient for the nociceptive

sensitization (Fig 7). Although it has been previously reported that BMP signaling in nociceptors is

required for nociceptive sensitization after tissue-injury in *Drosophila* (22), the mechanisms of the

regulation of nociception by BMP signaling was totally unknown. Our study provides the first

evidence that BMP signaling regulates calcium signaling in axon termini of nociceptors.

The BMP signaling pathway plays crucial roles in various developmental processes, such as

embryonic patterning, skeletal development, and the development of neuronal circuits (47, 48). The

roles of BMP signaling in the regulation of neuronal activity has also been extensively investigated in

the larval motoneuron system, where BMP signaling plays crucial roles in the homeostatic regulation

of synaptic morphology and transmission (49, 50). Since BMP signaling is important for synaptic

transmission in larval NMJ, an interesting hypothesis is that BMP signaling in nociceptors functions

to homeostatically regulate synaptic function (similar to that seen in motor neurons). However, our

study suggests that BMP signaling acts differently in nociceptors than in motor neurons. First,

although our data show that activated intracellular BMP signaling in nociceptors resulted in

hypersensitivity of nociception and nociceptor sensitivity, genetic manipulations that increase

intracellular BMP signaling in motor neurons does not increase evoked EJP in larval NMJ (24, 27-29).

Second, interfering with BMP signaling with dominant negative Mad did not cause nociception

insensitive phenotypes (Fig S3) (consistent with another study that found that nociceptor-specific

knockdown of BMP signaling components did not affect basal thermal nociception (22)), while loss of

BMP signaling components in motor neurons decreased evoked EJP by disturbing homeostatic

regulation of synapses (23, 35, 51). Finally, local pMad signals were detected at NMJ as well as at

nuclei in motor neurons. This is relevant in that local pMad at the NMJ functions through a

non-canonical signaling pathway to regulate synaptic maturation (25, 34). In the case of nociceptors,

however, pMad signals were undetectable at nociceptor axon termini (Fig 2G). Although a full

understanding of the mechanisms through which BMP signaling regulates nociceptor sensitivity

requires further investigation, these results indicate that BMP signaling may act differently in the

nociceptors and motor neurons to regulate neuronal outputs.

Potential conservation of Hiw-BMP pathway in regulating nociception in mammals

Hiw and BMP signaling pathway components are all evolutionarily well-conserved. The role of *hiw* in

the negative regulation of nociceptive signaling may be as well. A mammalian *hiw* orthologue

Phr1/MYCBP2 has been previously implicated in a negative regulation of nociception processing.

Specifically, it has been reported that *Phr1/MYCBP2* is expressed in DRG neurons, and that

intrathecal injection of an antisense oligonucleotide against *Phr1/MYCBP2* causes hypersensitivity in

formalin-induced nociceptive responses (52). Furthermore, nociceptive and thermoceptive

neuron-specific *Phr1/MYCBP2* knock-out mice show prolonged formalin-triggered sensitization in

thermal nociception, whereas no obvious phenotypes are observed for basal nociception in the

knock-out animals (53). Decreased internalization of the TRPV1 channel (which is mediated through

a p38 MAPK pathway) has been implicated in this prolonged nociceptive sensitization in *MYCBP2*

knock-out mice (53). In contrast, whether BMP signaling plays a role in regulating nociception in

mammals is unknown. Similarly, the degree to which the role of Hiw and BMP signaling is conserved

in the physiological regulation of mammalian nociceptors represents a fascinating topic for future

investigation.

Intriguingly, Hiw and BMP signaling have been implicated in nerve

regeneration/degeneration processes after axonal injury in both *Drosophila* and mammals (17, 54).

In flies, axonal injury leads to decrease of Hiw, which leads to the upregulation of Wnd that promotes

axonal degeneration in motor neurons (46). Phr1/MYCBP2 is also involved in promoting axonal

degeneration after sciatic or optic nerve axotomy (55). Smad1 is known to be activated and play an

important role for axonal regeneration after peripheral axotomy of DRG neurons (56-59). Because

nerve injuries are thought to be one of key contributors for neuropathic pain conditions and

peripheral axotomies are widely used to generate neuropathic pain models in mammals, it will be of

particular interest in the future to determine whether the Hiw-BMP signaling pathway and

up-regulation of intracellular BMP signaling in nociceptors play a role in the development of a

neuropathic pain state in mammals.

Materials and Methods

Fly strains

Canton-S and w^{1118} were used as control strains as indicated. The other strains used in this study

were as follows: *ppk1.9-GAL4* (60), *UAS-mCD8::GFP* (61), *UAS-ChR2::YFP* line C (4), $hiw^{ND8\,(18)}$,

$hiw^{\Delta N}$, $hiw^{\Delta C}$, UAS-*hiw*, UAS-*hiwΔRing* (20), UAS-*hiwNT,* UAS-*hiwCT,* UAS-*hiw△RCC*,

UAS-*hiw△HindIII*, UAS-*hiwCT1000 (38)*, wnd^1, wnd^2, *UAS-wnd* (30), *ppk1.9-GAL4*; *UAS>CD2*

stop>mCD8::GFP hs-flp, *UAS-tkv*^{QD} (33), *tub-GAL80*^{ts} (62), *ppk-CD4-tdGFP* (63) and

UAS-GCaMP6m (41). *UAS-mad*^{1} (36)

Thermal nociception assay

The thermal nociception assay was performed as described previously (3, 6, 10, 16, 64). NEL latency

was measured as initial contact of the thermal probe on the lateral side of the larval body wall to the

completion of NEL (a 360° roll). Stimulation was ceased at 11 seconds. A thermal probe heated to

46°C was used to examine the insensitive phenotype since it usually evokes NEL in less than 3

seconds (3, 6, 10, 16, 65). A 39°C probe, which usually results in NEL in 9-10 seconds, was used to

examine thermal hypersensitivity, as using a lower temperature probe is important to detecting the

hypersensitive phenotype (16).

Optogenetic nociception assay

383 The optogenetic nociception assay was performed as described previously (5) with slight

384 modifications. 3.8 klux was used to test for optogenetic hypersensitivity, but 76 klux blue light was

385 used in the analysis of acute *tkv*^{QD} induction (Fig 7). Because male larvae show a lower

386 responsiveness to optogenetic nociceptor activation than females (Honjo, unpublished), male larvae

387 were used to allow for more easily detectable hypersensitivity.

Immunohistochemistry

389 Antibodies used in this study were as follows: rabbit anti-GFP (Invitrogen, 1:1000), mouse anti-GFP

390 (Invitrogen, 1:250), mouse anti-rat CD2 (AbD Serotec, 1:200), rabbit anti-pMad (gift from Ed Laufer,

391 1:1000), goat anti-rabbit Alexa488 (Invitrogen, 1:1000), goat anti-rabbit Alexa568 (Invitrogen,

392 1:1000), goat anti-mouse Alexa488 (Invitrogen, 1:1000) and goat anti-mouse Alexa568 (Invitrogen,

393 1:1000). Larvae were filleted, fixed in 4% paraformaldehyde for 30 minutes and then stained

394 according to standard protocols.

395 **pMad staining and image analysis**

396 Wandering third instar larvae expressing mCD8::GFP in nociceptors were filleted and

397 immunostained as described above. To minimize variation due to processing controls, experimental

398 specimens were processed side-by-side within the same staining solutions. The dorsal Class IV

399 mutidendritic neuron (ddaC) was imaged in segments A4-6 (Zeiss LSM 710 with a 100x/1.4

400 Plan-Apochromat oil immersion or Olympus FV1200 with a 100x/1.4 UPLSAPO oil immersion).

401 Z-stack images were converted to maximum intensity projections. To quantify nuclear pMad signals,

402 nociceptor nuclei were identified based on the absence of GFP signal, and a region of interest (ROI)

403 outlining the nucleus was delineated. The average signal intensity of nuclear pMad staining in the

404 ROI was then calculated. Background signal intensity was determined as the mean from ROIs

405 (identical size and shape of the nucleus from the image) drawn in the four corners of each image.

406 The calculated background signal intensity was then subtracted from the nuclear pMad signal level.

407 Data are plotted as nuclear pMad levels normalized to that of the co-processed control specimens.

408 Image analyses were performed in Adobe Photoshop.

409 **Dendrite imaging and quantification**

Wandering third instar larvae expressing mCD8::GFP in nociceptors under the control of

ppk1.9-GAL4 were anesthetized by submersion in a drop of glycerol in a chamber that contained a

cotton ball soaked by a few drops of ether. ddaC neurons in segments A4-6 were imaged on Zeiss

LSM 5 Live with a 40x/1.3 Plan-Neofluar oil immersion objective lens. A series of tiled images were

captured and assembled to reconstruct the entire dendritic field of the three A4-6 ddaC neurons.

Z-stack images were then converted to maximum intensity projections. Dendritic field coverage was

quantified as described previously (16).

Flip-out clone analysis of axon termini

A *ppk1.9-GAL4; UAS>CD2 stop>mCD8::GFP hs-flp* strain was used to induce single cell flip-out

clones in order to sparsely label nociceptors. Six virgin females and three males were used to seed

vials containing a cornmeal molasses medium for a period of 2 days. The seeded vials were then

heat-shocked in a 35°C water bath for 30 minutes. After an additional 3 to 5 days, wandering third

instar larvae were harvested from the vials and dissected. In order to precisely identify the neurons

responsible for the axons labeled in the CNS, the incision made in filleting the larvae was along the

dorsal side, and the CNS remained attached to the fillet prep during immunostaining. mCD8::GFP

and rat CD2 were detected using rabbit anti-GFP and mouse anti-rat CD2 primary antibodies, and

visualized by anti-rabbit Alexa488 and anti-mouse Alexa568 secondary antibodies, respectively.

Axon terminal branches of single cell flip-out clones were imaged in the abdominal ganglion using a

Zeiss LSM 5 Live with a 40x/1.3 Plan-Neofluar oil immersion objective. The cell body of origin for

each flip-out clone was then determined by inspecting the body wall of the corresponding fillet.

Flip-out clones belonging to A1-7 segments were imaged and analyzed.

To analyze the projection patterns for axon terminals, the presence or absence of terminal

branches in each neuromere and longitudinal tract was manually identified for each single nociceptor

clone. In order to align clones projecting to different segments, positions relative to the entry

neuromere were used. The neurons that aligned were then used to calculate the percentage

projecting to each neuromere and longitudinal tract. Heat-maps were color-coded according to these

percentages using Microsoft Excel and Adobe Illustrator.

The quantification of axon terminal area was performed in Matlab. Z-stack images of axon

termini were converted to maximum intensity projections and manually cropped to exclude signals

from other clones in the same sample. The green channel (GFP) and red channel (CD2) of the

cropped images were separately binarized using Otsu's method (66). The number of GFP-positive

pixels were counted to calculate the area innervating the termini. To compensate for differences in

the size and shape of the ventral nerve cord, the number of GFP-positive pixels was normalized to

the average size of a single neuromere, which was calculated as the number of CD2-positive pixels

divided by the number of neuromeres in the cropped image.

Acute induction of tkv^{QD} by tub-GAL80^{ts}

Larvae raised in normal fly vials for 5 or 6 days at 25°C, or larvae raised on apple juice plates

containing ATR for 4 days at 25°C, were transferred to 30°C for 24 hours. In every experiment,

experimental genotypes and control animals were treated side-by-side to minimize the effect of

potential variations in temperature.

Calcium imaging

The ppk1.9-GAL4 UAS-GCaMP6m strain was crossed to either a control strain (w^{1118}) or *UAS-tkv^{QD}*

strain. Activity of larval nociceptors were monitored at their axon terminals in the larval ventral nerve

cord (VNC), which was exposed for imaging by a partial dissection as follows: wandering third instar

larvae expressing GCaMP6m in Class IV md neurons were immobilized in ice cold hemolymph-like

saline 3.1 (HL3.1) (70 mM NaCl, 5mM KCl, 1.5 mM $CaCl_2$, 4 mM $MgCl_2$, 10 mM $NaHCO_3$, 5 mM

Trehalose, 115 mM Sucrose, and 5 mM HEPES, pH 7.2)(67). The outer cuticle of each larvae was

cut at segment A2/A3 to expose the central nervous system from which intact ventral nerves

innervate the posterior larval body. The partially dissected animals were transferred to an imaging

chamber containing HL3.1 equilibrated to the room temperature (23-25 °C). A strip of parafilm was

placed over the larval VNC and was used to gently press the nerve cord down onto a coverslip for

imaging. A Zeiss LSM5 Live confocal microscope and a 20x/0.8 Plan-Apochromat objective with a

piezo focus drive were used to perform three-dimensional time-lapse imaging. Z-stacks consisting of

10-11 optical slices (Z depth of 63 to 70 µm) of 256 x 128 pixel images were acquired at

approximately 4 Hz. During imaging, and using a custom-made thermal probe, a heat ramp stimulus

was applied locally to one side of the A5 to A7 segments. The temperature of the thermal probe was

regulated using a variac transformer. 10V was used to generate a 0.1 °C/sec heat ramp stimulation

and no voltage was applied during cooling. A thermocouple probe (T-type) wire was placed inside of

the thermal probe to monitor the probe temperature, and the data were acquired at 4 Hz through a

digitizer USB-TC01 (National Instruments) and NI Signal Express software (National Instruments).

The acquired images and temperature data were analyzed using Matlab software (Mathworks).

Maximum intensity projections were generated from the time-series Z-stacks. Region of interest

(ROI) was selected as a circular area with a diameter of 6 pixels, whose center was defined as the

centroid of the A6 neuromere. Averaged fluorescent intensities (F) were calculated for the ROI for

each time point. The average of Fs from the first 30 frames was used as a baseline (F_0), and the

percent change in fluorescent intensity from baseline ($\Delta F/F_0$) was calculated for each time point.

Since acquisitions of images and probe temperatures were not synchronized, probe temperature for

each time point was estimated by a linear interpolation from the raw probe temperature reading. For

a comparison of controls and tkv^{QD} OE, $\Delta F/F_0$, data were binned and averaged in 1°C intervals.

Statistical analyses

To statistically compare proportional data, Fisher's exact test was used. Multiple comparisons of

proportional data were corrected by the Bonferroni method. For non-proportional data,

Mann-Whitney's U-test was used for pair-wise comparisons, and Steel's test (non-parametric

equivalent of Dunnet's test) was used for multiple comparisons. Statistical analyses were performed

in R software and Kyplot.

Acknowledgements

We thank TRiP at Harvard Medical School (NIH/NIGMS R01-GM084947), Aaron DiAntonio, Michael

B. O'Connor, Gary Struhl, Chunlai Wu, and the Bloomington Stock Center for fly stocks. We are

grateful to Dan Vasiliauskas, Susan Morton, Tom Jessell, and Ed Laufer for kindly providing the

pMad antibody. We also thank Dr. Emiko Suzuki for her mentorship and helpful advice to KH. We

acknowledge Hisako Honjo for technical assistance in dendrite analysis. Stephania Mauthner,

Andrew Bellemer, and Melissa Christiansen made helpful suggestions on this manuscript. This work

was supported by a grant from the National Institutes of Health (R01GM086458, W.D.T.) and JSPS

KAKENHI (26890025, K.H.). K.H. was supported by postdoctoral fellowships from the Uehara

Memorial Foundation, the Ruth. K. Broad Biomedical Research Foundation, the Japan Society for

the Promotion of Science, and the National Institute of Genetics.

References

1.   Woolf CJ, Ma Q. Nociceptors--noxious stimulus detectors. Neuron. 2007;55(3):353-64.

2.   Gold MS, Gebhart GF. Nociceptor sensitization in pain pathogenesis. Nat Med.

2010;16(11):1248-57.

3.   Tracey WD, Jr., Wilson RI, Laurent G, Benzer S. painless, a Drosophila gene essential for

nociception. Cell. 2003;113(2):261-73.

4.   Hwang RY, Zhong L, Xu Y, Johnson T, Zhang F, Deisseroth K, et al. Nociceptive neurons

protect Drosophila larvae from parasitoid wasps. Curr Biol. 2007;17(24):2105-16.

5.   Honjo K, Hwang RY, Tracey WD, Jr. Optogenetic manipulation of neural circuits and

behavior in Drosophila larvae. Nat Protoc. 2012;7(8):1470-8.

6.   Zhong L, Bellemer A, Yan H, Honjo K, Robertson J, Hwang RY, et al. Thermosensory and

non-thermosensory isoforms of Drosophila melanogaster TRPA1 reveal heat sensor domains of a

thermoTRP channel. Cell Rep. 2012;1(1):43-55.

7.   Kim SE, Coste B, Chadha A, Cook B, Patapoutian A. The role of Drosophila Piezo in

mechanical nociception. Nature. 2012;483(7388):209-12.

8.   Neely GG, Keene AC, Duchek P, Chang EC, Wang QP, Aksoy YA, et al. TrpA1 regulates

thermal nociception in Drosophila. PLoS One. 2011;6(8):e24343.

9.   Babcock DT, Shi S, Jo J, Shaw M, Gutstein HB, Galko MJ. Hedgehog signaling regulates

nociceptive sensitization. Curr Biol. 2011;21(18):1525-33.

10.   Zhong L, Hwang RY, Tracey WD. Pickpocket is a DEG/ENaC protein required for

mechanical nociception in Drosophila larvae. Curr Biol. 2010;20(5):429-34.

11.   Neely GG, Hess A, Costigan M, Keene AC, Goulas S, Langeslag M, et al. A genome-wide

Drosophila screen for heat nociception identifies alpha2delta3 as an evolutionarily conserved pain

gene. Cell. 2010;143(4):628-38.

12.   Kang K, Pulver SR, Panzano VC, Chang EC, Griffith LC, Theobald DL, et al. Analysis of

Drosophila TRPA1 reveals an ancient origin for human chemical nociception. Nature.

2010;464(7288):597-600.

13.   Babcock DT, Landry C, Galko MJ. Cytokine signaling mediates UV-induced nociceptive

sensitization in Drosophila larvae. Curr Biol. 2009;19(10):799-806.

14.   Neely GG, Rao S, Costigan M, Mair N, Racz I, Milinkeviciute G, et al. Construction of a

global pain systems network highlights phospholipid signaling as a regulator of heat nociception.

PLoS Genet. 2012;8(12):e1003071.

15.   Im SH, Takle K, Jo J, Babcock DT, Ma Z, Xiang Y, et al. Tachykinin acts upstream of

autocrine Hedgehog signaling during nociceptive sensitization in Drosophila. eLife. 2015;4:e10735.

16.   Honjo K, Mauthner SE, Wang Y, Skene JH, Tracey WD, Jr. Nociceptor-Enriched Genes

Required for Normal Thermal Nociception. Cell Rep. 2016;16(2):295-303.

17.   Grill B, Murphey RK, Borgen MA. The PHR proteins: intracellular signaling hubs in neuronal

development and axon degeneration. Neural Dev. 2016;11:8.

18.   Wan HI, DiAntonio A, Fetter RD, Bergstrom K, Strauss R, Goodman CS. Highwire regulates

synaptic growth in Drosophila. Neuron. 2000;26(2):313-29.

19.   Husson SJ, Costa WS, Wabnig S, Stirman JN, Watson JD, Spencer WC, et al. Optogenetic

analysis of a nociceptor neuron and network reveals ion channels acting downstream of primary

sensors. Curr Biol. 2012;22(9):743-52.

20.   Wu C, Wairkar YP, Collins CA, DiAntonio A. Highwire function at the Drosophila

neuromuscular junction: spatial, structural, and temporal requirements. J Neurosci.

2005;25(42):9557-66.

21.   Deshaies RJ, Joazeiro CA. RING domain E3 ubiquitin ligases. Annu Rev Biochem.

2009;78:399-434.

22.   Follansbee TL, Gjelsvik KJ, Brann CL, McParland AL, Longhurst CA, Galko MJ, et al.

Drosophila Nociceptive Sensitization Requires BMP Signaling via the Canonical SMAD Pathway. J

Neurosci. 2017;37(35):8524-33.

23.   McCabe BD, Hom S, Aberle H, Fetter RD, Marques G, Haerry TE, et al. Highwire regulates

presynaptic BMP signaling essential for synaptic growth. Neuron. 2004;41(6):891-905.

24.   Li W, Yao A, Zhi H, Kaur K, Zhu YC, Jia M, et al. Angelman Syndrome Protein Ube3a

Regulates Synaptic Growth and Endocytosis by Inhibiting BMP Signaling in Drosophila. PLoS Genet.

2016;12(5):e1006062.

25.   Sulkowski MJ, Han TH, Ott C, Wang Q, Verheyen EM, Lippincott-Schwartz J, et al. A Novel,

Noncanonical BMP Pathway Modulates Synapse Maturation at the Drosophila Neuromuscular

Junction. PLoS Genet. 2016;12(1):e1005810.

26.	Zhang X, Rui M, Gan G, Huang C, Yi J, Lv H, et al. Neuroligin 4 Regulates Synaptic Growth

via the Bone Morphogenetic Protein (BMP) Signaling Pathway at the Drosophila Neuromuscular

Junction. J Biol Chem. 2017.

27.	Zhao G, Wu Y, Du L, Li W, Xiong Y, Yao A, et al. Drosophila S6 Kinase like inhibits

neuromuscular junction growth by downregulating the BMP receptor thickveins. PLoS Genet.

2015;11(3):e1004984.

28.	Nahm M, Lee MJ, Parkinson W, Lee M, Kim H, Kim YJ, et al. Spartin regulates synaptic

growth and neuronal survival by inhibiting BMP-mediated microtubule stabilization. Neuron.

2013;77(4):680-95.

29.	Zhao L, Wang D, Wang Q, Rodal AA, Zhang YQ. Drosophila cyfip regulates synaptic

development and endocytosis by suppressing filamentous actin assembly. PLoS Genet.

2013;9(4):e1003450.

30.	Collins CA, Wairkar YP, Johnson SL, DiAntonio A. Highwire restrains synaptic growth by

attenuating a MAP kinase signal. Neuron. 2006;51(1):57-69.

31.	Vanlandingham PA, Fore TR, Chastain LR, Royer SM, Bao H, Reist NE, et al. Epsin 1

Promotes Synaptic Growth by Enhancing BMP Signal Levels in Motoneuron Nuclei. PLoS One.

2013;8(6):e65997.

32.   Shi W, Chen Y, Gan G, Wang D, Ren J, Wang Q, et al. Brain tumor regulates neuromuscular

synapse growth and endocytosis in Drosophila by suppressing mad expression. J Neurosci.

2013;33(30):12352-63.

33.   Nellen D, Burke R, Struhl G, Basler K. Direct and long-range action of a DPP morphogen

gradient. Cell. 1996;85(3):357-68.

34.   Sulkowski M, Kim YJ, Serpe M. Postsynaptic glutamate receptors regulate local BMP

signaling at the Drosophila neuromuscular junction. Development. 2014;141(2):436-47.

35.   McCabe BD, Marques G, Haghighi AP, Fetter RD, Crotty ML, Haerry TE, et al. The BMP

homolog Gbb provides a retrograde signal that regulates synaptic growth at the Drosophila

neuromuscular junction. Neuron. 2003;39(2):241-54.

36.   Takaesu NT, Herbig E, Zhitomersky D, O'Connor MB, Newfeld SJ. DNA-binding domain

mutations in SMAD genes yield dominant-negative proteins or a neomorphic protein that can activate

WG target genes in Drosophila. Development. 2005;132(21):4883-94.

37.   Wu C, Daniels RW, DiAntonio A. DFsn collaborates with Highwire to down-regulate the

Wallenda/DLK kinase and restrain synaptic terminal growth. Neural Dev. 2007;2:16.

38.   Tian X, Li J, Valakh V, DiAntonio A, Wu C. Drosophila Rae1 controls the abundance of the

ubiquitin ligase Highwire in post-mitotic neurons. Nat Neurosci. 2011;14(10):1267-75.

39.   Xiong X, Hao Y, Sun K, Li J, Li X, Mishra B, et al. The Highwire ubiquitin ligase promotes

axonal degeneration by tuning levels of Nmnat protein. PLoS Biol. 2012;10(12):e1001440.

40.   Wang X, Kim JH, Bazzi M, Robinson S, Collins CA, Ye B. Bimodal control of dendritic and

axonal growth by the dual leucine zipper kinase pathway. PLoS Biol. 2013;11(6):e1001572.

41.   Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, et al. Ultrasensitive

fluorescent proteins for imaging neuronal activity. Nature. 2013;499(7458):295-300.

42.   McGuire SE, Mao Z, Davis RL. Spatiotemporal gene expression targeting with the TARGET

and gene-switch systems in Drosophila. Sci STKE. 2004;2004(220):pl6.

43.   Wang LX, Wang ZJ. Animal and cellular models of chronic pain. Adv Drug Deliv Rev.

2003;55(8):949-65.

44.   Leung C, Wilson Y, Khuong TM, Neely GG. Fruit flies as a powerful model to drive or

validate pain genomics efforts. Pharmacogenomics. 2013;14(15):1879-87.

45.   DiAntonio A, Haghighi AP, Portman SL, Lee JD, Amaranto AM, Goodman CS.

Ubiquitination-dependent   mechanisms   regulate   synaptic   growth   and   function.   Nature.

2001;412(6845):449-52.

46.   Xiong X, Wang X, Ewanek R, Bhat P, Diantonio A, Collins CA. Protein turnover of the

Wallenda/DLK   kinase   regulates   a   retrograde   response   to   axonal   injury.   J   Cell   Biol.

2010;191(1):211-23.

47.   Bragdon B, Moseychuk O, Saldanha S, King D, Julian J, Nohe A. Bone morphogenetic

proteins: a critical review. Cell Signal. 2011;23(4):609-20.

48.   Hodge LK, Klassen MP, Han BX, Yiu G, Hurrell J, Howell A, et al. Retrograde BMP signaling

regulates trigeminal sensory neuron identities and the formation of precise face maps. Neuron.

2007;55(4):572-86.

49.   Bayat V, Jaiswal M, Bellen HJ. The BMP signaling pathway at the Drosophila

neuromuscular junction and its links to neurodegenerative diseases. Curr Opin Neurobiol.

2011;21(1):182-8.

50.   Marques G. Morphogens and synaptogenesis in Drosophila. J Neurobiol.

2005;64(4):417-34.

51.   Eaton BA, Davis GW. LIM Kinase1 controls synaptic stability downstream of the type II BMP

receptor. Neuron. 2005;47(5):695-708.

52.   Ehnert C, Tegeder I, Pierre S, Birod K, Nguyen HV, Schmidtko A, et al. Protein associated

with Myc (PAM) is involved in spinal nociceptive processing. J Neurochem. 2004;88(4):948-57.

53.   Holland S, Coste O, Zhang DD, Pierre SC, Geisslinger G, Scholich K. The ubiquitin ligase

MYCBP2 regulates transient receptor potential vanilloid receptor 1 (TRPV1) internalization through

inhibition of p38 MAPK signaling. J Biol Chem. 2011;286(5):3671-80.

54.   Zhong J, Zou H. BMP signaling in axon regeneration. Curr Opin Neurobiol. 2014;27:127-34.

55.   Babetto E, Beirowski B, Russler EV, Milbrandt J, DiAntonio A. The Phr1 ubiquitin ligase

promotes injury-induced axon self-destruction. Cell Rep. 2013;3(5):1422-9.

56.   Saijilafu, Hur EM, Liu CM, Jiao Z, Xu WL, Zhou FQ. PI3K-GSK3 signalling regulates

mammalian axon regeneration by inducing the expression of Smad1. Nat Commun. 2013;4:2690.

57.   Parikh P, Hao Y, Hosseinkhani M, Patil SB, Huntley GW, Tessier-Lavigne M, et al.

Regeneration of axons in injured spinal cord by activation of bone morphogenetic protein/Smad1

signaling pathway in adult neurons. Proc Natl Acad Sci U S A. 2011;108(19):E99-107.

58.   Zou H, Ho C, Wong K, Tessier-Lavigne M. Axotomy-induced Smad1 activation promotes

axonal growth in adult sensory neurons. J Neurosci. 2009;29(22):7116-23.

59.   Ma CH, Brenner GJ, Omura T, Samad OA, Costigan M, Inquimbert P, et al. The BMP

coreceptor RGMb promotes while the endogenous BMP antagonist noggin reduces neurite

outgrowth and peripheral nerve regeneration by modulating BMP signaling. J Neurosci.

2011;31(50):18391-400.

60.   Ainsley JA, Pettus JM, Bosenko D, Gerstein CE, Zinkevich N, Anderson MG, et al.

Enhanced locomotion caused by loss of the Drosophila DEG/ENaC protein Pickpocket1. Curr Biol.

2003;13(17):1557-63.

61.   Lee T, Luo L. Mosaic analysis with a repressible cell marker for studies of gene function in

neuronal morphogenesis. Neuron. 1999;22(3):451-61.

62.   McGuire SE, Le PT, Osborn AJ, Matsumoto K, Davis RL. Spatiotemporal rescue of memory

dysfunction in Drosophila. Science. 2003;302(5651):1765-8.

63.   Han C, Jan LY, Jan YN. Enhancer-driven membrane markers for analysis of

nonautonomous mechanisms reveal neuron-glia interactions in Drosophila. Proc Natl Acad Sci U S A.

2011;108(23):9673-8.

64.   Hwang RY, Stearns NA, Tracey WD. The Ankyrin Repeat Domain of the TRPA Protein

Painless Is Important for Thermal Nociception but Not Mechanical Nociception. PLoS One.

2012;7(1):e30090.

65.   Tracey WD, Jr. Genetics Can Be Painless: Molecular Genetic Analysis of Nociception in

Drosophila. 2007.

66.   Otsu N. Threshold Selection Method from Gray-Level Histograms. Ieee T Syst Man Cyb.

1979;9(1):62-6.

67.   Feng Y, Ueda A, Wu CF. A modified minimal hemolymph-like solution, HL3.1, for

physiological recordings at the neuromuscular junctions of normal and mutant Drosophila larvae. J

Neurogenet. 2004;18(2):377-402.

 Figure legends

Fig 1. *hiw* is involved in both desensitizing and sensitizing pathways in nociceptors.

(A) Insensitive thermal nociception in hiw^{ND8} mutant and nociceptor-specific rescue of the insensitivity.

In comparison to the control w^{1118} (n = 119, 1.8 ± 0.2), hiw^{ND8} (n = 114, 5.1 ± 0.3), no driver control

(hiw^{ND8}; UAS-*hiw*/+, n = 108, 3.7 ± 0.2) and hiw^{ND8} with GAL4 driver (hiw^{ND8}; *ppk-GAL4*

UAS-mCD8::GFP/+, n = 101, 3.9 ± 0.2) all showed significantly delayed nociceptive responses to a

46°C probe, while the rescue genotype (hiw^{ND8}; *ppk-GAL4 UAS-mCD8::GFP/UAS-hiw*, n = 122, 1.7 ±

0.1) had a normal response. *** p < 0.001 (Steel's test versus control). (B) A schematic of thermal

and optogenetic stimulation of a nociceptor. While heat stimuli activate nociceptors via sensory

transduction, ChR2 triggers nociceptor activation independently of sensory transduction. (C) The *hiw*

genetic mutant expressing ChR2 in nociceptors was more responsive than the control to

optogenetically triggered nociceptor activation, and the nociceptor-specific expression of *hiw* rescued

optogenetic nociception responses to levels similar to control. Control (w^{1118}/Y; *ppk-GAL4*

UAS-ChR2::YFP/+, n = 154, 31 ± 4%), hiw^{ND8} (hiw^{ND8}/Y; *ppk-GAL4 UAS-ChR2::YFP/+*, n =191, 75 ±

3%) and *hiw* rescue (hiw^{ND8}/Y; *ppk-GAL4 UAS-ChR2::YFP/UAS-hiw,* n = 112, 35 ± 5%). *** p < 0.001

(Fisher's exact test with Bonferroni correction). (D) *hiwΔRing* expression (*hiwΔRing* OE) in

nociceptors resulted in thermal hypersensitivity. *hiwΔRing* OE animals (*ppk-GAL4* x *UAS-hiwΔRing,*

n = 90, 5.7 ± 0.4) showed a significantly shortened latency to respond to a 39°C thermal probe

compared to controls (ppk-GAL4 x w^{1118}, n = 104, 8.8 ± 0.3). *** p < 0.001 (Mann-Whitney's U-test).

(E) *hiwΔRing* expression in nociceptors induced hypersensitivity to optogenetic nociceptor

stimulation. *hiwΔRing*-expressing animals (*ppk-GAL4 UAS-ChR2::YFP x UAS-hiwΔRing,* n = 71, 55

± 6%) exhibited increased responsiveness to optogenetic stimulation of nociceptors compared to

control animals (*ppk-GAL4 UAS-ChR2::YFP x w^{1118}*, n = 64, 23 ± 5%). *** $p < 0.001$ (Fisher's exact

test). All error bars represent standard error.

Fig 2. BMP signaling in nociceptors is negatively regulated at the downstream of *hiw*

(A-E) Representative images of pMad immunoreactivity in nociceptor cell bodies. Green represents

mCD8::GFP and magenta shows pMad signals. (A'-E') Split images for pMad signals. (F)

Quantification of nuclear pMad signals in nociceptors. *hiw^{ND8}* mutants (*hiw^{ND8}*; *ppk-GAL4*

UAS-mCD8::GFP/+, n = 21) and *hiwΔRing* OE (*ppk-GAL4 UAS-mCD8::GFP/+*; *UAS-hiwΔRing/+*, n

= 18) had 33 ± 7% and 40 ± 8% increases in nuclear pMad signals, respectively. No significant

difference in nuclear pMad level compared to controls was detected in *hiw rescue* animals (*hiw^{ND8}*;

ppk-GAL4 UAS-mCD8::GFP/UAS-hiw, n = 24). Nociceptors expressing *tkv^{QD}* (*ppk-GAL4*

UAS-mCD8::GFP x UAS-tkv^{QD}, n = 24) also showed significantly increased nuclear pMad levels (84

± 7%). Control (*ppk-GAL4 UAS-mCD8::GFP/+*, n > 24) * $p < 0.05$, *** $p < 0.001$ (Mann-Whitney's

U-test). (G) A projection image of axon terminal of Class IV neurons at A4 and A5 segments in the

larval ventral ganglia. Green represents mCD8::GFP and magenta shows pMad signals. pMad

signals at axon termini in nociceptors were not distinguishable from the background. (G') Split image

for mCD8::GFP. (G") Split image for pMad signals. (H) Expression of mad[1] suppressed the thermal

hypersensitivity in *hiwΔRing*-expressing animals. Control (*w^1118 x ppk-GAL4,* n = 73, 10.1 ± 0.3),

hiwΔRing OE (*ppk-GAL4 x UAS-hiwΔRing*, n = 59, 6.7 ± 0.4) and *hiwΔRing* & *mad1* OE (*ppk-GAL4*

UAS-hiwΔRing x UAS-mad^1, n = 54, 10.6 ± 0.2). *** $p < 0.001$ (Steel's test versus control). Error bars

represent standard error.

Fig 3. Activated BMP signaling in *hiw* mutant does not depend on *wallenda*

(A-E) Representative images of pMad immunoreactivity in nociceptor soma. Green shows

mCD8::GFP and magenta represents pMad signals. (A'-E') Split images for pMad signals. (F)

Quantification of nuclear pMad signals in nociceptors. Similarly to *hiw^ND8* mutants (*hiw^ND8*; *ppk-GAL4*

UAS-mCD8::GFP/+, n = 36), *hiw^ND8* with heterozygous or transheterozygous *wnd* mutations

(*hiw^ND8*; *ppk-GAL4 UAS-mCD8::GFP*/+; *wnd^2*/+ and *hiw^ND8*; *ppk-GAL4 UAS-mCD8::GFP*/+;

wnd^1/wnd^2, n = 48 and 45) showed significantly increased nuclear pMad level relative to controls

(*ppk-GAL4 UAS-mCD8::GFP*/+, n = 48). The transheterozygous *wnd* mutants (*ppk-GAL4*

UAS-mCD8::GFP/+; *wnd^1/wnd^2*, n = 33) did not show a significant difference in nuclear pMad level

compared to controls ($p > 0.7$). ** $p < 0.01$, *** $p < 0.001$ (Mann-Whitney's U-test). Error bars

represent standard error.

Fig 4. pMad signals in nociceptors expressing various *hiw* deletion constructs.

(A) A schematic showing the structure of Hiw and Hiw deletion constructs. (B-G) Representative

images of pMad immunoreactivity in nociceptor cell bodies. Green represents mCD8::GFP and

magenta shows pMad signals. (B'-G') Split images for pMad signals. (H) Quantification of nuclear

pMad signals in nociceptors. Nociceptors expressing *hiwNT* OE (*ppk-GAL4 UAS-mCD8::GFP* x

UAS-hiwNT, n = 12), *hiwCT* OE (*ppk-GAL4 UAS-mCD8::GFP* x *UAS-hiwCT*, n = 12) and *hiwΔRCC1*

OE (*ppk-GAL4 UAS-mCD8::GFP* x *UAS-hiwΔRCC1*, n = 12) showed nuclear pMad signals

increased by 218 ± 26%, 99 ± 19% and 68 ± 18%, respectively. A significant difference in nuclear

pMad level compared to controls was not detected in *hiwΔHindIII* OE (*ppk-GAL4 UAS-mCD8::GFP* x

UAS-hiwΔHindIII, n = 12) or *hiwCT1000* OE (*ppk-GAL4 UAS-mCD8::GFP* x *UAS-hiwCT1000*, n =

12). Control (*ppk-GAL4 UAS-mCD8::GFP*/+, n = 12) ** $p < 0.01$, *** $p < 0.001$ (Mann-Whitney's

U-test). Error bars represent standard error.

Fig 5. Activation of BMP signaling in nociceptors induces nociceptive hypersensitivity.

(A) Animals expressing tkv^{QD} in Class IV neurons showed thermal hypersensitivity. Control

(*ppk-GAL4* x w^{1118}, n=102, 6.6 ± 0.3) and tkv^{QD} OE (*ppk-GAL4 x UAS-tkv^{QD}*, n=118, 5.0 ± 0.3). ** $p <$

0.01 (Mann-Whitney's U-test). (B) Expression of tkv^{QD} in Class IV neurons caused optogenetic

hypersensitivity. tkv^{QD} overexpressors expressing ChR2::YFP in nociceptors (*ppk-GAL4*

UAS-ChR2::YFP x UAS-tkv^{QD}, n = 74, 59 ± 6%) showed significantly elevated responsiveness to

blue light-triggered nociceptor activation compared to controls (*ppk-GAL4 UAS-ChR2::YFP* x w^{1118}, n

= 67, 21 ± 5%). *** p < 0.001 (Fisher's exact test). (C-E) Overexpression of *tkv^QD* in nociceptors did

not affect dendritic coverage. (C and D) Representative images of ddaC dendrites in control

(*ppk-GAL4 UAS-mCD8::GFP* x *w^1118*) and *tkv^QD* overexpression (*ppk-GAL4 UAS-mCD8::GFP x*

UAS-tkv^QD) animals. Scale bars represent 100 µm. (E) Quantification of dendritic coverage. Dendritic

coverage in *tkv^QD*-overexpressing animals was indistinguishable from that in controls (n = 6 and 6, p

> 0.3, Mann-Whitney's U-test). (F-H) Expression of *tkv^QD* in nociceptors resulted in overextension of

axon termini. (F) A representative image of a v'ada Class IV axon terminal in a *tkv^QD* overexpressor

(ppk1.9-GAL4; UAS>CD2 stop>mCD8::GFP hs-flp x UAS-*tkv^QD*). Scale bar represents 5 µm. (G)

Heat map of axonal projections. Animals with expression of *tkv^QD* showed a severe overextension

phenotype (n = 13) compared to controls (ppk1.9-GAL4; UAS>CD2 stop>mCD8::GFP hs-flp x *w^1118*,

n = 24). (H) Quantification of terminal size of the v'ada Class IV neuron. Terminal size of the v'ada

axon was significantly increased in *tkv^QD*-expressing animals (n = 13) compared to controls (n = 24).

*** p < 0.001 (Steel's test versus control). All error bars represent standard error.

Fig 6. Elevated BMP signaling increases Ca^{2+} responses in nociceptor axon terminals

(A) A cartoon showing the Ca^{2+} imaging preparation to monitor GCaMP6m signals from nociceptor

terminals during heat ramp stimuli. (B and B') Representative images showing thermal activation of

nociceptors in control animals during calcium imaging (*ppk-GAL4 UAS-GCaMP6m* x *w^1118*). In

comparison to the initial frame (B), the GCaMP6m signal monitored at nociceptor axon termini was

increased when the probe temperature reached 40°C (B'). (C and C') Images showing a

representative result of animals with nociceptor-specific up-regulation of BMP signaling (tkv^{QD} OE,

ppk-GAL4 UAS-GCaMP6m x *UAS-tkv^{QD}*). Compared to the baseline (C), increase of GCaMP

fluorescent intensity was observed when the probe temperature reached 40°C (C'). (D) Average

percent increase of GCaMP6m fluorescent intensity relative to baseline ($\Delta F/F_0$) during heat ramp

stimulations. $\Delta F/F_0$ is plotted to binned probe temperature (interval = 1 °C). In controls, GCaMP6m

fluorescence in nociceptors began increasing when the probe temperature reached 37 °C, peaked at

43°C, and returned to baseline at 47 °C. In comparison to controls, nociceptors of tkv^{QD} OE animals

showed a highly exaggerated fluorescent increase of GCaMP through 36°C to 50°C. n = 17 and 19

for controls and tkv^{QD} OE, respectively. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (Mann-Whitney's U-test).

(E) Basal GCaMP6m signals (F_0) did not differ significantly between controls and tkv^{QD} OE (n = 17

and 19). $p > 0.6$ (Mann-Whitney's U-test). All error bars represent standard error.

Fig 7. Acute up-regulation of BMP signaling sensitizes optogenetic nociception

(A) Acute expression of tkv^{QD} induced hypersensitivity in optogenetic nociception. After 24 hour

induction of tkv^{QD} in Class IV nociceptors (ppk-GAL4 UAS-ChR2::YFP/+; UAS-tkv^{QD}/tub-GAL80^{ts}

incubated at 30°C for 24 hours, n = 65, 32 ± 6%), larval nociceptive responses to optogenetic

activation of Class IV nociceptors were significantly increased compared to those in controls

(ppk-GAL4 UAS-ChR2::YFP/+; tub-GAL80^{ts}/+ incubated at 30°C for 24 hours, n = 73, 15 ± 4%). * p <

0.05 (Fisher's exact test). (B) Acute induction of tkv^{QD} increased nuclear pMad levels in nociceptors.

pMad levels in nociceptor nuclei were significantly elevated (64 ± 8%) in animals with 24 hour tkv^{QD}

induction (ppk-GAL4 UAS-mCD8::GFP/+; tub-GAL80^{ts}/UAS-tkv^{QD} incubated at 30°C for 24 hours, n

= 12) compared to control animals (ppk-GAL4 UAS-mCD8::GFP/+; tub-GAL80^{ts}/+ incubated at 30°C

for 24 hours, n = 12). When raised at 25°C, animals with UAS-tkv^{QD} (ppk-GAL4

UAS-mCD8::GFP/ppk-CD4-tdGFP; tub-GAL80^{ts}/UAS-tkv^{QD}, n = 48) and controls (ppk-GAL4

UAS-mCD8::GFP/ppk-CD4-tdGFP; tub-GAL80^{ts}/+, n = 48) showed comparable pMad levels. *** p <

0.001 (Mann-Whitney's U-test). (C and D) 24-hour induction of tkv^{QD} did not induce axonal

overgrowth. (C) A representative image of axon termini of a single v'ada neuron. Scale bar

represents 5 μm. (D) Heat map of v'ada axonal projection. 24-hour expression of tkv^{QD} did not cause

a severe overextension phenotype (n = 7). The heat map of the control is reused from Fig. 5G for

comparison. All error bars represent standard error.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7