1 **Enhancing resolution of natural methylome reprogramming behavior in plants**

3 Robersy Sanchez[†], Xiaodong Yang[†], Jose R Barreras, Hardik Kundariya and Sally A. Mackenzie[*]

5 Departments of Biology and Plant Science, The Pennsylvania State University, University Park, PA

6 16802

[†]Equal contributors

*Correspondence to: sam795@psu.edu

Sally Mackenzie

362 Frear North Bldg

Pennsylvania State University

University Park, PA 16802

Ph 814-863-8324, email sam795@psu.edu

Robersy Sanchez: rus547@psu.edu

Xiaodong Yang: xiaodongy86@gmail.com

Jose R Barreras: barreras@gmail.com

Hardik Kundariya: kundariyahardik@gmail.com

Sally Mackenzie: sam795@psu.edu

Abstract

Background
Natural methylome reprogramming within chromatin involves changes in local energy landscapes that are

subject to thermodynamic principles. Signal detection permits the discrimination of methylation signal

from dynamic background noise that is induced by thermal fluctuation. Current genome-wide methylation

analysis methods do not incorporate biophysical properties of DNA, and focus largely on DNA

methylation density changes, which limits resolution of natural, more subtle methylome behavior in

relation to gene activity.

41
Results
We present here a novel methylome analysis procedure, Methyl-IT, based on information

thermodynamics and signal detection. Methylation analysis involves a signal detection step, and the

method was designed to discriminate methylation regulatory signal from background variation.

Comparisons with commonly used programs and two publicly available methylome datasets, involving

stages of seed development and drought stress effects, were implemented. Information divergence

between methylation levels from different groups, measured in terms of Hellinger divergence, provides

discrimination power between control and treatment samples. Differentially informative methylation

positions (DIMPs) achieved higher sensitivity and accuracy than standard differentially methylated

positions (DMPs) identified by other methods. Differentially methylated genes (DMG) that are based on

DIMPs were significantly enriched in biologically meaningful networks.

53
Conclusions
Methyl-IT analysis enhanced resolution of natural methylome reprogramming behavior to reveal

network-associated responses, offering resolution of gene pathway influences not attainable with previous

methods.

Keywords

Epigenomics, DNA methylation, gene expression, information theory, Arabidopsis

Background

Most chromatin changes that are associated with epigenetic behavior are reprogrammed each generation, with the apparent exception of cytosine methylation, where parental patterns can be inherited through meiosis [1]. Genome-wide methylome analysis, therefore, provides one avenue for investigation of transgenerational and developmental epigenetic behavior. Complicating such investigations in plants is the dynamic nature of DNA methylation [2, 3] and a presently incomplete understanding of its association with gene expression. In plants, cytosine methylation is generally found in three contexts, CG, CHG and CHH (H=C, A or T), with CG most prominent within gene body regions [4]. Association of CG gene body methylation with changes in gene expression remains in question. There exist ample data associating chromatin behavior with plant response to environmental changes [5], yet, affiliation of genome-wide DNA methylation with these effects, or their inheritance, remains inconclusive [6, 7].

The epigenetic landscape is modulated by thermodynamic fluctuations that influence DNA stability [8, 9] [10]. Most genome-wide methylome studies have relied predominantly on statistical approaches that ignore fundamental biophysical properties of cytosine DNA methylation, offering limited resolution of those genomic regions with highest probability of having undergone epigenetic change. Jenkinson and colleagues [11] have implemented statistical physics and information theory to the analysis of whole genome methylome data to define sample-specific energy landscapes. Our group [12, 13] proposed an information thermodynamics approach to investigate genome-wide methylation patterning based on the statistical mechanical effect of methylation on DNA molecules. The information thermodynamics-based approach is postulated to provide greater sensitivity for resolving true signal from background variation within the methylome [12]. Because gene-associated biological signal created within the dynamic methylome environment characteristic of plants may be subtle and is not free from background noise, the approach, designated Methyl-IT, includes application of signal detection theory [14-18].

A basic requirement for the application of signal detection is a probability distribution for background noise. Probability distribution, as a Weibull distribution model, can be deduced on a statistical mechanical basis for DNA methylation induced by thermal fluctuations [12]. Assuming that this background methylation variation is consistent with a Poisson process, it can be distinguished from variation associated with methylation regulatory machinery, which is non-independent for all genomic regions [12]. An information-theoretic divergence to express the background variation will follow a Weibull distribution model, provided that it is proportional to minimum energy dissipated per bit of information from methylation change.

The information thermodynamics model was previously verified with more than 150 Arabidopsis and
more than 90 human methylome datasets [12]. To test application of Methyl-IT to methylome analysis,
and compare resolution to approaches used in programs DSS [19], and methylpy [20], we investigated
two Arabidopsis methylome datasets. For resolution of methylation signal during plant development, we
used previously reported datasets from globular stage (4 days after pollination [DAP]), linear cotyledon
stage (8 DAP), mature green stage (13 DAP), post-mature green stage (18 DAP), dry seed (Ws-0), and
leaf [21, 22] . To assess methylation signal during stress in plants, and association of methylation with
altered gene expression during stress, we investigated data from Ganguly et al. (2017), which involves
mild drought stress by withholding irrigation for 9 days [23, 24]. Direct comparison of outputs by
Methyl-IT with previous analyses by methylpy and DSS are presented.
106

Results

The information thermodynamics model and Methyl-IT workflow
Methylation level is generally the ratio of methylated cytosine read counts divided by the sum of
methylated and unmethylated cytosine read counts for a given cytosine site. This is a descriptive variable
that reflects uncertainty of methylation level at a given cytosine site. Most methylation analyses test
whether or not the difference between control (CT) and treatment (TT) methylation levels (the uncertainty
variation) is statistically significant. The approach measures the absolute value of the difference between
methylation levels $\left| p_i^{TT} - p_i^{CT} \right|$ from control (p_i^{CT}) and treatment (p_i^{TT}) at each cytosine site. The
magnitude of $\left| p_i^{TT} - p_i^{CT} \right|$ is known as total variation distance (TVD).
116

To improve resolution of methylation signal, we applied Hellinger divergence (*HD*), ([25], detailed
description included in Methods section). Both TVD and HD are information divergences that follow
asymptotic chi-square distribution [25]. However, HD converges faster and carries more information than
TVD and, consequently, has higher discrimination power [26]. The improvement in discrimination power
is visible in Fig. 1 By way of illustration, we used the drought stress data, where CTR designated
unstressed control group and STR designated stressed group. Fig. 1a shows that treatment methylation
signal on chromosome 1, expressed in terms of methylation level, was indistinguishable from control.
Higher resolutions are reached with TVD and HD, with HD providing highest discrimination power.
125

Ganguly et al. reported individual variation and pre-existing methylation differences in the drought stress
materials [24], which is reflected by HD in Fig. 1c. The improvement in resolution attributed to HD
derives from the fact that TVD takes into account only one dimension of the methylation change, while

HD is estimated in bi-dimensional space $(p_i, 1 - p_i)$, where the goodness-of-fit test to detect differences

is performed.

Genome-wide Hellinger divergence for background methylation variation can be modeled by a Weibull

distribution [12]. On the other hand, biologically meaningful methylation changes result in an increment

of Hellinger divergence distinguishable in the signal detection step (Fig. 2). For a given level of

significance α (Type I error probability, eg. α = 0.05), cytosine positions with $H_{\alpha=0.05}$ can be selected as

sites carrying potential biological signal (shown as the blue shade region under the curve in Fig. 2). True

signal is detected based on optimal cutpoint [27], which can be estimated by area under the curve (AUC)

from a receiver operating characteristic (ROC) built from logistic regression with potential signals from

control and treatment. The AUC is the probability to distinguish biological regulatory signal naturally

generated in the control from that induced by the treatment. Cytosine sites carrying methylation signal are

designated *differentially informative methylation positions* (DIMPs). The probability that a DIMP is not

induced by the treatment is designated probability of false alarm (P_{FA}, false positive, Fig. 2). As suggested

in Fig. 2, we define DIMPs as cytosine positions with high probability to carry signal created in response

to treatment.

Estimation of optimal cutoff from AUC is an additional step to remove any remaining potential

methylation background noise that still remains with probability α = 0.05 > 0. We define as methylation

signal (DIMP) each cytosine site with Hellinger divergence values above the cutpoint (shown in Fig. 2 as

$H_{33}^{D_T}$). Each DIMP-associated signal may or may not be represented within a DMP derived by Fisher's

exact test (or other current tests, Fig. 2). The difference in resolution by current methods versus Methyl-

IT is illustrated by positioning H value sensitivity for Fisher's exact test (FET) at greater than H_{min} for

cytosine sites that are DMPs and DIMPs simultaneously (Fig. 2).

Table 1 provides a critical but non-unique example; assume there is an experiment that yields read counts

with $n_i^{mC_c} = 8$, $n_i^{C_c} = 2$, $n_i^{mC_t} = 350$, and $n_i^{C_t} = 20$, where $n_i^{mC_c}$ and $n_i^{mC_t}$ refer to methylated cytosine

read counts in control and treatment, respectively, and $n_i^{C_c}$ and $n_i^{C_t}$ to non-methylated cytosine counts in

control and treatment, respectively. In the given example, it's clear that control and treatment have

different methylation pattern, but Fisher's exact test (including one tail test or Monte Carlo (MC)

simulations with 3000 resamplings (3k)) failed to detect the difference (for significance level α = 0.05).

Root-mean-square test (RMST) used in methylpy [20] and goodness-of-fit test based on Hellinger chi-

square test (HCT, with HD as statistic) [25, 28] proved the sensitivity but still failed to detect the

difference (for $\alpha= 0.05$). However, if the hypothetical methylation changes were to occur in the drought

stress experiment, then Weibull distribution modeling in Methyl-IT would yield *p*-values of 5.08E-04,

5.08E-04, and 3.20E-04 for each stressed plant (Table 1). Such methylation changes represent potential

DIMPs. The conclusions will remain the same even for a generalized situation with n_i^{mCt} running between

80 and 350 ($80 \leq n_i^{mC_t} \leq 350$). Considering that even a small genome like Arabidopsis contains millions

of cytosine sites, the situation presented in Table 1 is not rare, and the difference caused by statistical tests

listed in Table 1 would be significant. A flow chart integrating the main procedures of Methyl-IT and

optional downstream analysis is shown in Fig. 3.

Methyl-IT sensitivity and genomic regions targeted by DIMPs

To investigate the sensitivity of Methyl-IT, we applied DIMP detection to the drought stress dataset and

compared with the outputs from other methods. Fig. 4 shows a direct comparison of DIMPs to DMPs

estimated with Fisher's exact test, DMSs (differentially methylated sites) estimated with root mean square

test (RMST, approach implemented in methylpy [20, 21]), and DMCs (differentially methylated cytosines)

estimated with Hellinger chi-square test (HCT).

In all methylation contexts, 100% of DMPs (*TVD* > 0.25) found by Fisher's exact test, 98.63% of DMSs

(*TVD* > 0.25) found by RMST, and 98.45% of DMCs (*TVD* > 0.25) found by HCT are identified as

DIMPs. On the other hand, DMPs only account for 30.9% of DIMPs, DMSs account for 59.8% of DIMPs,

and DMCs account for 47% of DIMPs. These observations suggest a much higher sensitivity by Methyl-

IT than other methods. DMS and DMC classes were relatively close, which helps validate our use of HD.

Results also suggest that differences in outcome between Methyl-IT and methylpy stem from signal

detection limitations rather than implementation of RMST. Application of signal detection requires

knowledge of the distribution of methylation background noise, which is not a component of the

methylpy procedure.

To evaluate whether DIMPs target genomic features in agreement with published reports [21-24], we

assessed their distribution across the genome. Fig. 5 shows DIMP distribution pattern within three major

genomic contexts (Gene regions in shades of blue, TE region in shades of red and small RNAs in shades

of green). Because total cytosine number within CHH context is about 5 times higher than CG and CHG

contexts, we have normalized data by presenting DIMP density (ratio of DIMP number at a given region /

total cytosine context number at corresponding region) rather than absolute numbers.

Results showed general agreement with the Kawakatsu et al. original study [21]. Strong methylation

changes were identified in all three contexts during the seed development process, with DIMP signal

increasing from COT to MG to PMG to dry seed, and reaching its peak in leaf tissue. CHG and CHH

changes were associated predominantly with non-genic and TE regions, and CG DIMPs showed higher

density within gene regions, which agreed with the DMP distribution pattern reported in the original

study[21]. A surprising CHG peak was observed in leaf tissues relative to seed, which we did not pursue

in detail, but may reflect a pronounced tissue-specific transition. Similar DIMP patterns were observed in

the drought stress dataset relative to cytosine context, although with higher signal levels in each context.

Hierarchical clustering based on AUC criteria and built on the set of 9893 DIMP-associated genes (using

caTools R package) permitted classification of seed developmental stages into two main phases:

morphogenesis/maturation versus dormancy (Fig. 6). In this analysis, methylation signal was expressed as

the sum of Hellinger divergence within genes plus 2kb upstream. Within the 9893-dimensional metric

space generated by 9893 AUC-selected genes, the linear cotyledon (*COT*) and mature green (*MG*) stages

(morphogenesis-maturation phase) grouped into a cluster quite distant from post mature green (*PMG*) and

dry seed (*DRY*) stages (dormancy phase). These observations indicate a detectably greater similarity in

methylome patterns between cotyledon and mature green stages, transitioning to a distinguishable state

for post-mature green and dry seed. This transition may relate to the desiccation and dormancy shift that

occurs within this timing [29, 30].

DIMPs can be predicted using a machine learning approach

An important test of DIMPs detected by the Methyl-IT pipeline is whether or not DIMPs identified within

treatment samples can be discriminated from those in the control. To address this question, machine-

learning approaches were implemented.

Each DIMP was represented as a four-dimensional vector with variables HD, TV, Weibull probability,

and cytosine relative position. The classification result for simulated data and seed development data are

presented in Table 2. Simulation experiments suggested that classification accuracy mainly depended on

the distance separating Weibull distributions (noise plus signal) for control and treatment. Weibull model

parameter values (alpha.1 and scale.1) from the first simulation for control samples (S11 to S13) were

close to those estimated in the treatment group (S21 to S23), suggesting that corresponding distribution

functions were close as well. Although the classifier performance to predict DIMPs could be considered

acceptable (about 80% accuracy), discriminatory power to predict DIMPs from an external sample (not

included to build the model) was relatively low. If probabilistic models were sufficiently distant, even a

classifier trained with samples having an overall mean TVD (absolute values of methylation differences)

equal to 0.13 could achieve good discrimination of DIMPs from an external sample. Importantly, a given

DIMP with the same HD value in control and treatment groups could be discriminated from control group

if the Weibull probability distributions from control and treatment were different.

Classification of DIMPs was accomplished for the seed development dataset as well. Since each seed

development stage comprised only one sample, groups were formed according to the hierarchical cluster

presented in Fig. 6. The best classification accuracies were obtained for CG and CHH methylation

contexts (Table 2). These were binary classifications, where control samples were the reference class.

Thus, probability P(x) that a new DIMP x could be observed in the control class determined its

classification, and the probability that a given DIMP did not classify within the control class was 1 - P(x).

A classifier model built on the groups CT: COT and MG, and non-CT: PMG and DRY (Table 2) could be

applied to classify a DIMP from the leaf stage sample as non-CT. If a DIMP from the leaf stage classified

as 'CT', this would mean that, with probability $1 - P(x) > P(x)$, its current methylation status for the

corresponding cytosine position was not distinguishable from the status observed during early seed stages.

The classifier model does not provide information for whether or not methylation status of a given

cytosine position changed across the developmental stages.

Differentially methylated genes identified by DIMPs are biologically meaningful

To investigate DIMP-based resolution of differences between seed development stages or between

stressed vs non-stressed conditions, we defined differentially methylated genes (DMGs) based on group

comparison for DIMP counts by applying generalized linear regression model (GLM). Genes displaying

statistically significant difference in DIMP number relative to control were defined as DMGs. The DMG

is defined distinctly from differentially methylated regions (DMRs), which comprise regions of high

density methylation changes. In the original study of seed methylation data, enormous DMP numbers

were identified in CHH context, corresponding to 23,195 DMRs that largely associated with transposable

elements [21]. However, DMR association with gene regions was only scant. In the drought stress dataset,

only 49 DMRs corresponding to drought stress were identified by the DSS method [24].

A total of 1068 DMGs were identified for the group comparison of morphogenesis/maturation versus

dormancy phases for seed development (Additional file 1). To investigate the biological meaning of these

DMGs, we conducted a network enrichment analysis test (NEAT). A statistically significant network

enrichment of links between genes from the set of seed development DMGs and the set of *GO-biological*

process associated with seed functions was observed (Table 3). The list of 16 networks identified includes

positive and negative regulation of GA-mediated signaling, positive and negative regulation of seed

germination, regulation of seed dormancy, and raffinose family oligosaccharide biosynthesis, all well-

established seed processes (full gene list in Additional file 2: Table S2). GeneMANIA [31] identified

interaction networks within the data, indicating that many DMGs in the seed development dataset

function together (Additional file 3: Figure S1). To test the impact of different minimum cytosine

coverage on Methyl-IT output, the pipeline was run without minimum coverage limit (Table 3) and with a

minimum coverage of 10 reads (Additional file 4: Table S3). Results were similar with either setting.

In the drought stress experiment, analyses performed by the original authors detected 2141 CG, 1039

CHG and 718 CHH DMRs, which eventually led to identification of 49 drought stress-related DMRs

[24]. A very weak relationship between methylome changes and phenotype or gene expression patterns

was suggested in the original study [24]. With Methyl-IT, we identified 6669 DMGs (Additional file 5:

Table S4). To investigate whether associations between identified DMGs and gene expression were

evident, we compared the DMG list with the differentially expressed gene (DEG) dataset reported in the

original study with 4371 genes [23]. Fig. 7a shows that the two lists shared 842 genes, accounting for

19.25% DEGs and 12.6% DMGs. Applying NEAT and Network Based Enrichment Analysis (NBEA) to

DEG and DMG datasets, we identified 73 significantly enriched DEG and 23 DMG networks. Among

them, 11 were shared and all were related to plant stress response mechanisms. Fig. 8 shows four

examples within the 11 networks, with MAPK cascade (GO:0000165), response to osmotic stress

(GO:0006970), response to salt stress (GO:0009651), and response to abscisic acid (GO:0009737). Each

gene shown carried significant DIMP signal (Additional file 5: Table S4, Additional file 6: Table S5),

suggesting that a systematic methylation repatterning had occurred within these networks. At an

individual gene level, numerous genes showed both significant gene expression and methylation changes

associated with drought stress response. For example, *ABA INSENSITIVE 1* (*ABI1*, AT4G26080) encodes

a protein involved in abscisic acid signal transduction that negatively regulates ABA promotion of

stomatal closure [32]. The locus carries 5 DIMPs on average in the three drought stressed plants, and is

up-regulated 3.36 fold. *ABRE BINDING FACTOR 4* (*ABF4*, AT3G19290), encodes a bZIP transcription

factor with specificity for abscisic acid-responsive elements (ABRE), and mediates ABA-dependent stress

responses, acting through the SnRK2 pathway [33]. This gene has an average of 8.7 DIMPs and 2.6-fold

up-regulation. *ABSCISIC ACID RESPONSIVE ELEMENTS-BINDING FACTOR 3* (*ABF3*, AT4G34000)

encodes an ABA-responsive element-binding protein with similarity to transcription factors expressed in

response to stress and abscisic acid [34]. In our study, this gene displays 11 DIMPs and is up-regulated 11

fold. *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*, AT2G46830) encodes a transcriptional repressor that

performs overlapping functions with *LHY* in a regulatory feedback loop that is closely associated with the

circadian oscillator of Arabidopsis [35]. This gene shows an average of 7.7 DIMPs and is up-regulated

38.6 fold. Taken together, these data provide enticing indication that differential gene methylation is

subtle, goes undetected by common methodologies, and identifies gene networks that are compelling

candidates for more detailed subsequent investigation.

Discussion

Methyl-IT draws from the perspective that DNA methylation functions to stabilize DNA [8, 36, 37] and,

as such, may exist in "activated-signal" versus "maintenance" states with regard to bioenergetics. The

theoretical premise underlying our approach, and based on Landauer's principle, is detailed elsewhere [12,

13], while the present study compares resolution of this methodology to current methods for analysis of

whole-genome methylation datasets. To date, there has not been a statistical biophysics model to

simulate background methylome variation. Consequently, comparisons with other methylation analysis

procedures presented here were limited to published experimental datasets.

Methyl-IT permits methylation analysis as a signal detection problem. The model predicts that most

methylation changes detected, at least in Arabidopsis, represent methylation "background noise" with

respect to methylation regulatory signal, explainable within a statistical physical probability distribution.

Implicit in our approach is that DIMPs can be detected in the control sample as well. These DIMPs are

located within the region of false alarm in Fig. 1, and correspond to natural methylation signal not

induced by treatment. Thus, using the Methyl-IT procedure, methylation signal is not only distinguished

from background noise, but can be used to discern natural signal from that induced by treatment.

Whereas methods underlying RMST (methylpy approach) and DSS provide essential information about

methylation density, context and positional changes on a genome-wide scale, Methyl-IT provides

resolution of subtle methylation repatterning signals distinct from background fluctuation. Data derived

from analysis with FET, RMST, HCT or DSS alone could lead to an assumption that gene body

methylation plays little or no role in gene expression, or that transposable elements are the primary target

of methylation repatterning. Yet ample data suggest that this picture is incomplete [38]. Methyl-IT results

show that these conclusions more likely reflect inadequate resolution of the methylome system. GLM

analysis applied to the identification of DMR-associated genes by methylpy [21] and DSS indicates that

DMRs (or DMR associated genes) do not provide sufficient resolution to link them with gene expression.

Signal detected by Methyl-IT may reflect gene-associated methylation changes that occur in response to
local changes in gene transcriptional activity. Pathway-associated methylome changes detected in seed
development data suggest participation of methylation in gene expression stage transitions, particularly
prominent between mature green and post-mature green stages. Likewise, coincident patterns between
methylome-associated gene networks and gene expression networks during drought stress appear to be
strongly non-random.

Methyl-IT analysis of various stages in seed development and germination showed evidence of
methylation changes. Previous methylpy output [21] defined predominant changes in non-CG
methylation residing within TE-rich regions of the genome, whereas Methyl-IT data resolved statistically
significant methylation signal within gene regions. With the complementary resolution provided by
Methyl-IT, it becomes possible to investigate the nature of chromatin response within identified genes in
greater detail during the various stages of a seed's development. Several of the identified DMGs in this
study involved genes that interact within known seed-associated pathways.

A limitation to currently existing methylome data analysis platforms is that most require fairly advanced
coding skills and statistics knowledge, rendering them less directly accessible to most biologists. Methyl-
IT has been designed to be highly user friendly, accessible to any biologist with basic R knowledge.

Conclusions

Methyl-IT is an alternative and complementary approach to plant methylome analysis that discriminates
DNA methylation signal from background and enhances resolution. Analysis of publicly available
methylome datasets showed enhanced signal during seed development and germination or during drought
stress within genes belonging to related pathways, providing new evidence that DNA methylation
changes occur within gene networks. Whereas, previous methylome analysis protocols identify changes in
methylome density and landscape, predominantly non-CG, Methyl-IT reveals effects within gene space,
mostly CG and CHG, for elucidation of methylome linkage to gene effects.

Methods

Methylome analysis

The alignment of BS-Seq sequence data from *Arabidopsis thaliana* was carried out with Bismark 0.15.0
[39]. BS-Seq sequence data from tomato experiment were aligned using ERNE 2.1.1 [40]. The basic
theoretical aspects of methylation analysis applied in the current work are based on previous published
results [12]. Details on Methyl-IT steps are provided in the next sections.

Methylation level estimation

In Methyl-IT pipeline, it is up to the user whether to estimate methylation levels at each cytosine position

following a Bayesian approach or not. In a Bayesian framework assuming uniform priors, the methylation

level p_i can be defined as: $p_i = \left(n_i^{mC} + 1\right) / \left(n_i^{mC} + n_i^{C} + 2\right)$ (1), where n_i^{mC} and n_i^{C} represent the numbers

of methylated and non-methylated read counts observed at the genomic coordinate i, respectively. We

estimate the shape parameters α and β from the beta distribution $P(p|\alpha,\beta) = \dfrac{p^{\alpha-1}(1-p)^{\beta-1}}{B(\alpha,\beta)}$ (2)

minimizing the difference between the empirical and theoretical cumulative distribution functions (ECDF

and CDF, respectively), where $B(\alpha,\beta)$ is the beta function with shape parameters α and β. Since the

beta distribution is a prior conjugate of binomial distribution, we consider the *p* parameter (methylation

level p_i) in the binomial distribution as randomly drawn from a beta distribution. The hyper-parameters

α and β are interpreted as pseudo counts. Then, the mean $E[p_i|D] = \hat{p}_i$ of methylation levels p_i, given

the data *D*, is expressed by $\hat{p}_i = \dfrac{\alpha + n_i^{mC}}{\alpha + \beta + n_i^{mC} + n_i^{C}}$ (3). The methylation levels at the cytosine with

genomic coordinate i are estimated according to this equation. If the Bayesian framework is not selected,

then methylation levels are estimated as: $p_i = n_i^{mc} / (n_i^{mC} + n_i^{c})$.

Hellinger and Total Variation divergences of the methylation levels

To evaluate the methylation differences between individuals from control and treatment we introduce a

metric in the bidimensional space of methylation levels: $P_i = (p_i, 1 - p_i)$. Vectors P_i provide a

measurement of the uncertainty of methylation levels at position *i*. However, we do not perform a direct

comparison between the uncertainty of methylation levels from each group of individuals, control ($\hat{p}_i^c$)

and treatment ($\hat{p}_i^t$), but the uncertainty variation with respect to the same individual reference ($\hat{p}_i^r$) on

the mentioned metric space. The reason to measure the uncertainty variation with respect to the same

reference resides in that even sibling individuals follow an independent ontogenetic development. This a

consequence of the "omnipresent" action of the second law of thermodynamics in living organisms, at

molecular level manifested throughout the actions of Brownian motion and thermal fluctuations on DNA

molecules.

The difference between methylation levels from reference and treatment (control) experiments is

expressed in terms of information divergences of their corresponding methylation levels, $\hat{p}_i^r$ and $\hat{p}_i^t$ ($\hat{p}_i^c$),

respectively. The reference sample(s) can be additional experiment(s) fixed at specific conditions, or a

virtual sample created by pooling methylation data from a set of control experiments, e.g. wild type

individual or group.

If the read counts n_i^{mC} and n_i^C are provided and taken into account, then the Hellinger divergence

between the methylation levels from reference and treatment experiments is defined as:

393
$$H\left(\hat{p}_i^r, \hat{p}_i^t\right) = w_i \left[\left(\sqrt{\hat{p}_i^r} - \sqrt{\hat{p}_i^t} \right)^2 + \left(\sqrt{1-\hat{p}_i^r} - \sqrt{1-\hat{p}_i^t} \right)^2 \right] \qquad (4)$$

394   Where $w_i = 2 \dfrac{m_i^t m_i^r}{m_i^t + m_i^r}$, $m_i^t = n_i^{mC_t} + n_i^{C_t} + 1$ and $m_r = n_i^{mCr} + n_i^{Cr} + 1$. Otherwise, Hellinger

395   divergence between the methylation levels from reference and treatment experiments is defined as:

396
$$H\left(\hat{p}_i^r, \hat{p}_i^t\right) = 2\left(\sqrt{\hat{p}_i^t} - \sqrt{\hat{p}_i^r} \right)^2 + \left(\sqrt{1-\hat{p}_i^t} - \sqrt{1-\hat{p}_i^r} \right)^2 \qquad (5)$$

The total variation of the methylation levels $TV\left(\hat{p}_i^r, \hat{p}_i^t\right) = \hat{p}_i^t - \hat{p}_i^r$ (6) indicates the direction of the

methylation change in the treatment, hypo-methylated $TV < 0$ or hyper-methylated $TV > 0$. TV is

linked to a basic information divergence, the total variation distance, defined as:

$TVD\left(\hat{p}_i^r, \hat{p}_i^t\right) = \left| TV\left(\hat{p}_i^r, \hat{p}_i^t\right) \right|$ (7) [41]. Distance $TVD\left(\hat{p}_i^r, \hat{p}_i^t\right)$ and Hellinger divergence (as given in

Eq. 4) hold the inequality: $TVD\left(\hat{p}_i^r, \hat{p}_i^t\right) \leq \dfrac{1}{\lambda_i} H\left(\hat{p}_i^r, \hat{p}_i^t\right)$ (8), where $\lambda_i = w_i/2$, which is a direct

consequence of the Cauchy-Schwarz inequality. Under the null hypothesis of non-difference between

distributions $\hat{p}_i^r$ and $\hat{p}_i^t$, Eq. 4 asymptotically has a chi-square distribution with one degree of freedom,

which set the basis for a Hellinger chi-square test (HCT). The term w_i introduces a useful correction for

the Hellinger divergence, since the estimation of $\hat{p}_i^t$ and $\hat{p}_i^r$ are based on counts (see Table 1).

In Methyl-IT pipeline, the statistics mean, median, or sum of the read counts at each cytosine site of some

control samples can be used to create a virtual reference sample. It is up to the user whether to apply the

'row sum', 'row mean' or 'row median' of methylated and unmethylated read counts at each cytosine site

across individuals.

     Non-linear fit of Weibull distribution

411   The cumulative distribution functions (CDF) for $H_k\left(\hat{p}_k^r, \hat{p}_k^t\right)$ can be approached by a Weibull

412   distribution $P\left(H_k \leq H^0 \mid \alpha, \lambda, \mu\right) = 1 - e^{-\left(\frac{H_k - \mu}{\lambda(l)}\right)^{\alpha}}$ (9) [12]. Parameter $\hat{\alpha}, \hat{\lambda}$ and $\hat{\mu}$ were estimated by non-

413   linear regression analysis of the ECDF $\hat{F}_n\left(\hat{H}_k \leq H^0\right)$ versus $H_k\left(\hat{p}_k^r, \hat{p}_k^t\right)$ [12]. The ECDF of the

414   variable $\hat{H}_k$ is defined as:

415
$$\hat{F}_n\left(\hat{H}_k \leq H^0\right) = \frac{\text{number of CDMs in the samples with } \hat{H}_k \leq H}{n} = \frac{1}{n}\sum_{k=1}^{n} 1_{\hat{H}_k \leq H^0} \quad (10)$$

, where $1_{\hat{H}_k \leq H^0} = \begin{cases} 1 \text{ if } \hat{H}_k \leq H^0 \\ 0 \text{ if } \hat{H}_k > H^0 \end{cases}$ is the indicator function. Function $\hat{F}_n\left(\hat{H}_k \leq H^0\right)$ is easily computed

(for example, by using function "*ecdf*" of the statistical computing program "R"[42]).

A statistical mechanics-based definition for a potential/putative methylation signal (PMS)

Most methylation changes occurring within cells are likely induced by thermal fluctuations to ensure

thermal stability of the DNA molecule, conforming to laws of statistical mechanics [12]. These changes

do not constitute biological signals, but methylation background noise induced by thermal fluctuations,

and must be discriminated from changes induced by the treatment. Let $P\left(E_k^D \leq E_k^{D_0}\right)$ be the probability

that energy E_k^D, dissipated to create an observed divergence D between the methylation levels from two

different samples at a given genomic position k, can be lesser than or equal to the amount of energy $E_k^{D_0}$.

Then, a single genomic position k shall be called a PMS at a level of significance α if, and only if, the

probability $P\left(E_k^D > E_k^{D_0}\right) = 1 - P\left(E_k^D \leq E_k^{D_0}\right)$ to observe a methylation change with energy dissipation

higher than $E_k^{D_0}$ is lesser than α. The probability $P\left(E_k^D \leq E_k^{D_0}\right)$ can be given by a member of the

generalized gamma distribution family and, in most cases, experimental data can be fixed by the Weibull

distribution [12]. Based on this dynamic nature of methylation, one cannot expect a genome-wide

relationship between methylation and gene expression. A practical definition of PMS based on Hellinger

divergence derives provided that H_k is proportional to E_k^H and using the estimated Weibull CDF for

H_k given by Eq. 8. That is, a single genomic position k shall be called a PMS at a level of significance

α if, and only if, the probability $\hat{P}\left(H_k > H^0 \middle| \hat{\alpha}, \hat{\lambda}, \hat{\mu}\right) = 1 - \hat{P}\left(H_k \le H^0 \middle| \hat{\alpha}, \hat{\lambda}, \hat{\mu}\right)$ to observe a

methylation change with Hellinger divergence higher than H_k is lesser than α .

The PMSs reflect cytosine methylation positions that undergo changes without discerning whether they

represent biological signal created by the methylation regulatory machinery. The application of signal

detection theory is required for robust discrimination of biological signal from physical noise-induced

thermal fluctuations, permitting a high signal-to-noise ratio [18].

439   *Robust detection of differentially informative methylated positions (DIMPs)*

Application of signal detection theory is required to reach a high signal-to-noise ratio [43, 44]. To

enhance DIMP detection, the set of PMSs is reduced to the subset of cytosines with

$TVD\left(\hat{p}_i^r, \hat{p}_i^t\right) \le TVD_0$, where TVD_0 is a minimal total variation distance defined by the user, preferably

$TVD_0 > 0.1$. If we are interested not only in DIMPs but also in the full spectrum of biological signals,

this constraint is not required. Once potential DIMPs are estimated in the treatment and in the control

samples, a logistic regression analysis is performed with the prior binary classification of DIMPs, i.e., in

terms of PMSs (from treatment versus control), and a receiver operating curve (ROC) is built to estimate

the cutpoint of the Hellinger divergence at which an observed methylation level represents a true DIMP.

There are several criteria to estimate the optimal cutpoint, many of which are implemented in the R

package *OptimalCutpoints* [27]. The optimal cutpoint used in Methyl-IT corresponds to the *H* value that

maximizes Sensitivity and Specificity simultaneously [45, 46]. These analyses were performed with the R

package *Epi* [47].

Once all pairwise comparisons are done, a final decision of whether a DFMP is a DIMP is taken based on

the highest cutpoint detected in the ROC analyses (Fig. 1). That is, the decision is taken based on the

cutpoint estimated in the ROC analysis for the control sample with the closest distribution to treatment

samples. The position of the cutpoint will determine a final posterior classification for which we would

estimate the number of true positive, true negatives, false positives and false negatives. For each cutpoint

we would estimate, the accuracy and the risk of our predictions. We may wish to use different cutpoints

for different situations. For example, if our goal is the early detection of a terminal disease and high

values of the target variable indicates that a patient carries the disease, then to save lives we would prefer

the lowest meaningful cutpoint reducing the rate of false negative.

461   *DIMP simulation and machine learning classifier*

Methyl-IT pipeline was applied to seven random generated individual samples, each on with 2×10^5

simulated cytosine positions with their corresponding methylation levels. A reference individual sample

was generated with parameters $\alpha = 1.54$ and $\beta = 2$ with mean of methylation levels $E[\hat{p}] = 0.435$ and

variance $Var[\hat{p}] = 0.0541$. Two simulation experiments were performed. For the first simulation, total

variations values for three control samples (S11 to S13) were generated using normal distribution with

means (standard deviation): 0.297 (0.31), 0.297 (0.32), and 0.295 (0.34) and for three treatment (S21 to

S23) individual with means (standard deviation): 0.44 (0.3), 0.45 (0.33), and 0.43 (34). The overall mean

of all the pairwise differences of methylation levels between control and treatment sample is 0.03.

TV treatment means were increase in the second simulation with values: 0.54, 0.55, and 0.53. The overall

mean of all the pairwise differences of methylation levels between control and treatment sample is 0.13.

DIMPs were estimated according to Methyl-IT pipeline and a classifier model was inferred with the three

control samples and the first two treatment samples to classify DIMPs into two classes: control (CT) and

treatment (non-CT or 'TT'). Each cytosine site is represented as a four dimensional vector with variables:

HD, TV, Weibull probability, and cytosine relative position estimated as $(x - x_{min})/(x_{max} - x)$, where x_{min}

and x_{max} are the maximum and minimum positions for the corresponding chromosome.

The set of four dimensional vectors integrated by control and treatment was randomly split into two

subsets: training (60%, used to train the model) and test (40%, used to evaluate the classifier). The

classification performance was evaluated with Monte Carlo resampling and the classifier model was

applied to predict DIMPs from the third treatment sample not included in the construction of the classifier

model. In the case of Monte Carlo resampling, a new random split of the samples is performed for each

resampling.

Currently, there are seven classifiers available to use with Methyl-IT: logistic regression model (LRM),

linear discriminant analysis (LDA), quadratic discriminant analysis (QDA), support vector machine

(SVM), PCA-LRM using the principal component (PCA) as predictor variables in LMR, PCA-LDA and

PCA-QDA.

Estimation of differentially methylated genes (DMGs) using Methyl-IT

Our degree of confidence in whether DIMP counts in both control and treatment represent true biological

signal was set out in the signal detection step. To estimate DMGs, we followed similar steps to those

proposed in Bioconductor R package DESeq2 [48], but the test looks for statistical difference between the

groups based on gene body DIMP counts rather than read counts. The regression analysis of the

generalized linear model (GLMs) with logarithmic link was applied to test the difference between group

counts. The fitting algorithmic approaches provided by *glm* and *glm.nb* functions from the R packages

stat and MASS were used for Poisson (PR), Quasi-Poisson (QPR) and Negative Binomial (NBR) linear

regression analyses, respectively.

Likewise for DESeq2 we used the linear regression model $\log_2\left(q_{ij}\right) = \sum_k x_{jk}\beta_{ik}$, with design matrix

elements x_{jk}, coefficients β_{ik}, and mean $\mu_{kj} = s_j q_{kj}$, where s_j normalization constants are considered

constant within a group. Only two groups were compared at a time. The design matrix elements indicate

whether a sample j is treated or not, and the GLM fit returns coefficients indicating the overall

methylation strength at the gene and the logarithm base 2 of the fold change (log2FC) between treatment

and control [48]. In particular, in the case of NBR, the inverse of the variance was used as prior weight

($\sigma_{jk}^2 = \dfrac{1}{\mu_{ij} + \mu_{ij} disp}$, where *disp* is data dispersion computed by the *estimateDispersions* function from

DESeq2 R package).

To test difference between group counts we applied the fitting algorithmic approaches: PR and PQR if

$\rho < \dfrac{\mu_{ij}}{\sigma_{ij}} \leq 1\,(0.9 < \rho < 1)$, NBR and NBR with '*prior weights*'. Next, best model based on Akaike

information criteria (AIC). The Wald test for significance of the independent variable coefficient indicates

whether or not the treatment effect is significant, while the coefficient sign (*log2FC*) will indicate the

direction of such an effect.

Bootstrap goodness-of-fit test for 2x2 contingency tables

The goodness-of-fit RMST 2x2 contingency tables as implemented in methylpy [20] for the estimation of

DMSs (based on the root-mean-square (RMS) statistics) is explained in Perkins et al. in reference [49](a

complemental description is found at arXiv:1108.4126v2). The bootstrap heuristic to perform the test is

given in reference [50]. An analogous bootstrap goodness-of-fit test based on Hellinger divergence was

also applied to estimate DMCs. In this case, Hellinger divergence estimated according to the first statistic

given in Theorem 1 from reference [51].

Network enrichment analysis

Network based enrichment analysis (NBEA) was applied using the EnrichmentBrowser R package [52,

53] and the Network Enrichment Analysis Test (NEAT) was performed by using the R package "neat"

version 1.1.1[53].

Abbreviations

AUC: Area under the receiver operating characteristic curve

CDM: Cytosine DNA methylation

DAGs: DMR associated genes

DEG: Differentially expressed gene

DIMPs: Differentially informative methylated positions

DMGs: Differentially methylated genes

DMPs: Differentially methylated positions

DMRs: differentially methylated regions

DSS: Dispersion Shrinkage for Sequencing

FET: Fisher's exact test

GLM: generalized linear regression model

HD: Hellinger divergence

HCT: Hellinger chi-square test. Goodness-of-fit test based on Hellinger divergence

NEAT: Network Enrichment Analysis Test

NBEA: Network based enrichment analysis

RMST: Root-mean-square test

ROC: Receiver operating characteristic curve

SD: Signal detection

TVD: total variation distance

PMS: Potential/putative methylation signal

Declarations

Acknowledgments

We thank Diep Ganguly from Australian National University provides guidance on the use of drought

stress data. We thank Professor David Miller from Department of electrical engineering for helpful

discussions.

Funding

The work was supported by funding from the Bill and Melinda Gates Foundation (OPP1088661).

Availability of data and materials

The source code for all of analysis and visualization, including The Methyl-IT package source code,

Network enrichment analysis, R code for all figures are available at the GitLab:

https://git.psu.edu/genomath/MethylIT. Seed development methylome dataset, original studied by

Kawakatsu et al. (2017) [21], was obtained from the Gene Expression Omnibus (GEO) under accession

numbers GSE68132. Drought stress methylome dataset was obtained from the Gene Expression Omnibus

(GEO) under accession numbers GSE94075.The differential expressed gene list and express level was

obtained Crisp et al. [23].

Authors' contributions

RS developed the application of the information thermodynamic theory on cytosine DNA methylation

and conducted mathematical and computational biology analyses, XY participated in methylome data

analysis, JRB conducted computation, HK conducted the NBEA, NEAT analysis. SM designed

experiments, participated in data analysis and wrote manuscript.

Competing interests

Not applicable

Consent for publication

Not applicable

Ethics approval and consent to participate

Not applicable

Figures

577
578
579
Fig. 1 Comparison of three variables used to measure DNA cytosine methylation.

The heatmap for CG methylation distribution represented by **(a)** methylation level (percentage), **(b)** total

variation distance (TVD), and **(c)** Hellinger divergence (HD) on chromosome 1 for the drought stress

experimental data are shown. Chromosomes were split into 2-kb non-overlapping windows (regions).

The mean of methylation levels for each region i was estimated as: $p_i = \sum_{j=1}^{2kb} mC_{ij} / \sum_{ij=1}^{2kb}(mC_{ij} + uC_{ij})$,

while $TVD_i = \sum_{j=1}^{2kb} TVD_{ij}$ and $HD_i = \sum_{j=1}^{2kb} HD_{ij}$.

586
587

588
589
590
591   **Fig. 2** Schematic of the theoretical principle underlying Methyl-IT. Methyl-IT is designed to identify a

592   statistically significant cutoff between thermal system noise (conforming to laws of statistical physics)

593   and treatment signal (biological methylation signal), based on Hellinger divergence (H), to identify "true"

594   differentially informative methylation positions (DIMPs). Empirical comparisons allow the placement of

595   Fisher's exact test for discrimination of DMPs.

596
597
598
599

600
601
Fig. 3 Methyl-IT processing flowchart. Ovals represent input and output data, squares represent

processing steps, with signal detection processing steps highlighted in blue and DIMPs and DMGRs, as

main outputs of Methyl-IT, highlighted in yellow. The generalized linear model is incorporated for group

comparison of genomic regions (GRs) based on the number of DIMPs in the treatment group relative to

control group. DIMPs and DMGRs can be subjected to further statistical analyses to perform network

enrichment analysis and to identify potential signature genes, multivariate statistical analysis (and

machine learning applications) for individual and group classifications.

609
610
611
612

614   **Fig. 4** Venn diagrams of overlapping DMSs (RMST implemented in methylpy software), DMPs

615   (obtained with Fisher Exact Test), DMCs (obtained with HCT, see methods) and DIMPs (obtained with

616   Methyl-IT) for the drought experimental data. Only methylated cytosine positions with total variation

617   distance (TVD) greater than 0.25 (25% of methylation level difference) are shown for the three

618   methylation contexts. DIMPs carrying methylation signal are in the region within the dashed oval.

Fig. 5 Results of signal detection with Methyl-IT for genome-wide methylome data from seed development samples from Kawakatsu et al [21] at five seed stages (GLOB, COT, MG, PMG, DRY) and leaf (globular (GLOB) stage used as control), and drought stress experiment control (CTR) and stress (STR) samples from Ganguly et al. [24]. The experimental results provide a direct, scaled comparison of methylation signal between datasets. The relative frequency of DIMPs was estimated as the number of DIMPs divided by the number of cytosine positions.

637
638
639 **Fig.** 6 Classification of seed development stages based on identified DIMPs. A hierarchical cluster built
640 on the set of 7006 selected DIMP-associated genes, based on AUC criteria, classified the stages into two
641 groups: morphogenesis-maturation phase and dormancy phase.
642
643

644
645 **Fig. 7** Differential expressed genes(DEGs) vs differential methylated genes(DMGs) in unstressed plants
646 vs drought stressed plants comparison. **(a)** 4371 DEGs were identified by [24] and 6669 DMGs were
647 identified by Methyl-IT. **(b)** 73 and 23 significantly enriched networks were identified from 4371 DEGs
648 and 6669 DMGs, respectively. NEAT and NBEA analysis were used to identify enriched network (see
649 Method).
650

Fig. 8 Examples of enriched networks identified by NBEA using DMGs. Genes involved in **(a)** MAPK

cascade, **(b)** response to salt stress, **(c)** response to osmotic stress, and **(d)** response to abscisic acid are

in circles. Network graphs were generated by EnrichmentBrowser R package in R. Details of DIMPs

number for each gene could be found in additional file 6.

Tables

Table 1. Relative sensitivity differences between several statistical tests applied to identify differentially

methylated cytosines. P-values for the 2x2 contingency table with read counts $n_i^{mC_c} = 8$, $n_i^{C_c} = 2$,

$n_i^{mC_t} = 350$, and $n_i^{C_t} = 20$.

Approach	*p*-value	Significance(α = 0.05)
FET	0.108615	No
FET one tail	0.108615	No
FET *p*.value MC 3k[(1)]	0.1086	No
RMST Boot 3k[(2)]	0.051	No
HT Boot 3k [(3)]	0.050667	No
Weibull STR1 CG[(4)]	5.08E-04	Yes
Weibull STR2 CG[(4)]	3.20E-04	Yes
Weibull STR3 CG[(4)]	3.20E-04	Yes

[1]*p*.value simulated with Monte Carlo (MC) simulation with 3000 resamplings (3k). [2]Bootstrap goodness-of-fit

RMST as implemented in methylpy [20]. [3]Bootstrap goodness-of-fit test based on Hellinger divergence estimated

according to the first statistic given Theorem 1 from reference [51]. [3]*p*-value based on the Weibull distribution for

memory lines (STR 1 to 3). n_i^{mCc} refers to methylated cytosine counts in control, n_i^{Cc} refers to non-methylated

cytosine counts in control, n_i^{mCt} refers to methylated cytosine counts in treatment and n_i^{Ct} refers to non-methylated

cytosine counts in treatment.The R script to compute RMST and *H* MC estimation is provided in GitLab:

https://git.psu.edu/genomath/MethylIT

Table 2. Classification of DIMPs into two classes: control (CT) and non-control (non-CT)

Parameters from the best fitted Weibull model estimated for simulate data							
Parameters	**[1]S11**	**S12**	**S13**	**S21**	**S22**	**S23**	**TVD mean**
alpha.1[2]	0.645650	0.645195	0.645586	0.649747	0.651112	0.650272	0.03
scale.1	0.249118	0.253707	0.253919	0.257382	0.268151	0.258988	
alpha.2[3]	0.645650	0.645195	0.645586	0.656598	0.654718	0.652522	0.13
scale.2	0.249118	0.253707	0.253919	0.290850	0.300384	0.290910	

Performance of classifier models built on simulated data							
Classifier model	**Accuracy**	**[4]MC. Accuracy**	**Sensitivity**	**Specificity**	**Pos. Pred**	**Neg. Pred**	**CT/non-CT (S23)**
PCA-QDA.1	0.8088	0.8101	0.537	0.9763	0.9483	0.7592	1213/**378**
PCA-QDA.2	0.9997	0.9998	1	0.9994	0.9995	1	0/**2508**

Performance of classifier models built on CT: COT and MG, and non-CT: PMG and DRY							
Methylation Context	**Classif. Model**	**Accuracy**	**Sensitivity**	**Specificity**	**Pos Pred**	**Neg. Pred**	**Predictions for LEAF**
							CT/non-CT
CG	Logistic	0.9011	0.9984	0.7222	0.8685	0.996	18/**166186**
CHG	Logistic	0.7541	0.8842	0.5574	0.7512	0.7611	3174/**205463**
CHH	PCA-QDA	0.9074	0.9716	0.671	0.9158	0.865	69102/**3906**

674 [1]Simulated samples were denoted S11, S12… S23 (S11 to S13 are control, the remainder treatment). [2]1st

675 simulation experiment. [3] 2nd simulation experiment. [4] Accuracy mean for 500 Monte Carlos resamplings.

678 **Table 3.** Network enrichment analysis test (NEAT) on the set of GO-biological process (BP-GO) for the

679 differentially methylated genes in Ws-0 seed development dataset.

BP-GO	NAB	Expected NAB	Adj. *p*-value
GO:0000902 cell morphogenesis	3	0.2492	0.00280
GO:0006623 protein targeting to vacuole	4	0.299	< 0.001
GO:0006891 intra-Golgi vesicle-mediated transport	4	0.3323	< 0.001
GO:0009723 response to ethylene	8	2.9072	0.00873
GO:0009740 gibberellic acid mediated signaling pathway	5	0.9802	0.00375
GO:0009845 seed germination	6	1.3456	0.00301
GO:0009938 negative regulation of gibberellic acid mediated signaling pathway	4	0.2658	< 0.001
GO:0010162 seed dormancy process	5	1.03	0.00434
GO:0010187 negative regulation of seed germination	3	0.4319	0.00916
GO:0010325 raffinose family oligosaccharide biosynthetic process	5	0.3323	0.00102
GO:0016049 cell growth	3	0.3655	0.00640
GO:0016192 vesicle-mediated transport	5	0.3987	< 0.001
GO:0016197 endosomal transport	2	0.0665	0.00280
GO:0048444 floral organ morphogenesis	5	0.3323	< 0.001
GO:2000033 regulation of seed dormancy process	3	0.1994	0.0017
GO:2000377 regulation of reactive oxygen species metabolic process	4	0.4153	0.00154

682 Only over-enriched pathways are included

683 NAB: observed number of (network) links from DMG list to GO term gene list

684 Expected NAB: expected number of links from DMG list to GO term gene list (in absence of enrichment)

685 Enrichment Fold: the ratio of NAB (observed number of network links) / expected nab (expected number of links)

Table 4. Overlapped pathways between DEGs and DMGs in drought stress data

No.	Gene Ontology term
1	GO:0000165 MAPK cascade
2	GO:0006970 response to osmotic stress
3	GO:0009409 response to cold
4	GO:0009651 response to salt stress
5	GO:0009723 response to ethylene
6	GO:0009737 response to abscisic acid
7	GO:0009862 systemic acquired resistance, salicylic acid mediated signaling pathway
8	GO:0009863 salicylic acid mediated signaling pathway
9	GO:0009867 jasmonic acid mediated signaling pathway
10	GO:0031348 negative regulation of defense response

| 11 | GO:0042742 defense response to bacterium |

Additional files

Additional file 1: Table.S1   DMGs from Arabidopsis seed development dataset.

Additional file 2: Table.S2   List of seed develoment DMGs found_in networks based on NEAT.

Additional file 3: Figure.S1 Interaction network built for the seed development DMGs in networks identified with NEAT using GeneMNIA.

Additional file 4: Table.S3 Enriched network from seed development DMGs (with minimum coverage 10 reads).

Additional file 5: Table.S4 List of 6669 DMGs identifed in the drought stress experiment.

Additional File 6: Table. S5 DMGs in the enriched networks identifid by NBEA for the drought stress data.

References

1.   Calarco JP, Borges F, Donoghue MTA, Van Ex F, Jullien PE, Lopes T, Gardner R, Berger F, Feijo JA, Becker JD et al: Reprogramming of DNA Methylation in Pollen Guides Epigenetic Inheritance via Small RNA. Cell 2012, 151(1):194-205.
2.   Schmitz RJ, Schultz MD, Lewsey MG, O'Malley RC, Urich MA, Libiger O, Schork NJ, Ecker JR: Transgenerational Epigenetic Instability Is a Source of Novel Methylation Variants. Science 2011, 334(6054):369-373.
3.   Becker C, Hagmann J, Muller J, Koenig D, Stegle O, Borgwardt K, Weigel D: Spontaneous epigenetic variation in the Arabidopsis thaliana methylome. Nature 2011, 480(7376):245-U127.
4.   Matzke MA, Mosher RA: RNA-directed DNA methylation: an epigenetic pathway of increasing complexity (vol 15, 394, 2014). Nat Rev Genet 2014, 15(8).
5.   Crisp PA, Ganguly D, Eichten SR, Borevitz JO, Pogson BJ: Reconsidering plant memory: Intersections between stress recovery, RNA turnover, and epigenetics. Sci Adv 2016, 2(2).
6.   Kinoshita T, Seki M: Epigenetic Memory for Stress Response and Adaptation in Plants. Plant Cell Physiol 2014, 55(11):1859-1863.

7.   Colaneri AC, Jones AM: Genome-Wide Quantitative Identification of DNA
Differentially Methylated Sites in Arabidopsis Seedlings Growing at Different Water
Potential. Plos One 2013, 8(4).
8.   Severin PM, Zou X, Gaub HE, Schulten K: Cytosine methylation alters DNA
mechanical properties. Nucleic Acids Res 2011, 39(20):8740-8751.
9.   Osakabe A, Adachi F, Arimura Y, Maehara K, Ohkawa Y, Kurumizaka H: Influence of
DNA methylation on positioning and DNA flexibility of nucleosomes with pericentric
satellite DNA. Open Biol 2015, 5(10).
10.   Yusufaly TI, Li Y, Olson WK: 5-Methylation of cytosine in CG:CG base-pair steps: a
physicochemical mechanism for the epigenetic control of DNA nanomechanics. J
Phys Chem B 2013, 117(51):16436-16442.
11.   Jenkinson G, Pujadas E, Goutsias J, Feinberg AP: Potential energy landscapes identify
the information-theoretic nature of the epigenome. Nat Genet 2017, 49(5):719-+.
12.   Sanchez R, Mackenzie SA: Information Thermodynamics of Cytosine DNA
Methylation. Plos One 2016, 11(3).
13.   Sanchez R, Mackenzie SA: Genome-Wide Discriminatory Information Patterns of
Cytosine DNA Methylation. Int J Mol Sci 2016, 17(6).
14.   Greiner M, Pfeiffer D, Smith RD: Principles and practical application of the receiver-
operating characteristic analysis for diagnostic tests. Prev Vet Med 2000, 45(1-
2):23-41.
15.   Carter JV, Pan J, Rai SN, Galandiuk S: ROC-ing along: Evaluation and interpretation of
receiver operating characteristic curves. Surgery 2016, 159(6):1638-1645.
16.   Harpaz R, DuMouchel W, LePendu P, Bauer-Mehren A, Ryan P, Shah NH:
Performance of pharmacovigilance signal-detection algorithms for the FDA adverse
event reporting system. Clin Pharmacol Ther 2013, 93(6):539-546.
17.   Kruspe S, Dickey DD, Urak KT, Blanco GN, Miller MJ, Clark KC, Burghardt E, Gutierrez
WR, Phadke SD, Kamboj S et al: Rapid and Sensitive Detection of Breast Cancer Cells
in Patient Blood with Nuclease-Activated Probe Technology. Mol Ther Nucleic Acids
2017, 8:542-557.
18.   Kay SM: Fundamentals of Statistical Signal Processing, Volume II: Detection Theory,
1 edition edn; 1998.
19.   Wu H, Xu T, Feng H, Chen L, Li B, Yao B, Qin Z, Jin P, Conneely KN: Detection of
differentially methylated regions from whole-genome bisulfite sequencing data
without replicates. Nucleic Acids Res 2015, 43(21):e141.
20.   Schultz MD, He Y, Whitaker JW, Hariharan M, Mukamel EA, Leung D, Rajagopal N,
Nery JR, Urich MA, Chen H et al: Human body epigenome maps reveal noncanonical
DNA methylation variation. Nature 2015, 523(7559):212-216.
21.   Kawakatsu T, Nery JR, Castanon R, Ecker JR: Dynamic DNA methylation
reconfiguration during seed development and germination. Genome Biol 2017,
18(1):171.
22.   Schmitz RJ, Schultz MD, Urich MA, Nery JR, Pelizzola M, Libiger O, Alix A, McCosh RB,
Chen H, Schork NJ et al: Patterns of population epigenomic diversity. Nature 2013,
495(7440):193-198.
23.   Crisp PA, Ganguly DR, Smith AB, Murray KD, Estavillo GM, Searle I, Ford E,
Bogdanovic O, Lister R, Borevitz JO et al: Rapid Recovery Gene Downregulation

during Excess-Light Stress and Recovery in Arabidopsis. Plant Cell 2017,
29(8):1836-1863.
24. Ganguly DR, Crisp PA, Eichten SR, Pogson BJ: The Arabidopsis DNA Methylome Is
Stable under Transgenerational Drought Stress. Plant Physiol 2017, 175(4):1893-
1912.
25. Basu A, Mandal A, Pardo L: Hypothesis testing for two discrete populations based on
the Hellinger distance. Stat Probabil Lett 2010, 80(3-4):206-214.
26. Vaart A: Asymptotic Statistics: Cambridge University Press. ; 1998.
27. Mónica López-Ratón MXR-Á, Carmen Cadarso-Suárez, Francisco Gude-Sampedro:
OptimalCutpoints: An R Package for Selecting Optimal Cutpoints in Diagnostic Tests.
Journal of statistical software 2014, Vol 61 (2014)(Issue 8):4896.
28. Perkins W, Tygert M, Ward R: Computing the confidence levels for a root-mean-
square test of goodness-of-fit. Applied Mathematics and Computation 2011,
217(22):9072-9084.
29. Le BH, Cheng C, Bui AQ, Wagmaister JA, Henry KF, Pelletier J, Kwong L, Belmonte M,
Kirkbride R, Horvath S et al: Global analysis of gene activity during Arabidopsis seed
development and identification of seed-specific transcription factors. Proc Natl Acad
Sci U S A 2010, 107(18):8063-8070.
30. Bassel GW, Lan H, Glaab E, Gibbs DJ, Gerjets T, Krasnogor N, Bonner AJ, Holdsworth
MJ, Provart NJ: Genome-wide network model capturing seed germination reveals
coordinated regulation of plant cellular phase transitions. Proc Natl Acad Sci U S A
2011, 108(23):9709-9714.
31. Mostafavi S, Ray D, Warde-Farley D, Grouios C, Morris Q: GeneMANIA: a real-time
multiple association network integration algorithm for predicting gene function.
Genome Biol 2008, 9 Suppl 1:S4.
32. Krzywinska E, Bucholc M, Kulik A, Ciesielski A, Lichocka M, Debski J, Ludwikow A,
Dadlez M, Rodriguez PL, Dobrowolska G: Phosphatase ABI1 and okadaic acid-
sensitive phosphoprotein phosphatases inhibit salt stress-activated SnRK2.4 kinase.
Bmc Plant Biol 2016, 16(1):136.
33. Yoshida T, Fujita Y, Maruyama K, Mogami J, Todaka D, Shinozaki K, Yamaguchi-
Shinozaki K: Four Arabidopsis AREB/ABF transcription factors function
predominantly in gene expression downstream of SnRK2 kinases in abscisic acid
signalling in response to osmotic stress. Plant Cell Environ 2015, 38(1):35-49.
34. Yoshida T, Fujita Y, Sayama H, Kidokoro S, Maruyama K, Mizoi J, Shinozaki K,
Yamaguchi-Shinozaki K: AREB1, AREB2, and ABF3 are master transcription factors
that cooperatively regulate ABRE-dependent ABA signaling involved in drought
stress tolerance and require ABA for full activation. Plant J 2010, 61(4):672-685.
35. Yakir E, Hilman D, Hassidim M, Green RM: CIRCADIAN CLOCK ASSOCIATED1
transcript stability and the entrainment of the circadian clock in Arabidopsis. Plant
Physiol 2007, 145(3):925-932.
36. Lefebvre A, Mauffret O, el Antri S, Monnot M, Lescot E, Fermandjian S: Sequence
dependent effects of CpG cytosine methylation. A joint 1H-NMR and 31P-NMR study.
Eur J Biochem 1995, 229(2):445-454.
37. Nathan D, Crothers DM: Bending and flexibility of methylated and unmethylated
EcoRI DNA. J Mol Biol 2002, 316(1):7-17.

38.   Huang SC, Ecker JR: Piecing together cis-regulatory networks: insights from
814   epigenomics studies in plants. Wiley Interdiscip Rev Syst Biol Med 2017.
815   39.   Krueger F, Andrews SR: Bismark: a flexible aligner and methylation caller for
816   Bisulfite-Seq applications. Bioinformatics 2011, 27(11):1571-1572.
817   40.   Prezza N, Vezzi F, Kaller M, Policriti A: Fast, accurate, and lightweight analysis of BS-
818   treated reads with ERNE 2. Bmc Bioinformatics 2016, 17.
819   41.   Sason I, Verdu S: f-Divergence Inequalities. Ieee Transactions on Information Theory
820   2016, 62(11):5973-6006.
821   42.   R_Core_Team: A language and environment for statistical computing. 2016.
822   43.   Hippenstiel RD: Detection Theory: Applications and Digital Signal Processing. CRC
823   Press 2001.
824   44.   Stanislaw H, Todorov N: Calculation of signal detection theory measures. Behav Res
825   Meth Ins C 1999, 31(1):137-149.
826   45.   Youden WJ: Index for rating diagnostic tests. Cancer 1950, 3(1):32-35.
827   46.   Perkins NJ, Schisterman EF: The inconsistency of "optimal" cutpoints obtained using
828   two criteria based on the receiver operating characteristic curve. Am J Epidemiol
829   2006, 163(7):670-675.
830   47.   Carstensen B, Plummer, M., Laara, E. & Hills, M.: Epi:A Package for Statistical
831   Analysis in Epidemiology. R package version 27 2016.
832   48.   Love MI, Huber W, Anders S: Moderated estimation of fold change and dispersion
833   for RNA-seq data with DESeq2. Genome Biol 2014, 15(12).
834   49.   William Perkins MT, Rachel Ward: Computing the confidence levels for a root-mean-
835   square test of goodness-of-fit. Applied Mathematics and Computation 2011, Volume
836   217(issue 22):Pages 9072-9084.
837   50.   He Y, Gorkin DU, Dickel DE, Nery JR, Castanon RG, Lee AY, Shen Y, Visel A,
838   Pennacchio LA, Ren B et al: Improved regulatory element prediction based on
839   tissue-specific local epigenomic signatures. Proc Natl Acad Sci U S A 2017,
840   114(9):E1633-E1640.
841   51.   F. Liese IV: On Divergences and Informations in Statistics and Information Theory.
842   IEEE Transactions on Information Theory 2006, Volume: 52(Issue: 10):4394 - 4412.
843   52.   Geistlinger L, Csaba G, Zimmer R: Bioconductor's EnrichmentBrowser: seamless
844   navigation through combined results of set- & network-based enrichment analysis.
845   Bmc Bioinformatics 2016, 17.
846   53.   Signorelli M, Vinciotti V, Wit EC: NEAT: an efficient network enrichment analysis
847   test. Bmc Bioinformatics 2016, 17.
848