# Natural variation in stomata size contributes to the local adaptation of water-use efficiency in *Arabidopsis thaliana*

H. Dittberner[1], A. Korte[2], T. Mettler-Altmann[3], A.P.M. Weber[3], G. Monroe[4], J. de Meaux[1]

Affiliations: [1]Institute of Botany, University of Cologne, 50674 Cologne; [2]Center for Computational and Theoretical Biology, Julius-Maximilians-University Würzburg, 97074 Würzburg; [3]Institute of Plant Biochemistry & CEPLAS Plant Metabolism and Metabolomics Laboratory, Heinrich-Heine-University Düsseldorf, 40225 Düsseldorf; [4]College of Agricultural Sciences, Colorado State University, Fort Collins, CO 80523-1101

## Abstract

Stomata control gas exchanges between the plant and the atmosphere. How natural variation in stomata size and density contributes to resolve trade-offs between carbon uptake and water-loss in response to local climatic variation is not yet understood. We developed an automated confocal microscopy approach to characterize natural genetic variation in stomatal patterning in 330 fully-sequenced *Arabidopsis thaliana* accessions collected throughout the European range of the species. We compared this to variation in water-use efficiency, measured as carbon isotope discrimination ($\delta^{13}$C). We detect substantial genetic variation for stomata size and density segregating within *Arabidopsis thaliana*. A positive correlation between stomata size and $\delta^{13}$C further suggests that this variation has consequences on water-use efficiency. Genome-wide association analyses indicate a complex genetic architecture underlying not only variation in stomata patterning but also to its co-variation with carbon uptake parameters. Yet, we report two novel QTL affecting $\delta^{13}$C independently of stomata patterning. This suggests that, in *A. thaliana*, both morphological and physiological variants contribute to genetic variance in water-use efficiency. Patterns of regional differentiation and co-variation with climatic parameters indicate that natural selection has contributed to shape some of this variation, especially in Southern Sweden, where water availability is more limited in spring relative to summer. These conditions are expected to favor the evolution of drought avoidance mechanisms over drought escape strategies.

## Keywords

GWAS; stomata; water-use efficiency; Arabidopsis thaliana; $Q_{ST}$ $F_{ST}$ analysis; local adaptation to climate

# 1  Introduction

In plants, carbon uptake and water loss are intimately linked by a trade-off between growth

and water conservation (Cowan, 1986; Cowan & Farquhar, 1977; Field, Merino, & Mooney,

1983). Stomata, the microscopic pores embedded in the epidermis of plant leaves, play a key

role in the resolution of this trade-off.  Their density, distribution and regulation control the

rate of $CO_2$ and water exchange (Raven, 2002). As a result, they impact the ratio of

photosynthetic carbon assimilation to water loss *via* transpiration. This ratio defines water-use

efficiency (WUE), a physiological parameter that directly determines plant productivity when

the water supply is limited. Stomata may thus have played a pivotal role in shaping the

diversity of plant communities throughout the globe (Lambers, Chapin, & Pons, 1998;

McDowell et al., 2008).

The density of stomata on the leaf surface is expected to correlate positively with the rate of

gas exchanges between the leaf and the atmosphere, also called "conductance". Models based

on gas diffusion theory predict that small stomata in high density can best maximize

conductance (Franks & Beerling, 2009). A positive relationship between stomata density and

conductance has been reported in a majority of studies looking at natural variation between

species (Anderson & Briske, 1990; Pearce, Millard, Bray, & Rood, 2006) as well as within

species (Carlson, Adams, & Holsinger, 2016; Muchow & Sinclair, 1989; Reich, 1984).

Molecular mutants in genes promoting stomata development show that reduced stomata

density translates into decreased water loss and increased ability to survive after exposure to

drought (Franks, W. Doheny-Adams, Britton-Harper, & Gray, 2015; Yoo et al., 2010).

Yet, decreased stomata density does not necessarily associate with increased demands on

WUE imposed by water limitation. In the *Mimulus guttatus* species complex, accessions from

drier inland populations showed decreased stomatal density and increased WUE, compared to

accessions collected in humid coastal populations (Wu, Lowry, Nutter, & Willis, 2010). By

contrast, in 19 *Protea repens* populations measured in a common garden experiment, stomata

density increased with decreasing summer rainfall at the source location (Carlson et al.,

2016). In addition, higher stomata density does not always translate into higher rates of gas

exchanges: in a diversity panel of rice (Ohsumi, Kanemura, Homma, Horie, & Shiraiwa,

2007) or within several vegetable crop species (Bakker, 1991), for example, the relationship

was not observed.

In fact, stomata density is not the only parameter modulating the balance between water loss

and carbon uptake. Variation in stomata size also impacts the efficiency of stomata regulation

(Raven, 2014). Stomata open and close in response to environmental and internal signals

(Chater et al., 2011; Kinoshita et al., 2011). This ensures that plants do not desiccate when

water evaporation is maximal and spares water when photosynthesis is not active

(Daszkowska-Golec & Szarejko, 2013). The speed of stomata closure is higher in smaller

stomata (Drake, Froend, & Franks, 2013; Raven, 2014). Stomatal responses are an order of

magnitude slower than photosynthetic changes, so any increase in closure time lag may favor

unnecessary water loss and reduce WUE (T. Lawson, Kramer, & Raines, 2012; Raven, 2014).

However, it is often observed that decreases in stomata size occur at the expense of increased

stomata density (reviewed in Hetherington & Woodward, 2003). This leads to a correlation

that may at first be counter-intuitive: increased stomata density can contribute to improve

WUE because of indirect effects on stomata size. In *Eucalyptus globulus*, however, plants

from the drier sites had smaller stomata and higher WUE but no concomitant change in

stomata density (Franks, Drake, & Beerling, 2009). This suggested that the developmental

effect correlating stomata size and density may sometimes be alleviated. Altogether, these

studies highlight interconnections between stomata size, stomata density and WUE that

change across species or populations. How and whether variation in these traits and their

connections support or constrain adaptive processes, however, is not clearly established.

Eco-evolutionary studies, e.g. the analysis of evolutionary forces shaping genetic variation in

natural populations, can determine whether phenotypic variance has a significant impact on

the ecology of species (Carroll, Hendry, Reznick, & Fox, 2007; Hendry, 2016). By drawing

on the elaborate toolbox of population genetics and genomics, it is not only possible to

determine the genetic architecture of any given trait but also to ask whether it is optimized by

natural selection and to investigate the ecological determinants of selective forces at work

(Hendry, 2016; Weinig, Ewers, & Welch, 2014). In this effort, the annual species *Arabidopsis*

*thaliana*, which thrives as a pioneer species in disturbed habitats, has a privileged position

(Gaut, 2012). Genome-wide patterns of nucleotide variation can be contrasted to phenotypic

variation and both the genetic architecture and the adaptive history of the traits can be

reconstructed (Atwell et al., 2010; Fournier-Level et al., 2011; Alonso-Blanco et al., 2016).

Environmental variation has a documented impact on local adaptation in this species (Debieu

et al., 2013; Hamilton, Okada, Korves, & Schmitt, 2015; Hancock et al., 2011; Kronholm,

Picó, Alonso-Blanco, Goudet, & Meaux, 2012; Lasky et al., 2014; Postma & Ågren, 2016). In

addition, natural variation in stomatal patterning is known to segregate in *A. thaliana*

(Delgado, Alonso-Blanco, Fenoll, & Mena, 2011). This species thus provides the ideal

evolutionary context in which the adaptive contribution of variation in stomata patterning can

be dissected.

Here, we developed an automated confocal microscopy approach that overcomes the technical

limitations which have so far complicated the phenotyping of stomatal variation on larger

samples. We characterized genetic variation in stomatal patterning in 330 fully-sequenced

accessions, across a North-South transect of the European range. Additionally, we measured

$\delta^{13}C$, a commonly used estimate of water-use efficiency (WUE) (Juenger et al., 2005; Martin

& Thorstenson, 1988; McKay et al., 2008; Mojica et al., 2016), for all genotypes. Combined

with public genomic and environmental resources, this dataset allows us to ask: i) how

variable are natural *A. thaliana* accessions in stomata patterning? ii) does variation in stomata

patterning influence the carbon-water trade-off? iii) what is the genetic architecture of traits

describing stomata patterning? iv) is stomata patterning optimized by natural selection?

By combining a genome-wide association approach with $Q_{ST}/F_{ST}$ analyses and associations

with environmental parameters, we show that, in *A. thaliana*, variation in stomata patterning

plays a role in local adaptation. Our results further indicate that natural variation in stomata

size is one of the adaptive traits contributing to the optimization of WUE.

## Methods

### Plant material, plant genotypes and growth conditions

In total, 330 accessions, spanning a wide geographical range were selected from the 1001

collection of fully sequenced genotypes (Suppl. Table 1). Accessions were assigned to five

groups based on their geographic origin and genetic clustering (Alonso-Blanco et al., 2016):

Spain, Western Europe, Central Europe, Southern Sweden and Northern Sweden (Figure S1).

If genetic information contradicted geographic information, we prioritized to geographic

information since we are focusing on local adaptation and expect that geography reflects the

scale at which local adaptation proceeds. To avoid oversampling, we randomly reduced the

number of plants sampled at the same location to one for the analysis of heritability, regional

differentiation ($Q_{ST}$-$F_{ST}$) and climatic correlation.

The genome sequences of the 330 genotypes included in the analysis were downloaded from

the 1001 genome database (Alonso-Blanco et al., 2016) on May 12th, 2017. Single nucleotide

polymorphism (SNP) data was extracted using *vcftools* (Danecek et al., 2011). Genomic data

was thinned to 1 SNP picked randomly in each 1000bp window to reduce computational load.

In *A. thaliana*, linkage disequilibrium extends beyond 1kb (Nordborg et al., 2002). Thus, this

data-size reduction should not impact statistics describing the geographical structure of

genomic variation. Additionally, minimum minor allele frequency was set to 5% and sites

exceeding 5% missing data were removed, resulting in 70,410 SNPs among all genotypes.

SNP information was loaded into R using the *vcfR* package (Knaus, Grunwald, Anderson,

Winter, & Kamvar, 2017). For genome-wide association studies the full, unthinned SNP

dataset was used and missing SNPs were imputed using BEAGLE version 3.0 (Browning &

Browning, 2009).

Seeds were stratified on wet paper for 6 days at 4°C in darkness. Plants were grown on soil in

5x5 cm paper pots in 3 replicates with one plant per pot. Genotypes were randomized within

each of 3 blocks of 12 trays containing 8x4 pots. Plants were grown for 7 weeks in growth

chambers (one for each block) under the following conditions: 16 h light; 95 µmol s$^{-1}$ mm$^{-2}$

light intensity; 20 °C day- and 18 °C night-temperature. Plants were watered twice a week and

trays shuffled and rotated every two to three days to account for variable conditions within the

chambers.

## High throughput phenotyping

After 7 weeks, one fully-expanded, intact, adult leaf (one of the largest leaves developed after leaf 4) was selected from each plant for microscopic analysis. Stomata density and size as well as leaf size were measured using our high-throughput microscopy pipeline (for details, see Suppl. Document 1). Stomata density was also determined manually on a random set of 14 individuals as well as on a set of 32 independently-grown individuals. Automatic and manual measurements were strongly correlated (Pearson correlation coefficient $r^2$=0.88, p<<0.01and $r^2$=0.81, p<<0.01, for the 14 and 32 individuals Figures S2-3). The algorithm was conservative and tended to slightly under-estimate stomata numbers. This ensured that stomata area was also correctly quantified. Due to quality filters in our pipeline, the number of analyzed images differed between samples (Figure S4). We found a significant correlation between the number of images analyzed and stomata density (r=0.21, p<<0.01, Figure S5), but not stomata size (r=0.02, p>0.05). Thus, we included the number of images as a co-factor into all statistical models for stomata density (see below). Carbon isotope discrimination measurements ($\delta^{13}C$) of whole rosettes were performed for all plants in block 1 (for details see Suppl. Document 1).

## Heritability estimates

Broad sense heritability $H^2$, the proportion of the observed phenotypic variance that is genetic, was estimated as:

$$H^2 = VarG/(VarG + VarE)$$

where VarG is the genetic variance and VarE is the environmental variance. Because we worked with inbred lines, VarE and VarG can be estimated as the variance between replicates of a genotype and the variance between genotypes, respectively, with a linear-mixed-model

140 using block as fixed effect and genotype as random effect. We ran a linear mixed model using

141 the *lme* function from *nlme* package (Pinheiro, Bates, DebRoy, Sarkar, & R Core Team,

142 2015) (Suppl. Document 2). For $\delta^{13}C$, no replicates were available but a pseudo-heritability

143 estimate was extracted from the GWAS mixed model including the kinship matrix (Atwell et

144 al., 2010).

## Genome-Wide Association Study (GWAS)

For GWAS, markers with minor allele count <5 were removed, leaving a dataset of 2.8-3M

SNPs, depending on missing data for the phenotypes. Minor allele frequency spectra for all

three datasets show that the subset of 261 genotypes, for which all three phenotypes were

determined, has a lower proportion of rare SNPs (Figure S6). GWAS was performed with a

mixed model correcting for population structure using a kinship matrix calculated under the

assumption of the infinitesimal model. Markers were first analyzed with a fast approximation

(Kang et al., 2010) and the 1000 top-most associated SNPs were reanalyzed with the complete

model that estimates the respective variance components for each marker separately (Kang et

al., 2008).

For trait pairs measured on the same plant, a Multi-Trait Mixed Model (MTMM) was applied

to distinguish common and trait-specific SNP-phenotype association (Korte et al., 2012).

The MTMM performs three different statistical tests on a bivariate phenotype including each

trait pair. The first model tests whether a given marker has the same effect on both traits. This

model has increased power to detect significant associations, which may fall under the

significance threshold when traits are analyzed in isolation. The second model identifies

markers having distinct effects on the two traits. It is well suited to detect SNPs with

antagonistic effects on both traits. The last model combines both trait-specific and common

effects. This last model is particularly powerful for detecting markers affecting both traits

with different intensity. The MTMM analysis also provides estimates of the genetic and

environmental correlation for each pair of traits. The statistical details of the models are

described in (Korte et al., 2012).

For all analyses (GWAS and MTMM), the significance threshold was determined as a 5 %

Bonferroni threshold, e.g. 0.05 divided by the number of polymorphic SNPs in the dataset.

## Climatic data

Climatic data included average precipitation, temperature, water vapor pressure (humidity),

wind speed and solar radiation estimates with 2.5 min grid resolution (WorldClim2 database

(Fick & Hijmans, 2017) on May 30[th], 2017) and soil water content (Trabucco & Zomer,

2010). For each variable and accession, we extracted a mean over the putative growing

season, i.e. the months in the year with average temperature greater than 5 °C and average soil

water content over 25% (Suppl. Table 1). We further computed historical drought frequencies

at *A. thaliana* collection sites using 30+ years of the remotely-sensed Vegetative Health Index

(VHI). The VHI is a drought detection method that combines the satellite measured

Vegetative Health and Thermal Condition Indices to identify drought induced vegetative

stress globally at weekly $4km^2$ resolution (Kogan, 1995). This is a validated method for

detecting drought conditions in agriculture. Specifically, we used VHI records to calculate the

historic frequency of observing drought conditions (VHI<40) during the spring (quarter

surrounding spring equinox) and summer (quarter surrounding summer solstice). These are

the typical reproductive seasons of *Arabidopsis* populations (reviewed in Burghardt, Metcalf,

Wilczek, Schmitt, & Donohue, 2015). The drought regime in each location was quantified as

the log-transformed ratio of spring over summer drought frequency. Positive values of this

drought regime measure reflect environments where the frequency of drought decreases over

the typical reproductive growing season, and vice versa for negative values. This ratio

quantifies the seasonality of water availability. It correlates with the ratio of soil water content

of the first and third month of the reproductive season (r=0.54, p<0.01), which we defined as

the first and third growing month in the year, giving similar estimates as Burghardt, Metcalf,

Wilczek, Schmitt, & Donohue (2015).

Because the seven climate variables are correlated, we combined them in seven orthogonal

principal components for 316 *A. thaliana* collection sites (Figures S7-9, loadings described in

Suppl. Document 2). Fourteen genotypes with missing climate data were excluded. Climatic

distance between each region pair was estimated as the F-statistic of a multivariate analysis of

variance (MANOVA) with climatic PCs as response variables and region of origin as

predictor.

Population genomic analysis

Principal component analysis of genomic data (thinned to 1kb) was done using the *adegenet*

package (Jombart et al., 2016) with missing data converted to the mean (Figures S10-11).

Comparing phenotypic differentiation ($Q_{ST}$) to the distribution of $F_{ST}$ is a useful method to

reveal signatures of local adaptation (Leinonen, McCairns, O'hara, & Merilä, 2013; Whitlock

& Guillaume, 2009). Genome-wide, pairwise $F_{ST}$ estimates between regions were calculated

using the *hierfstat* package (function *basic.stats,* Nei's Fst) (Goudet, 2005). Negative $F_{ST}$

values were set to zero before the 95th percentile was calculated.

For stomata density, stomata size and WUE, the respective phenotypic differentiation

between regions, $Q_{ST}$, was estimated as:

214
$$Qst = VarB/(VarW + VarB)$$

where VarW is the genetic variance within regions and VarB the genetic variance between

regions as decribed in Kronholm et al. (2012). Briefly, variance components were estimated

with the *lme* function mentioned previously, including block as a fixed effect and genotype

nested within region as a random effect (Suppl. Document 2). We used the genotype effect as

an estimate for within-region variance and the region effect as an estimate for between-region

variance. Since we did not measure replicates for $\delta^{13}C$, we could not estimate the

environmental variance component. Thus, we adapted the model to include only region as a

random effect, which was our estimate of between-region variance and used the residual

variance as within-region variance. This approach underestimates $Q_{ST}$ and its use for

detecting signatures of local adaptation at the phenotypic level is conservative (Whitlock &

Guillaume, 2009).

To test whether $Q_{ST}$ estimates significantly exceed the 95th percentile of the $F_{ST}$ distribution,

we permuted the phenotypic data by randomizing genotype labels to keep heritability

constant. For each permutation and phenotype, we calculated the difference between each $Q_{ST}$

value and the 95th percentile of the $F_{ST}$ distribution. We used the 95th percentile of the

maximum $Q_{ST}$-$F_{ST}$ distance distribution as a threshold for determining if phenotypic

differentiation significantly exceeds neutral expectations. Since this test takes the maximum

$Q_{ST}$-$F_{ST}$ distance for all population combinations in each permutation, it does not require

multiple testing correction.

## Statistical analysis

Statistical analysis was conducted using R (R Development Core Team, 2008) (R Markdown documentation in Suppl. Document 2). Plots were created using the following libraries: *ggplot2* (Wickham, 2009)*, ggthemes* (Arnold et al., 2017), *ggmap* (Kahle & Wickham, 2013)*, ggbiplot* (Vu, 2011) and *effects* (Fox et al., 2016).

We used Generalized Linear Models (GLM) to test the effect of block, origin, pot position in tray (edge or center) and leaf size on each phenotype (stomata density, stomata size and $\delta^{13}C$). For stomata density we also included the number of analyzed images as a co-factor. The error distribution was a quasi-Poisson distribution for stomata density and size and Gaussian for $\delta^{13}C$. Stomata density was log-transformed to avoid over-dispersion. Significance of each predictor was determined via a type-II likelihood-ratio test (*Anova* function of the *car* package). Significant differences between regions were determined with Tukey's contrasts using the *glht* function of the *multcomp* package (Hothorn et al., 2017). GLMs were also used to test the impact of all climatic principal components on phenotypic traits, while accounting for population structure with the first 20 principal components for genetic variation, which explain 28% of genetic variation (see above). Additionally, for $\delta^{13}C$ we also tested a simpler model including climatic parameters but not population structure. From the resulting models, we created effect plots for significant environmental principal components using the *effects* package (Fox et al., 2016). Further, we used GLMs with binomial distribution to test whether any of the climatic PCs significantly predicts the allelic states of loci associated with WUE in GWAS.

# Results

## Substantial genetic variation in stomata density and size

We analyzed over 31,000 images collected in leaves of 330 *A. thaliana* genotypes and observed high levels of genetic variation in stomata patterning. Genotypic means ranged from 87 to 204 stomata/mm² for stomata density and from 95.0 µm² to 135.1 µm² for stomata size (see Suppl. Table 2 for raw phenotypic data). Leaf size was not significantly correlated with stomata density (r=-0.02, p=0.7, Figure S12) and stomata size (r=-0.08, p=0.15, Figure S13), as expected in fully developed leaves. Broad-sense heritability reached 0.41 and 0.30 for stomata size and density, respectively. Mean stomata density and stomata size were negatively correlated (r=-0.51, p<<0.001; Figure 1). Due to the strong correlation between stomata size and density, we focus primarily on stomata size in the following report, but results for stomata density are in the supplemental material.

## Stomata size correlates with water-use efficiency

We expected variation in stomatal traits to influence the trade-off between carbon uptake and transpiration. Thus, we measured isotopic carbon discrimination, $\delta^{13}C$, an estimator that increases with water-use efficiency (WUE) (Farquhar, Hubick, Condon, & Richards, 1989; McKay et al., 2008). $\delta^{13}C$ ranged from -38.7‰ to -30.8‰ (Suppl. Table 2) and was significantly correlated with stomata size (r=-0.18, p=0.004; Figure 2), indicating that accessions with smaller stomata have higher WUE. About ~4% of the total phenotypic variation (i.e. the sum of phenotypic and genetic variance) in $\delta^{13}C$ is explained by genetic

variance in stomata size. We found no significant correlation between stomatal density and

$\delta^{13}C$ (r=-0.007, p=0.9, Figure S14).

Common genetic basis of stomata size and $\delta^{13}C$

To identify the genetic basis of the phenotypic variance we observe, we conducted a genome-

wide association study (GWAS) for each phenotype. Based on the genome-wide diversity

between genotypes (i.e. the kinship matrix), a pseudo-heritability can be calculated. Pseudo-

heritability estimates were 0.59 for stomata density, 0.56 for stomata size and 0.69 for $\delta^{13}C$,

indicating that differences in stomata patterning and carbon physiology decreased with

increasing relatedness. Despite considerable levels of heritability, we did not detect any

variant associating with stomata density at a significance above the Bonferroni-corrected p-

value of 0.05 (log10(p)=7.78). For stomata size, we detected one QTL with two SNPs

significantly associating at positions 8567936 and 8568437 (Figure S15). These SNPs have an

allele frequency of 1.5% (5 counts) and 2.1% (7 counts), respectively and map to gene

*AT4G14990.1*, which encodes for a protein annotated with a function in cell differentiation.

The former SNP is a synonymous coding mutation while the latter is in an intron.

For $\delta^{13}C$, one genomic region on chromosome 2 position 15094310 exceeded the Bonferroni

significance threshold ($\log_{10}(p)$=7.97, Figure S16). Allele frequency at this SNP was 9.7% (30

counts) and all accessions carrying this allele, except four, were from Southern Sweden (3

Northern Sweden, 1 Central Europe). Southern Swedish lines carrying the allele showed

significantly increased $\delta^{13}C$ compared to the remaining Southern Swedish lines (W=1868, p-

value=6.569e-05, Figure S17). A candidate causal mutation is a non-synonymous SNP at

position 15109013 in gene *AT2G35970.1*, which codes for a protein belonging to the Late

Embryogenesis Abundant (LEA) Hydroxyproline-Rich Glycoprotein family. This SNP also

shows elevated association with the phenotype. However, its significance was below the

Bonferroni-threshold (log(p)=7). Since this SNP is not in linkage disequilibrium with the

highest associating SNP in the region (Figure S18), it is possible that another, independent

SNP in this region is causing the association.

We used Multi-Trait Mixed-Model (MTMM) analysis to disentangle genetic and

environmental determinants of the phenotypic correlations. We found that the significant

correlation between stomata density and stomata size (r=-0.5) had no genetic basis, but had a

significant (r=-0.9, p<0.05) residual correlation. This suggests that the correlation was not

determined by common loci controlling the two traits, but by other, perhaps physical,

constraints or by epistatic alleles at distinct loci. By contrast, the correlation between stomata

size and $\delta^{13}C$ (r= -0.18) had a significant genetic basis (kinship-based correlation, r=-0.58,

p<0.05). Thus, in contrast to the phenotypic variation, genetic variation in stomata size

roughly explains over 33% of the genetic variation in $\delta^{13}C$.

To further investigate the genetic basis for the correlation between stomata size and $\delta^{13}C$, we

performed MTMM GWAS, which tests three models: the first model tests whether a SNP has

the same effect on both traits; the second model tests whether a SNP has differing effects on

both traits and the third model is a combination of the first two to identify SNPs which have

effects of different magnitude on the traits (Korte et al., 2012). We did not observe variants

with same or differing effects on $\delta^{13}C$ and stomata size. However, with the combined model,

we observed a marginally significant association on chromosome 4, which had an effect on

$\delta^{13}C$ but not stomata size, de-correlating the two traits. GWAS of $\delta^{13}C$ restricted to the 261

individuals used for the MTMM analysis confirmed the QTL on chromosome 4. GWAS

applied to different but overlapping sets of accessions yield correlated results but can

sometimes differ in the set of significant associations, since marginal changes in SNP

frequency can affect significance levels (Figure S6). Indeed, the p-values of associations with

$\delta^{13}C$ for the two datasets (310 and 261 accessions) are highly correlated (r=0.87, p<<0.0001,

Figure S19). In this set of genotypes, two SNPS, at position 7083610 and 7083612, exceeded

the Bonferroni-corrected significance threshold ($\alpha$=0.05) (both p=4.8e-09, Figure S20)

although they were under the significance threshold in the larger dataset. Allele frequency is

14% (37 counts) at these two loci and explains 11% of the phenotypic variation. The

association is probably due to complex haplotype differences since it coincides with a

polymorphic deletion and contains several imputed SNPs. Thirty-five of the 37 accessions

carrying the minor allele originated from Southern Sweden and showed significantly higher

$\delta^{13}C$ compared to other Southern Swedish accessions (mean difference=1.34; W=1707,

p=1.15e-06; Figure S21). In summary, we detected two genetic variants significantly

associating with $\delta^{13}C$, independent of stomata size, despite the common genetic basis of the

two traits.

Stomata size and stomata density correlate with geographical patterns of climatic

variation

We used PCA to describe multivariate variation in climatic conditions reported for the

locations of origins of the genotypes. We tested the correlation of each measured phenotype

with climatic principal components (PC) using a GLM which accounted for genetic

population structure (see methods). We found a significant, negative relationship between

genetic variation in stomata size and climatic PC2 (Likelihood ratio test Chi-Square (LRT X²)

=9.2784, degrees of freedom (df)=1, p=0.005) and PC5 (LRT X²=5.7335, df=1, p=0.02,

Figure 3). Climatic PC 2 explained 23.8% of climatic variation and had the strongest loadings

(both negative) from temperature and water vapor pressure (humidity). Climatic PC 5

explained 9% of the climatic variation and mostly increased with increasing spring-summer

drought probability ratio and increasing solar radiation. We also found significant climatic

predictors for the distribution of genetic variation in stomata density (PC 2: LRT X²=8.6612,

df=1 p=0.003; PC 5: LRT X²=7.3773, df=1, p=0.007; PC 7: LRT X²=6.6033, df=1, p=0.01;

Figure S22). $\delta^{13}C$ did not correlate with any of the climatic PCs. However, removing

population structure covariates from the model revealed significant correlations of $\delta^{13}C$ with

climatic PC2 (+, LRT X²=7.3564, df=1, p=0.006), PC3 (-, LRT X²=3.8889, df=1, p=0.048)

and PC4 (+, LRT X²=6.6885, df=1, p= 0.009) (Figure S23). PC3 explained 13.7% of climatic

variation and principally increased with rainfall and decreased with spring-summer drought

probability ratio. PC4 explained 11.4% of the total variation and mostly increased with wind

speed. Therefore, the covariation of $\delta^{13}C$ with climatic parameters describing variation in

water availability and evaporation in *A. thaliana* is strong but confounded with the

demographic history of the species. To test whether alleles associating with increased $\delta^{13}C$ in

GWAS are involved in adaptation to local climate, we checked whether any climatic PC is a

significant predictor of the allelic state of Southern Swedish accessions. However, none of the

climatic PCs was a significant predictor for one of the two loci.

## Patterns of regional differentiation depart from neutral expectations

Genotypes were divided into five regions based on genetic clustering (Alonso-Blanco et al.,

2016) and their geographic origin (Figure S1, see Methods). We detected significant

phenotypic differentiation among these regions for stomata size (LRT $X^2$=52.852, df=4,

p=9.151e-11, Figure 4). Stomata size was significantly lower in Southern Sweden (mean=108

μm²) compared to Central Europe (mean=114μm², GLHT z=-6.24, p<0.001), Western Europe

(mean=111 μm², GLHT z=2.769, p=0.04) and Spain (mean=113 μm², GLHT z=6.709,

p<0.001), which did not significantly differ from each other. Northern Sweden showed an

intermediate phenotype and did not differ significantly from any region (mean=110 μm²).

Variation for stomata density, showed a similar but inverted pattern (Figure S24).

Furthermore, we found significant regional differentiation in $\delta^{13}$C measurements (LR $X^2$

=58.029, df=4 p=7.525e-12, Figure 4). Highest $\delta^{13}$C levels (highest WUE) were found in

accessions from Northern Sweden (mean=-34.8) and Southern Sweden (mean=-35.2), which

were significantly higher than in accessions from Spain (mean=-35.7; GLHT Southern

Sweden z=-3.472, p=0.008; GLHT North Sweden z=-3.49, p=0.001) and Western Europe

(mean=-36.06; GLHT Southern Sweden z= -2.8, p=0.03; GLHT Northern Sweden z=-3.28,

p=0.008). Lowest $\delta^{13}$C levels were found in lines from Central Europe (mean=-36.6), which

were significantly lower than in lines from Northern Sweden (GLHT z=5.676, p<0.001),

Southern Sweden (GLHT z=6.992, p<0.001) and Spain (GLHT z=3.714, p=0.002).

The observed regional differences result either from the demographic history of the regions or

from the action of local selective forces. To tease these possibilities apart, the phenotypic

differentiation ($Q_{ST}$) can be compared to nucleotide differentiation ($F_{ST}$) (Kronholm et al.,

2012; Leinonen et al., 2013). We examined each pair of regions separately, since they are not

equidistant from each other and calculated $F_{ST}$ distributions for over 70,000 SNP markers

(spaced at least 1kb apart, see methods). For each trait, $Q_{ST}$ exceeded the 95th percentile of the

$F_{ST}$ distribution in at least two pairs of regions (Table 1 A-C). We used permutations to

calculate a significance threshold for the $Q_{ST}/F_{ST}$ difference (see methods). Significant

regional differentiation was pervasive in our sample, with Central Europe and Southern

Sweden being significantly differentiated for all three phenotypes. This analysis suggests that

natural selection has contributed to shape the phenotypic differentiation between regions.

Regional differences in climate may have imposed divergence in stomatal patterning. Thus,

we estimated climatic distances between regions using estimates of regional effects extracted

from a MANOVA. We did not observe significant correlations between adaptive phenotypic

divergence ($Q_{ST}$-$F_{ST}$) and the climatic distance of the respective regions (Mantel test $p>0.05$

for each of the three traits). Regional divergence in $\delta C^{13}$, stomata density and stomata size

was therefore not proportional to climatic divergence.

## Discussion

### Genetic variation for stomata density and size segregates in A. thaliana

We used high-throughput confocal imaging to characterize stomata patterning in over 31,000

images from 870 samples collected from 330 genotypes. Our high-throughput pipeline could

characterize stomata density and stomata size with a reliable accuracy, confirmed by high

correlation with manual measurements. Broad-sense heritability and pseudo-heritability

estimates for stomata density, which are 30% and 58%, respectively, are slightly lower than in

a previous report of manually counted stomata diversity across a smaller sample chosen to

maximize genetic diversity (Delgado et al., 2011). Despite the clear impact of environmental

(random) variance on both observed phenotypes, stomata size and stomata density showed a

strong negative correlation. This is consistent with earlier reports of studies manipulating

regulators of stomata development (Doheny-Adams, Hunt, Franks, Beerling, & Gray, 2012;

Franks et al., 2015), but also with studies analyzing stomatal trait variation in a wide range of

species (Franks & Beerling, 2009; Hetherington & Woodward, 2003).

The extensive genomic resources available in *A. thaliana* enabled us to investigate the genetic

basis of trait variation and co-variation, with the help of GWAS (Atwell et al., 2010). Much is

known about the molecular pathways that control the differentiation of stomata in *Arabidopsis*

*thaliana*, providing a set of candidate genes expected to control genetic variation in stomata

patterning (Bergmann & Sack, 2007; Pillitteri & Torii, 2012). However, we did not detect any

genomic region that associated with stomata density at a p-value beyond the Bonferroni-

significance threshold. For stomata size, there was only one significant association on

chromosome 4, albeit with very low minor allele frequency in a gene that has not been

reported previously in stomata development. GWAS studies can detect small-effect loci only

if they segregate at high frequency, whereas rare alleles only give detectable signals when

they are of large effect (Korte & Farlow, 2013; Wood et al., 2014). Given that variance for

both stomata size and stomata density is clearly heritable, the genetic variants controlling

these traits are not causing strong association signals in GWAS. Theoretically, the presence of

a large effect QTL impacting local adaptation can be masked by correction for population

structure. However, not correcting for population structure is known to lead to a high number

of false-positives and is thus not a reliable alternative (Vilhjálmsson & Nordborg, 2012).

Nevertheless, we can conclude that variation in stomata patterning is controlled by a

combination of i) alleles of moderate effect size segregating at frequencies too low to be

detected by GWAS, and/or ii) alleles segregating at high frequency but with effect size too

small to be detected and/or iii) rare alleles of small effect. In addition, it is possible that the

effect of associated loci is weakened by epistatic interactions among loci. In *A. thaliana*, the

genetic architecture of natural variation in stomata traits is therefore not caused by a handful

of large effect variants but complex and polygenic.

Using MTMM analysis (Korte et al., 2012), we further investigated the impact of genetic

variation on the negative co-variation between stomata size and density. This analysis

revealed that genetic similarity does not influence the pattern of covariation. It implies that

either multiple alleles act epistatically on the covariation, or that physical or environmental

factors explain the correlation.

Natural variation in stomata patterning can contribute to optimize physiological

performance

Both stomata development and reactions to drought stress are being intensively investigated

in *A. thaliana* (Bergmann & Sack, 2007; Krasensky & Jonak, 2012; Pillitteri & Torii, 2012;

Verslues, Govinal Badiger, Ravi, & M. Nagaraj, 2013). Mutants in stomata density or size

have recently been shown to have a clear impact carbon physiology (Franks et al., 2015;

Hepworth, Doheny-Adams, Hunt, Cameron, & Gray, 2015; Hughes et al., 2017; S. S.

Lawson, Pijut, & Michler, 2014; Masle, Gilmore, & Farquhar, 2005; Yoo et al., 2010; Yu et

al., 2008).Yet, the relevance of natural variation in stomatal patterning on the local limitation

in water availability had not been documented in this species so far. We provide here

concomitant measures of morphological and physiological variation to examine the impact of

variation in stomatal patterning on natural variation in WUE. By including genome-wide

patterns of nucleotide diversity, our analysis presents two major findings: i) the decrease in

stomata size associates with an increase in WUE in *A. thaliana* and ii) this pattern of co-

variation has a genetic basis. This shows that, in *A. thaliana*, variation in stomata size has the

potential to be involved in the optimization of physiological processes controlling the trade-

off between growth and water loss. Interestingly, in the close relative *A. lyrata ssp. lyrata*,

stomata were observed to grow smaller when water was limiting, which coincided with

increased WUE (Paccard, Fruleux, & Willi, 2014). This suggest that the consequences of

decreased stomata size are conserved in the genus.

While variation for stomata size and density is likely shaped by a complex genetic

architecture that hindered QTL detection, we detected two regions in the genome that

associated significantly with carbon isotope discrimination. Three previous QTL mapping

analyses, including one between locally adapted lines from Sweden and Italy, identified 16

distinct QTLs controlling $\delta^{13}C$ (Juenger et al., 2005; McKay et al., 2008; Mojica et al., 2016).

One of these is caused by a rare allele in the root-expressed gene *MITOGEN ACTIVATED*

*PROTEIN KINASE 12* (*MPK-12)*, (Campitelli, Des Marais, & Juenger, 2016; Juenger et al.,

2005). While QTL-mapping approaches can only reveal the variance shown by the parental

lines, GWAS approaches fail to detect rare alleles unless they have a very strong impact. It is

therefore not surprising that the loci that stand out in GWAS do not overlap with the QTL

previously mapped. In fact, one of the mapping populations used the parental genotype Cvi-0,

a genotypic and phenotypic outlier.

The two QTL we report here on chromosomes 2 and 4 add two novel loci, raising the number

of genomic regions known to impact $\delta^{13}$C in *A. thaliana* to 18. The two QTL we report here

on chromosomes 2 and 4 add two novel loci, raising the number of genomic regions known to

impact $\delta^{13}$C in *A. thaliana* to 18. The novel loci we report are locally frequent. Individuals

carrying the minor alleles of both loci are almost exclusively from Southern Sweden and

display significantly higher $\delta^{13}$C than other Southern Swedish accessions. However, we did

not find any climatic factor significantly correlated with the allelic states of our QTLs. This

suggests that other factors, like soil composition, play a role in drought adaptation.

Alternatively, locally adapted alleles may not yet be fixed within the region.

Interestingly, the accessions with the minor allele associating with high $\delta^{13}$C in both QTL did

not show decreased stomata size compared to other accessions. Multi-trait GWAS confirmed

that these QTL are associated with $\delta^{13}$C variants that are independent of genetic variation for

stomata patterning. We therefore can conclude that, stomata patterning is only one of the traits

contributing to the optimization of WUE. A large array of molecular and physiological

reactions is indeed known to contribute to tolerance to drought stress (Krasensky & Jonak,

2012; Verslues et al., 2013). The close vicinity of the chromosome 2 QTL to a non-

synonymous mutation in a gene encoding an LEA protein, known to act as a chaperone when

cells dehydrate, suggests one possible mechanism by which WUE might be optimized

independently of stomata size and density (Candat et al., 2014; Eriksson, Kutzer, Procek,

Gröbner, & Harryson, 2011; Reyes et al., 2005). Variation in rates of proline accumulation in

the presence of drought stress or in nutrient acquisition in the root are also among the

physiological mechanism that appear to have contributed to improve drought stress tolerance

in this species (Campitelli et al., 2016; Kesari et al., 2012).

Adaptive evolution of stomata patterning is suggested by the geographic distribution

of genetic variation

Phenotypic variation for stomata patterning and carbon uptake is not uniformly distributed

throughout the species range. All three phenotypes we report in this study were significantly

differentiated between the five broad regions defined in our sample of 330 genotypes. We

performed a comparison of phenotypic and nucleotide levels of divergence to evaluate the

putative role of past selective events in shaping the distribution of diversity we report

(Leinonen et al., 2013; Whitlock & Guillaume, 2009). Because these regions are not equally

distant, $F_{ST}/Q_{ST}$ comparisons averaged over all populations may mask local patterns of

adaptation (Leinonen et al., 2013). We therefore measured $Q_{ST}$ between pairs of regions and

compared them to the distribution of pairwise $F_{ST}$, using permutations to establish the

significance of outlier $Q_{ST}$. This analysis showed that, for all three traits, differentiation

between regions was stronger than expected from genome-wide patterns of diversity,

suggesting local adaptation. This is further supported by our finding that stomata density and

stomata size correlated with climatic principal components, which are most strongly driven by

temperature, humidity, solar radiation, and historic drought regimen.

The strongest $Q_{ST}$-$F_{ST}$ differences are found across regional pairs including Central Europe.

Particularly, WUE is significantly differentiated between Central Europe and Spain as well as

both Swedish regions, due to low WUE in Central Europe. It is tempting to speculate that the

significantly lower WUE observed in Central Europe results from selection for life cycling at

latitudes where two life cycles can be completed each year, as high WUE is usually associated

with a reduction in photosynthetic rate (Blum, 2009; Field et al., 1983; Kimball et al., 2014).

Interestingly, Central Europe and Southern Sweden are significantly differentiated for all

three traits and Southern Sweden and Spain are significantly differentiated for both stomata

traits. Combined with the fact that Swedish genotypes show the highest values for WUE, this

suggests that stomata size is involved in drought adaptation of Swedish accessions. This result

is somewhat counterintuitive because Sweden is not known to be a region experiencing

intense drought. However, our result is supported by an independent study showing that

Northern and Southern Swedish genotypes maintain photosynthetic activity under terminal

drought stress for a longer time than other, especially Central and Western European,

accessions (Exposito-Alonso et al., 2017). Additionally, locally adapted genotypes from

Northern Sweden (which showed high WUE in our study, as well) have been shown to

display higher WUE than Italian genotypes (Mojica et al., 2016).

This regional difference in *A. thaliana* further coincides with the satellite measurements of

historic drought regimen, which show that Sweden is a region where drought frequency is

changing along the season: it is relatively more frequent in the early growing season (spring)

than in the late growing season (summer). Drought episodes occurring earlier in the growth

season may favor the evolution of drought avoidance traits (e.g. morphological or

physiological stress adaptations) over that of escape strategies mediated by e.g. seed

dormancy (Kooyers, 2015; Passioura, 1996). Indeed, in Northern Europe, increased negative

co-variation between flowering time and seed dormancy suggested that the narrow growth

season imposes a strong selection on life-history traits (Debieu et al., 2013). In Southern

European regions, *A. thaliana* appeared to rely on escape strategies provided by increased

seed dormancy (Kronholm et al., 2012). Taken together, this suggests that decreased stomata

size and, consequently, increased $\delta^{13}$C have contributed to adaptation to water limitations in

spring in a region where the narrow growth season leaves no room for escape strategies.

Indeed, both stomata size and $\delta^{13}$C associate with historic drought regimen. For $\delta^{13}$C,

however, this association disappears when genetic population structure is included as a

covariate. This indicates that local adaptation for WUE might have also contributed to shape

the current population structure.

Finally, the coarse regional contrasts used in the present study cannot resolve patterns of local

adaptation occurring at a fine-grained scale within regions (as e.g. local adaptation to specific

soil patches). In fact, we observe most variation for all three phenotypes within regions. It is

therefore possible that we underestimate the magnitude of adaptive differentiation across the

species' European range, which could further explain why $Q_{ST}$ / $F_{ST}$ differences did not co-

vary with environmental divergence in our dataset.

Conclusion

This work provides a comprehensive description of the variation in stomata size and density

that segregates throughout the European range of *A. thaliana*. It shows that stomata size

covaries with water-use efficiency and may contribute to local adaptation. Several reports

indicate that plants can also change stomatal development in water-limiting conditions

(Fraser, Greenall, Carlyle, Turkington, & Friedman, 2009; Paccard et al., 2014; Xu & Zhou,

2008). Future work will have to investigate whether this variation in stomata size and number

also contributes to adaptive plasticity to drought stress.

## Acknowledgements

We thank Swantje Prahl and Hildegard Schwitte for technical support in microscopy, Maria

Graf for technical assistance in $\delta^{13}C$ analysis, Anja Linstädter for advice in statistical analysis

and Angela Hancock for critical comments on the manuscript. This research was supported by

the Deutsche Forschungsgemeinschaft (DFG) through grant INST 211/575-1 for the OPERA

microscope, and grant ME2742/6-1 in the realm of SPP1529 "Adaptomics", as well as by the

European Research Council with Grant 648617 "AdaptoSCOPE".

## References

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M., …
Zhou, X. (2016). 1,135 Genomes Reveal the Global Pattern of Polymorphism in
Arabidopsis thaliana. *Cell*, *166*(2), 481–491.
https://doi.org/10.1016/j.cell.2016.05.063
Anderson, V. J., & Briske, D. D. (1990). Stomatal Distribution, Density and Conductance of
Three Perennial Grasses Native to the Southern True Prairie of Texas. *The
American Midland Naturalist*, *123*(1), 152–159.
https://doi.org/10.2307/2425768
Arnold, J. B., Daroczi, G., Werth, B., Weitzner, B., Kunst, J., Auguie, B., … London, J. (2017).
ggthemes: Extra Themes, Scales and Geoms for "ggplot2" (Version 3.4.0).
Retrieved from https://cran.r-project.org/web/packages/ggthemes/index.html
Atwell, S., Huang, Y. S., Vilhjálmsson, B. J., Willems, G., Horton, M., Li, Y., … Nordborg, M.
(2010). Genome-wide association study of 107 phenotypes in Arabidopsis
thaliana inbred lines. *Nature*, *465*(7298), 627–631.
https://doi.org/10.1038/nature08800
Bakker, J. C. (1991). Effects of humidity on stomatal density and its relation to leaf
conductance. *Scientia Horticulturae*, *48*(3), 205–212.
https://doi.org/10.1016/0304-4238(91)90128-L
Bergmann, D. C., & Sack, F. D. (2007). Stomatal Development. *Annual Review of Plant
Biology*, *58*(1), 163–181.
https://doi.org/10.1146/annurev.arplant.58.032806.104023
Blum, A. (2009). Effective use of water (EUW) and not water-use efficiency (WUE) is the
target of crop yield improvement under drought stress. *Field Crops Research*,
*112*(2), 119–123. https://doi.org/10.1016/j.fcr.2009.03.009

Browning, B. L., & Browning, S. R. (2009). A Unified Approach to Genotype Imputation
and Haplotype-Phase Inference for Large Data Sets of Trios and Unrelated
Individuals. *The American Journal of Human Genetics*, *84*(2), 210–223.
https://doi.org/10.1016/j.ajhg.2009.01.005
Burghardt, L. T., Metcalf, C. J. E., Wilczek, A. M., Schmitt, J., & Donohue, K. (2015).
Modeling the Influence of Genetic and Environmental Variation on the Expression
of Plant Life Cycles across Landscapes. *The American Naturalist*, *185*(2), 212–227.
https://doi.org/10.1086/679439
Campitelli, B. E., Des Marais, D. L., & Juenger, T. E. (2016). Ecological interactions and the
fitness effect of water-use efficiency: Competition and drought alter the impact of
natural MPK12 alleles in Arabidopsis. *Ecology Letters*, *19*(4), 424–434.
https://doi.org/10.1111/ele.12575
Candat, A., Paszkiewicz, G., Neveu, M., Gautier, R., Logan, D. C., Avelange-Macherel, M.-H.,
& Macherel, D. (2014). The Ubiquitous Distribution of Late Embryogenesis
Abundant Proteins across Cell Compartments in Arabidopsis Offers Tailored
Protection against Abiotic Stress. *The Plant Cell*, *26*(7), 3148–3166.
https://doi.org/10.1105/tpc.114.127316
Carlson, J. E., Adams, C. A., & Holsinger, K. E. (2016). Intraspecific variation in stomatal
traits, leaf traits and physiology reflects adaptation along aridity gradients in a
South African shrub. *Annals of Botany*, *117*(1), 195–207.
https://doi.org/10.1093/aob/mcv146
Carroll, S. P., Hendry, A. P., Reznick, D. N., & Fox, C. W. (2007). Evolution on ecological
time-scales. *Functional Ecology*, *21*(3), 387–393. https://doi.org/10.1111/j.1365-
2435.2007.01289.x
Chater, C., Kamisugi, Y., Movahedi, M., Fleming, A., Cuming, A. C., Gray, J. E., & Beerling, D.
J. (2011). Regulatory mechanism controlling stomatal behavior conserved across
400 million years of land plant evolution. *Current Biology: CB*, *21*(12), 1025–
1029. https://doi.org/10.1016/j.cub.2011.04.032
Cowan, I. R. (1986). Economics of carbon fixation in higher plants. *On the Economy of
Plant Form and Function : Proceedings of the Sixth Maria Moors Cabot Symposium,
Evolutionary Constraints on Primary Productivity, Adaptive Patterns of Energy
Capture in Plants, Harvard Forest, August 1983*. Retrieved from
http://agris.fao.org/agris-search/search.do?recordID=US201301399273
Cowan, I. R., & Farquhar, G. D. (1977). Stomatal function in relation to leaf metabolism
and environment. *Symposia of the Society for Experimental Biology*, *31*, 471–505.
Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R.
(2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158.
https://doi.org/10.1093/bioinformatics/btr330
Daszkowska-Golec, A., & Szarejko, I. (2013). Open or Close the Gate – Stomata Action
Under the Control of Phytohormones in Drought Stress Conditions. *Frontiers in
Plant Science*, *4*. https://doi.org/10.3389/fpls.2013.00138
Debieu, M., Tang, C., Stich, B., Sikosek, T., Effgen, S., Josephs, E., … Meaux, J. de. (2013). Co-
Variation between Seed Dormancy, Growth Rate and Flowering Time Changes
with Latitude in Arabidopsis thaliana. *PLOS ONE*, *8*(5), e61075.
https://doi.org/10.1371/journal.pone.0061075
Delgado, D., Alonso-Blanco, C., Fenoll, C., & Mena, M. (2011). Natural variation in
stomatal abundance of Arabidopsis thaliana includes cryptic diversity for

673   different developmental processes. *Annals of Botany*, *107*(8), 1247–1258.
674   https://doi.org/10.1093/aob/mcr060

675 Doheny-Adams, T., Hunt, L., Franks, P. J., Beerling, D. J., & Gray, J. E. (2012). Genetic
676   manipulation of stomatal density influences stomatal size, plant growth and
677   tolerance to restricted water supply across a growth carbon dioxide gradient.
678   *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1588),
679   547–555. https://doi.org/10.1098/rstb.2011.0272

680 Drake, P. L., Froend, R. H., & Franks, P. J. (2013). Smaller, faster stomata: scaling of
681   stomatal size, rate of response, and stomatal conductance. *Journal of*
682   *Experimental Botany*, *64*(2), 495–505. https://doi.org/10.1093/jxb/ers347

683 Eriksson, S. K., Kutzer, M., Procek, J., Gröbner, G., & Harryson, P. (2011). Tunable
684   membrane binding of the intrinsically disordered dehydrin Lti30, a cold-induced
685   plant stress protein. *The Plant Cell*, *23*(6), 2391–2404.
686   https://doi.org/10.1105/tpc.111.085183

687 Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A. A., & Weigel, D.
688   (2017). Genomic basis and evolutionary potential for extreme drought
689   adaptation in Arabidopsis thaliana. *BioRxiv*, 118067.
690   https://doi.org/10.1101/118067

691 Farquhar, G. D., Hubick, K. T., Condon, A. G., & Richards, R. A. (1989). Carbon Isotope
692   Fractionation and Plant Water-Use Efficiency. In P. W. Rundel, J. R. Ehleringer, &
693   K. A. Nagy (Eds.), *Stable Isotopes in Ecological Research* (pp. 21–40). New York,
694   NY: Springer New York. https://doi.org/10.1007/978-1-4612-3498-2_2

695 Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate
696   surfaces for global land areas. *International Journal of Climatology*.

697 Field, C., Merino, J., & Mooney, H. A. (1983). Compromises between Water-Use Efficiency
698   and Nitrogen-Use Efficiency in Five Species of California Evergreens. *Oecologia*,
699   *60*(3), 384–389.

700 Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., & Wilczek, A. M.
701   (2011). A Map of Local Adaptation in Arabidopsis thaliana. *Science*, *334*(6052),
702   86–89. https://doi.org/10.1126/science.1209271

703 Fox, J., Weisberg, S., Friendly, M., Hong, J., Andersen, R., Firth, D., & Taylor, S. (2016).
704   effects: Effect Displays for Linear, Generalized Linear, and Other Models (Version
705   3.1-2). Retrieved from https://cran.r-
706   project.org/web/packages/effects/index.html

707 Franks, P. J., & Beerling, D. J. (2009). Maximum leaf conductance driven by CO2 effects on
708   stomatal size and density over geologic time. *Proceedings of the National Academy*
709   *of Sciences*, *106*(25), 10343–10347. https://doi.org/10.1073/pnas.0904209106

710 Franks, P. J., Drake, P. L., & Beerling, D. J. (2009). Plasticity in maximum stomatal
711   conductance constrained by negative correlation between stomatal size and
712   density: an analysis using Eucalyptus globulus. *Plant, Cell & Environment*, *32*(12),
713   1737–1748. https://doi.org/10.1111/j.1365-3040.2009.002031.x

714 Franks, P. J., W. Doheny-Adams, T., Britton-Harper, Z. J., & Gray, J. E. (2015). Increasing
715   water-use efficiency directly through genetic manipulation of stomatal density.
716   *New Phytologist*, *207*(1), 188–195. https://doi.org/10.1111/nph.13347

717 Fraser, L. H., Greenall, A., Carlyle, C., Turkington, R., & Friedman, C. R. (2009). Adaptive
718   phenotypic plasticity of Pseudoroegneria spicata: response of stomatal density,

leaf area and biomass to changes in water supply and increased temperature.
*Annals of Botany*, *103*(5), 769–775. https://doi.org/10.1093/aob/mcn252
Gaut, B. (2012). Arabidopsis thaliana as a model for the genetics of local adaptation.
*Nature Genetics*, *44*(2), 115–116. https://doi.org/10.1038/ng.1079
Goudet, J. (2005). hierfstat, a package for r to compute and test hierarchical F-statistics.
*Molecular Ecology Notes*, *5*(1), 184–186. https://doi.org/10.1111/j.1471-
8286.2004.00828.x
Hamilton, J. A., Okada, M., Korves, T., & Schmitt, J. (2015). The role of climate adaptation
in colonization success in Arabidopsis thaliana. *Molecular Ecology*, *24*(9), 2253–
2263. https://doi.org/10.1111/mec.13099
Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G., …
Bergelson, J. (2011). Adaptation to Climate Across the Arabidopsis thaliana
Genome. *Science*, *334*(6052), 83–86. https://doi.org/10.1126/science.1209244
Hendry, A. P. (2016). *Eco-evolutionary Dynamics*. Princeton University Press.
Hepworth, C., Doheny-Adams, T., Hunt, L., Cameron, D. D., & Gray, J. E. (2015).
Manipulating stomatal density enhances drought tolerance without deleterious
effect on nutrient uptake. *The New Phytologist*, *208*(2), 336–341.
https://doi.org/10.1111/nph.13598
Hetherington, A. M., & Woodward, F. I. (2003). The role of stomata in sensing and driving
environmental change. *Nature*, *424*(6951), 901–908.
https://doi.org/10.1038/nature01843
Hothorn, T., Bretz, F., Westfall, P., Heiberger, R. M., Schuetzenmeister, A., & Scheibe, S.
(2017). multcomp: Simultaneous Inference in General Parametric Models
(Version 1.4-7). Retrieved from https://cran.r-
project.org/web/packages/multcomp/index.html
Hughes, J., Hepworth, C., Dutton, C., Dunn, J. A., Hunt, L., Stephens, J., … Gray, J. E. (2017).
Reducing Stomatal Density in Barley Improves Drought Tolerance without
Impacting on Yield. *Plant Physiology*, *174*(2), 776–787.
https://doi.org/10.1104/pp.16.01844
Jombart, T., Kamvar, Z. N., Lustrik, R., Collins, C., Beugin, M.-P., Knaus, B., … Calboli, F.
(2016). adegenet: Exploratory Analysis of Genetic and Genomic Data (Version
2.0.1). Retrieved from https://cran.r-
project.org/web/packages/adegenet/index.html
Juenger, T. E., Mckay, J. K., Hausmann, N., Keurentjes, J. J. B., Sen, S., Stowe, K. A., …
Richards, J. H. (2005). Identification and characterization of QTL underlying
whole-plant physiology in Arabidopsis thaliana: δ13C, stomatal conductance and
transpiration efficiency. *Plant, Cell & Environment*, *28*(6), 697–708.
https://doi.org/10.1111/j.1365-3040.2004.01313.x
Kahle, D., & Wickham, H. (2013). ggmap: Spatial Visualization with ggplot2. *The R
Journal*, *5*(1), 144–161.
Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S., Freimer, N. B., … Eskin, E.
(2010). Variance component model to account for sample structure in genome-
wide association studies. *Nature Genetics*, *42*(4), 348–354.
https://doi.org/10.1038/ng.548
Kang, H. M., Zaitlen, N. A., Wade, C. M., Kirby, A., Heckerman, D., Daly, M. J., & Eskin, E.
(2008). Efficient control of population structure in model organism association

mapping. *Genetics*, *178*(3), 1709–1723.
https://doi.org/10.1534/genetics.107.080101
Kesari, R., Lasky, J. R., Villamor, J. G., Marais, D. L. D., Chen, Y.-J. C., Liu, T.-W., … Verslues,
P. E. (2012). Intron-mediated alternative splicing of Arabidopsis P5CS1 and its
association with natural variation in proline and climate adaptation. *Proceedings*
*of the National Academy of Sciences*, *109*(23), 9197–9202.
https://doi.org/10.1073/pnas.1203433109
Kimball, S., Gremer, J. R., Barron-Gafford, G. A., Angert, A. L., Huxman, T. E., & Venable, D.
L. (2014). High water-use efficiency and growth contribute to success of non-
native Erodium cicutarium in a Sonoran Desert winter annual community.
*Conservation Physiology*, *2*(1). https://doi.org/10.1093/conphys/cou006
Kinoshita, T., Ono, N., Hayashi, Y., Morimoto, S., Nakamura, S., Soda, M., … Shimazaki, K.
(2011). FLOWERING LOCUS T Regulates Stomatal Opening. *Current Biology*,
*21*(14), 1232–1238. https://doi.org/10.1016/j.cub.2011.06.025
Knaus, B. J., Grunwald, N. J., Anderson, E. C., Winter, D. J., & Kamvar, Z. N. (2017). vcfR:
Manipulate and Visualize VCF Data (Version 1.4.0). Retrieved from
https://cran.r-project.org/web/packages/vcfR/index.html
Kogan, F. N. (1995). Application of vegetation index and brightness temperature for
drought detection. *Advances in Space Research*, *15*(11), 91–100.
https://doi.org/10.1016/0273-1177(95)00079-T
Kooyers, N. J. (2015). The evolution of drought escape and avoidance in natural
herbaceous populations. *Plant Science*, *234*, 155–162.
https://doi.org/10.1016/j.plantsci.2015.02.012
Korte, A., & Farlow, A. (2013). The advantages and limitations of trait analysis with
GWAS: a review. *Plant Methods*, *9*, 29. https://doi.org/10.1186/1746-4811-9-29
Korte, A., Vilhjálmsson, B. J., Segura, V., Platt, A., Long, Q., & Nordborg, M. (2012). A
mixed-model approach for genome-wide association studies of correlated traits
in structured populations. *Nature Genetics*, *44*(9), 1066–1071.
https://doi.org/10.1038/ng.2376
Krasensky, J., & Jonak, C. (2012). Drought, salt, and temperature stress-induced
metabolic rearrangements and regulatory networks. *Journal of Experimental*
*Botany*, *63*(4), 1593–1608. https://doi.org/10.1093/jxb/err460
Kronholm, I., Picó, F. X., Alonso-Blanco, C., Goudet, J., & Meaux, J. de. (2012). Genetic Basis
of Adaptation in Arabidopsis Thaliana: Local Adaptation at the Seed Dormancy
Qtl Dog1. *Evolution*, *66*(7), 2287–2302. https://doi.org/10.1111/j.1558-
5646.2012.01590.x
Lambers, H., Chapin, F. S., & Pons, T. L. (1998). Photosynthesis, Respiration, and Long-
Distance Transport. In *Plant Physiological Ecology* (pp. 10–153). Springer, New
York, NY. https://doi.org/10.1007/978-1-4757-2855-2_2
Lasky, J. R., Marais, D., L, D., Lowry, D. B., Povolotskaya, I., McKay, J. K., … Juenger, T. E.
(2014). Natural Variation in Abiotic Stress Responsive Gene Expression and Local
Adaptation to Climate in Arabidopsis thaliana. *Molecular Biology and Evolution*,
*31*(9), 2283–2296. https://doi.org/10.1093/molbev/msu170
Lawson, S. S., Pijut, P. M., & Michler, C. H. (2014). The cloning and characterization of a
poplar stomatal density gene. *Genes & Genomics. 36(4): 427-441.*, *36*(4), 427–441.
https://doi.org/10.1007/s13258-014-0177-x

Lawson, T., Kramer, D. M., & Raines, C. A. (2012). Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. *Current Opinion in Biotechnology*, *23*(2), 215–220. https://doi.org/10.1016/j.copbio.2011.12.012

Leinonen, T., McCairns, R. S., O'hara, R. B., & Merilä, J. (2013). QST–FST comparisons: evolutionary and ecological insights from genomic heterogeneity. *Nature Reviews Genetics*, *14*(3), 179–190.

Martin, B., & Thorstenson, Y. R. (1988). Stable Carbon Isotope Composition (δ13C), Water Use Efficiency, and Biomass Productivity of Lycopersicon esculentum, Lycopersicon pennellii, and the F1 Hybrid. *Plant Physiology*, *88*(1), 213–217. https://doi.org/10.1104/pp.88.1.213

Masle, J., Gilmore, S. R., & Farquhar, G. D. (2005). The ERECTA gene regulates plant transpiration efficiency in Arabidopsis. *Nature*, *436*(7052), 866. https://doi.org/10.1038/nature03835

McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., … Yepez, E. A. (2008). Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New Phytologist*, *178*(4), 719–739. https://doi.org/10.1111/j.1469-8137.2008.02436.x

McKay, J. K., Richards, J. H., Nemali, K. S., Sen, S., Mitchell-Olds, T., Boles, S., … Juenger, T. E. (2008). Genetics of Drought Adaptation in Arabidopsis Thaliana Ii. Qtl Analysis of a New Mapping Population, Kas-1 × Tsu-1. *Evolution*, *62*(12), 3014–3026. https://doi.org/10.1111/j.1558-5646.2008.00474.x

Mojica, J. P., Mullen, J., Lovell, J. T., Monroe, J. G., Paul, J. R., Oakley, C. G., & McKay, J. K. (2016). Genetics of water use physiology in locally adapted Arabidopsis thaliana. *Plant Science*, *251*(Supplement C), 12–22. https://doi.org/10.1016/j.plantsci.2016.03.015

Muchow, R. C., & Sinclair, T. R. (1989). Epidermal conductance, stomatal density and stomatal size among genotypes of Sorghum bicolor (L.) Moench. *Plant, Cell & Environment*, *12*(4), 425–431. https://doi.org/10.1111/j.1365-3040.1989.tb01958.x

Nordborg, M., Borevitz, J. O., Bergelson, J., Berry, C. C., Chory, J., Hagenblad, J., … Weigel, D. (2002). The extent of linkage disequilibrium in Arabidopsis thaliana. *Nature Genetics*, *30*(2), 190–193. https://doi.org/10.1038/ng813

Ohsumi, A., Kanemura, T., Homma, K., Horie, T., & Shiraiwa, T. (2007). Genotypic Variation of Stomatal Conductance in Relation to Stomatal Density and Length in Rice (Oryza sativa L.). *Plant Production Science*, *10*(3), 322–328. https://doi.org/10.1626/pps.10.322

Paccard, A., Fruleux, A., & Willi, Y. (2014). Latitudinal trait variation and responses to drought in Arabidopsis lyrata. *Oecologia*, *175*(2), 577–587. https://doi.org/10.1007/s00442-014-2932-8

Passioura, J. B. (1996). Drought and drought tolerance. In *Drought Tolerance in Higher Plants: Genetical, Physiological and Molecular Biological Analysis* (pp. 1–5). Springer, Dordrecht. https://doi.org/10.1007/978-94-017-1299-6_1

Pearce, D. W., Millard, S., Bray, D. F., & Rood, S. B. (2006). Stomatal characteristics of riparian poplar species in a semi-arid environment. *Tree Physiology*, *26*(2), 211–218. https://doi.org/10.1093/treephys/26.2.211

Pillitteri, L. J., & Torii, K. U. (2012). Mechanisms of Stomatal Development. *Annual Review*
*of Plant Biology*, *63*(1), 591–614. https://doi.org/10.1146/annurev-arplant-
042811-105451
Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2015). *nlme: Linear and*
*Nonlinear Mixed Effects Models*. Retrieved from http://CRAN.R-
project.org/package=nlme
Postma, F. M., & Ågren, J. (2016). Early life stages contribute strongly to local adaptation
in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences*, *113*(27),
7590–7595. https://doi.org/10.1073/pnas.1606303113
R Development Core Team. (2008). *R: A language and environment for statistical*
*computing*. Retrieved from http://www.R-project.org
Raven, J. A. (2002). Selection pressures on stomatal evolution. *New Phytologist*, *153*(3),
371–386. https://doi.org/10.1046/j.0028-646X.2001.00334.x
Raven, J. A. (2014). Speedy small stomata? *Journal of Experimental Botany*, *65*(6), 1415–
1424. https://doi.org/10.1093/jxb/eru032
Reich, P. B. (1984). Leaf Stomatal Density and Diffusive Conductance in Three
Amphistomatous Hybrid Poplar Cultivars. *New Phytologist*, *98*(2), 231–239.
https://doi.org/10.1111/j.1469-8137.1984.tb02733.x
Reyes, J. L., Rodrigo, M.-J., Colmenero-Flores, J. M., Gil, J.-V., Garay-Arroyo, A., Campos, F.,
… Covarrubias, A. A. (2005). Hydrophilins from distant organisms can protect
enzymatic activities from water limitation effects in vitro. *Plant, Cell &*
*Environment*, *28*(6), 709–718. https://doi.org/10.1111/j.1365-
3040.2005.01317.x
Seren, Ü., Grimm, D., Fitz, J., Weigel, D., Nordborg, M., Borgwardt, K., & Korte, A. (2017).
AraPheno: a public database for Arabidopsis thaliana phenotypes. *Nucleic Acids*
*Research*, *45*(D1), D1054–D1059. https://doi.org/10.1093/nar/gkw986
Trabucco, A., & Zomer, R. (2010). Global soil water balance geospatial database. *CGIAR*
*Consortium for Spatial Information, Published Online, Available from the CGIAR-CSI*
*GeoPortal at: Http://Www. Cgiar-Csi. Org (Last Access: January 2013)*.
Verslues, P. E., Govinal Badiger, B., Ravi, K., & M. Nagaraj, K. (2013). Drought tolerance
mechanisms and their molecular basis. In  tthew A. Jenks & P. M. Hasegawa
(Eds.), *Plant Abiotic Stress* (pp. 15–46). John Wiley & Sons, Inc.
https://doi.org/10.1002/9781118764374.ch2
Vilhjálmsson, B. J., & Nordborg, M. (2012, November 20). The nature of confounding in
genome-wide association studies [Comments and Opinion].
https://doi.org/10.1038/nrg3382
Vu, V. Q. (2011). A ggplot2 based biplot (Version 0.55).
Weinig, C., Ewers, B. E., & Welch, S. M. (2014). Ecological genomics and process modeling
of local adaptation to climate. *Current Opinion in Plant Biology*, *18*(Supplement C),
66–72. https://doi.org/10.1016/j.pbi.2014.02.007
Whitlock, M. C., & Guillaume, F. (2009). Testing for Spatially Divergent Selection:
Comparing QST to FST. *Genetics*, *183*(3), 1055–1063.
https://doi.org/10.1534/genetics.108.099812
Wickham, H. (2009). *Ggplot2: elegant graphics for data analysis*. New York: Springer.
Wood, A. R., Esko, T., Yang, J., Vedantam, S., Pers, T. H., Gustafsson, S., … Frayling, T. M.
(2014). Defining the role of common variation in the genomic and biological

architecture of adult human height. *Nature Genetics*, *46*(11), 1173.
https://doi.org/10.1038/ng.3097
Wu, C. A., Lowry, D. B., Nutter, L. I., & Willis, J. H. (2010). Natural variation for drought-
response traits in the Mimulus guttatus species complex. *Oecologia*, *162*(1), 23–
33. https://doi.org/10.1007/s00442-009-1448-0
Xu, Z., & Zhou, G. (2008). Responses of leaf stomatal density to water status and its
relationship with photosynthesis in a grass. *Journal of Experimental Botany*,
*59*(12), 3317–3325. https://doi.org/10.1093/jxb/ern185
Yoo, C. Y., Pence, H. E., Jin, J. B., Miura, K., Gosney, M. J., Hasegawa, P. M., & Mickelbart, M.
V. (2010). The Arabidopsis GTL1 Transcription Factor Regulates Water Use
Efficiency and Drought Tolerance by Modulating Stomatal Density via
Transrepression of SDD1. *The Plant Cell Online*, *22*(12), 4128–4141.
https://doi.org/10.1105/tpc.110.078691
Yu, H., Chen, X., Hong, Y.-Y., Wang, Y., Xu, P., Ke, S.-D., … Xiang, C.-B. (2008). Activated
Expression of an Arabidopsis HD-START Protein Confers Drought Tolerance with
Improved Root System and Reduced Stomatal Density. *The Plant Cell*, *20*(4),
1134–1151. https://doi.org/10.1105/tpc.108.058263
919
920

## Data accessibility

Raw image data and image analysis scripts are available upon request and will be stored in a

Dryad repository upon acceptance. Phenotypic data is provided in the supplemental material

and will be uploaded to the AraPheno database (https://arapheno.1001genomes.org, (Seren et

al., 2017) and stored in a Dryad repository upon acceptance. Additionally, we provide an R

Markdown file, which contains all figures (except GWAS and MTMM) and the

corresponding R code used to create the figures and statistics in the supplemental material.

GWAS scripts are available at https://github.com/arthurkorte/GWAS. MTMM scripts are

available at https://github.com/Gregor-Mendel-Institute/mtmm.

Genomic data used is publicly available in the 1001 genomes database (Alonso-Blanco et al.,

2016)

## Author contributions

JdM, AK, and HD conceived the study. HD conducted the experiment and produced

phenotypic data for stomata traits. TM and AW were responsible for $\delta^{13}C$ measurements. GM

provided data on historic drought regimen. JdM, AK and HD were responsible for the

statistical analyses of the data. JdM and HD wrote the manuscript with significant

contributions from AK, TM, AW and GM.

941
942 **Figure 1: Natural variation in stomata patterning**

943 Stomata density and size were measured for 330 natural genotypes of *A. thaliana*. The plot shows
944 genotypic means of stomata density and stomata size. Dots are colored based on the geographical
945 origin of each accession. The red line shows a linear fit and gray shadows indicate the error of the fit.
946 Pearson's product-moment correlation r=-0.5, p<0.001.
947

**Figure 2: Stomata size correlates with water-use efficiency**

$\delta^{13}$C was measured for all plants in block 1. Plots show correlation of stomata size (block 1 only) with $\delta^{13}$C. $\delta^{13}$C is expressed as ‰ against the Vienna Pee Dee Belemnite (VPDB) standard. The red line shows a linear fit and gray shadows indicate the error of the fit. Pearson's product-moment correlation: r=-0.18, p=0.004. Correlation of $\delta^{13}$C and stomata size is not only driven by the Spanish outlier (correlation without outlier: r=-0.16, p=0.009). Genetic correlation was calculated using the MTMM approach: r=-0.58, p<0.05.

**Figure 3: Stomata patterns correlate with geographical patterns of climatic variation**

Correlation between stomata patterns and seven climatic principal components (PCs) was tested for each phenotype using a Generalized Linear Model (GLM) including genetic population structure as described by the 20 first genetic PCs. Plots are effect plots based on the GLM (see methods), showing the correlation between stomata size two climatic PCs. Black arrows indicate correlation with the climatic variables showing the strongest loadings for the respective principal component. Plots show the linear fit (red solid line) and the smoothed fit of partial residuals (gray) of the specific predictor. Gray dots are partial residuals. The red shade shows the error of the linear fit. Both principal components shown here are significant predictors of the respective response variable (p<0.05).

**Figure 4: Significant regional differentiation of stomata size and δ¹³C**

*A. thaliana* accessions were grouped based on their geographical origin. Boxplots show regional differentiation of stomata size (top) and δ¹³C (bottom). Significance of differentiation was tested using Generalized Linear Models followed by a post-hoc test. Statistical significance is indicated by letters on top: Groups that do not share a common letter are significantly different. Significance levels: top) a-c, a-bc: p<0.001; ab-c: p<0.05; bottom) a-c, a-b: p<0.001, b-c: p<0.01, ab-c: p<0.05.

**A) Stomata size**

| $Q_{ST}$ \ $Q_{ST}$-$F_{ST}$ | Central Europe | North. Sweden | South. Sweden | Spain | West. Europe |
|---|---|---|---|---|---|
| Central Europe | | -0.32 | **0.29** | -0.17 | -0.13 |
| Northern Sweden | 0.15 | | -0.31 | -0.38 | -0.51 |
| Southern Sweden | 0.41 | 0.09 | | **0.12** | -0.03 |
| Spain | 0.01 | 0.06 | 0.32 | | -0.18 |
| West. Europe | 0.02 | <0.01 | 0.21 | <0.01 | |

**B) Stomata density**

| $Q_{ST}$ \ $Q_{ST}$-$F_{ST}$ | Central Europe | North. Sweden | South. Sweden | Spain | West. Europe |
|---|---|---|---|---|---|
| Central Europe | | -0.37 | **0.31** | -0.16 | **0.17** |
| Northern Sweden | 0.09 | | -0.24 | -0.44 | -0.49 |
| Southern Sweden | 0.44 | 0.16 | | **0.17** | -0.19 |
| Spain | 0.01 | 0.01 | 0.36 | | 0.07 |
| West. Europe | 0.32 | 0.02 | <0.01 | 0.26 | |

**C) $\delta^{13}$C**

| $Q_{ST}$ \ $Q_{ST}$-$F_{ST}$ | Central Europe | North. Sweden | South. Sweden | Spain | West. Europe |
|---|---|---|---|---|---|
| Central Europe | | **0.21** | **0.28** | **0.13** | -0.07 |
| Northern Sweden | 0.7 | | -0.40 | -0.16 | -0.01 |
| Southern Sweden | 0.4 | 0.01 | | -0.08 | -0.01 |
| Spain | 0.30 | 0.28 | 0.11 | | -0.12 |
| West. Europe | 0.07 | 0.40 | 0.17 | 0.05 | |

 **Table 1 A-C: Patterns of regional differentiation depart from neutral expectations**

Pairwise $Q_{ST}$ estimates were derived from linear mixed models for all regions. Genome-wide, pairwise $F_{ST}$ distribution was calculated based on 70,000 SNPs for all regions. In the top half of each table, the difference $Q_{ST}$-$F_{ST}$ for each pair of regions is shown. In the bottom half of each table the $Q_{ST}$ estimate for each pair of regions is shown. Each table represents one phenotype as indicated by table headlines. Significant $Q_{ST}$-$F_{ST}$ differences are written in bold. The significance threshold is based on the 95[th] percentile of a distribution of maximum $Q_{ST}$-$F_{ST}$ values from 1000 random permutations of phenotypic data.