1
8 **Dendritic spikes in hippocampal granule cells are necessary for long-term**
9 **potentiation at the perforant path synapse**
10
11
12

13 Sooyun Kim*, Yoonsub Kim, Suk-Ho Lee and Won-Kyung Ho*

14
15
16

17 Department of Physiology, Seoul National University College of Medicine, 103

18 Daehak-ro, Jongno-gu, Seoul, 110-799, Korea

*For correspondence:
Drs. Sooyun Kim and Won-Kyung Ho
Department of Physiology
Seoul National University College of Medicine
103 Daehak-ro, Jongno-gu,
Seoul, 110-799, South Korea
Tel: +82-2-3668-7981
Fax: +82-2-763-9667

e-mail: sooyun.kim@snu.ac.kr, wonkyung@snu.ac.kr

Abstract

Long-term potentiation (LTP) of synaptic responses is essential for hippocampal memory function. Perforant-path (PP) synapses on hippocampal granule cells (GCs) contribute to the formation of associative memories, which are considered the cellular correlates of memory engrams. However, the mechanisms of LTP at these synapses are not well understood. Due to sparse firing activity and the voltage attenuation in their dendrites, it remains unclear how associative LTP at distal synapses occurs. Here we show that NMDA receptor-dependent LTP can be induced at PP-GC synapses without backpropagating action potentials (bAPs) in acute rat brain slices. Dendritic recordings reveal substantial attenuation of bAPs as well as local dendritic Na$^+$-spike generation during PP-GC input. Inhibition of Na$^+$-spikes impairs LTP suggesting that LTP at PP-GC synapse requires local Na$^+$-spikes. Thus, dendritic spikes are essential for LTP induction at PP-GC synapse and may constitute a key cellular mechanism for memory formation in the dentate gyrus.

Introduction

The cortico-hippocampal circuit is implicated in the formation, storage, and retrieval of spatial and episodic memories (Lisman, 1999). The dentate gyrus (DG), the first stage in the hippocampal circuitry, receives abundant excitatory projections from the entorhinal cortex via the perforant-path (PP) synapses. Theoretical models of hippocampal function propose that the DG is critically involved in pattern separation and that synaptic transmission and plasticity at PP-

granule cell (GC) synapses in the DG is required to remove redundant memory

representations (Marr, 1971; McNaughton and Morris, 1987; Treves & Rolls,

1994). In agreement with theoretical predictions, knockout of *N*-methyl-D-

aspartate (NMDA) receptors in GCs impairs long-term potentiation (LTP) and the

ability to rapidly form a contextual representation and discriminate it from previous

similar memories in a contextual fear conditioning task (McHugh et al., 2007).

Moreover, GCs that were activated by contextual fear conditioning, referred to as

memory engram cells, present clear signatures of synaptic potentiation such as

a larger AMPA-NMDA ratio and a greater density of dendritic spines (Ryan et al.,

2015). Thus, knowledge of plasticity at PP-GC synapses is essential for

understanding the hippocampal function.

Attempts to understand synaptic plasticity rules at PP-GC synapses date

to studies of Bliss and Lomo (1973), who first demonstrated LTP in the

hippocampus by high-frequency stimulation of the PP fibers *in vivo*. More recently,

theta-burst high-frequency stimulation (TBS) of the PP has been used to induce

LTP at PP-GC synapses (Schmidt-Hieber et al., 2004; McHugh et al., 2007; Ge

et al., 2007). However, the underlying mechanisms for the induction of LTP with

TBS at these synapses remain unclear. Associative forms of synaptic plasticity

depend on a presynaptic activity (e.g. excitatory postsynaptic potential, EPSP)

and a postsynaptic signal (e.g. action potential, AP). Classically, a

backpropagating AP (bAP) provides the associative postsynaptic signal at the

synaptic site for the induction of LTP (Hebb, 1949; Magee & Johnston, 1997; Dan

& Poo, 2006; Feldman, 2012). However, axosomatic APs are poorly propagated

back into the dendrites of the GCs (Krueppel et al., 2011), and are unlikely to

contribute to the induction of LTP. In addition, mature GCs are relatively silent

during exploration (Schmidt-Hieber et al., 2014; Diamantaki et al., 2016) and fire

with a low number of APs. Thus, spike-timing dependent plasticity (STDP) that

depends on an axosomatic postsynaptic AP will rarely occur under natural

conditions (Feldman, 2012). The pronounced attenuation of AP backpropagation,

the low occurrence of APs, along with the distinct intrinsic features of mature GCs,

such as a hyperpolarized resting membrane potential and reduced excitability

(Scharfman & Schwartzkroin, 1990; Mongiat et al., 2009; Pernía-Andrade &

Jonas, 2014), cannot explain how distal synaptic PP inputs can be potentiated.

Resolving this question has critical implications for understanding both the

mechanism and function of memory encoding and retrieval (Ryan et al., 2015).

Understanding LTP induction at PP synapses innervating distal dendrites

of GCs requires a detailed analysis of dendritic electrogenesis (Sjöström et al.,

2008). Synaptic plasticity at distal synapses may occur in the absence of

axosomatic APs via local dendritic spikes (Golding et al., 2002; Kim et al., 2015).

In GCs, linear somatic responses can be seen when glutamate is applied in the

dendrites (Krueppel et al., 2011). However, without directly accessing the

electrical properties of the distal GC dendrites, it is still not known whether

synaptic stimulation can generate local dendritic spikes in GCs.

To address this question, we performed subcellular patch-clamp

recordings on the thin dendrites of GCs. We found that inhibition of dendritic Na$^+$

channels prevented LTP induction via TBS at PP-GC synapses. Because

dendritic Na$^+$ channels could generate local spikes that were independent of

axosomatic AP generation and were not affected by voltage attenuation, we

suggest that Na$^+$ spikes in the dendrites provide the postsynaptic signal

necessary for the induction of LTP at PP-GC synapses.

Results

We examined GCs with an input resistance (R_{in}) < 200 MΩ (R_{in} = 108.4 ± 2.6 MΩ;

resting membrane potential: –81.3 ± 0.2 mV; see 'Materials and Methods') which

corresponds to the mature GC population in the acute hippocampal slices from

rats (Schmidt-Hieber et al., 2004).

TBS-induced LTP at the PP-GC synapses does not require postsynaptic

bAPs

We first induced long-term potentiation (LTP) in GCs by theta-burst stimulation

(TBS) of the PP synapses in the outer third of the molecular layer (**Figure 1A**).

To activate the PP synapses, the tip of a stimulation electrode was placed in close

proximity to the dendrite (< 50 μm) under the visual guidance of fluorescent image

of the dendrite ('Materials and Methods'). To ensure that no axosomatic AP

initiation and backpropagation occur during TBS, we locally applied tetrodotoxin

(TTX) to the GC axon, soma, and proximal dendrites in a subset of experiments

(6 out of 13 experiments). TBS-induced LTP caused an increase in the amplitude

of the excitatory postsynaptic potentials (EPSPs; **Figure 1B,C**; from 6.96 ± 0.40

mV to 9.96 ± 0.86 mV, n = 13, *P* < 0.01). During TBS, activation of the PP

synapses often produced voltage responses with a fast depolarizing phase

similar to somatic events observed during dendritic spike generation in other

types of neurons (**Figure 1D**; Golding & Spruston, 1998; Golding et al., 2002;

Jarsky et al., 2005; Losonczy et al., 2006; Kim et al., 2012). Consistent with

previous results (Golding & Spruston, 1998; Golding et al., 2002; Losonczy et al.,

2006; Kim et al., 2015), various shapes of putative dendritic spikes were observed

during TBS (**Figure 1D**). By examining the temporal derivative (dV/dt) of somatic

voltage responses, we identified weak (dV/dt < 10 mV/ms) and strong (dV/dt >

10 mV/ms) putative dendritic spikes that were distinguishable from EPSPs

without any local regenerative events (Without dendritic spikes, black, 1.3 ± 0.01

mV/ms, n = 2498; weak dendritic spike, orange, 4.1 ± 0.4 mV/ms, n = 18; strong

dendritic spike, red, 14.5 ± 0.8 mV/ms, n = 11; Kruskal-Wallis test: P < 0.0001;

Figure 1D). These putative dendritic spikes were accompanied by a sustained

plateau potential (**Figure 1D**; see also **Figure 2B**). Interestingly, the presence of

these weak and strong putative dendritic spikes during TBS was correlated with

LTP induction (**Figure 1E**; in the presence of putative dendritic spikes, 173.2 ±

12.1%, n = 7; in the absence of putative dendritic spikes 114.1 ± 12.8%, n = 6, P

< 0.005). Indeed, we found the strong correlation between the number of putative

dendritic spikes observed during TBS and the magnitude of LTP (**Figure 1F**; r =

0.77; P < 0.005; n = 13). To test the contribution of bAPs in this form of LTP, we

applied strong synaptic stimulation without perisomatic TTX application (**Figure**

1–figure supplement 1A). The presence of axosomatic spikes did not

significantly affect the magnitude of LTP indicating that bAPs are not critical for

this form of LTP (**Figure 1–figure supplement 1B,C**; 167.9 ± 26.8%, n = 8, P =

0.87, compared to control in **Figure 1I**). These results suggest that dendritic

spikes but not axosomatic APs contribute to LTP induction at PP-GC synapses.

To further test whether axosomatic spikes contribute to the induction of

LTP at the PP-GC synapse, we used associative pairing protocols. Presynaptic

activity was either followed (pre-postsynaptic sequence) or preceded (post-

presynaptic sequence) at a 10-ms interval by single or AP bursts (2 APs at 100

Hz). Both protocols failed to induce significant changes in EPSP amplitude

(**Figure 1G,H**; pre-postsynaptic sequence, control: 4.97 ± 0.38 mV, after

induction: 5.83 ± 0.86 mV, n = 15, P = 0.23; post-presynaptic sequence, control:

4.78 ± 0.54 mV, after induction: 4.91 ± 0.99 mV, n = 9, P = 0.54), suggesting that

APs of axonal origin are not important for potentiation. We further performed

similar experiments at the medial PP–GC synapses by stimulating axons in the

middle third of the molecular layer (**Figure 1–figure supplement 2A**). Pairing

EPSPs and APs in both pre-postsynaptic and post-presynaptic sequences did

not induce LTP of EPSPs (**Figure 1–figure supplement 2B–E**; pre-postsynaptic

sequence, control: 4.86 ± 0.40 mV, after induction: 5.09 ± 0.99 mV, n = 7, P =

0.9015; post-presynaptic sequence, control: 3.86 ± 0.26 mV, after induction: 4.27

± 0.57 mV, n = 6, P = 0.56), consistent with the previous studies that reported no

synaptic potentiation after similar pairing protocols (Yang & Dani, 2014; Lopez-

Rojas et al., 2016). Together, these results imply that dendritic spikes, rather than

axosomatic APs, are the essential signal for LTP induction at distal GC synapses.

LTP by TBS at PP–GC synapses requires NMDARs and Na$^+$ channels

We next examined the receptors involved in LTP expression at the PP-GC

synapses. Bath application of the NMDAR antagonist DL-AP5 (50 µM) abolished

LTP (**Figure 2A,D,E**; control: 8.89 ± 0.95 mV; after induction: 9.88 ± 1.21 mV, n

= 9, P = 0.50). While sustained plateau potentials during LTP induction were

inhibited by DL-AP5 in the external solution, fast rising events were remained

unchanged (**Figure 2B**). We thus tested whether dendritic voltage-gated Na^+

channels are involved in LTP. Including 5 mM QX-314 in the whole-cell patch

pipette abolished both plateau potentials and TBS-induced LTP (**Figure 2B–D**;

control: 11.5 ± 2.5 mV, after induction: 10.6 ± 2.4 mV, n = 8, P = 0.38). These

results indicate that TBS-induced LTP in GCs depends on the activation of

NMDARs and voltage-gated Na^+ channels on the postsynaptic dendritic

membrane, reinforcing the idea that dendritic Na^+ spikes may play a pivotal role

in TBS induction at distal synapses.

Backpropagation of axosomatic APs in the dendrites of GCs

To directly dissect the dendritic mechanism that determines the induction rule of

synaptic plasticity in GCs, we employed subcellular patch-clamp techniques to

analyze AP backpropagation and initiation in GC dendrites (**Figure 3A**). First, we

characterized backpropagation in GCs with somatic current injection evoking

trains of APs at the soma while simultaneously recording in the dendrite (**Figure**

3B). The AP always appeared first in the somatic recording and then in the

dendrites (**Figure. 3C**). Similar to the previous report (Krueppel et al., 2011), the

peak amplitude of the bAPs attenuated as a function of distance from the soma

(**Figure 3D,E**). At dendritic distances beyond 150 μm from the soma, the peak

amplitude of bAPs was reduced to ~36% of the amplitude of somatic APs (soma:

93.6 ± 1.7 mV; dendrite: 33.7 ± 3.1 mV; n = 10, P < 0.005; **Figure 3D**). The

attenuation per dendritic length is much more pronounced in GCs compared to

the other neocortical (Stuart et al., 1997; Nevian et al., 2007) and hippocampal

pyramidal neurons (Spruston et al., 1995; Kim et al., 2012). It is interesting to

note that the extent of bAP attenuation when normalized to the overall length of

GC dendrites (mean dendritic length: 278 ± 7.4 µm; n = 11) is quite similar to

layer 5 pyramidal neuron dendrites (cf. Figure S3B in Nevian et al., 2007; see

also Krueppel et al., 2011; **Figure 3F**), suggesting that the failure of LTP induction

by pairing protocol at distal GC synapses (**Figure 1G–I**) might be explained by

the voltage attenuation of bAPs, as was reported in layer 5 pyramidal neurons

(Letzkus et al., 2006; Sjöström et al., 2008; Feldman; 2012). We also estimated

the conduction velocity of bAPs by analyzing the AP latencies measured at the

half-maximal amplitude of the rising phase. APs were initiated axonally and

propagated back into the dendrites with a velocity of 226 µm/ms (**Figure 3G**; n =

56; corresponding to 0.2–0.3 m/s, Senzai & Buzsaki, 2017). This conduction

velocity is slower than those measured in the apical and basal dendrites of layer

5 pyramidal neurons (apical dendrites, 508 µm/ms; basal dendrites, 341 µm/ms;

Nevian et al., 2007 and Stuart et al., 1997) and is similar or lower than the velocity

estimated in the apical dendrite of other hippocampal principal neurons (Spruston

et al., 1995; Kim et al., 2012). In summary, these experiments show that bAPs in

GCs propagate into the dendrites with substantial voltage attenuation and

moderate conduction velocity.

Ionic mechanisms of AP backpropagation

To determine the ionic mechanisms underlying the strong voltage attenuation and

the moderate conduction velocity of bAPs, we assayed the somatic and dendritic

distribution of voltage-gated Na^+ and K^+ conductances in outside-out patches

isolated at various locations using pipettes of similar open tip resistance (soma:

17.9 ± 0.5 MΩ, n = 24; dendrite: 19.4 ± 0.5 MΩ, n = 36; **Figure 4**). To calculate

conductance density from the measured current amplitude of each component,

the surface area of outside-out patches was determined by capacitance

measurements (**Figure 4–figure supplement 1**; Schmidt-Hieber &

Bischofberger, 2010; Hu & Jonas, 2014). Depolarizing voltage pulses from −120

mV to 0 mV evoked TTX-sensitive inward Na^+ currents in the majority of outside-

out patches excised from both soma and dendrites (**Figure 4A** and **Figure 4–**

figure supplement 2&3). Pooled data demonstrated that a moderate density of

Na^+ channels is distributed over the dendritic membrane (**Figure 4D**). On average,

the peak Na^+ conductance density was not significantly different between somatic

and dendritic patches (**Figure 4D**, soma, 184.5 ± 36.5 $pS/\mu m^2$, n = 19; proximal

dendrite (within 100 μm; PD), 136.4 ± 39.5 $pS/\mu m^2$, n = 8; Distal dendrite (beyond

100 μm; DD), 206.2 ± 38.6 $pS/\mu m^2$, n = 21; soma vs PD and soma vs DD, $P >$

0.99; PD vs DD, $P = 0.98$, Kruskal-Wallis test with Dunn's multiple comparisons).

Surprisingly, a very high density of transient K^+ currents that appeared to activate

and inactivate rapidly in response to voltage pulses from −120 mV to +70 mV was

found in patches obtained from the dendrites (**Figure 4B**). These inactivating A-

type components were reduced by the A-type K^+ channel blocker 4-

aminopyridine (4-AP, **Figure 4B** and **Figure 4–figure supplement 2&3**). Plotting

the conductance density along the dendrite demonstrated that GC dendrites

show a significantly higher density of A-type K^+ components than the soma

(**Figure 4E,** soma, 384.8 ± 54.3 $pS/\mu m^2$, n = 21; PD, 784.9 ± 107.9 $pS/\mu m^2$, n =

7; DD, 1319.9 ± 181.6 pS/µm^2, n = 21; soma vs. PD, P = 0.07; soma vs. DD, P <

0.0001; PD vs. DD, P = 0.76, Kruskal-Wallis test with Dunn's multiple

comparisons). Consistent with this finding, we found that bath application of 5

mM 4-AP in simultaneous somatic and dendritic voltage recordings results in a

significant enhancement of the peak amplitude and duration of bAPs in the

dendrites, indicating that dendritic A-type K$^+$ current (I$_A$) contributes to voltage

attenuation of bAPs in GCs (**Figure 4–figure supplement 4**). Finally, the delayed

rectifier K$^+$ current components showed a low and uniform expression over the

entire somatodendritic axis (**Figure 4C,F**; soma, 247.9 ± 28.0 pS/µm^2, n = 21;

PD, 231.5 ± 44.3 pS/µm^2, n = 7; DD, 280.5 ± 46.7 pS/µm^2, n = 22; P > 0.99 for

all cases, Kruskal-Wallis test with Dunn's multiple comparisons). Taken together,

these voltage-clamp data reveal that a markedly high density of K$^+$ channels and

a moderate and uniform density of Na$^+$ channels are present in the dendrites of

GCs, suggesting the specific biophysical mechanisms underlying both the strong

dendritic AP attenuation and the moderate AP propagation velocity.

Dendritically initiated Na$^+$ spikes are required for TBS-induced LTP at PP–

GC synapses

Voltage-gated Na$^+$ channels may contribute to the generation of

dendritically initiated local spikes in regions where the dendrites are very small

(Holmes, 1989). Indeed, depolarizing dendrites with brief current pulse injection

evoked local spikes in distal dendrites of GCs (55 of 63 recordings; **Figure 5A,B**).

In the proximal domain (within 100 µm from the soma), dendritic spikes were not

detected and depolarizing dendrites resulted in an axosomatic AP that

propagated back to the dendritic recording site (**Figure 5C**). At dendritic recording

locations 70 to 150 µm from the soma, dendritic spikes were associated with

axosomatic spikes. For distances larger than 150 µm from the soma

(approximately corresponding to the outer molecular layer), current injections

robustly initiated isolated dendritic spikes (**Figure 5C**). Pharmacological analysis

revealed that dendritic spikes were resistant to 200 µM $CdCl_2$ and 50 µM $NiCl_2$

but were inhibited by 0.5 µM TTX, indicating that they were mediated by dendritic

voltage-gated Na^+ channels rather than by Ca^{2+} channels (TTX, $11.3 \pm 3.3\%$, n =

3; $CdCl_2$, $93.9 \pm 7.8\%$, n = 4; $NiCl_2$, $97.9 \pm 10.2\%$, n = 4; **Figure 5–figure**

supplement 1). Because our results indicate that GC dendrites contain a high

density of transient, A-type K^+ channels (**Figure 4B**), we further explored how

these channels influence dendritic spike initiation. We combined dual

somatodendritic recordings and focal application of 4-AP (10 mM) to the dendritic

patch (**Figure 5–figure supplement 2A**). Local block of I_A significantly decreased

the current threshold for dendritic spike initiation, suggesting that dendritic I_A

affect the generation of dendritic spikes (**Figure 5–figure supplement 2B,C**).

We finally examined whether Na^+ spikes in GC dendrites can be evoked by

synaptic stimulation of PP inputs. Extracellular synaptic stimulation was

combined with simultaneous double patch-clamp recordings from the soma and

dendrites (**Figure 5D**). Dendritic spikes could be triggered by stimulating the PP

inputs with TBS protocol (**Figure 5D**) and were inhibited by the application of 10

nM TTX in the bath (**Figure 5E**; Kim et al., 2015). Repeated trials of a high-

frequency PP stimulation using the same stimulus intensity produced variable

dendritic responses with subthreshold EPSP, weak dendritic spikes, or strong

dendritic spikes that appeared either in isolation or associated with axosomatic

APs (**Figure 5–figure supplement 3A,B**). When dendritic spikes were present,

the dV/dt of the corresponding somatic voltages showed variable peak

amplitudes that were similar to those of the putative dendritic spikes during TBS

induction shown in **Figure 1D** (**Figure 5–figure supplement 3C,D**, EPSPs with

a dendritic spike: $dV/dt = 7.2 \pm 1.1$ mV/ms, n = 25; $P = 0.77$ compared to EPSPs

with a putative dendritic spike: $dV/dt = 8.0 \pm 1.0$ mV/ms, n = 29, **Figure 1D**).

Finally, low concentration of TTX (10 nM) prevented TBS-induced LTP (**Figure**

5E–H; control: 6.58 ± 0.51 mV; after induction: 7.04 ± 0.88 mV, n = 8; $P = 0.64$),

without affecting the baseline EPSP amplitude (**Figure 5–figure supplement 4**;

control: 5.21 ± 0.78 mV; after TTX: 5.06 ± 0.68 mV, n = 11; $P = 0.83$). All together,

these results suggest that dendritically generated local Na^+ spikes in response to

TBS of the PP are necessary for LTP induction in GCs.

Discussion

In summary, the present study demonstrates several major findings. First, our

results show that conventional STDP protocols (Dan & Poo, 2006; Feldman, 2012)

do not trigger synaptic potentiation at PP-GC synapses. Second, a

physiologically relevant TBS paradigm can efficiently induce LTP in GCs without

axosomatic APs. Finally, direct dendritic recordings revealed that LTP induction

requires dendritic spikes. To our knowledge, these studies show the first direct

evidence for dendritic spike generation in GCs and the role of these spikes during

induction of LTP in the dentate gyrus network. However, whether our findings

also hold for GCs at early stages of development remains to be determined.

Considerable evidence has shown that the electrical properties of neurons

at the cellular and subcellular levels are brain region-specific and cell type-

specific and endow unique rules and characteristics on circuit function (Sjöström

et al., 2008; Stuart & Spruston, 2015). Our findings support this idea.

Hyperpolarized resting potential, strong dendritic voltage attenuation, and low

occurrence of APs are the known electrophysiological characteristics of mature

GCs (Scharfman & Schwartzkroin, 1990; Schmidt-Hieber et al., 2004; Mongiat et

al., 2009; Krueppel et al., 2011; Pernía-Andrade & Jonas, 2014), which are

distinct from other types of hippocampal principal neurons (Spruston et al., 1995;

Kim et al., 2012; Spruston & Stuart, 2015). While backpropagated APs are an

important associative signal for triggering plasticity at the synaptic site via the

voltage-dependent relief of Mg^{2+} block of NMDARs (Magee & Johnston, 1997;

Dan & Poo, 2006; Feldman, 2012; Mishra et al., 2016), the above features of

mature GCs are highly unfavorable for LTP induction at distal synaptic contacts.

Accordingly, we found that GCs do not show LTP at PP-GC synapses during

standard STDP (**Figure 1G,H** and **Figure 1–figure supplement 2**), which is

consistent with recent reports (Yang & Dani, 2014; Lopez-Rojas et al., 2016).

However, several studies have demonstrated EPSP-AP pairing protocol-

induced synaptic potentiation at the same synapses (Levy & Steward, 1983; Lin

et al., 2006). Given that Levy and Steward (1983) and Lin et al. (2006) employed

in vivo and *in vitro* field recording configurations, respectively, the discrepant

results could be attributed to GCs at different stages of maturation as immature

GCs exhibit a lower threshold for LTP induction (Schmidt-Hieber et al., 2004; Ge

et al., 2007) or under different recording circumstances that were exposed to

various neuromodulators. For example, Yang and Dani (2014) reported that the

pairing protocol that showed no synaptic potentiation could induce reliable LTP

at PP-GC synapses after D1-type dopamine receptor activation by which

dendritic A-type K$^+$ currents are suppressed. Because I_A is known to limit the

backpropagation of APs (Hoffman et al., 1997), suppresion of I_A can boost AP

backpropagation, allowing sufficient dendritic depolarization for LTP induction at

distal synapses. Our observations of a high density of dendritic I_A and their effects

on AP backpropagation in GCs directly support the finding of Yang and Dani

(2014). Therefore, under physiological conditions, long-lasting dendritic

depolarization during theta oscillation (Buzsaki, 2002) or activation of

neuromodulatory systems (Hamilton et al., 2010; Yang & Dani, 2014) may cause

inactivation of dendritic I_A and trigger pairing-induced LTP (Lin et al., 2006;

Brunner & Szabadics, 2016).

Although the presence of A-type K$^+$ channels in GC dendrites had been

shown in earlier immunocytochemical studies (Birnbaum et al., 2004; Monaghan

et al., 2008; Menegola et al., 2008), it has been proposed that dendritic A-type K$^+$

channels have no significant impact on bAP-induced Ca^{2+} signals (Krueppel et

al., 2011). Krueppel et al.'s (2011) results appear to be inconsistent with our

present data, which show the strong expression of functional A-type K$^+$ channels

in GC dendrites. In contrast to that study, we directly tested the effect of 4-AP in

simultaneous soma-dendrite recordings by using both global and local application

methods. Our results show that the effect of local puff application of 4-AP to the

dendritic recording site is much smaller than that of global application. As

Krueppel et al. (2011) used local application halfway between the soma and the

Ca^{2+} imaging site, the impact of 4-AP on bAPs in their study could be

underestimated.

We further found that high-frequency stimulation of PP synapses is efficient

in eliciting dendritic spikes required for LTP induction. As reported in a recent *in*

vivo study, mature GCs are exposed to abundant functional glutamatergic inputs

from the entorhinal cortex during theta rhythm (Pernía-Andrade & Jonas, 2014;

Schmidt-Hieber et al., 2014). Moreover, several lines of evidence demonstrated

that mature GCs are highly innervated by PP synaptic connections while

receiving a powerful perisomatic inhibition (Dieni et al., 2013, 2016; Temprana et

al., 2015). Under these *in vivo* network conditions, strong dendritic excitation by

high-frequency PP inputs and strong perisomatic inhibition may promote

amplification of dendritic responses without axosomatic APs. Therefore, dendritic

spikes are likely physiologically relevant signals for the induction of LTP. Support

for the idea of cooperative LTP at PP-GC synapses has also come from

McNaughton et al. (1978), who demonstrated that high-frequency stimulation of

PP synapses could induce synaptic enhancement in the absence of GC

discharges.

This specific high-frequency stimulation-dependent synaptic potentiation in

GCs presumably stems from distinct functional and geometrical features of GC

dendrites. A moderate density of dendritic Na^+ channels together with higher input

impedance of the distal dendrites (Hama et al., 1989; Schmidt-Hieber et al., 2007;

Holmes, 1989) suffices for initiating spikes locally in distal dendrites that have

small capacitive load but not for supporting active backpropagation of axosomatic

APs. Although GCs have relatively short dendrites, a high K^+ to Na^+ current ratio

in these thin-caliber dendrites imposes a strong distance-dependent attenuation

of axosomatic APs, leading to a lack of pairing-induced LTP at distal synapses.

Thus, Na^+ spikes in the dendrites contribute the postsynaptic depolarization

necessary for the induction of associative plasticity (Golding et al., 2002; Kim et

al., 2015). However, it should be noted that dendritic Na^+ spikes were

accompanied by NMDAR-mediated plateau potentials and therefore these two

dendritic events could act in concert to trigger LTP in GCs (Schiller et al., 2000).

Consequently, our findings suggest that in the absence of axonal firing

(Alme et al., 2010; Diamantaki et al., 2016), dendritic spikes would allow a silent

GC to participate in the storage of memories via LTP induction. It would further

permit a functional separation between a storage phase mediated by Na^+

channels in GC dendrites, and a recall phase (O´Neil et al., 2008) that effectively

activates the CA3 neurons (Vyleta et al., 2016) via Na^+ channels in GC axons.

Materials and Methods

Slice preparation and electrophysiology

Acute hippocampal slices (thickness, 350 μm) were prepared from the brains of 17- to 25-day-old Sprague-Dawley rats of either sex. Rats were anesthetized (isofluorane, Forane®; Abbott) and decapitated rapidly. All the experiments were approved by the University Committee Animal Resource in Seoul National University (Approval #: SNU-090115-7). All brains were sliced coronally, and dorsal slices displaying all the subregions of the hippocampal formation were used for the experiments (Coronal sections located between 4.3 mm and 5.7 mm from the posterior end of the right hemisphere). Slices were prepared in an oxygenated ice-cold sucrose-containing physiological saline using a vibratome (VT1200, Leica), incubated at ~36°C for 30 min, and subsequently maintained in the same solution at room temperature until the recordings. Recordings were performed at near-physiological temperature (33–35°C) in an oxygenated artificial cerebral spinal fluid (ACSF).

Patch pipettes were made from borosilicate glass capillaries (outer diameter = 1.5 mm, inner diameter = 1.05 mm) with a horizontal pipette puller (P-97, Sutter Instruments). The open-tip resistance of patch pipettes was 2.5–6.5 MΩ and 11–30 MΩ for somatic and dendritic recordings, respectively. Current-clamp recordings were performed with an EPC-10 USB Double amplifier (HEKA Elektronik). In current-clamp recordings, series resistance was 8–80 MΩ. Pulse protocols were generated, and signals were low-pass filtered at 3 or 10 kHz (Bessel) and digitized (sampling rate: 20–50 kHz) and stored using Patchmaster software running on a PC under Window 10. Resting membrane potential (RMP)

was noted immediately after rupture of the patch membrane. R_{in} was determined

by applying Ohm's law to the steady-state voltage difference resulting from a

current step (±50 pA). Pipette capacitance and series resistance compensation

(bridge balance) were used throughout current-clamp recordings. Bridge balance

was checked continuously and corrected as required. Experiments were

discarded if the resting membrane potential depolarized above –70 mV and were

stopped if the resting membrane potential or R_{in} changed by more than 20%

during the recording.

All experiments were performed on visually identified mature GCs on the

basis of the relatively large and round-shaped somata, and the location of the cell

body under DIC optics. GCs located at the superficial side of the GC layer in the

suprapyramidal blade were preferentially targeted. These cells had the average

RMP of –81.3 ± 0.2 mV and R_{in} of 108.4 ± 2.6 MΩ (n = 165), that is similar to

characteristic intrinsic properties of mature GC population (Schmidt-Hieber et al.,

2004). Cells were filled with a fluorescent dye, Alexa Fluor 488 (50 µM, Invitrogen)

and imaged with an epifluorescence system mounted on an upright microscope

equipped with a 60 x (1.1 N.A.) water immersion objective lens. Focal electrical

stimulation (100 µs pulses of 1–35 V) was applied on isolated dendrites in the

outer third of the molecular layer (within 100 µm of the hippocampal fissure) by

placing a glass microelectrode (0.5–3 MΩ) containing 1 M NaCl or ACSF in the

vicinity of the selected dendrite (typically at < 50 µm distance), guided by the

fluorescent image of the dendrite. All experiments were performed in the

presence of the GABA receptor antagonist picrotoxin (PTX, 100 µM) and

CGP52432 (1 µM).

To record voltage-gated Na$^+$ or K$^+$ currents, outside-out patches were

excised from the soma and the dendrite with pipettes of similar geometry and

open-tip resistances (18.8 ± 0.4 MΩ, n = 60, ranging from 12.5 to 24.4 MΩ) for

comparison of channel density between soma and dendrite (Kim et al., 2012).

Ensemble K$^+$ currents were evoked by a pulse protocol consisting of a 50–200

ms prepulse to –120 mV followed by a 200 ms test pulse to 70 mV. Na$^+$ currents

were generated by a pulse sequence comprised of a 100-ms prepulse to –120

mV and a 30-ms test pulse to 0 mV. In all cases, the holding potential was –90

mV before and after the pulse protocol. Voltage protocols were applied to outside-

out patches once every 3 and 5 s for Na$^+$ and K$^+$ current recordings, respectively.

Leak and capacitive currents were subtracted online using the pipette

capacitance compensation circuit of the amplifier and a 'P over –4' correction

procedure (Schmidt-Hieber & Bischofberger, 2010).

Subcellular dendritic patch-clamp recording

Dendritic recordings from GCs were obtained similarly as described previously

(Kim et al., 2012). First, Alexa Fluor 488 (50 µM, Invitrogen) diluted in an internal

solution was loaded into cells via a somatic recording pipette. Second, after ~10

min of loading, fluorescently labeled dendrites were traced from the soma into the

molecular layer using epifluorescence microscope. Finally, fluorescent and

infrared differential interference contrast (IR-DIC) images were compared, and

GC dendrites were patched under IR-DIC.

Stimulation protocols for the induction of long-term potentiation (LTP)

LTP was induced by either theta burst stimulation (TBS; Schmidt-Hieber et al.,

2004) or pairing protocols. TBS induction protocol consisted of burst of EPSPs

(10 stimuli at 100 Hz) repeated 10 times at 5 Hz. These episodes were repeated

four times every 10 seconds. The pairing protocol comprised 300 repetitions of a

presynaptic stimulation and one or two postsynaptic APs at the different time

intervals at 1 Hz. A postsynaptic AP was evoked by a brief current injection to the

soma (2 ms, 3 nA). For LTP experiments, baseline EPSPs evoked by stimulating

presynaptic axon fibers at 0.1 Hz were measured for ~10 min after whole-cell

recording. After LTP induction protocol, EPSPs were recorded for 30 min. LTP

magnitude was evaluated as the percentage of EPSP baseline (5 min) after 25

to 30 minutes after the induction protocol. In a subset of recordings, local

application of TTX (1 µM) to the perisomatic area was used to prevent axosomatic

AP initiation and backpropagation during TBS induction, without affecting evoked

EPSPs.

Solutions and chemicals

The extracellular solution for dissection and storage of brain slices was sucrose-

based solution (87 mM NaCl, 25 mM $NaHCO_3$, 2.5 mM KCl, 1.25 mM NaH_2PO_4,

7 mM $MgCl_2$, 0.5 mM $CaCl_2$, 10 or 25 mM glucose, and 75 sucrose). Physiological

saline for experiments was standard ACSF (125 mM NaCl, 25 mM $NaHCO_3$, 2.5

mM KCl, 1.25 mM NaH_2PO_4, 1 mM $MgCl_2$, 2 mM $CaCl_2$, and 25 mM glucose).

TTX and 4-AP were applied either via bath perfusion (0.5 µM and 10 mM,

respectively) or by local application (1 µM and 10 mM, respectively) with a

pressure application system (Picospritzer 3, General Valve). Pressure pulses had

durations of 0.2 s and amplitudes of ~10 psi. $CdCl_2$ and $NiCl_2$ were applied in the

bath at a concentration of 200 µM and 50 µM, respectively.

For whole-cell recording and K^+ current recording in outside out patches,

we used K^+ rich intracellular solution that contained 115 mM K-gluconate, 20 mM

KCl, 10 mM HEPES, 0.1 mM EGTA, 2 or 4 mM MgATP, 10 mM Na_2-

phosphocreatine, and 0.3 mM NaGTP, pH adjusted to 7.2–3 with KOH (~300

mOsm). In a subset of experiments, 50 µM Alexa 488 and 0.1–0.2 % biocytin

(wt/vol) were added to the internal solution for labeling during the experiment or

after fixation, respectively. All drugs were dissolved in physiological saline

immediately before the experiment and perfused on slices at a rate 4–5 ml min^{-1}.

These included: DL-AP5, PTX, CGP52432, TTX, 4-AP, and TEA. Intracellularly

delivered QX-314 (N-(2,6-dimethylphenylcarbamoylmethyl) triethylammonium

chloride) were directly added to the pipette solution before the experiment was

started. TTX and DL-AP5 (D,L-2-amino-5-phoshonovaleric acid; 50 µM) were

purchased from Tocris Bioscience; CGP52432 were from Abcam; All other drugs

were from Sigma-Aldrich.

Immunohistochemistry

Post hoc morphological analysis of GCs was performed as described previously

with slight modifications (Kim et al., 2012). Slices were fixed overnight at 4°C in

4% paraformaldehyde in 100 mM phosphate buffer solution (PBS), pH7.4. After

fixation, slices were rinsed several times with PBS and permeabilized with 0.3%

Triton X-100 in PBS. Subsequently, slices were treated with 0.3% Triton X-100

and 0.5% BSA in PBS to prevent any unwanted. Next, slices were treated with

0.3% Triton X-100 and 0.2% streptavidin-cy3 in PBS and were again incubated

overnight in 4°C. Finally, slices were mounted with DAKO S3023 medium.

Data analysis

Custom-made routines written in Igor Pro 6.3 (Wavemetrics), Stimfit 0.15

(Guzman et al., 2014) or Prism (Graphpad) were used for data analysis and

statistical testing. Spike threshold value was determined as the time point at

which the derivative of voltage exceeded 40 V/s at the soma or 20 V/s at the

dendrite. To determine the conduction velocity, latency–distance data were fit

with a linear regression, and the velocity was calculated from the linear regression

slope. Conductance density ($\overline{g}$) was calculated as $\overline{g} = I_{Peak} / ((V - V_{rev}) \cdot A)$, where

I_{Peak} is the corresponding peak current amplitude, V is the test pulse potential,

and A is the membrane surface area of outside-out patches. Membrane patch

area was estimated by capacitance measurement (**Figure 4–figure supplement**

1; Schmidt-Hieber & Bischofberger, 2010; Hu & Jonas, 2014). Reversal potentials

were assumed as +75 mV for Na^+ current and –95 mV for K^+ currents (Kim et al.,

2012).

All values indicate mean ± standard error of the mean (SEM), with "n"

denoting the number of experiments. To test statistical significance, a two-sided

nonparametric Wilcoxon signed-rank test or Wilcoxon rank sum test were used

(unless noted otherwise). For comparisons of more than two groups, a Kruskal-

Wallis test with Dunn's multiple comparisons for post-hoc testing was used.

Differences with P-value less than 0.05 were indicated with an asterisk and

considered significant. Membrane potential values were displayed without

correction for liquid junction potentials. Distances were measured from the soma

to the dendritic recording site along the dendritic trajectory (Kim, 2014).

References

1. Alme CB, Buzzetti RA, Marrone DF, Leutgeb JK, Chawla MK, Schaner MJ,
Bohanick JD, Khoboko T, Leutgeb S, Moser EI, Moser MB. McNaughton BL,
Barnes CA. 2010. Hippocampal granule cells opt for early retirement.
Hippocampus **20**:1109–1123. doi: 10.1002/hipo.20810.

2. Bliss TV, Lomo T. 1973. Long-lasting potentiation of synaptic transmission in
the dentate area of the anaesthetized rabbit following stimulation of the
perforant path. *The Journal of Physiology* **232**:331–356.

3. Birnbaum SG, Varga AW, Yuan LL, Anderson AE, Sweatt JD, Schrader LA.
2004. Structure and function of Kv4-family transient potassium channels.
Physiological Review **84**:803–833. doi: 10.1152/physrev.00039.2003.

4. Buzsaki G. 2002. Theta oscillations in the hippocampus. *Neuron* **33**:325–340.

5. Brunner J, Szabadics J. 2016. Analogue modulation of back-propagating
action potentials enables dendritic hybrid signalling. *Nature Communications*
7:13033. doi: 10.1038/ncomms13033.

6. Dan Y, Poo MM. 2006. Spike timing-dependent plasticity: from synapse to
perception. *Physiological Reviews* **86**:1033–1048.

7. Diamantaki M, Frey M, Berens P, Preston-Ferrer P, Burgalossi A. 2016.
Sparse activity of identified dentate granule cells during spatial exploration.
Elife **5**:e20252. doi: 10.7554/eLife.20252.

8. Dieni CV, Nietz AK, Panichi R, Wadiche JI, Overstreet-Wadiche L. 2013.
Distinct determinants of sparse activation during granule cell maturation. *The
Journal of Neuroscience* **33**:19131–19142. doi: 10.1523/JNEUROSCI.2289-
13.2013.

9. Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L.
2016. Low excitatory innervation balances high intrinsic excitability of
immature dentate neurons. *Nature Communications* **7**:11313. doi:
10.1038/ncomms11313.

10. Feldman DE. 2012. The spike-timing dependence of plasticity. *Neuron*
75:556–571. doi: 10.1016/j.neuron.2012.08.001.

11. Ge S, Yang CH, Hsu KS, Ming GL, Song H. 2007. A critical period for
enhanced synaptic plasticity in newly generated neurons of the adult brain.
Neuron **54**:559–566.

12. Golding NL, Spruston N. 1998. Dendritic sodium spikes are variable triggers
of axonal action potentials in hippocampal CA1 pyramidal neurons. *Neuron*
21:1189–1200.

13. Golding NL, Staff NP, Spruston N. 2002. Dendritic spikes as a mechanism for
cooperative long-term potentiation. *Nature* **418**:326–331.

14. Guzman SJ, Schlogl A, Schmidt-Hieber C. 2014. Stimfit: quantifying
electrophysiological data with Python. *Frontiers in Neuroinformatics* **8**:16. doi:
10.3389/fninf.2014.00016.

15. Hama K, Arii T, Kosaka T. 1989. Three-dimensional morphometrical study of
dendritic spines of the granule cell in the rat dentate gyrus with HVEM stereo
images. *Journal of Electron Microscopy Technique* **12**:80–87.

16. Hamilton TJ, Wheatley BM, Sinclair DB, Bachmann M, Larkum ME, Colmers
WF. 2010. Dopamine modulates synaptic plasticity in dendrites of rat and
human dentate granule cells. *Proceedings of the National Academy of
Sciences of the USA* **10**:18185–18190. doi: 10.1073/pnas.1011558107.

17. Hebb DO. 1949. The Organization of Behavior (John Wiley & Sons, 1949).

18. Hoffman DA, Magee JC, Colbert CM, Johnston D. 1997. K^+ channel regulation
of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature*
387:869–875.

19. Holmes WR. 1989. The role of dendritic diameters in maximizing the
effectiveness of synaptic inputs. *Brain research* **478**:127–137.

20. Hu H, Jonas P. 2014. A supercritical density of Na^+ channels ensures fast
signaling in GABAergic interneuron axons. *Nature Neuroscience* **17**:686–693.
doi: 10.1038/nn.3678.

21. Jarsky T, Roxin A, Kath WL, Spruston N. 2005. Conditional dendritic spike
propagation following distal synaptic activation of hippocampal CA1 pyramidal
neurons. *Nature Neuroscience* **8**:1667–1676.

22. Kim S, Guzman SJ, Hu H, Jonas P. 2012. Active dendrites support efficient
initiation of dendritic spikes in hippocampal CA3 pyramidal neurons. *Nature
Neuroscience* **15**:600–606. doi: 10.1038/nn.3060.

23. Kim S. 2014. Action potential modulation in CA1 pyramidal neuron axons
facilitates OLM interneuron activation in recurrent inhibitory microcircuits of
rat hippocampus. *PLoS One* **9**:e113124. doi: 10.1371/journal.pone.0113124.

24. Kim Y, Hsu CL, Cembrowski MS, Mensh BD, Spruston N. 2015. Dendritic
sodium spikes are required for long-term potentiation at distal synapses on
hippocampal pyramidal neurons. *Elife* **4**:e06414. doi: 10.7554/eLife.06414.

25. Krueppel R, Remy S, Beck H. 2011. Dendritic integration in hippocampal
dentate granule cells. *Neuron* **71**:512–528. doi:
10.1016/j.neuron.2011.05.043.

26. Levy WB, Steward O. 1983. Temporal contiguity requirements for long-term
associative potentiation ∕ depression in the hippocampus. *Neuroscience*
8:791–797.
27. Letzkus JJ, Kampa BM, Stuart GJ. 2006. Learning rules for spike timing-
dependent plasticity depend on dendritic synapse location. *The Journal of*
Neuroscience **26**:10420–10429. DOI: 10.1523/JNEUROSCI.2650-06.2006.
28. Lin YW, Yang HW, Wang HJ, Gong CL, Chiu TH, Min MY. 2006. Spike-timing-
dependent plasticity at resting and conditioned lateral perforant path
synapses on granule cells in the dentate gyrus: different roles of N-methyl-D-
aspartate and group I metabotropic glutamate receptors. *The European*
Journal of Neuroscience **23**:2362–2374.
29. Lisman JE. 1999. Relating hippocampal circuitry to function: recall of memory
sequences by reciprocal dentate–CA3 interactions. *Neuron* **22**:233–242.
30. Lopez-Rojas J, Heine M, Kreutz MR. 2016. Plasticity of intrinsic excitability in
mature granule cells of the dentate gyrus. *Scientific Reports* **6**:21615 doi:
10.1038/srep21615.
31. Losonczy A, Magee JC. 2006. Integrative properties of radial oblique
dendrites in hippocampal CA1 pyramidal neurons. *Neuron* **50**:291–307.
32. Magee JC, Johnston D. 1997. A synaptically controlled, associative signal for
Hebbian plasticity in hippocampal neurons. *Science* **275**:209–213.
33. Marr D. 1971. Simple memory: a theory for archicortex. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **262**:23–81.
34. McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK,
Lowell BB, Fanselow MS, Wilson MA, Tonegawa S. 2007. Dentate gyrus
NMDA receptors mediate rapid pattern separation in the hippocampal network.
Science **317**:94–99.
35. McNaughton BL, Douglas RM, Goddard GV. 1978. Synaptic enhancement in
fascia dentata: cooperativity among coactive afferents. *Brain Research*
157:277–293.
36. McNaughton BL, Morris RGM. 1987. Hippocampal synaptic enhancement
and information storage within a distributed memory system. *Trends in*
Neuroscience **10**:408–415.
37. Menegola M, Misonou H, Vacher H, Trimmer JS. 2008. Dendritic A-type
potassium channel subunit expression in CA1 hippocampal interneurons.
Neuroscience. **154**:953–964. doi: 10.1016/j.neuroscience.2008.04.022

38. Mishra RK, Kim S, Guzman SJ, Jonas P. 2016. Symmetric spike timing-
dependent plasticity at CA3-CA3 synapses optimizes storage and recall in
autoassociative networks. *Nature Communications* **7**:11552. doi:
10.1038/ncomms11552.

39. Monaghan MM, Menegola M, Vacher H, Rhodes KJ, Trimmer JS. 2008.
Altered expression and localization of hippocampal A-type potassium channel
subunits in the pilocarpine-induced model of temporal lobe epilepsy.
Neuroscience. **156**:550–562. doi: 10.1016/j.neuroscience.2008.07.057

40. Mongiat LA, Espósito MS, Lombardi G, Schinder AF. 2009. Reliable activation
of immature neurons in the adult hippocampus. *PLoS One* **4**:e5320. doi:
10.1371/journal.pone.0005320.

41. Nevian T, Larkum ME, Polsky A, Schiller J. 2007. Properties of basal
dendrites of layer 5 pyramidal neurons: a direct patch-clamp recording study.
Nature Neuroscience **10**:206–214. doi: 10.1038/nn1826

42. O'Neill J, Senior TJ, Allen K, Huxter JR, Csicsvari J. 2008. Reactivation of
experience-dependent cell assembly patterns in the hippocampus. *Nature*
Neuroscience **11**:209–215.

43. Pernía-Andrade AJ, Jonas P. 2014. Theta-gamma-modulated synaptic
currents in hippocampal granule cells *in vivo* define a mechanism for network
oscillations. *Neuron* **81**:140–152. doi: 10.1016/j.neuron.2013.09.046.

44. Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. 2015. Engram cells
retain memory under retrograde amnesia. *Science* **348**:1007–1013. doi:
10.1126/science.aaa5542.

45. Schiller J, Major G, Koester HJ, Schiller Y. 2000. NMDA spikes in the basal
dendrites of cortical pyramidal neurons. *Nature* **404**:285–289. doi:
10.1038/35005094.

46. Schmidt-Hieber C, Bischofberger J. 2010. Fast sodium channel gating
supports localized and efficient axonal action potential initiation. *The Journal*
of Neuroscience **30**:10233–10242. doi: 10.1523/JNEUROSCI.6335-09.2010.

47. Schmidt-Hieber C, Jonas P, Bischofberger J. 2004. Enhanced synaptic
plasticity in newly generated granule cells of the adult hippocampus. *Nature*
429:184–187.

48. Schmidt-Hieber C, Jonas P, Bischofberger J. 2007. Subthreshold dendritic
signal processing and coincidence detection in dentate gyrus granule cells.
The Journal of Neuroscience **27**:8430–8441.

49. Schmidt-Hieber C, Wei H, Häusser M. 2014. Synaptic mechanisms of sparse
activity in hippocampal granule cells during mouse navigation. *Society for*
Neuroscience Abstract.
50. Scharfman HE, Schwartzkroin PA. 1990. Responses of cells of the rat fascia
dentate to prolonged stimulation of the perforant path: sensitivity of hilar cells
and changes in granule cell excitability. *Neuroscience* **35**:491–504.
51. Senzai Y, Buzsáki G. 2017. Physiological Properties and Behavioral
Correlates of Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**:691–
704. doi: 10.1016/j.neuron.2016.12.011.
52. Sjöström PJ, Turrigiano GG, Nelson SB. 2001. Rate, timing, and cooperativity
jointly determine cortical synaptic plasticity. *Neuron* **32**:1149–1164.
53. Sjöström PJ, Rancz EA, Roth A, Häusser M. 2008. Dendritic excitability and
synaptic   plasticity.   *Physiological   Reviews*   **88**:769–840.   doi:
10.1152/physrev.00016.2007.
54. Spruston N, Schiller Y, Stuart G, Sakmann B. 1995. Activity-dependent action
potential invasion and calcium influx into hippocampal CA1 dendrites. *Science*
268:297–300.
55. Stuart GJ, Häusser M. 2001. Dendritic coincidence detection of EPSPs and
action potentials. *Nature Neuroscience* **4**:63–71.
56. Stuart GJ, Schiller J, Sakmann B. 1997. Action potential initiation and
propagation in rat neocortical pyramidal neurons. *The Journal of Physiology*
505: 617–632.
57. Stuart GJ, Spruston N. 2015. Dendritic integration: 60 years of progress.
Nature Neuroscience **18**:1713–1721. doi: 10.1038/nn.4157.
58. Temprana SG, Mongiat LA, Yang SM, Trinchero MF, Alvarez DD, Kropff E,
Giacomini D, Beltramone N, Lanuza GM, Schinder AF. 2015. Delayed
coupling to feedback inhibition during a critical period for the integration of
adult-born   granule   cells.   *Neuron*   **85**:116–130.   doi:
10.1016/j.neuron.2014.11.023.
59. Treves A, Rolls ET. 1994. Computational analysis of the role of the
hippocampus in memory. *Hippocampus.* **4**:374–391.
60. Vyleta NP, Borges-Merjane C, Jonas P. 2016. Plasticity-dependent, full
detonation at hippocampal mossy fiber-CA3 pyramidal neuron synapses. *Elife*
5:e17977. doi: 10.7554/eLife.17977.
61. Yang K, Dani, JA. 2014. Dopamine d1 and d5 receptors modulate spike
timing-dependent plasticity at medial perforant path to dentate granule cell

synapses. *The Journal of Neuroscience* **34**:15888–15897. doi:
10.1523/jneurosci.2400-14.2014.

Acknowledgments

We thank Jose Guzman and Hua Hu for critically reading the manuscript.

Figure 1. Putative dendritic spikes during theta-burst stimulation (TBS) induction

are required for long-term potentiation (LTP) at the perforant-path (PP) to granule

cell (GC) synapse.

(**A**) Maximum intensity projection of confocal stack fluorescence images of a GC

indicating the medial molecular layer (MML) and the outer molecular layer (OML).

Synaptic responses of the PP were evoked by electrical stimulation in the OML.

Scale bar is 50 µm.

(**B**) Time course of excitatory postsynaptic potentials (EPSPs) before (average of

30 EPSP traces, gray) and after (average of 30 EPSPs, black) TBS of the PP

synapses. Red line denotes average EPSP baseline value. Increment in EPSP

amplitude denotes LTP.

(**C**) Bar plot of average EPSP amplitude before and after TBS indicating a
significant increment in synaptic responses 25–30 minutes after TBS stimulation
(*n* = 13, **P* < 0.01). Data points from the same experiment are connected by
lines.

(**D**) (Left) Representative somatic voltage traces evoked by high-frequency burst
stimulation to the PP synapses. The arrows indicate weak (orange) and strong
(red) putative dendritic spikes and accompanying plateau potentials (blue arrow).
(Middle) Somatic voltages (top row) and corresponding dV/dt (bottom row) of
weak (orange) and strong (red) putative dendritic spikes in the burst responses
(left) on an expaned time scale. Note that dV/dt values of the putative dendritic
spikes increase sharply in a non-linear manner. (Right) Summary bar graphs of
dV/dt peak values of somatically recorded EPSPs with or without dendritic spikes.

(**E**) Average time course of EPSPs when TBS stimulation evokes putative
dendritic spikes and in the absence of putative dendritic spikes during TBS (black).
LTP is induced only if putative dendritic spikes are present during TBS stimulation.

(**F**) The number of putative dendritic spikes are significantly correlated with the
magnitude of LTP. Black lines represent linear regressions (n = 13).

(**G**) Representative time courses of EPSP amplitudes before and after pairing
presynaptic stimulation of the PP synapses and postsynaptic action potentials
with short time intervals (+10ms, Pre–post sequence, black, top left inset; –10 ms,
Post–pre sequence, green, top right inset). Both pairing protocols induce no
significant changes in EPSP, suggesting that action potential (AP)
backpropagation is not necessary for LTP induction. Right inset shows that LTP

was not induced after pairing of EPSPs and APs in both pre-post (top) and post-

pre sequences (bottom).

(**H**) The average EPSP time courses of pre-post (black) and post-pre (green)

induction protocols of pairings between synaptic responses and postsynaptic APs,

showing that low frequency pairing protocols failed to induce LTP.

(**I**) Bar summary graph and individual experiments (circles) indicating that

experiments showing the occurrence of putative dendritic spikes during TBS

induction induced a significant increment of LTP (i.e., percentage of EPSP

baseline), whereas pairings of synaptic stimulation with postsynaptic APs did not

show a statistically significant LTP induction, independent of temporal order.

Bars indicate mean ± SEM; circles represent data from individual cells. Lines

connect data points from the same experiment. ***$P < 0.005$. ****$P < 0.0001$.

Single-cell data (**B,E**) and mean data (**D,F**; mean ± SEM). Vertical gray bars in **B,**

D, E, and **F** indicate the time point of the induction protocol.

The following figure supplements are available for figure 1:

Figure supplement 1. TBS-induced LTP at PP-GC synapses is independent of

axosomatic APs.

Figure supplement 2. Pairing protocols did not induce LTP at the medial PP-GC

synapses.

Figure 2. Induction of LTP at PP–GC synapses requires activation of NMDARs

and the involvement of Na$^+$ channels.

(**A**) An example of the time course of EPSP amplitude when TBS protocols was

applied in the presence of the NMDAR antagonist, DL-AP5 (50 µM). Red line

indicates the average EPSP amplitude. Insets show that EPSP did not increase

in the presence of DL-AP5 (orange) after TBS, but a robust LTP was induced

when TBS was applied in standard saline (ACSF, black).

(**B**) (Left) Long-duration plateau potentials are mediated by NMDA receptor

channels. Somatically recorded voltages in response to high-frequency burst

sitmulation (green, 10 shocks, 100 Hz) under control (black) and in the presence

of DL-AP5 (50 µM; orange). The inset shows putative dendritic spikes (indicated

by the red arrow) before and after the addition of DL-AP5; Note that putative

dendritic spikes are resistant to the NMDAR blockers. (Right) Summary of the

effects of DL-AP5 on plateau potentials. Peak amplitdue of plateau potentials

were measured after the stimulus (indicated by dashed lines; Control: 36.6 ± 2.2

mV; DL-AP5: 25.6 ± 3.2 mV; n = 6, *$P < 0.05$).

(**C**) A representative time course of EPSP amplitude before and after TBS when

the cells were dialyzed with a sodium-channel blocker, QX-314 (5 mM). Inset

shows that the averaged EPSP amplitude did not change when blocking sodium

channels with QX-314 (blue) despite TBS induction. In contrast, when cells were

dialyzed with the standard intracellular solution, the amplitude of EPSP increased

after TBS (i.e., LTP).

(**D**) Summary plot of TBS-induced LTP experiments in the presence of DL-AP5

(orange) and with dialysis of QX-314 in the recording pipette (blue). Both

treatments prevented the induction of LTP at the PP to GC synapse.

(**E**) Summary bar graph and individual average EPSP amplitudes after TBS in

control (standard saline, black), bath application of DL-AP5 (orange) and dialysis

of QX-314 (blue). Treatments with DL-AP5 and QX-314 showed a significant

difference compared to the standard LTP induction (DL-AP5, **$P < 0.01$; QX-314,

***$P < 0.005$; compared to control in **Figure 1I**).

Representative traces in **A** and **C** correspond to the numbers (1 & 2) n the time-

course plot. Bars indicate mean ± SEM; circles represent data from individual

cells. Lines connect data points from the same experiment. Single-cell data (**A,C**)

and mean data (**D**; mean ± SEM). Vertical gray bars in **A, C,** and **D** indicate the

time point of the induction protocol.

Figure 3. Properties of backpropagating APs in the dendrites of GCs.

(**A**) Morphological reconstruction of a GC with representative double somatic and

dendritic whole-cell recording configuration used to analyze the AP

backpropagation. Scale bar is 50 μm.

(**B**) A train of APs elicited by a 1-s current pulse applied at the soma. Black traces

indicate somatic voltage and corresponding current; red traces indicate dendritic

voltage and corresponding current.

(**C**) First AP in the train displayed at expanded time scale. Voltage traces (soma

in black, dendrite in red) indicate that the AP is initiated first near the soma and

propagated back into the dendrites with a lower amplitude.

(**D**) Summary graph to compare somatic (black) and dendritic (red) AP peak

amplitude. Bars indicate mean ± SEM; circles represent data from individual cells.

Lines connect data points from the same experiment. ***$P < 0.005$

(**E**) Scatter plot of peak amplitude of the backpropagating AP against the absolute

physical distance of the recording site from the soma (56 somatodendritic

recordings). The blue curve represents a mono-exponential fit to the data points

between 0 and 212 µm.

(**F**) Scatter plot of the bAP amplitude normalized to the corresponding axosomatic

AP amplitude plotted against the distance of the recording site scaled to the total

dendritic length (278 ± 7.4 µm; n = 11). The blue curve is a mono-exponential fit

to the data.

(**G**) Scatter plot of AP latency as a function of the distance from the soma (56

somatodendritic recordings) together with a linear regression (blue line) to

compute the average conduction velocity of the AP into the dendrites; dendritic

AP propagation velocity was 226 µm/ms.

Single-cell data (**E–G**, red) and mean data (**E**, black; mean ± SEM).

Figure 4. Differential Na$^+$ and K$^+$ conductance gradients in the dendrites of GCs.

(**A**) Averages of Na$^+$ current recorded from outside-out patches from soma (black, averages of 25–27 sweeps) and dendrite (red, 110 µm, averages of 20 sweeps) in response to a test pulse potential to 0 mV (bottom). Na$^+$ currents were recorded in the presence of 4-AP (5 mM) and TEA (20 mM). Left, soma; right, dendrite; Top, control; bottom, currents in the presence of 0.5 µM TTX in the bath. Leak and capacitive currents were subtracted by a 'P over –4' correction procedure. Note that the remaining outward current is the resistant K$^+$ current component to 5 mM 4-AP (**Figure 4–figure supplement 2,3**; Hoffman et al., 1997). Blockade of outward K$^+$ currents by extracellular 4-AP (5 mM) only had a negligible effects on the peak amplitude of Na$^+$ currents (**Figure 4–figure supplement 3**).

(**B**) Averages of A-type K$^+$ current evoked in outside-out patches excised from

soma (black, averages of 6–8 sweeps) and dendrite (red, 110 μm averages of

15–19 sweeps) in response to a test pulse potential to +70 mV (top). Transient

A-current was measured by subtraction of traces with a −40 mV prepulse from

those with a −120 mV prepulse. Left column, soma; Right column, dendrite; Top

row, control; Bottom row, currents in the presence of 5 mM 4-AP in the bath.

(**C**) Averages of delayed rectifier K$^+$ current evoked in outside-out patches

excised from soma (black, averages of 6–8 sweeps) and dendrite (red, 124 μm,

averages of 10–18 sweeps) in response to a test pulse potential to +70 mV (top).

Delayed rectifier K$^+$ current was measured by a −40 mV prepulse. Left column,

soma; Right column, dendrite; Top row, control; Bottom row, currents in the

presence of 20 mM TEA in the bath. See also **Figure 4–figure supplement 2**.

(**D, E, F**) (Top) Plot of Na$^+$ channel (**D**), A-type K$^+$ channel (**E**), and delayed

rectifier K$^+$ conductance density (**F**) as a function of distance from the soma,

demonstrating that density of various conductances is differentially expressed

across the length of GC dendrites. Data from 19, 21, and 21 somatic (black circles)

and 29, 28, and 29 dendritic patches (red circles). Blue circles represent the

average of somatic recordings. (Bottom) Summary bar graph showing Na$^+$ (**D**),

A-type K$^+$ (**E**) and delayed rectifier K$^+$ (**F**) conductance density in the soma,

proximal dendrite (< 100 μm) and distal dendrite (≥ 100 μm). Bars indicate

mean ± SEM; circles represent data from individual experiments. n.s., not

significant; ****P < 0.0001 by Kruskal Wallis test with *post hoc* multiple

comparison using Dunn's test.

The following figure supplements are available for figure 4:

Figure supplement 1. Estimation of the surface area of outside-out patches by

capacitance measurements.

Figure supplement 2. Pharmacological analysis of voltage-dependent Na^+ and

K^+ currents.

Figure supplement 3. The dose-dependent effect of 4-AP on transient outward

currents and Na^+ currents.

Figure supplement 4. Effect of A-type K^+ channel blockade on AP

backpropagation.

Figure 5. Blockade of Na$^+$ channels prevent dendritic spikes and LTP induction

by TBS.

(**A**) (Left) Schematic recording configuration of a simultaneous somatic (black)

and dendritic (red) patch-clamp recording on a GC. (Right) Local dendritic spikes

in GCs evoked by dendritic current injection pulses of increasing amplitude (black,

voltage in the soma; red, voltage in the dendrite).

(**B**) The relationship between voltage amplitude (black, soma; red, dendrite) and

dendritic injection resembles a step-function, suggesting the all-or-none nature of

the dendritic spike. Peak values of dendritic spikes were measured after

subtraction of scaled subthreshold dendritic responses.

(**C**) A summary plot showing whether a spike was evoked first in the dendrite

(right, red) or in the soma (left, black) for an increasing current pulse injection at

the dendrite. Red squares indicate the cells showing a dendritic spike followed

by an axosomatic spike. Red triangles show the cells showing isolated dendritic

spike. Black circles indicate the cells showing only an axosomatic spike. The

green shaded area approximately corresponds to the outer molecular layer. The

black circle and red lines in the plot indicate the soma and the dendrite,

respectively.

(**D**) (Left) Schematic recording configuration of a triple pipette consisting of

electrical stimulation of the PP synapses (10 shocks, 100 Hz, green), dendritic

patch-clamp recording (120 μm from the soma, red) and somatic whole-cell

recording (black). (Right) Dendritic spikes can be identified as larger spikes in the

dendrite (red) with the corresponding small spike at the soma (black).

(**E**) Bath application of TTX (10 nM) prevents the generation of dendritic spikes

evoked by synaptic stimulation (black), as seen on the somatic whole-recording

trace (blue). Note that decreasing availability of Na^+ channels by 10 nM TTX had

a negligible effect on the $EPSP_1$ (see also **Figure 5–figure supplement 4**).

(**F**) Time course of EPSP amplitudes in the presence of TTX (10 nM). The

induction of LTP by TBS can be prevented by bath application of TTX, since the

average EPSP amplitude does not change.

(**G**) Average EPSP before and after TBS shows no changes in amplitude in the

presence of TTX.

(**H**) Summary bar graph and single experiments (circles) indicating that the TTX

application does not produce an increment in the EPSP amplitude and therefore

no statistically significant LTP induction. Bars indicate mean ± SEM; circles

represent data from individual cells. Data points from the same experiment are

connected by lines.

Single-cell data (**F**) and mean data (**G**; mean ± SEM). Scale bar in **A** and **D** is 50

µm. Vertical gray bars in **F** and **G** indicate the time point of the induction protocol.

The following figure supplements are available for figure 5:

Figure supplement 1. Dendritic spikes are mediated by voltage-gated Na^+

channels.

Figure supplement 2. Dendritic A-type K^+ channels control dendritic spike

initiation.

Figure supplement 3. Relationship between the dV/dt of somatically and

dendritically recorded voltages during dendritic spikes generation.

Figure supplement 4. Low concentration of TTX does not affect glutamate

release from presynaptic nerve terminals.

Figure supplement

Figure 1–figure supplement 1. TBS-induced LTP at PP-GC synapses is

independent of axosomatic action potentials (APs).

(**A**) Time course of excitatory postsynaptic potentials (EPSPs) before (average of

30 EPSP traces, gray) and after (average of 30 EPSPs, black) theta-burst

stimulation (TBS) of the PP synapses. Red line denotes average EPSP baseline

value. Inset: example of the first burst of TBS responses (red) showing initiation

of multiple axosomatic APs during TBS induction upon increasing the stimulus

intensity.

(**B**) Average time course of EPSPs when TBS stimulation evokes axosomatic

APs (red) and in the absence of axosomatic APs during TBS (black). Note that a

similar degree of potentiation in both cases is shown.

(**C**) Summary bar plot of average EPSP amplitude before and after TBS indicating

that there are no significant differences in the magnitude of LTP between two

groups in **B**. Bars indicate mean ± SEM; circles represent data from individual

cells.

Single-cell data (**A**) and mean data (**B**; mean ± SEM). Vertical bars (gray) in **A**

and **B** indicate the time point of the pairing protocol.

Figure 1–figure supplement 2

Figure 1–figure supplement 2. Pairing protocols did not induce LTP at the

medial perforant path (MPP)-GC synapses.

(**A**) An example of reconstructed granule cells and diagram illustrating the

experimental configuration. Note that a stimulating electrode was placed in the

middle third of the molecular layer. Scale bar is 50 μm.

(**B**) Representative time course of EPSP amplitude, showing that repeated

pairing of EPSPs and following postsynaptic APs (pre-post sequence; Δt = +10

ms, inset) induced no significant LTP at MPP-GC synapses. (Right) EPSPs

before (gray) and 25–30 min after induction (black).

(**C**) Representative time course of EPSP amplitude, showing that post-

presynaptic pairing (Δt = −10 ms, inset) induces no LTP at MPP-GC synapses.

(Right) EPSPs before (gray) and 25–30 min after induction (green).

(**D**) Summary graph of pairing-induced LTP experiments at the MPP-GC

synapses demonstrating the absence of LTP.

(**E**) Summary graph and scatter (individual cells) plot of the change in EPSP

amplitude before and after the pairing protocols. Bars indicate mean ± SEM;

circles represent data from individual cells.

Single-cell data (**B,C**) and mean data (**D**; mean ± SEM). Vertical gray bars in **B**,

C, and **D** indicate the time point of the induction protocol.

1000

1001

Figure 4–figure supplement 1. Estimation of the surface area of outside-out

patches by capacitance measurements.

(**A**) Experimental configuration showing the patch pipette near the silicone

elastomer ball (Sylgard 184, Dow Corning). After obtaining excised outside-out

patches, the pipette tip was gently pressed against the silicone ball (< 5 µm),

resulting in complete sealing of the tip. All the pipettes were coated with dental

wax for this set of experiments (Kimsdent).

(**B**) The traces show the difference current (evoked by a test pulse from –60 to –

110 mV, average of 100 individual capacitive current traces) before and after

pressing the pipette tip into the silicone ball. The integral under the curve (grey

area) represents the capacitive charge used for membrane surface (Schmidt-

Hieber & Bischofberger, 2010). Membrane surface area was determined from the

capacitance assuming a specific membrane capacitance of 1.0 µF·cm^{-2}. Dashed

lines indicate leakage current.

(**C**) Plot of membrane surface area of outside-out patches against pipette open-

tip conductance. Data (n = 10) were fit by linear regression according to $A(g_P) =$

$4.7124 \times g_P + 0.21421$, where A is membrane patch area (µm^2) and g_P is pipette

conductance (µS).

Figure 4–figure supplement 2

1020

Figure 4–figure supplement 2. Pharmacological analysis of voltage-dependent

Na$^+$ and K$^+$ currents.

(**A**) Bar graph summarizing the effects of the I_{Na} channel blocker TTX (0.5 µM)

on the transient Na$^+$ currents excised from the soma (left, black, n = 6) and the

dendrite (right, red, n = 6).

(**B**) Bar graph summarizing the effects of the I_A channel blocker 4-AP (5 mM) on

the transient outward K$^+$ currents excised from the soma (left, black, n = 10) and

the dendrite (right, red, n = 8).

(**C**) Bar graph summarizing the effects of the I_K channel blocker TEA (20 mM) on

the steady-state outward K$^+$ currents excised from the soma (left, black, n = 10)

and the dendrite (right, red, n = 6).

*$0.01 \leq P < 0.05$; **$p < 0.01$; ***$P < 0.005$; ****$P < 0.0001$ by paired t-test. Bars

indicate mean ± SEM; circles represent data from individual cells. Lines connect

data points from the same experiment.

1035

Figure 4–figure supplement 3. The dose-dependent effect of 4-AP on transient

outward currents and Na$^+$ currents.

(**A**) Representative traces showing dendritic A-type K$^+$ currents (113 μm from the

soma, averages of 10 sweeps) evoked as in **Figure 4B** in control (black) and

subsequently in 5 mM 4-AP (red), followed by 10 mM 4-AP (purple) in the bath.

Inset shows an expanded traces corresponding to the dashed box in the left panel.

(**B**) Summary plot indicating that 4-AP reduced the transient K$^+$ current in a dose-

dependent manner. Colored circles represent data from individual dendritic

recordings at different distances (right). Lines connect data points from the same

experiment. Filled circles (blue) represent average data of the normalized peak

currents, showing the percentage block of transient outward K$^+$ currents in the

presence of 5 and 10 mM 4-AP (right axis; 5 mM 4-AP, n = 8, ***P < 0.005 by

paired t-test).

(**C**) Representative traces showing averages of Na$^+$ currents (top, 110 µm from

the soma, averages of 10 sweeps) evoked as in **Figure 4A** in control (black) and

subsequently in 5 mM 4-AP (red), followed by 0.5 µM TTX and 5 mM 4-AP (blue)

in the bath. Inset shows an expanded traces corresponding to the dashed box on

the left.

(**D**) Summary graph showing the effects of 5 mM 4-AP and 0.5 µM TTX on Na$^+$

currents. Colors as in **B**. Filled blue circles show the average of the normalized

peak currents demonstrating that 5 mM 4-AP did not significantly affect the peak

of transient inward currents (n = 6, P = 0.69; n.s., not significant, by paired t-test).

These inward currents were blocked by 0.5 µM extracellular TTX (n = 5, ***P <

0.005 by paired t-test).

Open circles represent data from individual cells and filled circles represent mean

data (mean ± SEM).

1062

Figure 4–figure supplement 4

Figure 4–figure supplement 4. Effect of 4-AP on AP backpropagation.

(**A**) Diagram of the experimental setup displaying simultaneous recordings from the soma (black) and the dendrite (red) of a GC in the presence of 5 mM 4-AP. Scale bar is 50 μm.

(**B**) Sample traces of somatic (black) and dendritic (red) APs evoked by somatic current injection in the absence (top; control) or the presence of 4-AP (5 mM; bottom) in the bath.

(**C, D**) First and second APs (**C**) in the train displayed at expanded time scale and overlay of dendritic voltages (**D**) under control and in the presence of 4-AP (dashed line). The blue line illustrates an exponential fit to the repolarizing phase of the 1st dendritic AP to calculate the duration at half-maximal amplitude.

(**E**) Summary graph of the effects of 5 mM 4-AP on AP peak amplitude (Left; Soma: 102.6 ± 0.8%; Dendrite: 142.8 ± 2.6 %; n = 6, ***P < 0.005) and duration at half-maximal amplitude (right; soma: 144.4 ± 7.1%; dendrite: 377.6 ± 59.0%; n

1078   = 6, ***P < 0.005) in 6 somatodendritic recordings. Bars indicate mean ± SEM;

1079   circles represent data from individual cells.

1080

1081

Figure 5–figure supplement 1. Dendritic spikes are mediated by voltage-gated

Na$^+$ channels.

(**A**) Schematic diagram illustrating the recording configuration of a simultaneous

somatic (black amplifier) and dendritic (red amplifier) patch-clamp recording on a

GC combined with bath application of blockers of voltage-gated Na$^+$ and Ca^{2+}

channel. Scale bar is 50 μm.

(**B**) Blockade of voltage-gated Na$^+$ channels by a bath application of TTX (0.5 μM)

eliminated dendritic spikes elicited by a 5-ms current pulse injected into the

dendritic electrode.

(**C**) Application of 200 μM CdCl$_2$ did not abolish dendritic spikes. Bath application

of CdCl$_2$ did not affect the late depolarizing phase of the dendritic spikes (S.K.,

unpublished observations).

(**D**) Application of 50 μM NiCl$_2$ did not abolish dendritic spikes.

(**E**) Summary of the effects of the indicated pharmacological agents on the peak

of dendritic spikes.

The dendritic recording sites are 165 μm (**B**), 137 μm (**C**), and 100 μm (**C**) from

the soma, respectively. Bars indicate mean ± SEM; ***P < 0.005 by Student's t-

test; Black traces: somatic voltage; red traces: dendritic voltage.

1100

Figure 5–figure supplement 2. Dendritic A-type K$^+$ channels control dendritic

spike initiation.

(**A**) Diagram of the experimental setup illustrating the recording configuration of

a simultaneous somatic (black amplifier) and dendritic (red amplifier) patch-clamp

recording on a GC combined with focal application of 10 mM 4-AP directly to the

dendritic patch. Scale bar is 50 μm.

(**B**) Somatic and dendritic voltage responses to dendritic current injection pulses

with increasing amplitude (left: 400 pA; right; 500 pA). Local application of 4-AP

to the dendrite near the recording pipette tip decreased the threshold for initiating

dendritic spikes (arrows). Black traces represent somatic signal and red traces

represent dendritic signal. Dashed line indicates the voltage response after puff

application of 4-AP. The dendritic recording site is 102 μm from the soma.

(**C**) Summary of the effects of local application of 4-AP on the threshold required

to initiate dendritic spikes (Control, 0.84 ± 0.15 nA; 4-AP, 0.76 ± 0.14 nA; n = 8,

***$P < 0.005$, Paired t-test). Bars indicate mean ± SEM. Lines connect data points

from the same experiment.

1117

The table within the figure:

	①	②	③	④	⑤
dV/dt (mV/ms)	8.9	17.2	22.4	30.2	24.1
dV/dt (mV/ms)	2.8	4.3	5.5	16.5	551.1

Figure 5–figure supplement 3

1118

Figure 5–figure supplement 3. Relationship between the dV/dt of somatically

and dendritically recorded voltages during dendritic spikes generation.

(**A**) (Left) Diagram illustrating the experimental configuration of a triple pipette

consisting of electrical stimulation of the PP synapses in the OML (green),

dendritic patch-clamp recording (147 µm from the soma, red) and somatic whole-

cell recording (black). Scale bar is 50 µm. (Right) Somatic and dendritic voltages

in response to a high-frequency PP stimulation with a constant stimulus intensity

are shown superimposed for comparison.

(**B**) Traces of somatic (black) and dendritic (red) voltage responses (top row) in

A (red box) and corresponding dV/dt (Bottom). dV/dt peak amplitudes of each

traces were summarized in the table (bottom). The red arrows indicate the

dendritic spikes. When dendritic spikes were present, the corresponding somatic

voltage changes were used for analysis of dV/dt peaks. For comparison, somatic

and dendritic membrane voltages and corresponding dV/dt traces during

axosomatic AP generation are also shown on the right (dashed box). Encircled

numbers indicate correspondence between traces in **B** and data points in **C**.

(**C**) (Left) Summary plot of peaks in somatic dV/dt against corresponding peaks

in dendritic dV/dt for 4 simultaneous somatodendritic recordings in response to

high-frequency burst stimulation of the PP synapses (Dendritic recording sites

are from 136 μm to 175 μm from the soma). (Right) An enlarged view of the box

(blue) in the left panel. Black circles, in the absence of dendritic spikes; red circles,

in the presence of the dendritic spikes; gray circles, data points from the first

EPSPs (EPSP$_1$). Note that somatic dV/dt peak amplitudes of subthreshold

EPSP1s were comparable to those of the somatic traces when dendritic spikes

were present.

(**D**) Summary bar graphs of dV/dt peak amplitudes of somatically recorded

dendritic spikes in **C** (red) and in **Figure 1D**. Bars indicate mean ± SEM; circles

represent data from individual cells.

1147

Figure 5–figure supplement 4

1148

Figure 5–figure supplement 4. Low concentration of TTX does not affect

glutamate release from presynaptic nerve terminals.

(**A**) Plot of EPSP peak amplitude against time during perfusion of 10 nM TTX

(blue horizontal bar).

(**B**) Summary graph of the effect of 10 nM TTX on EPSP peak amplitude (control:

5.21 ± 0.78 mV, TTX: 5.06 ± 0.68, n = 11; $P = 0.83$). Data shown in **A** and example

traces in **Figure 5E** were taken from the same cell. Filled circles represent data

from individual experiments. Bars indicate mean $\pm$ SEM. Lines connect data

points from the same experiment.

1158