Beyond synthetic lethality: multiple gene interaction types play a key functional role in

cancer

Assaf Magen, Avinash Das, Joo Sang Lee, Mahfuza Sharmin, Alexander Lugo, Silvio Gutkind,

Eytan Ruppin*, Sridhar Hannenhalli*

Center for Bioinformatics and Computational Biology

University of Maryland, College Park, Maryland, USA

*Corresponding authors

Sridhar Hannenhalli & Eytan Ruppin

3104G Biomolecular Sciences Building

University of Maryland, College Park, MD

sridhar@umiacs.umd.edu

The phenotypic effect of perturbing a gene's activity depends on the activity state of other

genes, reflecting the fundamental notion that genotype to phenotype linkage is mediated by a

network of functionally interacting genes. The vast majority of contemporary investigations have

focused on just one type of genetic interactions (GI) – synthetic lethality (SL). However, there

may be additional types of GIs whose systematic identification may markedly enrich the

molecular and functional characterization of cancer. Here, based on a novel data-driven

approach, we identify ~72K GIs of 11 new types, shared across cancers. These GIs are highly

predictive of patient survival, stratify breast cancer tumors into refined subtypes, and explain

differences in patients' response to drugs and cancer driver genes' tissue-specificity. These

results markedly expand the scope of cancer GIs and lay a strong conceptual and

computational basis for future studies of additional types of GIs and for their translational

applications.

Introduction

Many cellular functions are mediated by functionally interacting networks of genes. Genetic

interactions (GIs), whereby the phenotypic effects of a gene's activity are modified by the

activity of another gene, are thus a key to understanding complex diseases, including cancer,

which involves an interplay among a myriad of genes[1–6]. GIs are of particular interest in cancer

because, on one hand the dependence of one gene's phenotypic effect on the activity state of

another gene provides opportunities for selective killing of cancer cells with minimal side

effects[7], and on the other hand, it can render a drug targeting a particular gene ineffective

because of unfavorable state of another gene, and it can result in emergence of resistance[8,9].

Three types of GIs have been studied so far, showing significant roles in cancer

progression and patient survival, and suggesting novel therapeutic avenues. First and foremost

are Synthetic Lethal (SL) interactions, describing the relationship between two genes whose

individual inactivation results in a viable phenotype while their combined inactivation is lethal to

the cell[1,9–11]. The vast majority of GI studies up until now have focused on the SL, which have

long been considered as important for developing selective anticancer drugs[9]. Such drugs

inhibit the SL partner of a gene that is inactivated by genomic alterations in the particular

cancer[1,6,12], thus primarily killing the cancer cells with few cytotoxic effects on healthy cells.

Another related class of GIs are Synthetic Dosage Lethal (SDL) interactions, where the

underactivity of one gene together with the over-activity of another gene is lethal but not each

event by itself[10,13,14]. SDLs are promising for targeting tumors with over-activated oncogenes,

many of which are difficult to target directly, by targeting their SDL partners[15–17]. A third class of

GIs are Synthetic Rescues (SR), where a change in the activity of one gene is lethal to the cell

but an alteration in its SR partner 'rescues' cell viability. SRs may play a key role in tumor

relapse and emergence of resistance to therapy[18–20]. Indeed, previous investigations have

shown that the overall numbers of functionally active SLs and SDLs in a given tumor sample is

highly predictive of patient survival[13]. SLs, and even SDLs and SRs whose investigation in

cancer has just began, represent just the 'tip of the GI iceberg', though, as there are many

additional types of GI that can be conceived at a conceptual level, but have remained

unchartered, and whose systematic exploration may have important functional ramifications for

cancer therapy.

Here we present a novel data-driven computational pipeline, called ''EnGIne"

(Encyclopedia of clinically significant GIs in cancer) for the identification of multiple types of GIs.

The application of EnGine to analyze 5288 TCGA samples (Methods) belonging to 18 different

cancer types reveals 71,946 significant GIs of 12 distinct types, having either positive or

negative associations with patient survival. A GI between two genes is considered functionally

active in a given tumor if the activity states of the two genes correspond to the definition of that

GI. We show that a tumor-specific 'fitness' score based on the active GIs in the tumor is highly

predictive of patient survival, comparing favorably with the gene-level approach, and that it can

be utilized for predicting a tumor's drug response. Furthermore, the GI activity profile of a tumor

can be used for stratifying breast cancer into functionally and clinically relevant subtypes. In

sum, these results provide the first comprehensive characterization of clinically relevant pan-

cancer GIs of 12 distinct basic types, substantively expanding the current knowledge of GIs in

cancer, and laying a strong conceptual and computational foundation for future studies of

additional GI types.

Results

1. Overview of the Encyclopedia of Genomic Interactions (EnGIne) Pipeline

The overall EnGine pipeline is summarized in Fig. 1 and the technical details are provided in the

Methods section. Here we provide an overview of the approach. Given a large set of tumor

transcriptomes (Fig. 1A), we first partition the expression level of each gene into low, medium,

and high activity state. Thus, for a pair of genes, there are 9 combinations, or bins, of possible

activity states for the two genes (Fig. 1B). For a given ordered pair of genes, each tumor sample

maps to exactly one of the 9 bins. Our goal is to identify GI pairs of the form $(x, y, b, \pm\alpha)$ such

that for the specific gene pair (x, y), the tumors in which the joint activity of (x,y) maps to bin b

have a significant fitness advantage (+) or disadvantage (-) with effect size α, relative to all other

tumors where the activity of (x,y) maps to a different bin. The effect size α is estimated by

measuring the difference in the survival curves between those patients where the activity of (x,

y) is in bin b in their tumors and those where it is not, as depicted in Fig. 1C. We term a specific

GI pair $(x, y, b, \pm\alpha)$ as *functionally active* in a given tumor if the activity states of (x,y) fall in bin b

in that tumor. Notably, we expect that the patients having a large number of functionally active

interactions with negative effects on the tumor and a small number of such positive effect

interactions are likely to have an overall higher survival and vice versa.

Our goal is to detect the genome-wide GIs in each of the 9 bins (Fig. 1B) searching over

a space of ~163M gene pairs corresponding to ~18,000 genes. We analyzed 5288 TCGA

samples for 18 cancer types (see Methods for initial filtering of TCGA samples). Underline{First}, as an

initial screening, we performed a Log-Rank test (depicted in Fig. 1C) for each gene pair in each

of the 9 bins and retained the 223,946 genes pairs that exhibit a significant association with

survival in any of the 9 bins (Methods). Second, if a potential GI in bin b has a differential effect

on tumor fitness, we expect the number of tumors that map to bin b to be relatively enriched (for

'+' interaction positively affecting tumor survival), or depleted (for '-' interaction negatively

affecting tumor survival). We thus applied an additional filter (Fig. 1D) to retain only GIs with a

molecular depletion/enrichment signal (Methods), yielding 179,444 gene pairs overall. Third, for

each retained gene pair, in each of the 9 bins, we implemented a Cox proportional hazards

model, specifically controlling for age, cancer-type, gender, and race, to assess whether being

in a particular bin is associated with patient survival, either positively or negatively (Fig. 1E);

while the second step aims to identify interactions that are selected for or against at the tumor

level, the third step aims to identify interactions that also potentially affect survival – only

interactions that have a consistent effect (positive on tumor growth and negative on survival, or

vice versa) are retained. Further refining the initial filtering based on the log rank test, the Cox

proportional hazards model enables us to specifically control for cancer type, race, gender, age

and individual gene effects in the interactions identification process (Methods). Finally, to

identify the most significant GIs in each bin independently, we applied an empirical False

Discovery Rate (FDR) correction based on the significance of the Cox interaction term of the

179,444 gene pairs (identified in the previous step) relative to those obtained for randomly

shuffled gene pairing as the NULL control. At FDR <= 0.01%, this resulted in 71,946 predicted

GIs across the 9 bins, of the form $(x, y, b, \pm\alpha)$, which form the final set of TCGA inferred GIs (Fig.

1E). Considering the symmetry among bins (bin 2 ~ bin 4 corresponding to low-medium

expression interaction; bin 3 ~ bin 7 corresponding to low-high expression interaction; bin 6 ~

bin 8 corresponding to medium-high expression interaction), there are 6 unique types of

interaction bins, and considering the two directions of the effect size yields a total of 12 basic

types of GIs.

Figure 1. Overview of the EnGIne pipeline. Given a large set of tumor transcriptomes (A), we
first partition the expression level of each gene into low, medium, and high activity state,
resulting in 9 joint activity state bins for any two genes (B). Each gene pair and bin *b* induces a
bipartition of the set of tumor samples based on whether the activity levels of the gene pair
examined is in bin *b* in the specific tumor. The first step of EnGIne is a preparatory one, aiming
to screen for the gene pairs that show distinct survival trends in the two sets of tumors in any of

the bins, based on log-rank test (C). Next, for a gene pair and a bin identified in the previous
step, we test whether the putative gene interaction in bin *b* has a differential effect on tumor
fitness, by testing for depletion/enrichment of samples in the bin *b* relative to expectation based
on individual genes (D). Finally, for each retained gene pair, in each of the 9 bins separately, we
fit a Cox proportional hazards model to assess whether being in a particular bin is associated
with a distinct (positive or negative) pattern of patient survival, followed by correction for multiple
hypotheses testing (E). The output of this computational pipeline is (a) a list of GIs of each of
the 12 types studied, and (b) GI profile in each of the individual tumor samples, defined as
activity state of each GI in the tumor sample (F).

EnGIne TCGA analysis reveals survival-associated GIs involving known cancer drivers

Applying EnGIne to 5288 tumor samples spanning 18 types of cancers results in the selection of

71,946 GIs of 12 different GI types (~0.02% of all the possible candidate gene pairs and GI

types tested). Considering the expectation that neighboring genes in the protein interaction

network (PIN) are more likely to be involved in a GI[21], we aimed to additionally construct a PIN-

supported GI network. Knowing that the a great many of the interactions may not correspond to

a direct PPI interaction[22], we filter the GIs that are separated by more than two edges in the

PIN, resulting in a PIN-supported GI network composed of 1704 gene pairs. Among the 1786

genes involved in these 1704 PIN GIs (Supplementary Table S1) we identified 133 known

cancer driver genes (Cosmic dataset[23]) that are associated with various cancer types

(enrichment P-value < 2.5E-22) and 50 breast cancer specific (Intogen dataset[24]) driver genes

(enrichment P-value < 7.7E-13).

Fig. 2A shows the distribution of the detected 1704 GIs across the 12 GI types. Notably,

SL interactions, whose functional activity has a negative effect on tumor fitness and are the

most experimentally studied type of GI, are surprisingly one of the least abundant types of GI,

and so are SDLs (1% of all GIs). Remarkably, the positive "anti-symmetric" type of SLs, which is

composed of interactions where the joint low activity of the two interacting genes is associated

with a higher tumor fitness, are 3 times more abundant than SLs. The Interaction between the

cancer genes GNAQ and JAK2 is one example of such a positive interaction in bin-1

(Supplementary Fig. S1). Our analysis reveals several other types of GIs to be much more

abundant than the SL and SDL-type GIs. In fact, in the 2 most abundant types of pan-cancer

GIs (Bin-2 and Bin-6), one of the genes has medium level of activity and only the extreme

activity of its partner gene reveals a phenotypic effect. For most GI bins we see a higher

proportion of GIs exerting a positive effect on tumor fitness, which corresponds to a negative

association with patient survival. Importantly, it is consistent with the hypothesis that the GIs

uncovered during the evolution of cancer are under positive selection.

We tested and validated EnGIne by capitalizing on a collection of 7297 interaction pairs

that were tested experimentally in SL *in vitro* screens, collected from the literature[25–30]. Each of

the three filtering steps of EnGIne (Fig. 1C-E), could discriminate the experimentally determined

positive SLs from the negative ones, with ROC-AUCs of 0.63 (P-value < 0.0005), 0.62 (P-value

< 0.001), and 0.59 (P-value < 0.012), respectively. The SL screening was done *in vitro*, a

platform known to exhibit major discrepancy with in-vivo data. Given that our goal is to identify

clinically-relevant effects and thus the EnGine predicted GIs are inferred directly from patient

data, this significant recapitulation of the SL signal in pan-caner heterogeneous tumor data,

while of modest accuracy, is encouraging and supports the contribution of each of the individual

steps in EnGIne. As shown in Supplementary Fig. S2 and detailed in Supplementary Results,

we found that known cancer drivers are enriched among GIs of various types, and interestingly,

relative to tumor suppressors, oncogenes show a greater enrichment and are especially

enriched among GIs in bin-3, having a positive effect on tumor fitness as expected.

We investigated the biological processes involved in each of the 12 broad types of GIs

shown in Fig. 2A. We found an enrichment (FDR <= 0.01) of several processes with well-

established links to cancer among the genes involved in these GIs (Methods). These include

cell cycle control, cell division, cell adhesion, extracellular matrix organization, angiogenesis,

hypoxia, the Wnt signaling pathway, and more (Supplementary File 2). An illustrative example is

a genetic interaction involving a breast tumor suppressor ERCC2, a transcription-coupled DNA

excision repair gene[31,32], and a breast cancer oncogene KLF4, a zinc finger transcription

factor[33]. Fig. 2B show survival patterns for different activity state combinations of these two

genes. It reveals two interesting trends: First, the over-activation of the oncogene and under-

activation of the tumor suppressor (bin 3) results in poorer survival than expected from the

individual gene effects (bins, 1, 2, 6, and 9). Second, and surprisingly, the survival curve reveals

a reversal of the effect of the tumor suppressor ERCC2 inactivity on survival when the

oncogene KLF4 has medium activity (bin 2), whose individual activity is actually associated with

better survival. This and several other examples of GIs involving a cancer driver gene

(Supplementary File 1) demonstrate that the context-specific effects of driver genes may show

very different trends than their previously established individual gene effects. Supplementary

Fig. S3 shows the extended GI-network involving the known driver genes.

Fig. 2C shows the predicted GIs involving the well-established oncogene Myc and tumor

suppressor BRCA1 obtained by target-specific FDR (Methods), highlighting their number and

variety. On one hand, this richness of GIs involving these driver genes might account for the

lack of efficacy in targeting cancer drivers in some cases, and on the other hand, our analysis

suggests combination therapies jointly targeting GI partners of these driver genes. Consistent

with its Myc's role as an oncogene, the GIs occurring when Myc has low activity mostly have

negative effect on tumor fitness. However, when Myc is activated, we find that the low

expression of PUF60, a repressor of Myc[34,35], is associated with higher tumor fitness (type 3+

interaction). Fig. 2D demonstrates that high expression of Myc does not significantly contribute

to poorer prognosis (P-value $\sim$ 0.9), unless its suppressor PUF60 is lowly expressed (HR =

1.37, P-value < 5.0E-04), underscoring the importance of molecular context in developing anti-

Myc treatments. Although PUF60 is known to suppress Myc, we provide evidence for a

functional relationship that deviates from a simple epistatic interaction.

Figure 2 – Broad distribution and characteristics of the detected GIs and context-specific effect of cancer driver genes on survival. (A) Distribution of the 1704 significant PIN-supported GIs across 9 joint activity bins. The fractions of GIs in each bin are shown for GIs with positive (blue) and negative (red) effect on tumor fitness. Only the data in the lower half of the matrix are shown as the GIs are symmetric relative to the genes in a pair. (B) The Kaplan-Meier (KM) survival curve of GI involving ERCC2, a transcription-coupled DNA excision repair gene, known to be a breast cancer tumor suppressor, and KLF4, a zinc finger transcription factor known to be oncogenic in breast cancer, reveals increasingly poor survival by over-activation of the oncogene and under-activation of the tumor suppressor (bin 3). Strikingly, the effect of ERCC2 inactivity on survival is reversed when KLF4 has medium activity level (bin 2). (C) The predicted GIs involving the oncogene Myc and the tumor-suppressor BRCA1. (D) KM survival curve of GI involving Myc and its suppressor PUF60. High expression of Myc is associated with poor prognosis specifically at low activation of PUF60.

Finally, we quantified the ability of tumor-specific GIs to predict patient survival relative to

a comparable gene-wise approach, both in cross-validation setting in TCGA[15], as well as in an

independent breast cancer METABRIC dataset[36] (Fig. S4A, supplementary note 3 for additional

datasets discussion). As detailed in supplementary results, overall, the GI-based approach

achieves higher concordance than the gene-wise approach. An even bigger improvement is

observed in the independent dataset, testifying that the GI-based approach is generalizable,

while the individual-gene approach fails (concordance $\approx$ 0.51). Supplementary Fig. S4B depicts

the survival prediction accuracy of each GI type. As evident, interactions involving both genes in

their wild type mid-activity levels (i.e. bin 5) have negligible predictive power on survival,

testifying that more extreme levels of gene expression tend to be involved in functional GIs

affecting survival.

Differential GI activation between drug-response groups
236
To assess the extent to which tumor-specific GIs are associated with the tumor's response to a

drug we studied whether the detected GIs exhibit differential activity between patients who are

responsive to a drug and those who are not. For a given drug and cancer type having data on

responders and non-responders, we first identify drug-specific GIs based only on the samples

that do not have drug response information to avoid circularity (Methods); note that this network

is different from the generic GI networks presented in the previous sections, as focusing on

specific drug target genes requires a smaller multiple hypotheses burden, resulting in a greater

number of drug target-specific GIs than in the TCGA global pan-cancer network above. Also,

because the drugs considered are inhibitory, it suffices to focus on GI bins 1, 2, and 3, where

one of the genes (the drug's target) is presumed inactivated. For each identified GI, we test,

using Fisher exact test[37] whether the GI exhibits differential activation between the responders

and the non-responders (Methods). Note that, for positive GIs, we expect a lower activation

among responders and for negative GIs we expect a higher activation among responders.

However, owing to very small and unbalanced numbers of responders and non-responders

(5~35 samples per response group per drug), the Fisher test is underpowered and despite high

effect sizes, the p-values for many individual tests are not significant. Given that, we compared

the Fisher test p-values of the GIs with those obtained using randomly shuffled drug-response

labels, using paired Wilcoxon test[38] (Methods), performed separately for each drug-cancer type

pair and segregated by GI type.

We considered 12 drug-cancer type pairs that have tumor size reduction information following

treatment for at least 10 patients (at least 5 responders and 5 non-responders) in TCGA. Fig. 3A

shows the significance of differential activation of the drug-specific GIs, segregated by the 6 GI

types. Overall, in 18 of the 72 tests of drug-cancer type combinations, GIs of a particular type

exhibit differential activation between responders and non-responders, consistent with the

predicted effects of the GIs (5 fold enrichment for $P \leq 0.05$). Reassuringly, several drugs come

up as significant using SLs (GI type -1), consistent with previous reports showing the role of SLs

in mediating drug response[39]. Among the drugs, Gemcitabine, Lomustine and Paclitaxel exhibit

differential GI activation for most GI types (aggregate P-values = 4E-16 ~ 2E-11).

To further explore the most differentially activated GIs, we imposed an empirical FDR threshold

of 0.01 on the Fisher test P-value yielding a total of 521 GIs for the 12 drug-cancer type

combinations (Methods, Supplementary Table S2). Individual genes comprising the 521 GIs are

closer to each other's interacting gene in the PPI network relatively to shuffled pairs (Wilcoxon P

< 0.001, Methods) and have a significantly increased number of direct PPI interactions between

them (Fisher P < 0.02, Methods).

As an illustrative test case, we explored the GIs associated with the response to *Paclitaxel* in

Head and Neck Squamous Cell (HNSC) Carcinoma. We identified a GI involving the inactivation

of BCL-2 (known to suppress apoptosis, indirectly inhibited through phosphorylation[40]) and the

over-activation of ITPR1 (Inositol 1,4,5-trisphosphate receptor type 1, coding for IP3 Receptor)

negatively affecting tumor fitness. Interestingly, our analysis shows that this GI is activated

among the responders at a significantly higher ratio than among non-responders (odds-ratio $\approx$

11.1). Interaction between the IP3 receptor and BCL-2 is well characterized[41–43]. BCL-2 exerts

its oncogenic effect by inhibiting IP3-dependent channel opening and Ca^{2+} release from the ER

(by inhibiting mitochondrial Ca^{2+} overload), and thus preventing cancer cell apoptosis. Our

analysis strongly suggests that BCL-2 inhibition by *Paclitaxel* is especially effective when the

IP3 receptor is abundant and therefore increased Ca^{2+} release is possible. We find additional

interactions with literature evidence[44] among the *Paclitaxel* targets (Fig. 3B), however without

complete experimental characterization, suggesting promising avenues for additional studies.

287
288
GIs explain cancer type-specific effects of cancer driver genes

Many of the known cancer driver genes affect tumor initiation and development in a tissue-

specific manner. We assessed the extent to which the molecular context, characterized by the

functionally active GIs, may explain the tissue-specific effects of cancer drivers. Toward this, we

identified 15 oncogenes and 20 tumor suppressors whose effects are likely to be restricted to

specific cancer types, based on preferentially high mutation rates in those cancer types,

including breast, bladder, and gastric cancer (Methods, Supplementary Table S3). We then

assessed the risk-score associated with each of these driver gene's GI partners in each of the

TCGA samples (Methods). We hypothesized that the specific activity states of the GI partners of

tumor-suppressors and oncogenes may explain their functional significance (as estimated by

their effects on patient survival) in the specific tissue(s) where the driver's effect is known to

manifest itself. Indeed, for 15 out of 35 (~43%) driver genes, we see a higher GI risk score

(measuring the functional activity of the predicted GI partners) in the specific cancer samples

belonging to the type they are known to be associated with in the literature, compared to their GI

risk scores in the other cancer types (Wilcoxon rank-sum test, FDR < 0.1, Supplementary Table

S3). For instance, HLF, a bZIP transcription factor, has been linked to lung and breast cancer

based on its significantly greater missense mutation frequency in those cancer types[24]. We

observed a significant difference (FDR < 1.07E-12) in GI activation between breast and lung

cancer relative to the other tissues (Fig. 3C,D), where positive interactions are preferentially

activated in these two tissues while negative interactions are preferentially activated in the

background tissues, consistent with the increased tumor fitness in the foreground tissues.

Among the GI partners, JAMIP1 is a member of the Integrated Breast Cancer Pathway and

CLPTM1 is associated with Estrogen-Receptor Negative Breast Cancer. Overall, these results

imply that cancer type-specific effects of many driver genes may be explained by their tissue-

specific GI network activity (note that mutation data used here was not used in the inference of

these GIs, so there is no circularity).

Figure 3 – (A-B) differential GI activation between drug response groups. (A) For each drug (left row labels) and each cancer type (right row label) combination, and for each GI type (columns), the heat plot shows the significance of differential activation of GIs in responders and non-responders consistent with expectation. The last column shows the significance when all GI types are aggregated. (B) The network shows the inferred functional interactions (based on STRING[44]) among the genes interacting with *Paclitaxel* targets, as well as inferred GI types. **(C-D) Differential GI activation between tissues in gene-specific GI-network.** For HLF-specific GI network the figure shows the activity states of GIs in breast and lung cancers (C - foreground tissues) and in other cancer types (D - background tissues). The edge weight (thickness) represents the fraction of samples in which the GI was functionally active. Several GIs can be

328 seen as differentially active in the two sets of cancer types. The figure also depicts cell surface
329 proteins among the HLF's GI partners. The GI network-based sample-specific risk score is
330 significantly higher (q-value < 1.07E-12) in breast and lung cancer relative to other cancer
331 types, potentially mediated by a selective activation of positive GIs in the foreground tissues and
332 negative GIs in the background tissues.
333

Stratifying breast cancer tumors based on their GI profiles

We set out to study whether functionally active GIs in a tumor can provide an interaction-based

alternative to gene expression or sequence data conventionally used for tumor stratification into

various sub-types. Given the cancer type specificity of many of the GIs studied above, we asked

whether GI-based characterization of breast cancer is effective in providing a functionally

relevant stratification of tumors to their subtypes. We used the raw GI profile of a tumor (which

consists of 1704 binary values representing the functional activity of all PIN-supported GIs) to

cluster the 1981 breast cancer samples in an independent METABRIC dataset [36] (recall that the

GIs were identified in TCGA data) using a conventional Non-Negative Matrix Factorization

(NMF) (Methods). Since NMF requires a predetermined number of clusters as input, we applied

the method for a range of 2-15 clusters and assess the clustering quality using Dunn's index,

repeating the clustering procedure used in the original work[36]. The Dunn's index achieved the

maximum value for 10 clusters (Supplementary Fig. S7A) and upon closer inspection of the

distributions of known breast cancer subtypes in these clusters (Supplementary Fig. S7B) we

merged two of the clusters, thus yielding 9 clusters for further analyses.

As shown in Fig. 4A, the 9 clusters have distinct survival characteristics with an overall

mean hazard ratio (HR) difference of 1.94 (P-value ~ 0). Supplementary Fig. S8A shows the

survival characteristics obtained for the previously published clustering of the METABRIC

samples[36]. As evident, both approaches obtain similar survival separation levels, but exhibit

differences in their histopathological composition.

Currently Breast cancer has 5 well-established clinically distinct subtypes based on the

tumors' histopathological attributes. Fig. 4B shows the fraction of these known subtypes among

the 9 GI-based clusters. Several clusters are highly associated with specific subtypes such as

Basal [triple-negative] (cluster 5), Luminal A (cluster 3,4) etc. Others show association with

several subtypes, e.g., Luminal A and B are clustered together in cluster 8. Interestingly, the

basal subtype, which are largely triple-negative and have poor prognosis, correspond to a

distinct cluster in our analysis (cluster 5), consistent with their distinct clinical status. In the

original METABRIC publication 50% of the samples were left unassigned to any of their 10

clusters in METABRIC while our GI-based clustering covers all samples. In order to provide an

equivalent comparison, we selected a matching number of samples (50%) such that we retain

the samples that are the most confidently associated with a cluster. Supplementary figure

S8B,C compare the GI-based clusters and the clusters in the original METABRIC publication in

terms of their histopathological subtype composition.

Fig. 4C,D illustrate the added survival predictive value of GI based characterization over

the classical histopathological ones. Our evaluation proceeds along two different perspectives:

(1) cases where known histopathological breast cancer subtypes are split across multiple GI-

based clusters (e.g. Luminal B across clusters 1, 2 and 8), and conversely (2), cases where one

GI-based cluster harbors multiple known histopathological subtypes (e.g. cluster 2 contains Her-

2 and Luminal B subtypes). In the former we find, e.g., that the 1989 Luminal B tumors that are

split across different GI-based clusters exhibit very distinct survival trends (Fig. 4C). Likewise, in

the latter case, we find that the survival trends of Luminal A and Luminal B histopathological

subtype samples that are assigned to the same GI-based cluster 8 do not show a significant

difference in their survival trends (Fig. 4D). We systematically identified 6 additional cases of the

above two kinds where (1) a known tumor histopathological subtype was split across multiple

GI-based clusters or (2) multiple known histopathological subtypes were assigned to the same

GI-based cluster (and each cluster has at least 30 samples). In each case of the first kind we

tested for differences in survival and in each of the second case we tested for lack thereof. As

shown in Supplementary Fig. S9, in 5 out of 6 cases we found that the GI-based clusters

provided a more accurate survival difference than the known histopathological subtypes. In

contrast, we identified 4 cases of GI-based clusters of the second kind and in none of them the

survival trends of the samples with different histopathology markers were significantly different

(Supplementary Fig. S10).

Additionally, we assessed whether the samples in GI-based clusters harbor distinct gene

mutations. Based on the TCGA mutation data, we identified 196 genes (Supplementary Table

S4) with significantly greater mutation frequency in one or more of the clusters, relative to their

overall mutation frequency in breast cancer (Methods). Fig. 4E shows the mutational frequency

profiles of these genes across the 9 clusters. Overall, the differentially mutated genes across

the GI based clusters include 10 cancer drivers: CDK12, CDKN1B, DNAJB1, ERBB2, EXT2,

FCGR2B, FNBP1, HOXC13, SEC24D, and PDGFRB. Cyclin dependent kinase Cdk12, involved

in DNA damage response, is known to affect BRAC1 transcription [45]. Likewise, cyclin CDKN1B

is known to be mutated in luminal breast cancer [46]. Our analysis suggests a more specific

enrichment of this mutation in subset of Luminal-A tumors (cluster 9 in Fig. 4B). The HER2

breast cancer subtype is conventionally characterized by overexpression of HER2 gene (also

known as ERBB2), but rare activating mutations of this gene have also been implicated in the

disease, independent of overexpression [47]. Our analysis reveals an enrichment of HER2

mutations in cluster 1, which includes only a small subset of annotated HER2 subtype tumors.

Most of the HER2 mutations (11 out of 13) are predicted to have a significant impact on the

protein structure (SIFT and PolyPhen measures, Methods). This may suggest a distinct

oncogenic mechanism underlying this subclass of tumors, which may express a HER2-like

subtype including a few samples where HER2 is over-expressed and many others where it is

not over-expressed, but mutated. As another example, the FCGR2B mutation modulates HER2

breast tumor's response to Trastuzumab [48]. Our analysis reveals a specific enrichment of

FCGR2B mutations in cluster 6, which includes several other breast cancer subtypes besides

HER2, pointing to the potential relevance of these mutations in tumors belonging to other

cancer subtypes as well (we emphasize that no mutation data was used at all in the GIs

inference via EnGIne).

We quantified the fraction of each of the 12 types of functionally active GIs among the

samples in each cluster (see Methods). Fig. 4F shows the active GI profiles of each cluster, and

reveals two broad subgroups, one including clusters 4, 3, 8, and 1 and another including

clusters 2, 5, 6, and 7. Interestingly, the two subgroups clearly segregate in terms of their

survival, testifying that the GI profiles capture a simplified yet robust characterization of the

clinical prognosis. Supplementary Fig. S11 compares the GI-profiles of clusters revealed in the

TCGA and the METABRIC breast cancer data and shows a high degree of consistency. A

global comparison of the GI profiles of the 9 clusters in the two datasets shows a Spearman

correlation of 0.67 (P-value < 2.4E-14) between the GI types composition of these clusters,

implying that GI-profiles are a robust characteristic of breast cancer tumors across different

tumor collections.

Figure 4 – (A) Mean survival curves of the individuals in the 9 inferred GI-based breast cancer subtypes. **(B) Cluster subtype composition based on PAM50 breast cancer sub typing**[49]. **(C-D) Survival trends of tumors of known histopathological cancer subtypes within and across GI-based clusters.** (C) Luminal B samples that are split across GI-based clusters 1, 2 and 8 show significant survival differences. (D) Her2 and Luminal-B type tumors that are included within the GI-based cluster 2 exhibit similar survival trends. **(E) Mutational profile of GI-based breast cancer subtypes**. Mutation profiles of 196 genes (rows) across the 9 GI-based clusters (columns). For each gene and each cluster, the figure depicts the fraction of samples in the cluster in which the gene is mutated. **(F) GI types composition of the GI-based breast cancer subtypes in the METABRIC dataset**. In clustering the samples based on GI profile, each GI is probabilistically assigned to a single cluster, based on which, the composition of GIs assigned to each cluster is obtained. The x-axes represent the 12 GI types (6 activity bins and 2 directional effects on survival), and the y-axes represent the clusters. The colors represent the fraction of cluster-assigned GIs of each GI type.

Discussion

Analyzing molecular and clinical data across thousands of tumors of dozens of types, here, for the first time, we comprehensively map the landscape of 12 basic GI types in cancer. Our work substantially extends previous investigations of gene interactions in cancer, which have been almost entirely focused on synthetic lethality (SL: corresponding to positive effect on survival in bin 1)[26,27,50], with a few studies of synthetic dosage lethality (SDL: corresponding to positive effect on survival in bin 1)[11,12], to a total of 12 types of interactions. The identified GIs are predictive of patient survival and drug response based on the tumor transcriptomes, and reveal novel functionally and clinically relevant breast cancer subtypes. The set of functionally active GIs thus provides a complimentary molecular characterization of tumors to those obtained by histology and contemporary individual gene-centric transcriptomic and sequence based profiles. A better concordance of the GI-based breast cancer subtypes with survival trends may be partly due to the fact that GIs were inferred based on their impact on survival. However, interestingly, the detected subtypes are marked by distinct mutational profiles, which were not utilized in inferring the GIs. Overall, these results underscore the importance of molecular context represented by functionally active GIs.

In this first conceptual foundation and application of EnGIne, we have analyzed the TCGA compendium to identify GIs interactions shared across many cancer types. Identifying pairwise genes interaction is only a first step toward capturing the true complexity of cellular networks. Future work can go beyond the 9 basis GI types studied here to investigate more complex GI types that involve different compositions of these basic types; for instance, a given interacting gene pair can confer tumor fitness benefit in multiple co-activity bins and reduce tumor fitness in others. Thus, while the results presented here go markedly beyond previous definitions and analyses of GIs, they only begin to explore the full scope and clinical potential of GI-based analyses of cancer, awaiting future investigations.

Methods and Materials

Cancer datasets

The Cancer Genome Atlas (TCGA)[15] consists of 8749 tumor samples across 28 cancer types, 3 races and 2 genders (data quality testing is described in supplementary note 1). We do gene expression binning specifically for each combination of cancer type, race and gender. Furthermore, we control for various clinical and demographic group specific effects. Because using small groups of samples may result in insufficiently robust models, we filtered rare combinations of clinical and demographic groups, resulting in 5288 mRNA samples derived from patients spread across 18 cancer types, 3 races and 2 genders. The RSEM normalized expression data was downloaded from the TCGA data portal and quantile normalization was applied within each sample. METABRIC breast cancer dataset[36] consists of 1989 microarray samples and was used for independent validation. Similarly, quantile normalization was applied in each sample.

Protein interaction Network (PIN)

The PIN was obtained from a previously published resource called the HIPPIE which aggregates physical protein interaction data from 10 source databases and 11 studies[21]. This network consists of 15673 proteins and 203159 edges.

Identification of gene interactions (GI) associated with cancer patient survival

As shown in Fig. 1, we divide the range of each gene's expression across tumor samples into 3 equal-sized bins that correspond to the 3 activity states: low, medium and high expression. Given a gene pair, each tumor sample is thus mapped into one of the 9 joint activity states of the two genes. The choice of dividing a gene's activity into three classes, while somewhat arbitrary, was made in consideration of interpretability of functional states and robustness of inference. However, to account for differences in expression distributions across clinical and demographic confounders, we apply subpopulation specific binning approach. We considered the following categorical confounders: cancer type, race, and gender. We considered the combination of confounder states for which there were at least 100 tumor samples (Supplementary Table S5). Our binning is thus not confounded by various clinical and demographic variables.

The GI pipeline consists of three steps that successively refine the predictions to arrive at high-

confidence set of predicted GIs. As such, the number of all pair-wise combinations of genes is

excessively large to apply a comprehensive Cox regression model.

Step 1: Log-Rank. In the first step, for each of the ~163 million gene pairs, say (X,Y), we

compute the Log-rank statistics [51] estimating the survival difference between the samples that

map to one of the 9 activity bins and the other 8 bins. We implemented the log-rank test in C++

for computational speed. To control for gene-wise effect, we compare the Log-rank statistics of

the gene pair (X,Y) (in a bin) with those for (X,U) and separately with those for (V,Y), where U

and V represent all other genes. We consider Log-rank(X,Y) to be significant if it among the top

0.1% relative to all (X,U) and top 0.1% among all (V,Y) gene pairs. We retain a gene pair if it is

deemed significant in any of the 9 bins. This procedure retained 210,612 gene pairs of the total

of ~163M.

Step 2: Molecular enrichment and depletion. For a gene pair, we expect that its positive

(respectively, negative) effect on survival should reflect in negative (respectively, positive)

selection on the cancer cell's viability. We therefore expect the number of samples mapped to

the corresponding activity bin relative to the gene pair to be depleted (respectively, enriched).

We only retained the gene pairs which the number of samples in a particular bin were lower

(bottom 45 percent) or respectively, greater (top 45 percent among all gene pairs) than

expectation, reducing the number of GIs to 271,096 across 179,444 gene pairs (recall that a

gene pair can participate in multiple GIs corresponding to multiple bins and effect on survival).

Step 3: Cox proportional hazard test. Cox proportional hazards model is the most widely

acceptable approach for modeling survival while accounting for censored data as well as

confounding factors. We modeled each of the retained gene pairs' effect on survival in each of

the 9 bins using the interaction status λ (active if the sample mapped to the bin and inactive

otherwise), along with the confounders. Specifically, we introduced the two individual genes σ_1,

σ_2 expression to model each gene's independent effect on survival, and additionally, clinical and

demographic confounders, namely, cancer type, race, gender, and age. The model is stratified

based on the discrete confounders, to account for differences in the baseline hazard (risk)

characteristics. We did not control for tumor stage and grade as these classifications reflect the

very same tumor characteristics our model aims to capture and such control would prevent us

from learning an important element of the disease. Control for genomic-stability and tumor purity

as a potential confounder is described in supplementary note 2.

$$527 \qquad risk \sim \sigma_1 + \sigma_2 + \lambda + age + strata(type, gender, race)$$

Cox modeling provided a P-value representing the significance of the effect of joint gene pair

activity on survival. To obtain a NULL distribution for the P-values, we repeated this process for

corresponding list of randomly shuffled pairs. We retained ~71K gene pairs in the top 1% of the

NULL P-values, corresponding to a nominal FDR of ~3.8%.

Optional Step 4: Filtering by protein interactions. To gain additional confidence in the

predicted GIs, given the greater tendency (and expectation) for neighbors in the Protein

Interaction Networks (PIN) to exhibit functional interactions (see Results section), we further

refined the GI set by retaining the pairs that are found within distance of 2 in the PIN. Overall we

obtain a set of 1704 GIs that exhibit molecular and clinical evidence in cancer as well as

evidence from the PIN network.

GO enrichment of hub genes

We conducted Gene Enrichment analysis on genes involved in GIs using GOrilla tool[52] taking all

17990 genes as background set. For enrichment analysis, we considered the genes into

multiple categories and the obvious categories can be the genes separated by 12 broad GI

types. However, the GI bins (Fig 1B) suggests that unlike in bin 1, 5, and 9, in bin 2, 3, and 9,

one of the 'x' gene and 'y' gene has higher or lower expression/activity than the other.

Therefore, considering bin number, patient survival, and gene activity states, we have total 18

(taking bin symmetricity into account) kinds of gene sets. For each gene sets, we conducted

Gene-Ontology enrichment analysis using Gorilla tool[52]. For the enrichment analysis, we have

used all 17990 genes as background set.

Patient Survival Risk Prediction

We applied a semi-supervised approach to assign a risk score to each patient according to the

functionally active interactions in the sample. Consider a GI involving genes X and Y conferring

a positive effect on the tumor fitness in a particular joint-activity bin B (Fig. 1). If in a sample

genes X and Y fall in bin B then then the GI is said to be 'functionally active' in the sample, and

a score of +1 is contributed to the overall tumor 'fitness'. Likewise if the GI has negative effect

on the tumor fitness then a score of -1 is contributed. For each sample, we compute the overall

score conferred by functionally active GIs (either in a bin-specific and effect-specific fashion, or

overall) in the sample. The higher the tumor 'fitness' score the lower the survival potential.

However, to make our approach comparable to gene-wise approaches[53], we assigned each

gene the sum of the contributions by all active GIs involving that gene. The estimated gene-wise

GI score is used as a predictor variable in a Cox model along with the confounding factors

discussed above to predict patient's survival.

This model is trained on the same data used for the GIs training and then validated on its

cross-validation counterpart, or in the case of independent validation, the model is trained on

the full TCGA dataset, and tested on an independent METABRIC breast cancer data with 1989

samples[36]. The prediction accuracy is estimated in terms of widely used C-index[54]. Several

previous publications have assessed survival risk prediction accuracy based on dichotomized

analysis where samples are separated into distinct low- and high-survival groups and their

survival curves the compared[51], which is prone to overestimating prediction accuracy. For

comparison, we have also performed accuracy estimation following the dichotomized

comparison of survival risks between the extreme cases of predicted survival risk groups, for

variable threshold to define the extreme (such as top versus bottom 10% or top versus bottom

20% and so on).

To compare the GI-based survival prediction with individual gene approach, we implemented an

analogous scheme for individual genes where the gene expression values were discretized into

3 expression levels (low, medium and high), and the discretized representation was used as a

predictor variable in a controlled Cox regression model to obtain the significance (P-value) of

each gene with respect to survival prediction. The most significant predictors (top 5%) were

chosen and precisely used as described for the GIs survival prediction procedure. An analogous

procedure was used to estimate the prediction accuracy based on both individual gene effects

and GIs.

Characterization of differential GI activation between drug-response groups

We inferred the GIs involving each drug's target gene based on the TCGA samples that do not

have that drug's response information to avoid circularity, and filtered based on FDR as for

global pan-cancer GI detection as above, but restricted to the target-specific GIs. For each of

the drug-specific GIs, we compare its activation frequency (whether the GI was functionally

activate or inactivate) among the responders (stable disease, partial response and complete

response categories) and non-responders group (clinical progressive disease categories), using

one-sided Fisher's exact test[37], where the alternative hypothesis is that negative (respectively,

positive) GIs are more frequently active among responders (respectively, non-responders).

However, given the extremely small and imbalanced sample sizes, and the conservative nature

of Fisher's exact test[55], we test whether the overall distribution of the obtained Fisher test p-

values are lower than those obtained using randomly shuffled drug-response labels, using one-

sided Wilcoxon test[38]. We thus obtain a p-value for each drug-cancer type pair, segregated by

GI type.

Then, for each gene pair in the inferred GIs list and, as a control, in a shuffled list of size 10x as

the original GIs list size, we compute the distance between the genes in the PPI network[21]. We

then use one-sided Wilcoxon test[38] to assess whether GI gene pairs are closer to each other

than random expectation. Alternatively, we also compare the number of directly GI gene pairs

having direct interaction using one-sided Fisher's exact test[37].

Characterization of tissue-specific effect of cancer driver genes

A recent study[56] of genes' somatic mutation profile across cancer types has identified and

characterized the tissue specificity of 459 candidate drivers. For each such candidate, we

matched the driver role annotation (oncogene or tumor-suppressor) obtained from the Census[23]

cancer genes dataset (genes with missing annotation were discarded) for 33 tumor suppressors

and 25 oncogenes. For each gene, we calculated the significant GI interactions involving this

gene, where the significance threshold is defined over the shuffled background interactions

involving this gene (90th percentile of the interaction confidence score). We further excluded

genes with 5 or fewer interactions with 300 or fewer samples where they are expressed,

resulting in 20 tumor suppressors and 15 oncogenes spanning 10 cancer types (Supplementary

Table S3). For each gene, the cancer types are partitioned into affected types (cancer types

affected by the driver) and the other unaffected cancer types. We hypothesized that higher-risk

will be associated with lower expression of a tumor-suppressor and higher expression of an

oncogene, much more in the affected cancer types relative to the unaffected cancer types.

Thus, we assess the associated risk-score of the samples found in the low (respectively, high)

expression bin for a tumor-suppressor (respectively, oncogene). Using a one-sided Wilcoxon

rank-sum test, we tested for higher risk-score in the samples in the affected cancer type in

comparison to those in unaffected types. After multiple hypotheses testing, 15 out of the 35

(~43%) driver genes were found to have significant tissue-specific GI-based risk score (FDR q-

value < 0.1, Supplementary Table S3).

Breast cancer tumor stratification

We represent a tumor sample as a vector indicating the functional activity status of each

predicted GI. This provides a survival-cognizant alternative to the widely-used gene expression

profile representation of a sample. We used this representation to stratify the METABRIC breast

cancer patients using Non-Negative Matrix Factorization (NMF)[57,58], which has suitable

statistical properties and has been shown to be effective in a variety of contexts[59]. Since NMF

requires a predetermined number of clusters, we performed the analysis for 2-15 clusters, and

assessed their fitness using Dunn's index[60], which quantifies compactness within and

separation across clusters. Similar analysis was performed on the TCGA breast cancer cohort.

The Hazard-ratio significance values were computed for each pair of clusters, while the P-

values were generated using multi-class log-rank test. For comparison purposes, to match our

estimated clusters' sizes to previously published METABRIC cluster sizes (~900 samples), we

constrained the number of samples in each cluster to the 100 samples that were found to be

most highly associated with the cluster.

Each cluster's GI profiles were constructed as follows. Our clustering approach – NMF,

assigned each GI to a single cluster. For each cluster, and for each of the 12 GI-types (6 bins in

Fig. 1 and the two directional effects on survival), we obtain the frequency of GIs of that type,

relative to all GI assigned to the clusters.

Mutation frequency analysis was performed on the TCGA clusters, where we first tested

whether the mutation frequency distribution of each gene differs significantly across clusters

using Chi-square test. The selected genes (Chi-Square test to ascertain the dependence of

mutation abundance on the interaction type) were then used to illustrate the mutational profiles

of the clusters.

The prediction values of a mutation's effect on the protein were obtained from the cBioPortal

repository. Out of the 196 differentially mutated genes found, 138 genes had matching extended

mutation information indicating their SIFT (sorts intolerant from tolerant amino acid

substitutions) and PolyPhene (polymorphism phenotyping) predictions. We calculated a gene-

wise ratio of mutations predicted to have a significant effect on the protein. 68 (49%) and 63

(45%) of which had at least 50% of the mutations belonging to the tested category in SIFT and

PolyPhene, respectively.

The breast cancer subtypes were derived using the widely accepted PAM50 algorithm[49]. The

METABRIC PAM50 subtypes were annotated in the original publication[36], while the TCGA

breast cancer subtypes were calculated using the original published class centroids[49].

References

1.   Ashworth, A., Lord, C. J. & Reis-Filho, J. S. Genetic interactions in cancer progression and treatment. *Cell* **145,** 30–38 (2011).

2.   Kelley, R. & Ideker, T. Systematic interpretation of genetic interactions using protein networks. *Nat. Biotechnol.* **23,** 561–566 (2005).

3.   Wong, S. L. *et al.* Combining biological networks to predict genetic interactions. *Proc. Natl. Acad. Sci. U. S. A.* **101,** 15682–15687 (2004).

4.   Zhong, W. & Sternberg, P. W. Genome-wide prediction of C. elegans genetic interactions. *Science* **311,** 1481–4 (2006).

5.   Lu, X., Kensche, P. R., Huynen, M. a & Notebaart, R. a. Genome evolution predicts genetic interactions in protein complexes and reveals cancer drug targets. *Nat. Commun.* **4,** 2124 (2013).

6.   Jerby-Arnon, L. *et al.* Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. *Cell* **158,** 1199–1209 (2014).

7.   Kaelin, W. G. The concept of synthetic lethality in the context of anticancer therapy. *Nat. Rev. Cancer* **5,** 689–98 (2005).

8.   Fong, C. Y. *et al.* BET inhibitor resistance emerges from leukaemia stem cells. *Nature* **525,** 538–+ (2015).

9.   Miyamoto, D. T. *et al.* RNA-Seq of single prostate CTCs implicates noncanonical Wnt signaling in antiandrogen resistance. *Science* **349,** 1351–6 (2015).

10.   Stuhlmiller, T. J. *et al.* Inhibition of lapatinib-induced kinome reprogramming in ERBB2-positive breast cancer by targeting BET family bromodomains. *Cell Rep.* **11,** 390–404 (2015).

11.   Sajesh, B. V., Guppy, B. J. & McManus, K. J. Synthetic genetic targeting of genome instability in cancer. *Cancers* **5,** 739–761 (2013).

12.   Kroll, E. S., Hyland, K. M., Hieter, P. & Li, J. J. Establishing genetic interactions by a synthetic dosage lethality phenotype. *Genetics* **143,** 95–102 (1996).

13.   Megchelenbrink, W., Katzir, R., Lu, X., Ruppin, E. & Notebaart, R. a. Synthetic dosage lethality in the human metabolic network is highly predictive of tumor growth and cancer patient survival. *Proc. Natl. Acad. Sci. U. S. A.* **112,** 12217–12222 (2015).

14.   Szappanos, B. *et al.* An integrated approach to characterize genetic interaction networks in yeast metabolism. *Nat. Genet.* **43,** 656–62 (2011).

15.   Chang, K. *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* **45,** 1113–1120 (2013).

16.   Rathert, P. *et al.* Transcriptional plasticity promotes primary and acquired resistance to BET inhibition. *Nature* **525,** 543–547 (2015).

17.   Luo, J., Solimini, N. L. & Elledge, S. J. Principles of Cancer Therapy: Oncogene and Non-oncogene Addiction. *Cell* **136,** 823–837 (2009).

18.   Hartwell, L. H., Szankasi, P., Roberts, C. J., Murray, A. W. & Friend, S. H. Integrating genetic approaches into the discovery of anticancer drugs. *Science (80-.).* **278,** 1064–1068 (1997).

19.   Brough, R., Frankum, J. R., Costa-Cabral, S., Lord, C. J. & Ashworth, A. Searching for synthetic lethality in cancer. *Current Opinion in Genetics and Development* **21,** 34–41 (2011).

20.   McLornan, D. P., List, A. & Mufti, G. J. Applying Synthetic Lethality for the Selective Targeting of Cancer. *N. Engl. J. Med.* **371,** 1725–1735 (2014).

21.   Schaefer, M. H. *et al.* Hippie: Integrating protein interaction networks with experiment

based quality scores. *PLoS One* **7,** (2012).

22. Costanzo, M. *et al.* A global genetic interaction network maps a wiring diagram of cellular
function. *Science (80-.).* **353,** aaf1420-aaf1420 (2016).

23. Futreal, P. A. *et al.* A census of human cancer genes. *Nat. Rev. Cancer* **4,** 177–83
(2004).

24. Gonzalez-Perez, A. *et al.* IntOGen-mutations identifies cancer drivers across tumor types.
Nat. Methods **10,** 1081–1082 (2013).

25. Lord, C. J., McDonald, S., Swift, S., Turner, N. C. & Ashworth, A. A high-throughput RNA
interference screen for DNA repair determinants of PARP inhibitor sensitivity. *DNA*
Repair (Amst). **7,** 2010–2019 (2010).

26. Martin, S. A. *et al.* Methotrexate induces oxidative DNA damage and is selectively lethal
to tumour cells with defects in the DNA mismatch repair gene MSH2. *EMBO Mol. Med.* **1,**
323–337 (2009).

27. Turner, N. C. *et al.* A synthetic lethal siRNA screen identifying genes mediating sensitivity
to a PARP inhibitor. *EMBO J.* **27,** 1368–77 (2008).

28. Luo, J. *et al.* A Genome-wide RNAi Screen Identifies Multiple Synthetic Lethal
Interactions with the Ras Oncogene. *Cell* **137,** 835–848 (2009).

29. Steckel, M. *et al.* Determination of synthetic lethal interactions in KRAS oncogene-
dependent cancer cells reveals novel therapeutic targeting strategies. *Cell Res.* **22,**
1227–45 (2012).

30. Bommi-Reddy, A. *et al.* Kinase requirements in human cells: III. Altered kinase
requirements in VHL-/- cancer cells detected in a pilot synthetic lethal screen. *Proc. Natl.*
Acad. Sci. U. S. A. **105,** 16484–16489 (2008).

31. Benhamou, S. & Sarasin, A. ERCC2/XPD gene polymorphisms and cancer risk.
Mutagenesis **17,** 463–469 (2002).

32. Bernard-Gallon, D. *et al.* DNA repair gene ERCC2 polymorphisms and associations with
breast and ovarian cancer risk. *Mol. Cancer* **7,** 36 (2008).

33. Akaogi, K. *et al.* KLF4 suppresses estrogen-dependent breast cancer growth by inhibiting
the transcriptional activity of ERalpha. *Oncogene* **28,** 2894–902 (2009).

34. Rahmutulla, B. *et al.* Alternative splicing of FBP-interacting repressor coordinates c-Myc,
P27Kip1/cyclinE and Ku86/XRCC5 expression as a molecular sensor for bleomycin-
induced DNA damage pathway. *Oncotarget* **5,** 2404–2417 (2013).

35. Matsushita, K. *et al.* Non-transmissible Sendai virus vector encoding c-myc suppressor
FBP-interacting repressor for cancer therapy. *World J. Gastroenterol.* **20,** 4316–4328
(2014).

36. Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours
reveals novel subgroups. *Nature* **486,** 346–52 (2012).

37. Fisher, R. A. On the Interpretation of χ 2 from Contingency Tables, and the Calculation of
P. *J. R. Stat. Soc.* **85,** 87 (1922).

38. Wilcoxon, F. Individual Comparisons by Ranking Methods. *Biometrics Bull.* **1,** 80 (1945).

39. Jerby, L., Waldman, Y., Weinstock, A., Geiger, T. & Ruppin, E. Genome-wide detection of
synthetically-lethal genes uncovers a novel repository of selective cancer targets. 1–9

40. Ruvolo, P. P., Deng, X. & May, W. S. Phosphorylation of Bcl2 and regulation of
apoptosis. *Leuk. Off. J. Leuk. Soc. Am. Leuk. Res. Fund, U.K* **15,** 515–522 (2001).

41. Oakes, S. A. *et al.* Proapoptotic BAX and BAK regulate the type 1 inositol trisphosphate
receptor and calcium leak from the endoplasmic reticulum. *Proc. Natl. Acad. Sci.* **102,**
105–110 (2005).

42. Chen, R. *et al.* Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate receptors to

regulate calcium release from the ER in response to inositol 1,4,5-trisphosphate. *J. Cell*
Biol. **166,** 193–203 (2004).

43.   Rong, Y.-P. *et al.* The BH4 domain of Bcl-2 inhibits ER calcium release and apoptosis by
binding the regulatory and coupling domain of the IP3 receptor. *Proc. Natl. Acad. Sci.*
106, 14397–14402 (2009).

44.   Szklarczyk, D. *et al.* STRING v10: Protein-protein interaction networks, integrated over
the tree of life. *Nucleic Acids Res.* **43,** D447–D452 (2015).

45.   Blazek, D. *et al.* The Cyclin K/Cdk12 complex maintains genomic stability via regulation
of expression of DNA damage response genes. *Genes Dev.* **25,** 2158–72 (2011).

46.   Stephens, P. J. *et al.* The landscape of cancer genes and mutational processes in breast
cancer. *Nature* **486,** 400–4 (2012).

47.   Sun, Z. *et al.* Analysis of different HER-2 mutations in breast cancer progression and
drug resistance. *J. Cell. Mol. Med.* **19,** 2691–2701 (2015).

48.   Norton, N. *et al.* Association Studies of Fcγ Receptor Polymorphisms with Outcome in
HER2+ Breast Cancer Patients Treated with Trastuzumab in NCCTG (Alliance) Trial
N9831. *Cancer Immunol. Res.* **2,** 962–9 (2014).

49.   Bernard, P. S. *et al.* Supervised risk predictor of breast cancer based on intrinsic
subtypes. *J. Clin. Oncol.* **27,** 1160–1167 (2009).

50.   Farmer, H. *et al.* Targeting the DNA repair defect in BRCA mutant cells as a therapeutic
strategy. *Nature* **434,** 917–21 (2005).

51.   Harrington, D. P. & Fleming, T. R. A class of rank test procedures for censored survival
data. *Biometrika* **69,** 553–566 (1982).

52.   Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: a tool for discovery
and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10,** 48
(2009).

53.   Yuan, Y. *et al.* Assessing the clinical utility of cancer genomic and proteomic data across
tumor types. *Nat. Biotechnol.* **32,** 644–652 (2014).

54.   Harrell, F. E., Lee, K. L. & Mark, D. B. in *Tutorials in Biostatistics, Statistical Methods in*
Clinical Studies **1,** 223–249 (2005).

55.   Berkson, J. In dispraise of the exact test. Do the marginal totals of the 2X2 table contain
relevant information respecting the table proportions? *J. Stat. Plan. Inference* **2,** 27–42
(1978).

56.   Rubio-Perez, C. *et al.* In Silico Prescription of Anticancer Drugs to Cohorts of 28 Tumor
Types Reveals Targeting Opportunities. *Cancer Cell* **27,** 382–396 (2015).

57.   Paatero, P. & Tapper, U. Positive Matrix Factorization - A Nonnegative Factor Model With
Optimal Utilization of Error-Estimates of Data Values. *Environmetrics* **5,** 111–126 (1994).

58.   Lee, D. D. & Seung, H. S. Algorithms for Non-negative Matrix Factorization. in *NIPS* 556–
562 (2000). doi:10.1109/ICASSP.2006.1661352

59.   Lee, D. D. & Seung, H. S. Learning the parts of objects by non-negative matrix
factorization. *Nature* **401,** 788–91 (1999).

60.   Dunn, J. C. & Dunnf, J. C. Well-Separated Clusters and Optimal Fuzzy Partitions. *J.*
Cybern. **4,** 95–104 (1974).

797