GENERAL INFORMATION

Title

Seasonal plasticity for anti-predatory strategies: matching colour and colour preference for effective

crypsis

Authors

Erik van Bergen[1*] and Patrícia Beldade[1,2]

Affiliations

[1] *Instituto Gulbenkian de Ciência, Oeiras, Portugal*

[2] *UMR5174 - CNRS, Evolution et Diversité Biologique, Université Paul Sabatier, Toulouse,*

France

Corresponding author

[*] Correspondence to Erik van Bergen (erikvanbergen.science@gmail.com)

ABSTRACT

Effective anti-predatory strategies typically require matching morphology and behaviour in prey and there are many compelling examples of behavioural strategies that enhance the effect of morphological defences. When protective adult morphologies are induced by a developmental environment predictive of future predation risk, adult behaviour should be adjusted accordingly to maximize predator avoidance. Behaviour is typically strongly affected by the adult environment but an effective match with a developmentally plastic morphology might be ensured when behaviour is affected the same pre-adult environmental cues. We study early and late life environmental effects on a suite of traits that are expected to work together for effective crypsis. We focused on body colour and colour preference in *Bicyclus anynana*, a model of developmental plasticity that relies on crypsis as a seasonal strategy for predator avoidance. Using a full-factorial design, we disentangle effects of developmental and adult thermal environments on both appearance and behaviour. We show that developmental conditions affect both adult colour and colour preference, with temperatures that simulate dry season conditions leading to browner butterflies with a perching preference for brown backgrounds. Our results reveal that coordinated environmentally-induced responses can increase effective seasonal anti-predatory strategies in species coping with environmental heterogeneity.

KEYWORDS

animal colouration – background matching – seasonal environments – phenotypic plasticity

1. INTRODUCTION

Predation is an important selective pressure and organisms have evolved a wide range of adaptations to improve chances of survival in the presence of predators. Such adaptations typically include a suite of traits working together to maximize predator avoidance. For example, spider-mimicking moths both look and act like their predators to improve their chances of survival [1], and butterflies bearing large vertebrate-eye-like eyespots generally only expose these when threatened to increase their chances of startling predators [2,3]. While defensive mechanisms are often constitutive properties of prey species (e.g. the spines of the hedgehog), they can also be induced by specific environmental conditions. For example, the protective morphology of *Daphnia* is induced by chemical cues that signal predator presence [4,5]. Plasticity in anti-predator strategies also occurs in response to abiotic factors such as those that affect the visual background under which species need to escape predators. For example, in many polyphenic butterflies, seasonal strategies to avoid predation are associated with seasonal fluctuations in vegetation cover [6]. While such environmental regulation of development, or developmental plasticity, often leads to changes in adult morphology that are irreversible, adult behaviour is generally more responsive to environmental conditions experienced during adulthood. The question of whether plastic responses of morphological and behavioural anti-predator traits are coordinated, to ensure that animals use a behavioural repertoire that matches their appearance, remains under explored.

The tropical butterfly *Bicyclus anynana* is well suited to test hypotheses about the integration of multiple environmentally-induced anti-predator traits. This species exhibits seasonal polyphenism as an adaptive response to dry-wet seasonal environments, with crypsis and deflection as seasonally alternating anti-predator strategies [6]. Dry season individuals with small inconspicuous wing pattern elements are camouflaged against the brown background of dry leaves, while wet season individuals have large conspicuous marginal eyespots that deflect predators' attention away from the more vulnerable body. This seasonal variation in wing patterns is induced by the temperature during development, with cooler temperatures leading to small pattern elements [7]. Mark-release-

recapture experiments in the field [8] and experimental work with different putative predators have

provided support for the adaptive advantage of seasonal variation in eyespot size [9,10].

Here we use thermal plasticity in *B. anynana* and a full factorial design to study the combined

effects of developmental and adult environments on anti-predator pigmentation and behaviour.

While thermal plasticity for *B. anynana* eyespot size is well characterized, other traits that could

enhance the crypsis of dry season individuals, such as actual wing colour and colour preference,

have largely been ignored (but see [6,9]). We investigated thermal plasticity in the colours of wing

patterns as well as in behaviours that are predicted to make crypsis more effective. These

behaviours are background matching, by which animals choose to rest on backgrounds that

resemble their appearance [11,12], and motionlessness, since camouflage is known to be less

effective when animals are in motion [13]. We test the hypotheses that the cryptic dry season forms,

that are produced when development occurs in cooler temperatures, will have not only smaller

pattern elements, but also colours that are less conspicuous. Moreover, we hypothesise that these

cryptic individuals will be less active and rest preferentially on brown backgrounds reminiscent of

the natural dry foliage. Globally, this work explores the extent to which seasonal strategies for

avoiding predation involve synchronized responses to environmental information, to prevent a

potentially fatal mismatch between anti-predatory morphology and behaviour.

2. METHODS

(a) Butterflies and rearing conditions

Eggs from a laboratory population of *B. anynana* were allocated to one of two climate-controlled

rooms (65% RH, 12L:12D) differing in ambient temperature to simulate the conditions of the dry

(Td=19°C) and wet seasons (Td=27°C). Groups of 250 larvae were reared in large population cages

and fed with young maize plants. Upon eclosion, males and females were separated and allocated to

one of two adult temperature regimes (Ta; 19°C and 27°C); yielding four treatments with different

combinations of Td and Ta (electronic supplementary material, figure S1). Adults were kept at

densities of 25 same-sex individuals per cage, and fed with moist banana.

(b) Wing pigmentation analysis

Using a colour-calibrated digital scanner (Epson V600), we scanned the ventral surfaces of

hindwings of 16 female and 16 male butterflies per treatment (N=128). The resulting images were

analysed with custom-made interactive Mathematica notebooks [14]. We first drew two contiguous

transects defined by five landmarks on the wing compartment bearing the eyespot that is often used

to characterize seasonal polyphenism in *Bicyclus* and related genera [15]. We then specified the

limits of the central band and each of the eyespot colour rings along the transect. The colours of the

wing background and different pattern elements were quantified by extracting the mean RGB value

of 3×3 pixel squares centred on the transect (figure 1*a*). For each individual, we then calculated the

Euclidean distance in RGB colour space between the colour of the different pattern elements and

that of the background of the wing. These distances were used as a proxy for colour

conspicuousness. We also calculated the distance from each pixel along the transect to the brown

patches used in our behavioural assay and used the mean of these as a proxy for crypsis in our

experimental setup (electronic supplementary material, figure S2).

(c) Behavioural assays

The behavioural assays were performed in large flight cages (150×50×54 cm) that allow expression

of a more natural behaviour [16]. The bottoms, backs, and sides of the cages were covered with

canvas depicting a mosaic of green (Hue=130°, Saturation=60%, Brightness=50%) and brown

(HSB 25°, 60%, 50%) squares (16.7 cm^2). The fronts were covered with mesh netting and the tops

had three fluorescent lamps (Osram L58W/840). Digital cameras (Canon EOD 1000D), connected

to a Wireless Remote Control (Hähnel - Giga T Pro), were positioned in front of the cage. Cohorts

of 20 same-sex, 6-8 day old individuals were released into the flight cage and left to habituate for

two hours. Subsequently, about 8 hours after the lights were switched on, 15 time-lapse photos were

taken at four-minute intervals for a period of 60 minutes. For each photo, we registered the number

of butterflies at rest on each of the two background colours, as well as the number of individuals

resting at exactly the same position as in the previous time point (i.e. no re-location). For perching

preference we used the number of new arrivals to brown patches and the number of new arrivals to

green patches, for each time point, as a 2-vector dependent variable. Likewise, for activity levels,

we used the number of inactive individuals and the number relocated individuals. Sixteen cohorts

were tested for each experimental treatment, eight for each of the sexes ($N_{cohort} = 64$).

(d) Statistical analyses

All statistical tests were done using R [17]. We examined the effects of developmental temperature

(Td), adult temperature (Ta), and sex on wing pigmentation using linear mixed models (electronic

supplementary material, table S1). The effects of Td, Ta, sex on perching preference and on levels

of activity were assessed using generalized linear mixed models with a binomial distribution and a

logit link. The cohort, and the time point (i.e. consecutive photos) for the behavioural data, were

included as random effects in the models. For both behavioural variables, minimum adequate

models were obtained by backward elimination, starting from the full model and using the Bayesian

Information Criterion (BIC). P-values were calculated via likelihood ratio tests and by parametric

bootstrapping (nsim=999), using the package *afex* [18]. Post-hoc pairwise comparisons

(alpha=0.05) were conducted using the package *lsmeans* (electronic supplementary material, tables

S2 and S3).

3. RESULTS

We used a full factorial design, with two developmental temperatures (Td; 19°C and 27°C) and two

adult temperatures (Ta; 19°C and 27°C) that simulate the natural dry and wet seasons, respectively,

to study thermal plasticity in pigmentation and behaviour and partition environmentally-induced

variation into early and late life effects.

(a) Thermal plasticity in wing colours

Using a custom image analysis tool, we quantified the colour of different wing pattern components

(central band and eyespot rings; figure 1*a*) and of the wing background of adults from each of the

four combinations of Td and Ta. We found that developmental temperature affected many of these

aspects of wing pigmentation in both males and females. Pattern elements of individuals reared at

cooler temperatures were not only smaller, as had been shown previously [15,19], but they were

also of less conspicuous colours. Specifically, the colours of the eyespot's external ring (Td;

$F=278.38$, $P<0.001$) and central focus (Td; $F=757.57$, $P<0.001$) and, in females, the central band

(Td; $F=47.65$, $P<0.001$) were closer to the background colour of their respective wings (figure

1*b,c*). Moreover, overall colouration of individuals reared at 19°C was also closer to the brown

background patches used in our behavioural assays (figure 1*d*), with the colours of males being

darker and closer to the brown patches (Sex; $F=171.40$, $P<0.001$). Adult temperature had, as

expected, no effect on wing pigmentation.

(b) Thermal plasticity in colour preference

The behaviour of males and females from the four temperature treatments was monitored in large

flying cages containing a mosaic of green and brown patches. Using time-lapse photographs, we

monitored preference to rest on brown versus green, as well as overall levels of activity (electronic

supplementary material, figure S3). We found that developmental temperature affected adult colour

preference (figure 2*a*). Compared to their more conspicuous wet-season counterparts (Td=27°C),

individuals with more cryptic colouration (Td=19°C) were more likely to alight in brown patches

(Td; $\chi^2=20.66$, $P<0.001$). This was true for both sexes and in both adult temperatures. The

preference for brown backgrounds was stronger in females (Sex; $\chi^2=62.53$, $P<0.001$) and these sex-

specific colour preferences were stronger when animals were tested at low adult temperatures

(Ta:Sex; $\chi^2=14.26$, $P<0.001$). Activity levels of females were not affected by either Td or Ta while,

in contrast to our prediction, males were more active when reared (Td:Sex; $\chi^2=7.79$, $P=0.005$) or

kept (Ta:Sex; $\chi^2=22.91$, $P<0.001$) at cooler temperatures (figure 2*b*).

4. DISCUSSION

Using a full factorial design that allows separating the effects of developmental and adult environments [20], we showed that both appearance and behaviour are affected by thermal conditions during development in *B. anynana*. We demonstrated for the first time that developmental temperatures affect wing pigmentation beyond the size of the ornamental pattern elements. For better camouflage against the dry brown foliage, dry-season individuals not only develop smaller pattern elements but the colours of these contrast less with their wings' brown background colour. Moreover, we also showed that individuals reared at cooler temperatures (i.e. the browner ones) also had stronger preferences for brown backgrounds. Such a concerted response in adult colour and colour preference can help ensure that the development of a cryptic adult appearance is matched by a behavioural repertoire that enhances crypsis.

The integrated background colour preferences that we demonstrate here may reduce the risk of detection by predators when cryptic individuals are confronted with visually heterogeneous environments [21]. For B. a*nynana* this is especially relevant during seasonal transitions when dead brown leaves and green vegetation co-occur and butterflies can choose among a variety of backgrounds to rest on. We found that the preference for brown backgrounds was stronger in females. In fact, overall, males favoured green backgrounds which may be related to sex-specific trade-offs between survival and reproduction in the wild. Males of *B. anynana* pursue a perch-and-chase strategy to procure mates and are typically found in patches of green sunlit vegetation that improve the probability of detecting potential mates [16,22]. In contrast, females tend to fly more closely to the ground searching for suitable host plants for oviposition. While our data supports the hypothesis that cryptic individuals show stronger preference for the background colour that presumably maximizes camouflage, we found no evidence for responses in another type of behaviour that could enhance crypsis; motionlessness. Field observations had suggested that cryptic

adults spend the dry season being relatively inactive, likely to conserve energy and avoid detection

[6], but we found no evidence for reduced activity in dry-season-like individuals in this study.

We demonstrated that environmental conditions during early life can affect wing colouration and

lead to "fixed" changes in adult behaviour, as had been shown in insects and vertebrates (e.g.

[23,24]). The mechanisms underlying the environmentally-induced differences in colour and colour

preference remain unclear, though traits responding in an integrated manner to environmental cues

typically share underlying regulators. Traits making up the *B. anynana* seasonal syndrome,

including eyespot size [19], life-history traits [25] and courtship behaviour [23], are known to be

mediated by ecdysteroid hormone dynamics during the late-larval and early pupal stages. Moreover,

sex differences in ecdysteroid titres during development have been shown to influence patterns of

sex-specific pigmentation in dorsal eyespots [26], which are not analysed here nor involved in

predator avoidance [27,28]. Whether and how ecdysone dynamics can have a direct effect on the

biosynthetic pathways for pigment production [29–31] is unclear, but effects of developmental

temperatures on these pathways have been studied in detail in *Drosophila melanogaster* [32], a

well-described model for thermal plasticity in body pigmentation.

Temperature-induced changes in ecdysteroid signalling may also alter neuronal activity and

connectivity [33,34] and/or mediate visual sensitivity of adults through the regulation of eye

development [35]. In fact, the size of the *B. anynana* compound eye is known to be

developmentally plastic and sexually dimorphic [36]. Wet-season individuals and males have larger

eyes than dry-season individuals and females, respectively, which could affect colour vision and

preferences. These published data on thermal plasticity in eye size [36] correlate strongly with our

behavioural data; individuals reared at lower temperatures have smaller eyes and stronger

preferences for brown patches (electronic supplementary material, figure S4).

Our data clearly show that morphological and behavioural anti-predator traits are affected in concert

by the conditions experienced during development and underscore the tight integration of form and

function necessary for survival under heterogeneous environments.

Data accessibility

Raw data will be archived in the Dryad Digital Repository upon acceptance of the manuscript.

Authors' contributions

EvB and PB conceived and designed the experiments. EvB performed the experiments and analysed

the data. EvB and PB wrote the manuscript.

Competing interests

The authors declare no competing financial interest.

Funding

Financial support was provided by the Portuguese science funding agency, Fundação para a Ciência

e Tecnologia, FCT (PTDC/BIA-EVF/0017/2014); French research funding agency, Agence

Nationale de la Recherche, ANR (Laboratory of Excellence TULIP, ANR-10-LABX-41) and

French research centre, Centre National de la Recherche Scientifique, CNRS (International

Associated Laboratory, LIA BEEG-B).

Acknowledgements

We are grateful to Carolina Peralta for technical assistance and Filipa Alves for developing a set of

interactive Mathematica notebooks we based our scripts on.

237   REFERENCES

1. Wang M-Y, Vasas V, Chittka L, Yen S-H. 2017 Sheep in wolf's clothing: multicomponent traits enhance the success of mimicry in spider-mimicking moths. *Anim. Behav.* **127**, 219–224. (doi:10.1016/j.anbehav.2017.03.020)

2. Olofsson M, Eriksson S, Jakobsson S, Wiklund C. 2012 Deimatic Display in the European Swallowtail Butterfly as a Secondary Defence against Attacks from Great Tits. *PLOS ONE* **7**, e47092. (doi:10.1371/journal.pone.0047092)

3. De Bona S, Valkonen JK, López-Sepulcre A, Mappes J. 2015 Predator mimicry, not conspicuousness, explains the efficacy of butterfly eyespots. *Proc. R. Soc. B Biol. Sci.* **282**. (doi:10.1098/rspb.2015.0202)

4. Grant JWG, Bayly IAE. 1981 Predator induction of crests in morphs of the *Daphnia carinata* King complex. *Limnol. Oceanogr.* **26**, 201–218. (doi:10.4319/lo.1981.26.2.0201)

5. Weiss LC, Leimann J, Tollrian R. 2015 Predator-induced defences in *Daphnia longicephala*: location of kairomone receptors and timeline of sensitive phases to trait formation. *J. Exp. Biol.* **218**, 2918. (doi:10.1242/jeb.124552)

6. Brakefield PM, Reitsma N. 1991 Phenotypic plasticity, seasonal climate and the population biology of *Bicyclus* butterflies (Satyridae) in Malawi. *Ecol. Entomol.* **16**, 291–303. (doi:10.1111/j.1365-2311.1991.tb00220.x)

7. Kooi RE, Brakefield PM. 1999 The critical period for wing pattern induction in the polyphenic tropical butterfly *Bicyclus anynana* (Satyrinae). *J. Insect Physiol.* **45**, 201–212. (doi:10.1016/S0022-1910(98)00093-6)

8. Brakefield PM, Frankino WA. 2009 Polyphenisms in Lepidoptera: multidisciplinary approaches to studies of evolution and development. In *Phenotypic plasticity of insects: mechanisms and consequences*, pp. 337–368. Enfield, USA: Science Publishers, Inc.

9. Lyytinen A, Brakefield PM, Lindström L, Mappes J. 2004 Does predation maintain eyespot plasticity in *Bicyclus anynana*? *Proc. R. Soc. B Biol. Sci.* **271**, 279–283.

10. Prudic KL, Stoehr AM, Wasik BR, Monteiro A. 2015 Eyespots deflect predator attack increasing fitness and promoting the evolution of phenotypic plasticity. *Proc. R. Soc. B Biol. Sci.* **282**. (doi:10.1098/rspb.2014.1531)

11. Michalis C, Scott-Samuel NE, Gibson DP, Cuthill IC. 2017 Optimal background matching camouflage. *Proc. R. Soc. B Biol. Sci.* **284**. (doi:10.1098/rspb.2017.0709)

12. Endler JA. 1984 Progressive background in moths, and a quantitative measure of crypsis. *Biol. J. Linn. Soc.* **22**, 187–231. (doi:10.1111/j.1095-8312.1984.tb01677.x)

13. Ioannou CC, Krause J. 2009 Interactions between background matching and motion during visual detection can explain why cryptic animals keep still. *Biol. Lett.* **5**, 191. (doi:10.1098/rsbl.2008.0758)

14. Rodrigues YK, van Bergen E, Alves F, Duneau D, Beldade P. 2017 Complex effects of day and night temperature fluctuations on thermally plastic traits in an experimental model of adaptive seasonal plasticity. *bioRxiv* (doi:10.1101/207258)

15. van Bergen E, Osbaldeston D, Kodandaramaiah U, Brattström O, Aduse-Poku K, Brakefield PM. 2017 Conserved patterns of integrated developmental plasticity in a group of polyphenic tropical butterflies. *BMC Evol. Biol.* **17**, 59. (doi:10.1186/s12862-017-0907-1)

16. van Bergen E, Brakefield PM, Heuskin S, Zwaan BJ, Nieberding CM. 2013 The scent of inbreeding: a male sex pheromone betrays inbred males. *Proc. R. Soc. B Biol. Sci.* **280**. (doi:10.1098/rspb.2013.0102)

17. R Core Team. 2017 *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing. See www.R-project.org.

18. Singmann H, Bolker B, Westfall J, Aust F. 2018 *afex: Analysis of Factorial Experiments.* See https://CRAN.R-project.org/package=afex.

19. Mateus ARA, Marques-Pita M, Oostra V, Lafuente E, Brakefield PM, Zwaan BJ, Beldade P. 2014 Adaptive developmental plasticity: Compartmentalized responses to environmental cues and to corresponding internal signals provide phenotypic flexibility. *BMC Biol.* **12**, 97. (doi:10.1186/s12915-014-0097-x)

20. Groothuis TGG, Taborsky B. 2015 Introducing biological realism into the study of developmental plasticity in behaviour. *Front. Zool.* **12**, S6. (doi:10.1186/1742-9994-12-S1-S6)

21. Merilaita S, Lyytinen A, Mappes J. 2001 Selection for cryptic coloration in a visually heterogeneous habitat. *Proc. R. Soc. Lond. B Biol. Sci.* **268**, 1925. (doi:10.1098/rspb.2001.1747)

22. Vande Velde L, Turlure C, van Dyck H. 2011 Body temperature and territory selection by males of the speckled wood butterfly (*Pararge aegeria*): what makes a forest sunlit patch a rendezvous site? *Ecol. Entomol.* **36**, 161–169. (doi:10.1111/j.1365-2311.2010.01257.x)

23. Bear A, Prudic KL, Monteiro A. 2017 Steroid hormone signaling during development has a latent effect on adult male sexual behavior in the butterfly *Bicyclus anynana*. *PLOS ONE* **12**, e0174403. (doi:10.1371/journal.pone.0174403)

24. Fischer S, Bohn L, Oberhummer E, Nyman C, Taborsky B. 2017 Divergence of developmental trajectories is triggered interactively by early social and ecological experience in a cooperative breeder. *Proc. Natl. Acad. Sci.* **114**, E9300–E9307. (doi:10.1073/pnas.1705934114)

25. Oostra V, Mateus ARA, van der Burg KRL, Piessens T, van Eijk M, Brakefield PM, Beldade P, Zwaan BJ. 2014 Ecdysteroid Hormones Link the Juvenile Environment to Alternative Adult Life Histories in a Seasonal Insect. *Am. Nat.* **184**, E79–E92. (doi:10.1086/677260)

26. Bhardwaj S, Prudic KL, Bear A, Dasgupta M, Wasik BR, Tong X, Cheong WF, Wenk MR, Monteiro A. 2018 Sex Differences in 20-Hydroxyecdysone Hormone Levels Control Sexual Dimorphism in *Bicyclus anynana* Wing Patterns. *Mol. Biol. Evol.* **35**, 465–472. (doi:10.1093/molbev/msx301)

27. Breuker CJ, Brakefield PM. 2002 Female Choice Depends on Size but Not Symmetry of Dorsal Eyespots in the Butterfly *Bicyclus anynana*. *Proc. Biol. Sci.* **269**, 1233–1239.

28. Robertson KA, Monteiro A. 2005 Female *Bicyclus anynana* butterflies choose males on the basis of their dorsal UV-reflective eyespot pupils. *Proc. R. Soc. B Biol. Sci.* **272**, 1541–1546. (doi:10.1098/rspb.2005.3142)

29. Zhang L, Martin A, Perry MW, van der Burg KRL, Matsuoka Y, Monteiro A, Reed RD. 2017 Genetic Basis of Melanin Pigmentation in Butterfly Wings. *Genetics* **205**, 1537. (doi:10.1534/genetics.116.196451)

30. Beldade P, Peralta CM. 2017 Developmental and evolutionary mechanisms shaping butterfly eyespots. *Curr. Opin. Insect Sci.* **19**, 22–29. (doi:10.1016/j.cois.2016.10.006)

31. Van Belleghem SM *et al.* 2017 Complex modular architecture around a simple toolkit of wing pattern genes. *Nat. Ecol. Amp Evol.* **1**, 0052.

32. Gibert J-M, Mouchel-Vielh E, Peronnet F. 2017 Modulation of yellow expression contributes to thermal plasticity of female abdominal pigmentation in *Drosophila melanogaster*. *Sci. Rep.* **7**, 43370.

33. Ishimoto H, Wang Z, Rao Y, Wu C-F, Kitamoto T. 2013 A Novel Role for Ecdysone in *Drosophila* Conditioned Behavior: Linking GPCR-Mediated Non-canonical Steroid Action to cAMP Signaling in the Adult Brain. *PLOS Genet.* **9**, e1003843. (doi:10.1371/journal.pgen.1003843)

34. Carvalho MJA, Mirth CK. 2015 Coordinating morphology with behavior during development: an integrative approach from a fly perspective. *Front. Ecol. Evol.* **3**, 5. (doi:10.3389/fevo.2015.00005)

35. Champlin DT, Truman JW. 1998 Ecdysteroids govern two phases of eye development during metamorphosis of the moth, *Manduca sexta*. *Development* **125**, 2009.

36. Everett A, Tong X, Briscoe AD, Monteiro A. 2012 Phenotypic plasticity in opsin expression in a butterfly compound eye complements sex role reversal. *BMC Evol. Biol.* **12**, 232. (doi:10.1186/1471-2148-12-232)

 FIGURE LEGENDS

 Figure 1 Environmentally-induced wing pigmentation. (*a*) Examples of ventral hindwing

 surfaces from females reared at Td=19°C (left) and Td=27°C (right; with section highlighted).

 Section of a hindwing showing landmarks (white circles) that defined two contiguous transects

 (white line) passing through the centre of the eyespot. The colours of the background, central band,

 and eyespot rings were quantified by extracting the mean RGB values of 3×3 pixel squares. (*b*)

 Schematic representation of the colour of the background (square fill), central band (stripe on top

 left corner) and eyespot rings (circles), as well as the size of the wings and eyespot rings (cf. scale

 on left-most drawing), for individuals reared at Td=19°C (left) and Td=27°C (right). (*c*) Euclidian

 distance in RGB colour space between the colour of the background and the colour of different

 pattern elements for females are males from four thermal treatments. Colour conspicuousness was

 not affected by adult temperature (Ta), statistical significance for effect of developmental

 temperature (Td) is indicated as: * for $P<0.05$ and *** for $P<0.001$. (*d*) Mean Euclidian distance in

 RGB colour space, using all pixels along the transect, to the colour of the brown patches used in our

 behavioural assay. Significant differences between groups (Tukey's HSD, $P<0.05$) are indicated by

 different letters. Details of statistical tests in electronic supplementary material, table S1.

Figure 2 Environmentally-induced behaviours. (*a*) Probability of relocating to a brown patch

for females and males from different combinations of Td and Ta. Within the same testing

temperature and sex, cryptic individuals (Td=19°C; blue dots) are more likely to alight in brown

patches than individuals with a more conspicuous colouration (Td=27°C; red dots). Preference for

brown patches is stronger in females and sex-specific colour preferences are stronger when tested at

cooler temperatures (Ta=19°C). Asterisks indicate groups having probabilities of relocating to

brown that are significantly different from a 1:1 ratio. (*b*) Probability of relocating for females and

males from different treatments. Males were more active when reared or kept at cooler temperatures

while activity levels of females were not affected by either Td or Ta. Dots in both panels represent

the probability of relocating (to brown) for each cohort based on the total number of relocations (to

brown) observed throughout the assay. Significant differences between groups (Tukey's HSD,

P<0.05) are indicated by different letters. Details of statistical tests in electronic supplementary

material, table S2 and S3.

SUPPLEMENTARY MATERIALS

* Supplementary Table 4 and 5 will be uploaded to Dryad upon acceptance for publication.

Supplementary Figure 1. Schematic representation of the experimental design. Panels on the left-hand side depict the natural environment of *B. anynana* during the wet season (upper) and dry season (lower) while panels on the right-hand side provide a schematic representation of our experimental design. Individuals were reared, from egg to adult eclosion, at one of two developmental temperatures (Td; 19°C and 27°C) that simulate the ambient temperatures of the dry and wet season, respectively. Variation in rearing temperature is sufficient to induce seasonal polymorphism in this species, with cool and warm temperatures inducing dry season form (DS) and wet season form (WS) individuals, respectively. On the day of eclosion, the butterflies were allocated to one of two adult temperatures (Ta; 19°C and 27°C), yielding four experimental treatments with different combinations of Td and Ta. Behavioural assays were performed at the temperature of the adult regime using cohorts of 20 six to eight day old males or females ($N_{cohort} = 64$). The red colours in this scheme represent warm temperatures or individuals reared at warm temperatures. Cool temperatures, or individuals reared at cool temperatures, are represented by blue colours.

Supplementary Figure 2. Quantification of crypsis-enhancing pigmentation traits. A. We used a set of custom-made interactive Mathematica notebooks to analyse the ventral hindwing surfaces of 16 female and 16 male butterflies per experimental treatment (i.e. two individuals per cohort, N=128). Two contiguous transects were defined by five landmarks (red dots) in the wing compartment bearing the large fifth eyespot (Cu1). Secondary landmarks, represented by grey lines, were used to specify the edge of the central band and the limits of each of the colour rings along the transect line. **B.** The colours of the eyespot rings were quantified by extracting the mean RGB value of 3×3 pixel squares centred on the intersection between the midpoints of the secondary landmarks and the transect line. The mean RGB values of the background and central band were extracted from a 3×3 cluster located 10 pixels after the associated secondary landmarks on the marginal side. Moreover, for each pixels we calculated the Euclidean distance in RGB colour space to the colour of the brown patches used in our behavioural assays. These distances are presented on the y-axis of the graph. **C.** We determined the Euclidean distance in RGB colour space between the colour of the each pattern element and the colour of the wing's background. The 3D-plots represent the mean RGB values for the each pattern element for females reared at cool (Td=19°C; left) and warm (Td=27°C; right) temperatures and lines represent the Euclidean distances for the band, golden ring, black disc and the white focus. **D.** Detailed images of the wing scales that make up the white focus for females (left) and males (right). The blue coloured outline represents individuals reared at cool temperatures (Td=19°C) while red outlines represents individuals reared at warm temperatures (Td=27°C). Scale bars represent 200 μm.

Supplementary Figure 3. Behavioural data per cohort. Panels represent all behavioural data per cohort. The number of butterflies at rest on each of the two background colours is represented by the brown and green bars for each time point. The number of individuals resting at exactly the same position as in the previous time point (i.e. no re-location) is depicted by lighter colours inside each bar. Note that individuals that were resting on the mesh netting that covered the front of the cage or on the upper surface of the cage, or were airborne when the time-lapse photos were taken were denoted as 'unaccounted for' in the data. The number of individuals that was unaccounted for was higher at cooler temperatures (Ta=19°C) for both males and females.

Supplementary Figure 4. Correlations between eye size and perching preference. The size of the *B. anynana* compound eye is known to be developmentally plastic and sexually dimorphic. Individuals reared at warm temperatures (wet-season; red dots) have larger eyes than individuals reared at cold temperatures (dry-season; blue dots) and females (circles) have smaller eyes than males (squares); data from Everett *et al.* 2012. Our behavioural data on perching preferences (y-axis) correlates strongly with the published data on thermal plasticity in eye size (x-axis) such that individuals with smaller eyes have stronger preferences for brown patches. Left-hand panel gives the correlation between eye size and perching preferences measured at cool temperatures (Ta=19°C) while the right-hand panel shows the correlation with behavioural responses assessed at warm temperatures (Ta=27°C).

Supplementary Table 1. Colours and sizes of the wing pattern elements across treatments. Left-hand panels represent the Euclidian distance in RGB colour space from the colour of the brown patches used in the experiment to **A)** all pixels along the transect and **B)** the colour of background. Panels C to F represent the Euclidian distance from the background colour of each individual to **C)** the central band, **D)** the golden ring, **E)** the black disc and **F)** the white focus. Panels G to K are related to (relative) sizes and panels represent **G)** the length of the transect (i.e. proxy for wing size), **H)** the relative distance from the start of the transect to the proximal edge of the central band (i.e. inverse proxy for the width of the band, which is difficult to measure directly because of its indistinct distal edge), **I)** the relative width of the golden ring, **J)** the relative width of the black disc and **K)** the relative width of the white focus. Relative widths were calculated by dividing the absolute size by the length of the transect. Tables on the right-hand side of each panel give the effects of developmental temperature (Td), adult temperature (Ta), sex and their interactions on the phenotypic trait measured. Effects with P<0.01 are shown in bold The experimental cohort was included as a random factor and denominator degrees of freedom were determined with the Kenward-Roger method, using the package *afex*. Post-hoc pairwise comparisons (alpha=0.05) were conducted using the package *lsmeans*. Differences between experimental treatments and sexes are given in the graphs, with groups that are not significantly different from each other sharing a common letter.

A)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**706.86**	**1,56**	**<0.001**
T$_A$	1.67	1,56	0.201
Sex	**49.42**	**1,56**	**<0.001**
T$_D$:T$_A$	1.26	1,56	0.267
T$_D$:Sex	**8.48**	**1,56**	**0.005**
T$_A$:Sex	0.05	1,56	0.828
T$_D$:T$_A$:Sex	0.37	1,56	0.546

B)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	2.13	1,56	0.150
T$_A$	3.79	1,56	0.056
Sex	**171.40**	**1,56**	**<0.001**
T$_D$:T$_A$	0.60	1,56	0.441
T$_D$:Sex	3.49	1,56	0.067
T$_A$:Sex	0.01	1,56	0.914
T$_D$:T$_A$:Sex	0.03	1,56	0.861

Continuation of Supplementary Table 1.

C)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**47.65**	**1,56**	**<0.001**
T$_A$	0.09	1,56	0.765
Sex	4.062	1,56	0.049
T$_D$:T$_A$	0.417	1,56	0.521
T$_D$:Sex	0.453	1,56	0.503
T$_A$:Sex	0.173	1,56	0.678
T$_D$:T$_A$:Sex	0.330	1,56	0.568

D)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**278.38**	**1,56**	**<0.001**
T$_A$	0.47	1,56	0.494
Sex	**19.45**	**1,56**	**<0.001**
T$_D$:T$_A$	0.03	1,56	0.861
T$_D$:Sex	5.01	1,56	0.029
T$_A$:Sex	1.16	1,56	0.287
T$_D$:T$_A$:Sex	0.25	1,56	0.619

E)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	0.57	1,56	0.452
T$_A$	0.37	1,56	0.548
Sex	**59.72**	**1,56**	**<0.001**
T$_D$:T$_A$	0.03	1,56	0.860
T$_D$:Sex	**13.80**	**1,56**	**<0.001**
T$_A$:Sex	0.15	1,56	0.695
T$_D$:T$_A$:Sex	0.14	1,56	0.705

F)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**757.57**	**1,56**	**<0.001**
T$_A$	2.11	1,56	0.151
Sex	3.94	1,56	0.051
T$_D$:T$_A$	1.29	1,56	0.261
T$_D$:Sex	**9.13**	**1,56**	**0.004**
T$_A$:Sex	2.82	1,56	0.099
T$_D$:T$_A$:Sex	0.39	1,56	0.536

Continuation of Supplementary Table 1.

G)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**74.60**	**1,56**	**<0.001**
T$_A$	1.89	1,56	0.174
Sex	**662.76**	**1,56**	**<0.001**
T$_D$:T$_A$	1.97	1,56	0.166
T$_D$:Sex	0.83	1,56	0.366
T$_A$:Sex	0.11	1,56	0.740
T$_D$:T$_A$:Sex	0.12	1,56	0.731

H)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**142.75**	**1,56**	**<0.001**
T$_A$	1.58	1,56	0.214
Sex	**15.87**	**1,56**	**<0.001**
T$_D$:T$_A$	4.09	1,56	0.048
T$_D$:Sex	<0.01	1,56	0.975
T$_A$:Sex	1.92	1,56	0.171
T$_D$:T$_A$:Sex	0.01	1,56	0.934

I)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**741.30**	**1,56**	**<0.001**
T$_A$	0.85	1,56	0.361
Sex	6.22	1,56	0.016
T$_D$:T$_A$	0.02	1,56	0.883
T$_D$:Sex	**10.88**	**1,56**	**0.002**
T$_A$:Sex	0.12	1,56	0.727
T$_D$:T$_A$:Sex	0.02	1,56	0.878

J)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**784.99**	**1,56**	**<0.001**
T$_A$	0.01	1,56	0.941
Sex	**12.55**	**1,56**	**<0.001**
T$_D$:T$_A$	1.10	1,56	0.299
T$_D$:Sex	1.30	1,56	0.258
T$_A$:Sex	0.04	1,56	0.837
T$_D$:T$_A$:Sex	0.25	1,56	0.618

Continuation of Supplementary Table 1.

K)

	F-value	df$_{KR}$	P$_{KR}$
T$_D$	**633.18**	**1,56**	**<0.001**
T$_A$	3.54	1,56	0.065
Sex	2.95	1,56	0.091
T$_D$:T$_A$	0.06	1,56	0.807
T$_D$:Sex	0.02	1,56	0.894
T$_A$:Sex	0.01	1,56	0.932
T$_D$:T$_A$:Sex	0.30	1,56	0.584

Supplementary Table 2: GLMM for perching preference (related to Fig 2a in the main text). **A.** Minimum adequate models were obtained by backward elimination, starting from the full model and using Bayesian Information Criterion (BIC). **B.** P-values for the minimum adequate model were calculated via likelihood ratio tests and parametric bootstrap (nsim=999), using the package *afex*. **C.** Mean probabilities, as well as upper and lower confidence limits, are given for each sex and experimental treatment. Post-hoc pairwise comparisons (alpha=0.05), obtained using the package *lsmeans*, specify differences between experimental treatments and sexes.

A. Perching preference: cbind(newBrown,newGreen)			BIC
M$_{full}$:	T$_D$ * T$_A$ * Sex	+ (1|Cohort) + (1|Timepoint)	2846.96
M$_2$:	T$_D$ + T$_A$ + Sex + T$_D$:T$_A$ + T$_D$:Sex + T$_A$:Sex	+ (1|Cohort) + (1|Timepoint)	2842.95
M$_3$:	T$_D$ + T$_A$ + Sex + T$_D$:Sex + T$_A$:Sex	+ (1|Cohort) + (1|Timepoint)	2836.78
M$_{mam}$:	**T$_D$ + T$_A$ + Sex + T$_A$:Sex**	**+ (1|Cohort) + (1|Timepoint)**	**2829.95**
M$_5$:	T$_D$ + T$_A$ + Sex	+ (1|Cohort) + (1|Timepoint)	2837.34

B.	Chisq	P$_{LRT}$	P$_{PB (n=999)}$
T$_D$	**20.66**	**<0.001**	**0.001**
T$_A$	1.32	0.251	0.245
Sex	**62.53**	**<0.001**	**0.001**
T$_A$:Sex	**14.26**	**<0.001**	**0.001**

C.	Sex	T$_A$	T$_D$	mean	LCL	UCL	pairs
	F	19	DS	0.624	0.575	0.671	E
	F	19	WS	0.566	0.518	0.613	CD
	F	27	DS	0.590	0.553	0.627	DE
	F	27	WS	0.530	0.493	0.566	C
	M	19	DS	0.448	0.408	0.490	B
	M	19	WS	0.389	0.349	0.431	A
	M	27	DS	0.516	0.479	0.552	C
	M	27	WS	0.455	0.419	0.492	B

Supplementary Table 3: GLMM for activity levels (related to Fig 2b in the main text). **A.** Minimum adequate models were obtained by backward elimination, starting from the full model and using Bayesian Information Criterion (BIC). **B.** P-values for the minimum adequate model were calculated via likelihood ratio tests and parametric bootstrap (nsim=999), using the package *afex*. **C.** Mean probabilities, as well as upper and lower confidence limits, are given for each sex and experimental treatment. Post-hoc pairwise comparisons (alpha=0.05), obtained using the package *lsmeans*, specify differences between experimental treatments and sexes.

A. Activity level: cbind(leaving,staying)		BIC
M_{full} :	T_D * T_A * Sex	+ (1|Cohort) + (1|Timepoint)	4362.39
M_2 :	T_D + T_A + Sex + T_D:T_A + T_D:Sex + T_A:Sex	+ (1|Cohort) + (1|Timepoint)	4360.09
M_{mam} :	**T_D + T_A + Sex + T_D:Sex + T_A:Sex**	**+ (1|Cohort) + (1|Timepoint)**	**4354.53**
M_4 :	T_D + T_A + Sex + T_A:Sex	+ (1|Cohort) + (1|Timepoint)	4355.46
M_5 :	T_D + T_A + Sex + T_D:Sex	+ (1|Cohort) + (1|Timepoint)	4370.57

B.	Chisq	P_{LRT}	$P_{PB\ (n=999)}$
T_D	**8.54**	**<0.001**	**0.006**
T_A	**13.14**	**<0.001**	**0.001**
Sex	0.25	0.620	0.661
T_D:Sex	**7.79**	**0.005**	**0.013**
T_A:Sex	**22.91**	**<0.001**	**0.001**

C.	Sex	T_A	T_D	mean	LCL	UCL	pairs
	F	19	DS	0.554	0.472	0.634	B
	F	19	WS	0.551	0.470	0.629	B
	F	27	DS	0.588	0.511	0.661	B
	F	27	WS	0.585	0.508	0.657	B
	M	19	DS	0.738	0.669	0.797	C
	M	19	WS	0.617	0.540	0.689	B
	M	27	DS	0.544	0.467	0.619	B
	M	27	WS	0.406	0.333	0.482	A