2,4-Diaminothieno[3,2-*d*]pyrimidines, a new class of anthelmintic with activity against adult and egg stages of whipworm

Short title: Diaminothienopyrimidines, a new chemotype for the control of whipworm

Author List

Frederick A. Partridge[1,¶], Ruth Forman[2,¶], Nicky J. Willis[3,¶, #a], Carole J.R. Bataille[3,¶], Emma A. Murphy[2, #a], Anwen E. Brown[1], Narinder Heyer-Chauhan[1], Bruno Marinič[3], Daniel J.C. Sowood[3], Graham M. Wynne[3], Kathryn J. Else[2*], Angela J. Russell[3,4*] and David B. Sattelle[1*]

[1] Centre for Respiratory Biology, UCL Respiratory, Division of Medicine, University College London, London, United Kingdom

[2] Faculty of Biology, Medicine and Health, University of Manchester, Manchester, United Kingdom

[3] Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Oxford, United Kingdom

[4] Department of Pharmacology, University of Oxford, Oxford, United Kingdom

[#a] Current address: Liverpool School of Tropical Medicine, Liverpool, United Kingdom

* Corresponding authors:

Email: d.sattelle@ucl.ac.uk (DBS), kathryn.else@manchester.ac.uk (KJE), angela.russell@chem.ox.ac.uk (AJR)

[¶] These authors contributed equally.

Abstract

The human whipworm *Trichuris trichiura* is a parasite that infects around 500 million people globally, with consequences including damage to physical growth and educational performance. Current drugs such as mebendazole have a notable lack of efficacy against whipworm, compared to other soil-transmitted helminths. Mass drug administration programs are therefore unlikely to achieve eradication and new treatments for trichuriasis are desperately needed. All current drug control strategies focus on post-infection eradication, targeting the parasite *in vivo*. Here we propose developing novel anthelmintics which target the egg stage of the parasite in the soil as an adjunct environmental strategy. As evidence in support of such an approach we describe the actions of a new class of anthelmintic compounds, the 2,4-diaminothieno[3,2-*d*]pyrimidines (DATPs). This compound class has found broad utility in medicinal chemistry, but has not previously been described as having anthelmintic activity. Importantly, these compounds show efficacy against not only the adult parasite, but also both the embryonated and unembryonated egg stages and thereby may enable a break in the parasite lifecycle.

Author Summary

The human whipworm, *Trichuris trichiura,* infects around 500 million people globally, impacting on their physical growth and educational performance. There are currently huge mass drug administration (MDA) programs aiming to control whipworm, along with the other major soil transmitted helminths, *Ascaris* and hookworm. However single doses of albendazole and mebendazole, which are used in MDA, have particularly poor effectiveness against whipworm, with cure rates less than 40%. This means that MDA may not be able to control and eliminate whipworm infection, and risks the spread of resistance to albendazole and mebendazole in the parasite population.

We are attempting to develop new treatments for parasitic worm infection, particularly focused on whipworm. Herein we report the identification of a class of compounds, diaminothienopyrimidines (DATPs), which have not previously been described as anthelmintics. These compounds are effective against adult stages of whipworm, and also block the development of the model nematode *C. elegans.*

Our DATP compounds reduce the ability of treated eggs to successfully establish infection in a mouse model of human whipworm. These results support a potential environmental spray to control whipworm by targeting the infectious egg stage in environmental hotspots.

Introduction

Current anthelmintics

The benzimidazole anthelmintics albendazole and mebendazole are typically used to treat human

whipworm infection but are compromised by lack of single-dose efficacy and the risk of resistance.

Thus, existing drugs lack sufficient efficacy in mass drug administration (MDA) programs to

adequately control or potentially eradicate whipworm. This is a major stumbling block in the WHO

target to eliminate morbidity from soil transmitted helminthiases in children by 2020. The current

approach for controlling soil-transmitted helminths such as *Trichuris* is mass drug administration of a

single-dose of albendazole or mebendazole, typically repeated annually [1]. However for infection

with *T. trichiura*, single doses of benzimidazoles lead to low cure rates, only 28% and 36% for

albendazole and mebendazole respectively[2]. These cure rates are much lower than those of other

major human soil-transmitted helminths, *Ascaris lumbricoides* and hookworm, demonstrating the

need for improvements to therapy specifically targeting *Trichuris*. Indeed modelling studies have

demonstrated that, due to these low cure rates, MDA with benzimidazoles does not interrupt

whipworm transmission and thus cannot achieve eradication in many settings [3].

Furthermore, the experience from studies on veterinary parasites is that widespread usage of

anthelmintics can lead to rapid development of resistance. The discovery of isolates of two species of

gastrointestinal nematodes resistant to monepantel only four years after its introduction [4] underlies

the real threat imposed by emerging drug resistance to control programmes. Indeed, the combination

of MDA programs and low single-dose cure rates may facilitate the development of drug resistance in

populations of human parasites. For example, resistance to benzimidazole drugs is caused by point

mutations in β-tubulin. Such resistance mutations have been found in *T. trichiura* after mass drug

administration [5], and have been found to increase in frequency after MDA. High frequency of

resistance mutations in a population may be associated with lower egg-reduction rates after MDA [6].

Whilst there is no clear evidence yet of widespread anthelmintic resistance in human populations,

identification of new drugs with novel mechanisms of actions is warranted to slow the development of

drug resistance.

Trichuris lifecycle

A *T. trichiura* infection becomes patent when adult female worms, embedded in the gut of the host,

start to lay eggs. A single female worm can lay up to 20,000 eggs per day and these unembryonated

eggs pass out with the faeces and embryonate in the soil. Development only proceeds further if the

embryonated eggs are accidentally consumed via contact of the next host with contaminated food,

water or soil. Once ingested, signals for hatching are received when the eggs reach the large intestine

[7,8], the newly emerged first stage larvae invade the mucosal epithelium and development to the

adult stage of the parasite occurs through a succession of larval moults. Importantly, even when active

infections are successfully treated, hosts are constantly re-infected due to high levels of infective eggs

present within the water and soil, which can remain viable for years.

Current anthelmintic programmes, including those targeting *Trichuris*, focus on post-infection

eradication of existing infections. However, lifecycle stages outside of the host are also potential

viable targets for small molecule drugs. Thus, both preventing egg embryonation and reducing the

infectivity of embryonated eggs prior to ingestion offer targets that would break the parasite lifecycle.

Screening *ex vivo T. muris* adults for new anthelmintic chemotypes

The mouse whipworm, *T. muris,* is a convenient model of the human whipworm as it can be grown

routinely in the laboratory via infection of severe combined immune deficiency (SCID) mice.

Screening *ex vivo* adult *T. muris* has been used to test the anthelmintic activity of a variety of

compounds, including approved drugs with the potential for repurposing, and also plant extracts [9–

11]. We recently reported a small molecule screen utilising an automated assay for assessment of the

motility of *ex vivo T. muris* adults. This screen led to the identification of a class of molecules termed

dihydrobenzoxazepinone (DHB) which demonstrated encouraging activity in this assay, as well as the

ability to reduce *in vivo* infectivity of treated eggs [12]. Most of the active molecules identified from

that screen belonged to the dihydrobenz[*e*][1,4]oxazepin-2(3*H*)-one chemotype, but interestingly one

additional active was from a completely different structural class. Here we report the identification,

synthesis and characterisation of this second chemotype, which has not previously been described as

having anthelmintic activity, the 2,4-diamino thieno[3,2-*d*]pyrimidines (henceforth called

diaminothienopyrimidines or DATPs).

Materials and methods

In vivo culture of *Trichuris muris*

T. muris worms were cultured using severe combined immune deficiency (SCID) mice, at the Biological Services Facility at the University of Manchester. Male and female mice were infected with 200 infective embryonated *T. muris* eggs via oral gavage. Thirty-five days later, the mice were sacrificed. Adult *T. muris* were obtained from the intestine as previously described [12]. Worms were maintained in Roswell Park Memorial Institute (RPMI) 1640 media supplemented with penicillin (500 U/mL) and streptomycin (500 µg/mL) at approximately 37 °C and studied on the same day. All animal experiments were approved by the University of Manchester Animal Welfare and Ethical Review Board and performed under the regulation of the Home Office Scientific Procedures Act (1986) and the Home Office project licence 70/8127.

Ex vivo T. muris adult maintenance for motility screen

Individual adult worms were added to wells containing 75 µL of RPMI-1640 medium, penicillin (500 U/mL), streptomycin (500 µg/mL) plus 1% v/v final concentration of dimethylsulfoxide (DMSO) or compound dissolved in DMSO. Plates were incubated at 37 °C, 5% CO_2. Motility was determined after 24 hours.

Automated motility assay

An automated system was used to quantify worm movement. An earlier version of this system has been previously described [13,14]. Two hundred frame movies of the whole plate were recorded at 10 frames per second and then motility determined by an algorithm based on thresholding pixel variance over time [15]. Dose-response curves were calculated with the four factor log-logistic model using the R package *drc* [16] or using GraphPad Prism.

Chemical synthesis

Thin layer chromatography (TLC) was performed on aluminium sheets coated with 60 F_{254} silica. All

solvents are used anhydrous unless stated otherwise. NMR spectra were recorded on Bruker AV400

(400 MHz), Bruker AVII 500 (500 MHz) or AVIIIHD 600 (600 MHz) instruments in the deuterated

solvent stated. All chemical shifts (δ) are quoted in ppm and coupling constants (*J*), which are not

averaged, in Hz. Residual signals from the solvents were used as an internal reference using the stated

deuterated solvent. Infrared spectra were recorded on a Perkin-Elmer 1750 IR Fourier Transform

spectrophotometer using thin films on a diamond ATR surface (thin film). Only the characteristic

peaks are quoted. Melting points were determined using a Stanford Research Systems EZ-Melt. Low

resolution mass spectra (*m/z*) were recorded on an Agilent 6120 spectrometer and high resolution

mass spectra (HRMS *m/z*) on a Bruker microTOF mass analyzer using electrospray ionization

(ESI).Compounds were synthesised from commercially available starting materials, and fully

characterised by Infrared (IR) Spectroscopy, Mass Spectrometry (ESI-MS, HRMS-ESI) and Nuclear

Magnetic Resonance (^{1}H and ^{13}C NMR).

***2-Chloro-N-(2-(chlorophenoxy)ethyl)thieno[3,2-d]pyrimidin-4-amine* 2a**

To a 20 mL microwave vial containing 2,4-dichlorothieno[3,2-*d*]pyrimidine (1.50 g, 7.32 mmol, 1.0

equiv.) in 1,4-dioxane (15 mL) at RT was added 2-(2-chlorophenoxy)ethylamine (1.26 g, 7.32 mmol,

1.0 equiv.) and *N,N*-diisopropylethylamine (2.5 mL, 14.64 mmol, 2.0 equiv.) under an argon

atmosphere. The vessel was sealed and the reaction heated at 80 °C for 3 h. The mixture was cooled to

RT, concentrated *in vacuo* and the crude residue was purified by flash column chromatography (silica

gel) to afford the title compound as an off-white solid (1.59 g, 64%).

mp = 118−119 °C; R_f = 0.2; v_{max} (film)/cm^{-1} = 3398m (NH), 3088w (CH), 2970w (CH), 1586s

(arom.), 1539m (arom.), 1483m (arom.); ^{1}H NMR (500 MHz, CDCl$_3$) δ 7.75 (1H, d, *J* = 5.4 Hz), 7.39

(1H, dd, *J* = 8.0, 1.6 Hz), 7.36 (1H, d, *J* = 5.4 Hz), 7.23 (1H, ddd, *J* = 8.3, 7.5, 1.6 Hz), 7.02 (1H, dd, *J*

= 8.4, 1.4 Hz), 6.96 (1H, apparent td, *J* = 7.9, 1.3 Hz), 5.79 (1H, t, *J* = 5.4 Hz), 4.30 (2H, t, *J* = 5.0 Hz)

4.13 (2H, apparent q, *J* = 5.4 Hz); ^{13}C NMR (500 MHz, CDCl$_3$) δ 161.4, 158.1, 157.6, 153.9, 132.6,

130.5, 128.1, 124.9, 123.5, 122.6, 114.7, 114.3, 68.1, 40.5; LRMS (ESI$^-$) calculated for

[C$_{14}$H$_{11}$ON$_3$^{35}Cl$_2$^{32}S-H]$^-$ = 338.0, found 337.9, [M-H]$^-$, 100%, calculated for [C$_{14}$H$_{11}$ON$_3$^{35}Cl^{37}Cl ^{32}S-

H]$^-$ = 340.0, found, 339.9 [M-H]$^-$, 60%; HRMS (ESI$^+$) calculated for [C$_{14}$H$_{11}$ON$_3$^{35}Cl$_2$^{32}S+H]$^+$ =

340.0073, found 340.0071, [M+H]$^+$.

2-Chloro-N-(2-phenoxyethyl)thieno[3,2-d]pyrimidin-4-amine (**2b**)

To a 20 mL microwave vial containing 2,4-dichlorothieno[3,2-*d*]pyrimidine (1.0 g, 5.0 mmol, 1.0

equiv.) in 1,4-dioxane (10 mL) at RT was added 2-phenoxyethylamine (0.6 mL, 5.0 mmol, 1.0 eq.)

and *N,N*-diisopropylethylamine (1.7 mL, 10.0 mmol, 2.0 equiv.) under an argon atmosphere. The

vessel was sealed and the reaction heated at 80 °C for 3 h. The mixture was cooled to RT,

concentrated *in vacuo* and the crude residue was purified by flash column chromatography (silica gel)

to afford the title compound as an off-white solid (1.23 g, 80%).

mp = 115.5–116.9 °C; R$_f$ = 0.5 (EtOAc: Petroleum; 1:4); ν$_{max}$ (film)/cm^{-1} = 3228w (NH), 3041w

(CH), 2962w (CH), 1597s (arom.), 1581s (arom.), 1533m (arom.), 1511m (arom.), (arom.), 1496m

(arom.), 1469m (arom.), 1434m (arom.); ^{1}H NMR (500 MHz, CDCl$_3$) δ 7.72 (1H, d, *J* = 5.4 Hz), 7.34

(1H, d, *J* = 5.4 Hz), 7.32-7.28 (2H, m), 6.97 (1H, app t, *J* = 7.3 Hz), 6.96 -6.92 (2H, m), 5.78 (1H, t, *J*

= 4.8 Hz), 4.22 (2H, t, *J* = 5.1 Hz), 4.11-4.06 (2H, m); ^{13}C NMR (126 MHz, CDCl$_3$) δ 161.4, 158.4,

158.1, 157.5, 132.6, 129.7, 124.8, 121.5, 114.6, 114.1, 66.2, 40.9; LRMS (ESI$^+$) calculated for

[C$_{14}$H$_{12}$ON$_3$^{35}Cl$_2$^{32}S+H]$^+$ = 306.0, found 306.0, [M+H]$^+$, 100%, calculated for

[C$_{14}$H$_{12}$ON$_3$^{35}Cl^{37}Cl^{32}S+H]$^+$ = 308.0, found 308.0, [M+H]$^+$, 40%, calculated for

[C$_{14}$H$_{12}$ON$_3$^{35}Cl$_2$^{32}S+Na]$^+$ = 328.0, found 328.0, [M+H]$^+$, 60%, calculated for

[C$_{14}$H$_{12}$ON$_3$^{35}Cl^{37}Cl^{32}S+Na]$^+$ = 330.0, found 330.0, [M+H]$^+$, 20%; HRMS (ESI$^+$) calculated for

[C$_{14}$H$_{12}$ON$_3$^{35}Cl$_2$^{32}S+H]$^+$ = 306.0462, found 306.0462, [M+H]$^+$.

General Synthetic Procedure

To a 10 mL microwave vial containing 2-chlorothieno[3,2-*d*]pyrimidine (1.0 equiv) in ^{i}PrOH (10

µL/mg chloride) at room temperature was added the requisite amine (10.0 equiv.) under an argon

atmosphere. The vessel was sealed and the mixture heated at 100°C for 16-24 h. The reaction was

cooled to ambient temperature (RT), concentrated *in vacuo* and the crude residue was purified by

flash column chromatography (silica gel).

***N2-Methyl-N4-(2-phenoxyethyl)thieno[3,2-d]pyrimidine-2,4-diamine* (3a, OX02925)**

Following general procedure 1, the title compound was obtained from **2b** (600 mg, 1.96 mmol, 1.0

equiv.) and methylamine (2.0 M in THF, 9.8 mL, 19.6 mmol, 10.0 eq). Purification by flash column

chromatography (MeOH:CH$_2$Cl$_2$; 1:49 v/v) followed by trituration with cold Et$_2$O afforded the desired

product as a pale yellow viscous oil (526 mg, 89%).

R_f = 0.2 (MeOH:CH$_2$Cl$_2$; 1:49 v/v); ν_{max} (film)/cm^{-1} = 3418w (NH), 3232w (NH), 3038w (CH), 2936w

(CH), 1585s (arom.), 1532s (arom.), 1508s (arom.), 1460s (arom.), 1405m (arom.); ^{1}H NMR (500

MHz, CDCl$_3$) δ 7.55 (1H, d, J = 5.4 Hz), 7.33-7.28 (2H, m), 7.15 (1H, d, J = 5.4 Hz), 7.00-6.96 (1H,

m), 6.96-6.93 (2H, m), 5.16 (1H, brs), 4.83 (1H, brs), 4.21 (2H, t, J = 5.26 Hz), 4.02 (2H, m), 3.04

(3H, d, J = 5.04 Hz); ^{13}C NMR (126 MHz, CDCl$_3$) δ 161.9, 161.8, 158.5, 157.4, 130.4, 129.6, 124.1,

121.2, 114.5, 106.5, 66.6, 40.2, 28.7; LRMS (ESI$^+$) calculated for [C$_{15}$H$_{16}$ON$_4$^{32}S+H]$^+$ = 301.1, found

301.1 [M+H]$^+$, 100%; HRMS (ESI$^+$) calculated for [C$_{15}$H$_{16}$ON$_4$C^{32}S+H]$^+$ = 301.1119, found 301.1118

[M+H]$^+$.

***N4-(2-(2-Chlorophenoxy)ethyl)-N2-(2-methoxyethyl)thieno[3,2-d]pyrimidine-2,4-diamine* (3b,**

OX02926)

Following general procedure 1, the title compound was obtained from **2a** (600 mg, 1.76 mmol, 1.0

equiv.) and 2-methoxyethylamine (1.5 mL, 17.6 mmol, 10.0 eq). Purification by flash column

chromatography (MeOH:CH$_2$Cl$_2$; 1:49 v/v) followed by trituration with cold Et$_2$O afforded the desired

product as an off-white solid (380 mg, 57%).

mp = 69–97 °C (Et$_2$O); R_f = 0.1 (MeOH:CH$_2$Cl$_2$; 1:49 v/v); ν_{max} (film)/cm^{-1} = 3424w (NH),

3304w (NH), 3076w (CH), 2949w (CH), 1606m (arom.), 1532s (arom.), 1476m (arom.), 1460m

(arom.), 1444m (arom.), 1412m (arom.); ^{1}H NMR (400 MHz, CDCl$_3$) δ 7.56 (1H, d, J = 5.3 Hz), 7.39

(1H, dd, J = 7.9, 1.4 Hz), 7.23-7.21 (1H, m), 7.13 (1H, d, J = 5.3 Hz), 7.00-6.92 (2H, m), 5.29 (1H,

s), 5.17 (1H, s) 4.27 (2H, t, J = 5.2 Hz), 4.05 (2H, apparent q, J = 5.4 Hz), 3.68-3.59 (4H, m), 3.40

(3H, s); ^{13}C NMR (151 MHz, CDCl$_3$) δ 161.8, 161.1, 157.3, 154.0, 130.5, 130.3, 127.8, 124.0, 123.3,

122.1, 114.3, 106.8, 71.6, 68.2, 58.7, 41.4, 39.9; LRMS (ESI$^+$) calculated for [C$_{17}$H$_{19}$O$_2$N$_4$^{35}Cl^{32}S+H]$^+$

= 379.1, found 379.1, [M+H]$^+$, 100%, calculated for [C$_{17}$H$_{19}$O$_2$N$_4$^{35}Cl^{32}S+Na]$^+$ = 401.1, found 401.1,

[M+Na]$^+$, 10%; HRMS (ESI$^+$) calculated for [C$_{17}$H$_{19}$O$_2$N$_4$^{35}Cl^{32}S+H]$^+$ = [M+H]$^+$, 379.0990, found

379.0991 [M+H]$^+$.

N4-(2-(2-chlorophenoxy)ethyl)-N2-(2-methoxybenzyl)thieno[3,2-d]pyrimidine-2,4-diamine

(3c, OX03143)

Following general procedure 1, the title compound was obtained from **2a** (240 mg, 0.70 mmol, 1.0

eq.) and 2-methoxybenzylamine (0.92 mL, 7.0 mmol, 10.0 eq.). Purification by flash column

chromatography (MeOH:CH$_2$Cl$_2$; 3:37 v/v) afforded the desired product (189 mg, 61%) as a thick pale

yellow oil.

R$_f$ = 0.4 (MeOH:CH$_2$Cl$_2$; 3:22 v/v); ν$_{max}$ (film)/cm^{-1}= 3424w (NH), 3247w (NH), 2935w (CH), 1587m

(arom.), 1553 (arom.), 1487 (arom.), 1461 (arom.); ^{1}H NMR (600 MHz, CDCl$_3$) δ 7.54 (1H, d, J = 5.3

Hz), 7.38 (1H, dd, J = 8.1, 1.7 Hz), 7.36 (1H, s) 7.22 (1H, ddd, J = 9.5, 7.5, 1.6 Hz), 7.20 (1H, ddd, J

= 8.2, 7.5, 1.6 Hz), 7.12 (1H, d, J = 5.3 Hz), 6.93 (1H, dt, J = 7.5, 1.5 Hz), 6.91 (1H, ddd, J = 8.1, 7.1,

1.1 Hz), 6.89 (2H, d, J = 7.9 Hz), 5.34 (1H, br), 5.29 (1H, t, J = 5.3 Hz), 4.68 (2H, d, J = 6.2 Hz) 4.20

(2H, t, J = 5.3 Hz), 4.04 (2H, apparent q, J = 5.3 Hz), 3.87 (3H, s); ^{13}C NMR (151 MHz, CDCl$_3$) δ

161.9, 161.2, 157.6, 157.4, 154.1, 130.5, 130.3, 129.1, 128.1 (x2), 127.8, 123.9, 123.3, 122.1, 120.3,

114.3, 110.1, 106.6, 68.2, 55.3, 41.4, 40.0; LRMS (ESI$^+$) calculated for [C$_{22}$H$_{21}$O$_2$N$_4$^{35}Cl^{32}S+H]$^+$ =

441.1, found 441.2, [M+H]$^+$, 100%; HRMS (ESI$^+$) calculated for [C$_{22}$H$_{21}$O$_2$N$_4$^{35}Cl^{32}S+H]$^+$ = 441.1147,

found 441.1142 [M+H]$^+$.

N4-(2-(2-chlorophenoxy)ethyl)-N2-methylthieno[3,2-d]pyrimidine-2,4-diamine **(3d, OX03147)**

Following general procedure 1, the title compound was obtained from **2a** (237 mg, 0.70 mmol) and

methylamine (2.0 M in THF) (3.5 mL, 7.0 mmol, 10 eq). Purification by flash column

chromatography (MeOH:CH_2Cl_2; 1:19 v/v) afforded the desired product (218 mg, 93%) as a pale

brown oil.

R_f = 0.4 (MeOH:CH_2Cl_2; 1:19 v/v); ν_{max} (film)/cm^{-1}= 3247w (NH), 2940w (CH), 1588m (arom.),

1552m (arom.), 1510m (arom.), 1484m (arom.), 1461m (arom.), 1446m (arom.), 1406 (arom.); ^{1}H

NMR (600 MHz, CDCl$_3$) δ 7.56 (1H, d, J = 5.3 Hz), 7.38 (1H, dd, J = 7.9, 1.7 Hz) 7.21 (1H, ddd, J =

8.2, 7.5, 1.7 Hz), 7.15 (1H, d, J = 5.3 Hz), 6.98 (1H, dd, J = 8.2, 1.5 Hz), 6.95 (1H, dd, J = 7.5, 1.5

Hz) 5.31 (1H, t, J = 5.5 Hz), 4.84 (1H, s) 4.27 (2H, t, J = 5.4 Hz), 4.06 (2H, apparent q, J = 5.5 Hz),

3.04 (3H, d, J = 5.0 Hz); ^{13}C NMR (151 MHz, CDCl$_3$) δ 162.0, 161.9, 157.5, 154.2, 130.7, 130.5,

128.0, 124.2, 123.6, 122.3, 114.6, 106.9, 68.5, 40.1, 28.9; LRMS (ESI$^+$) calculated for

$[C_{15}H_{15}ON_4{}^{35}Cl{}^{32}S+H]^+$ = 335.1, found 335.0, [M+H]+, 100%, calculated for $[C_{15}H_{15}ON_4{}^{37}Cl{}^{32}S+H]^+$

= 337.1, found 337.0, [M+H]+, 30%, HRMS (ESI$^+$) calculated for $[C_{15}H_{15}ON_4{}^{35}Cl{}^{32}S+H]^+$ =

335.0728, found 335.0725 [M+H]$^+$, calculated for $[C_{15}H_{15}ON_4{}^{37}Cl{}^{32}S+H]^+$ = 337.0698, found

337.0695 [M+H]$^+$.

C. elegans growth assay

A mixed-stage *C. elegans* N2 population was obtained by liquid culture (20 °C) according to standard

methods [17]. It was then bleached to obtain an egg population with 1.5 mL 4M NaOH, 2.4 mL

NaOCl, 2.1 mL water, washed three times, and allowed to hatch in 50 mL S-basal buffer at 20 °C

overnight to obtain a synchronised L1 population. For the growth assay, 49 µL of S-complete buffer

and 1 µL of DMSO or DMSO plus compound were added to each well of 96-well plates. 50 µL of a

worm suspension (approximately 20 synchronised L1 worms, 1% w/v *E. coli* HB101 in S-complete

buffer) were then added to each well. Plates were incubated at 20 °C before imaging 5 days later.

Worm movement was stimulated by inserting and removing a 96-well PCR plate into/from the wells

of the assay plate, and then whole plate 200 frame movies were recorded at 30 frames per second.

Growth was quantified as a correlate of movement using the same automated system described earlier.

Cytotoxicity testing

The mouse rectal epithelial cell line CMT-93 was used for these studies. The WST-8 and neutral red

cytotoxicity assays were performed as described [12]. Briefly, cells were cultured with test

compounds, chlorpromazine positive control or DMSO alone (final compound concentrations of 0 to

100 µM) for 72 hours. The WST-8 assay was then carried out using the Cell Counting Kit – 8 (Sigma

Aldrich # 96992) with an incubation time of 2 hours. Following this assay, the medium was

exchanged, and the ability of the cells to take up the dye neutral red (concentration 33 µg/mL,

incubation time 2 hours was determined using a microplate reader (absorbance at 540 nm). Results

were analysed using GraphPad Prism and fitted using a log-logistic model.

In vitro and *in vivo* establishment of infection

100 infective embryonated eggs were incubated in deionised water with 1% v/v DMSO or test

compounds at a final concentration of 100 µM in 1% v/v DMSO for 14 days at room temperature in

the dark. Eggs were then washed and resuspended in deionised water. For *in vitro* hatching assays 100

eggs were added to 1 mL of *E. coli* bacterial culture grown in LB broth overnight at 37 °C shaking at

200 rpm. Egg-bacterial cultures were incubated for 24 hours at 37 °C, 5% CO_2 and hatching

determined following blinding by visual examination under a dissecting microscope. For *in vivo*

hatching assays, 40 eggs were counted under a dissecting microscope and given to a SCID mouse in

200 µL water. At day 15 post-infection mice were culled and the number of worms present in the

caecae and colon enumerated in a blinded manner.

Statistical analysis of *in vivo* establishment of infection data

The experiment was conducted in two 'experimental batches'. For batch one there were 5 mice in

each of the DMSO and **OX02926** groups. For batch two there were 9 mice in each of the DMSO and

OX02926 groups. The raw data (number of worms that established infection in each mouse) are

shown separated by batch and treatment in the S1 Figure.

To analyse the data we used a two-way ANOVA (worm number ~ treatment * batch). This showed a

significant effect of treatment [$F_{(1,24)}$ = 8.520, P = 0.00752]. It also showed a significant effect of

batch [$F_{(1,24)}$ = 10.956, P = 0.00294]. There was no significant interaction between treatment and

batch [$F_{(1,24)}$ = 0.296, P = 0.59153]. The significant effect of batch reflected that in both DMSO-

and **OX02926**-treated groups, the number of worms that established infection was generally lower in

mouse batch 1 than in batch 2 (S1 Figure). Variation in control worm establishment is commonplace

in *Trichuris* infections due to natural variation in egg infectivity from a standardised egg number, but

was within expected ranges. We therefore took the approach of normalising each data point by

dividing by the mean of the DMSO-treated group for that batch. This gave the % batch normalised

infection establishment.

We used a two-way ANOVA (% batch normalised infection establishment ~ treatment * batch) to

analyse the data. There was a significant effect of treatment [$F_{(1,24)}$ = 9.569, P = 0.00497] but no

effect of batch [$F_{(1,24)}$ = 0.083, P = 0.77618] or interaction [$F_{(1,24)}$ = 0.083 0.77618]. We therefore

conducted a post-hoc Tukey HSD test which showed that infection establishment in the **OX02926**-

treated group was significantly different from the DMSO-treated control group (P = 0.0050).

Embryonation assay

One hundred unembryonated eggs were treated with a water, 1% v/v DMSO in water or test

compounds at a final concentration of 100 µM (unless stated) with a final concentration of 1% v/v

DMSO for 56 days in the dark at 26 °C. Images were collected on an Olympus BX63 upright

microscope using a 60x / 1.42 PlanApo N (Oil) objective and captured and white-balanced using

an DP80 camera (Olympus) in monochrome mode through CellSens Dimension v1.16 (Olympus).

Images were then processed and analysed using Fiji [18].

Data availability

Structures of resynthesized compounds have been deposited in the PubChem database with CID

49790760, 49790669, 46948320 and 49778268 and SID 348479445, 348479446, 348479447 and

348479448. Assay results for resynthesized compounds have been deposited in the PubChem database

with assay ID 1259352 and 1259353.

Results

Ex vivo T. muris adult motility screen

We have recently described a small molecule screen for new anthelmintics, which used reduction or loss of motility of adult *ex vivo T. muris* as an endpoint for screening [12]. This screen was designed to identify compounds active on *Trichuris* as existing drugs are notably less efficacious against this nematode, and it is comparatively evolutionary distant to nematodes typically screened in anthelmintic-discovery efforts, such as *H. contortus, M. incognita* and *C. elegans.* From this primary screen, we found 13 members of the dihydrobenzoxazepinone chemotype, which had not previously been shown to have anthelmintic activity.

In this report we describe the identification of a second new anthelmintic chemotype from this screen. A single 2,4-diaminothieno[3,2-*d*]pyrimidine (DATP) compound was found in the primary screen (Fig 1A). This has been given the identifier **OX02926**. We confirmed this activity in a secondary screen using the same source sample (DMSO solution containing 10 mM compound), and also tested a number of structurally-related compounds from our small molecule collection using the same assay (Fig 1C). This led to the identification of three further active molecules in this series **OX02925**, **OX03143** and **OX03147** (Fig 1D).

Fig 1. Identification of a diaminothienopyrimidine series from an *ex vivo T. muris* motility

screen. (a) Volcano plot of the primary screen conducted as in [12], with the single DATP hit

highlighted by a red asterix. Movement change and P-value compared to DMSO-only controls. *n*=4.

(b) Structure of the hit compound that was given the identifier **OX02926**. **(c)** Hit expansion by testing

of structurally-related compounds using library material. Significance was determined by a two-sided

Mann-Whitney test compared to DMSO-only controls, adjusted for multiple comparisons using the

Bonferroni method (for test compounds n=5, each replicate on different assay plates, each point

indicates one assay well). Blue bar indicates mean movement score. Compounds X (Pubchem CID:

49790326) and Y (Pubchem CID: 49795120) were not significantly active in this assay. **(d)** Structures

and identifiers of additional active compounds from this class.

Resynthesis of active compounds

Having identified promising active DATPs from testing of DMSO solution samples of compounds,

these were then resynthesised to obtain authentic, unambiguously characterised samples from which

confirmatory screening could take place. Compound resynthesis is important since DMSO solution

samples can degrade over time, and this often leads to so-called 'false positive' hits [19]. These

compounds could be readily prepared in two steps from commercially available 2,4-

dichlorothieno[3,2-*d*]pyrimidine **1**, *via* two sequential nucleophilic aromatic substitution reactions.

Treatment of **1** with 2-(2-chlorophenoxy)ethylamine or 2-phenoxyethylamine gave exclusively

monosubstitution affording **2a** and **2b** as a single regioisomer in 64% and 80% yield respectively.

Aubsequent displacement reaction at C4 gave authentic samples of **OX02925**, **OX02926**, **OX03143**

and **OX03147** in 57 – 91% yield (Fig 2).

2a R^1 = Cl, 64%
2b R^1 = H, 80%

3a (OX02925); R^1 = H; R^2 = NHMe, 89%
3b (OX02926); R^1 = Cl; R^2 = NHCH$_2$CH$_2$OMe, 57%
3c (OX03143); R^1 = Cl; R^2 = NHCH$_2$(2-OMePh), 61%
3d (OX03147); R^1 = Cl; R^2 = NHMe, 93%

Fig 2. Synthetic route to putative hit compounds (a) Substituted 2-phenoxyethan-1-amine

(1.0 equiv.), DIPEA (2.0 equiv.), 1,4-dioxane, 80 ^{o}C, 3 h. **(b)** Alkyl amine (10.0 equiv.), ^{i}PrOH,

100 ^{o}C, 16-24 h.

Activity of resynthesised diaminothienopyrimidines in the *T. muris ex vivo* adult motility assay

The resynthesized hits were then tested in this screen and a concentration-response curve constructed,

thereby confirming the anthelmintic activity of several examples of this structural class. (Fig 3,

Table 1).

Fig 3. Concentration-response curves for resynthesized DATPs in the *T. muris ex vivo* adult

motility assay. n=5 wells per concentration per compound, each replicate on a different 96-well plate,

conducted on two experimental sessions using worms from different mice. Blue line indicates

concentration-response curve fitted with the 4-factor log-logistic model using *drc* [16]. Figure in

parenthesis indicates EC_{50} estimate ± standard error from this model. No confidence limit could be

calculated for **OX03143.**

Chemical properties of the hit series and synthetic suitability for further development

This class has 'lead-like' or 'drug-like' chemical properties [20], although it is important to note that in the contemporary medicinal chemistry literature this term is usually applied in the context of imparting oral bioavailability characteristics (Table 1.). For agents targeting the gastrointestinal located *Trichuris,* minimal systemic exposure of the host is desirable and therefore it is critical to differentiate between the conventionally used terminology and parameters for 'drug-like' molecules, which affect solubility and permeability, and properties that would be relevant to agents targeting other body compartments. Recent literature has described this important caveat for non-peripheral CNS drugs [21], and indeed for anti-parasitic drug development [22]. Importantly, there is considerable scope for generating the large number of structural variants of the DATPs needed for the iterative improvement of compound properties during the downstream lead optimisation process.

Active diaminothienopyrimidines block *C. elegans* development

Although we are focused on developing an anthelmintic with improved efficacy over existing drugs against *Trichuris,* activity across the nematode phylum is valuable, particularly as efficacy against economically significant agricultural animal parasites would make further development more economically viable.

We therefore wanted to test the activity of the DATP chemotype against the clade V nematode *Caenorhabditis elegans.* Using a quantitative development assay to measure the growth of synchronised L1 stage worms, we tested varying concentrations of the compounds to determine the concentration-response effects. As shown in Fig 4, all four DATP compounds were active in this assay with EC_{50} values from 7 – 87 µM.

Fig 4. Concentration-response curves for resynthesized DATPs in the *C. elegans* growth assay.
n=5 wells per concentration per compound, each replicate on a different 96-well plate. Blue line
indicates concentration-response curve fitted with the 4-factor log-logistic model using *drc* [16].
Figure in parenthesis indicates EC_{50} estimate ± standard error from this model. No confidence limit
could be calculated for **OX03143**.

Interestingly, the DATPs display differing trends in activity between the *Trichuris* and *C. elegans*
assays. At this stage we do not know whether this reflects different potency at the target or different
patterns of drug access between the species, but the findings highlight the importance of screening
against *Trichuris* in the search for novel anthelmintic agents targeting whipworm. The data from each
of these assays as well as structural descriptors and Lipinski rule assessment for the four DATP
compounds and other anthelmintics are summarised in Table 1.

 Table 1. Properties and activities of resynthesized diaminothienopyrimidines, and other anthelmintics

Compound	PubChem CID	Structure	EC$_{50}$ (µM)		RMM	cLogP	HBA	HBD	tPSA (Å^2)	ROTB
			T. muris paralysis assay	*C. elegans* growth assay						
		(*'Drug-like' guidelines*)			*<500*	*<5*	*<10*	*<5*	*(<140)*	*(≤10)*
OX02925	49790760		28 ± 18	82 ± 9	300	2.5	5	2	87	6
OX02926	49790669		26 ± 4	82 ± 37	379	3.0	6	2	97	9
OX03143	46948320		70	7 ± 1	440	4.5	6	2	97	9
OX03147	49778268		44 ± 7	11 ± 2	344	3.1	5	2	87	6
Levamisole	26879		8 ± 3[a]	5 ± 1[a]	204	1.7	2	0	41	1
Albendazole	2082		> 800[b]	n.d.	249	2.3	6	2	76	5
Mebendazole	4030		>600[b]	1.1 ± 0.2[a]	295	2.7	6	2	84	4
OX02983	71447449		50 ± 13[c]	n.d.	344	2.7	4	0	42	3

RMM: relative molecule mass. HBA: number of hydrogen bond acceptors. HBD: number of hydrogen bond donors. tPSA: topological polar surface area,

calculated using DataWarrior [23]. ROTB: number of rotatable bonds. [a] Data from [15]. [b] Data from [24]. [c] Data from [12].

Assessment of the cytotoxicity of the diaminothienopyrimidine series

It was critical to ensure that this series showed minimal cytotoxicity towards mammalian cells, and

showed selective activity against the parasite. For example, gut cytotoxicity may result in the

compounds having too narrow a therapeutic window. Selected examples of the DATPs were assessed

for cytotoxicity using the mouse gut epithelial cell line CMT-93 (Table 2). Although, the DATPs

exhibited increased *in vitro* cytotoxicity in these assays compared to the previously reported DHB

series [12], an encouraging overall profile was exhibited for these early stage molecules. Furthermore,

the nematode cuticle often limits drug access which reduces target engagement by small drug-like

molecules [25,26]. This means that compound optimisation to improve uptake through the cuticle may

be a fruitful route to improved anti-nematode selectivity, as well as improving the cytotoxicity profile.

It is interesting to note that the activity against *Trichuris* did not correlate with cytotoxicity, with the

most cytotoxic compound (**OX03143**) showing the lowest activity in the *T. muris* adult paralysis

assay, by a factor of over 10-fold. This suggests that either anti-*Trichuris* activity is distinct from

cytotoxic action, or that differential drug access can be exploited to achieve differential host-parasite

activity. Either possibility is encouraging and suggests that continued exploration and iterative

improvement of the DATP structure might be anticipated to deliver a more potent anthelmintic with

acceptable host toxicity.

Compound	WST-8 EC_{50} (µM)	Neutral red EC_{50} (µM)	Adult *Trichuris* paralysis assay EC_{50} (µM)
OX02925	75 (48-124)	29 (19-43)	28
OX02926	43 (28-67)	21 (15-31)	26
OX03143	15 (9-26)	5 (3-7)	70
OX03147	37 (24-57)	21 (14-30)	44

Table 2. Summary of the cytotoxicity in a mouse epithelial cell line of the DATP series. Mouse

CMT-93 rectal epithelial cells were used for this assay. Maximum tested concentration was 100 µM.

$n=8$, error range (in parentheses) shows 95% confidence interval. EC$_{50}$ values in the adult *Trichuris*

paralysis assay are shown for comparison.

Activity of diaminothienopyrimidines against the infective egg stages of *T. muris*

Developing novel anthelmintics to disrupt the *T. trichiura* life cycle at the egg stage represents an

exciting and complementary strategy to an oral therapy and is particularly attractive as *T. trichiura*

eggs are highly resistant to extreme temperature changes and ultraviolet radiation, thereby remaining

viable in the environment for many years [27]. We assessed whether the DATP derivatives were

capable of affecting either infection establishment or embryonation of eggs. We first explored whether

the compounds could alter the establishment of infection by soaking embryonated *T. muris* eggs in the

test compounds for 14 days, washing the eggs and then determining infectivity both *in vitro* and *in*

vivo (Fig 5A).

To determine effects on *in vitro* hatching, a protocol modified from that previously described [8] was

established whereby eggs were induced to hatch when incubated in a culture of *Escherichia coli* at 37

°C. The results are summarised in Fig 5B. Strikingly all DATPs were capable of significantly reducing

in vitro hatching compared to the DMSO control.

Diaminothienopyrimidines reduce the ability of *T. muris* eggs to infect mice

To extend this finding, we selected   **OX02926** to test in an *in vivo* hatching and infection

establishment assay, as this compound showed both a significant decrease in *in vitro* hatching and a

small standard deviation between samples. The eggs were soaked as for the *in vitro* experiment and

SCID mice were infected with 40 treated eggs (**OX02926** or DMSO) by oral gavage. Egg infectivity

was quantified at day 15 post infection by culling the mice and counting the number of established

worms in the gut.

This experiment was carried out in two batches and the raw data are shown in the S1 Figure. Because

variation in control worm establishment is commonplace in *Trichuris* infections due to natural

variation in egg infectivity from a standardised egg number, we took the approach of normalising data

for each batch relative to the mean of the DMSO-only control group for that batch. This allowed us to

determine the effects of **OX02926** treatment (a full statistical description is given in the Methods

section). The results are shown in Fig 5C. We used a two-way ANOVA (% batch normalised infection

establishment ~ treatment * batch) to analyse the data. There was a significant effect of treatment

$[F_{(1,24)} = 9.569, P = 0.00497]$ but no effect of batch $[F_{(1,24)} = 0.083, P = 0.77618]$ or interaction

$[F_{(1,24)} = 0.083\ 0.77618]$. We therefore conducted a post-hoc Tukey HSD test which showed that

infection establishment in the **OX02926**-treated group was significantly different from the DMSO-

treated control group (P = 0.0050). **OX02926** was able to significantly reduce worm establishment *in*

vivo by an estimated 40%.

Fig 5. _T. muris_ eggs treated with diaminothienopyrimidines are less infective. (a) Embryonated

eggs were soaked in compound for 14 days, washed in water and then used in either in vitro or in vivo

hatching assays. **(b)** Treatment with DATPs reduced the ability of embryonated eggs to hatch in _E._

coli bacterial suspension after 24 hours. A one-way ANOVA showed a significant difference between

treatment groups (F(5,26)=25.95 p<0.0001) with a post-hoc Dunnett's compared to DMSO control

(****= p<0.0001) _n_=7 (DMSO), _n_=5 (DATP compounds) **(c)** SCID mice were infected with 40 eggs

and worm burden assessed at day 15 post infection. The experiment was carried out in two batches,

with n=5 and n=9 mice respectively in each of the control and treatment groups. Data were normalised

for each batch relative to the mean of the DMSO-only control group for that batch. Blue line indicates

mean for each treatment group. A two-way ANOVA showed a significant effect of treatment [F(1,24)

= 9.569, P = 0.00497] but no effect of batch [F(1,24) = 0.083, P = 0.77618] or interaction [F(1,24) =

0.083 0.77618]. A post-hoc Tukey HSD test showed that the **OX02926**-treated group was significantly

different from the DMSO control group (P = 0.0050).

Activity of diaminothienopyrimidines against the embryonation of _T. muris_ eggs

The ability of the DATPs to alter the embryonation of _T. muris_ eggs was investigated by soaking

unembryonated _T. muris_ eggs collected overnight from live adult _T. muris_ in the test compounds at 26

°C for the duration of the embryonation process (56-60 days). During embryonation the first larval

stage of the parasite develops within the egg shell (Fig 6A) from a ball of cells (Fig 6B). Treatment

with the DATPs **OX02925** and **OX03147** resulted in a significant increase in the percentage of

unembryonated eggs present compared to the DMSO control (Fig 6D). Importantly, although the other

DATPs did not alter the percentage of eggs unable to undergo the embryonation process, the larvae

that developed were atypical (Fig 6D-I). These atypical larvae were morphologically altered with the

granules present within the larvae appearing less distinct.

As **OX03147** had the clearest phenotype with a significant increase in the number of unembryonated

eggs, a concentration response study was performed to determine if an effect could be seen at lower

treatment doses. Additionally, we repeated the experiment at room temperature to allow for more

physiological conditions rather than the constant 26 °C utilised in the initial study to standardise

conditions across experiments. Although the increased number of unembryonated eggs was only

detected at the highest drug dose tested (100 µM) at both 26 °C and room temperature (Fig 6C, J)

striking effects on egg morphology was detectable at concentrations as low as 1 µM with significant

larvae stunting observed (Fig 6K).

To the known range of applications of DATPs in medicinal chemistry we can now add anthelmintic

activity. This study suggests they have significant potential for further development into dual-acting

therapeutic agents for both the reduction of *Trichuris* egg infectivity, and embryonation in the

environment. Thus, their actions on both the embryonated and unembryonated egg stages may enable

a break in the parasite lifecycle.

Fig 6. Unembryonated *T. muris* eggs treated with diaminothienopyrimidines have altered embryonation. Unembryonated eggs were soaked in 100 µM compound (unless specified otherwise) at 26 °C (unless specified otherwise) for the duration of the embryonation process (56-60 days) and

then embryonation determined and eggs imaged using an Olympus BX63 microscope. Scale bar

indicates 10 µm. **(a)** Typical embryonated egg and **(b)** unembryonated egg. **(c)** treatment with DATPs

increased the incidence of unembryonated eggs. Representative pictures of **(d)** DMSO, **(e) OX02925**,

(f) OX02926, **(g) OX03143** and **(h) OX03147** 100 µM and **(i) OX03147** 1 µM soaked *T. muris* eggs.

(j) Unembyronated eggs soaked in **OX03147** at room temperature for 56 days and embryonation

determined. **(k)** Unembyronated eggs soaked in **OX03147** at 26 °C for 56 days and larval length

calculated using ImageJ.

Discussion

Gastrointestinal nematode parasites remain a significant human health burden. Current anthelmintics lack efficacy and achieve low cure rates, threatening the targets set by the World Health Organisation for control of soil-transmitted helminths [2,28]. In particular, existing drugs have notably low efficacy against *T. trichiura,* the human whipworm. *T. trichiura* may be especially difficult to target as it inhabits the large intestine and is in part intracellular [29]. Thus, the metabolically active head end, the stichosome, is buried in the host epithelial cells lining the gut, affording some protection from orally delivered anthelmintics.

Diaminothienopyrimidines (DATPs), a new anthelmintic chemotype

We recently reported a small molecule screen for new anthelmintics targeting the gastro-intestinal (GI) nematode parasite *Trichuris muris* that identified the dihydrobenzoxazepinone (DHB) chemotype. The DHBs had not previously been ascribed anthelmintic activity [10]. Here, we describe a second class of novel anthelmintic, the diaminothienopyrimidines (DATPs). The potential for this early stage series is significant; the chemical synthesis of this series is facile and lends itself to iterative optimisation, will facilitate structural modifications aiming, for example, to increase local epithelial penetrance and hence improve efficacy during future development. Furthermore, this straightforward production imparts a favourable cost-economic aspect onto the series.

Activity against the egg stage of *T. muris*

In addition to activity against the adult stage of whipworm, the DATPs were also able to significantly reduce egg hatching, both *in vitro* and *in vivo*. These data are in keeping with members of the DHB series, which also were able to inhibit parasite egg hatching. However, unlike the DHB series, we identified members of the DATPs that also significantly reduced the percentage of eggs embryonating *ex vivo,* with other members of the DATP series appearing to disrupt the embryonation process, resulting in defects in embryonic elongation and abnormal egg shape. *Trichuris* egg embryonation occurs gradually and the mechanism by which it occurs is currently a poorly understood process. A detailed characterisation of the morphological changes which occur with the *Trichuris suis* egg during

embryonation has been described and other *Trichuris* species appear to undergo the same process.

Once the unembryonated, unsegmented eggs are deposited the two clear, nuclei-like areas move

together and fuse. Cellular division then begins, initially occurring asymmetrically with two

blastomeres of unequal size. The larger blastomere then divides again and then subsequently each

blastomere divides in two until a blastula forming of many small blastomeres develops. The initial

larval differentiation then occurs with the appearance of a motile cylindrical embryo, which gradually

turns into an infective larva with its characteristic oral spear. The fully developed larva is no longer

motile and is thought to be a L1 larva as no moult is observed within the egg [30]. The embryonation

process is temperature sensitive. The effect of temperature on egg embryonation has been

characterised in detail in recent years for *T. suis* eggs with the embryonation process accelerated at 30-

32 °C compared to 18 °C, with degeneration of the eggs rather than embryonation observed at higher

temperatures (40 °C). At low temperatures (5-10 °C) no embryonation occurs, however once these

eggs are then transferred to optimal embryonation temperatures normal embryonation proceeds [31].

Similar temperature sensitivity has been described for other *Trichuris* species including *Trichuris*

trichiura with different species embryonating with different kinetics [32,33]. More research is required

to understand the mechanisms behind this embryonation process, which may then allow an even more

targeted approach to breaking the life cycle.

Humans become infected with *Trichuris* via a faecal oral route. Adult parasites in the intestine shed

unembryonated eggs, which pass out with the faeces and embryonate in the external environment over

a period of five weeks. Parasite eggs are only infective if fully embryonated upon ingestion. Thus, the

ability of the DATPs to disrupt both embryonated egg hatching and the embryonation provides an

exciting alternative strategy, environmental control, to decrease *Trichuris* infection rates in the field

without the need to deliver anthelmintics to the infected host.

Other diaminothienopyrimidines – their applications and targets

Thienopyrimidines have received much interest in medicinal chemistry as they are bioisosteres for

purines, such as the nucleic acid components adenine and guanine. They are also related to

quinazolines, an important class of kinase inhibitors, including gefitinib and erlotinib, which act by

recognizing the ATP-binding site of the enzyme [34]. Thieno[2,3-*d*]pyrimidines are a particularly

important scaffold, with many reported examples of protein kinase inhibitors, as well as inhibitors of

dihydrofolate reductase, kainite receptor agonists, and α_1-adrenoreceptor antagonists [35]. The

thieno[3,2-*d*]pyrimidine scaffold, found in the compounds reported in this study, has also been

investigated, leading to the identification of a series of phosphatidylinositol-3-kinase inhibitors [36]. A

series of 2,4-diaminothieno[3,2-*d*]pyrimidines (the same core scaffold as the compounds reported in

this study) have been described as orally active antimalarial agents [37], with activity in the low

nanomolar range against *Plasmodium falciparum.* This series was later improved by systematic

modification giving improved antimalarial activity, but unfortunately continued hERG inhibition [38].

A series of 2,4-diaminothieno[3,2-*d*]pyrimidines has also recently been reported as active against the

endosymbiotic bacterium *Wolbachia,* with potential use against filarial nematodes [39]. In neither of

these latter two cases is the molecular target of the compounds known.

Conclusions

In summary we report the discovery of a new class of anthelmintic, the DATPs, which possesses

activity directed against adult stage *T. muris* parasites and the egg stage. Importantly, as a chemical

series the DATPS are notable, since they are relatively facile to produce synthetically thereby

presenting considerable scope for structural modifications to improve efficacy and deliver an

optimised agent.

Acknowledgements

The Bioimaging Facility microscopes used in this study were purchased with grants from BBSRC, Wellcome and the University of Manchester Strategic Fund. Special thanks goes to Roger Meadows for his help with the microscopy. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

References

1. World Health Organization. Helminth control in school age children: a guide for managers of control programmes [Internet]. 2nd ed. 2011. Available: http://www.who.int/intestinal_worms/resources/9789241548267/en/

2. Keiser J, Utzinger J. Efficacy of current drugs against soil-transmitted helminth infections: Systematic review and meta-analysis. JAMA. 2008;299: 1937–1948. doi:10.1001/jama.299.16.1937

3. Turner HC, Truscott JE, Bettis AA, Hollingsworth TD, Brooker SJ, Anderson RM. Analysis of the population-level impact of co-administering ivermectin with albendazole or mebendazole for the control and elimination of *Trichuris trichiura*. Parasite Epidemiol Control. 2016;1: 177–187. doi:10.1016/j.parepi.2016.02.004

4. Scott I, Pomroy WE, Kenyon PR, Smith G, Adlington B, Moss A. Lack of efficacy of monepantel against *Teladorsagia circumcincta* and *Trichostrongylus colubriformis*. Vet Parasitol. 2013;198: 166–171. doi:10.1016/j.vetpar.2013.07.037

5. Diawara A, Drake LJ, Suswillo RR, Kihara J, Bundy DAP, Scott ME, et al. Assays to Detect β-Tubulin Codon 200 Polymorphism in *Trichuris trichiura* and *Ascaris lumbricoides*. PLoS Negl Trop Dis. 2009;3: e397. doi:10.1371/journal.pntd.0000397

6. Diawara A, Halpenny CM, Churcher TS, Mwandawiro C, Kihara J, Kaplan RM, et al. Association between Response to Albendazole Treatment and β-Tubulin Genotype Frequencies in Soil-transmitted Helminths. PLoS Negl Trop Dis. 2013;7: e2247. doi:10.1371/journal.pntd.0002247

7. Koyama K. Bacteria-induced hatching of *Trichuris muris* eggs occurs without direct contact between eggs and bacteria. Parasitol Res. 2016;115: 437–440. doi:10.1007/s00436-015-4795-2

8. Hayes KS, Bancroft AJ, Goldrick M, Portsmouth C, Roberts IS, Grencis RK. Exploitation of the intestinal microflora by the parasitic nematode *Trichuris muris*. Science. 2010;328: 1391–1394. doi:10.1126/science.1187703

9. Hurst RJ, Hopwood T, Gallagher AL, Partridge FA, Burgis T, Sattelle DB, et al. An antagonist of the retinoid X receptor reduces the viability of *Trichuris muris* in vitro. BMC Infect Dis. 2014;14: 520. doi:10.1186/1471-2334-14-520

10. Silbereisen A, Tritten L, Keiser J. Exploration of novel in vitro assays to study drugs against *Trichuris* spp. J Microbiol Methods. 2011;87: 169–175. doi:10.1016/j.mimet.2011.08.009

11. Wangchuk P, Pearson MS, Giacomin PR, Becker L, Sotillo J, Pickering D, et al. Compounds Derived from the Bhutanese Daisy, *Ajania nubigena,* Demonstrate Dual Anthelmintic Activity against *Schistosoma mansoni* and *Trichuris muris*. PLOS Negl Trop Dis. 2016;10: e0004908. doi:10.1371/journal.pntd.0004908

12. Partridge FA, Murphy EA, Willis NJ, Bataille CJR, Forman R, Heyer-Chauhan N, et al. Dihydrobenz[*e*][1,4]oxazepin-2(3*H*)-ones, a new anthelmintic chemotype immobilising whipworm and reducing infectivity *in vivo*. PLoS Negl Trop Dis. 2017;11: e0005359. doi:10.1371/journal.pntd.0005359

13. Buckingham SD, Sattelle DB. Fast, automated measurement of nematode swimming (thrashing) without morphometry. BMC Neurosci. 2009;10: 84. doi:10.1186/1471-2202-10-84

14. Buckingham SD, Partridge FA, Sattelle DB. Automated, high-throughput, motility analysis in
Caenorhabditis elegans and parasitic nematodes: Applications in the search for new
anthelmintics. Int J Parasitol Drugs Drug Resist. 2014;4: 226–232.
doi:10.1016/j.ijpddr.2014.10.004

15. Partridge FA, Brown AE, Buckingham SD, Willis NJ, Wynne GM, Forman R, et al. An
automated high-throughput system for phenotypic screening of chemical libraries on *C. elegans*
and parasitic nematodes. Int J Parasitol Drugs Drug Resist. 2018;8: 8–21.
doi:10.1016/j.ijpddr.2017.11.004

16. Ritz C, Baty F, Streibig JC, Gerhard D. Dose-Response Analysis Using R. PLOS ONE. 2015;10:
e0146021. doi:10.1371/journal.pone.0146021

17. Stiernagle T. Maintenance of *C. elegans*. WormBook. 2006; doi:10.1895/wormbook.1.101.1

18. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.
doi:10.1038/nmeth.2019

19. Dahlin JL, Walters MA. The essential roles of chemistry in high-throughput screening triage.
Future Med Chem. 2014;6: 1265–1290. doi:10.4155/fmc.14.60

20. Oprea TI, Allu TK, Fara DC, Rad RF, Ostopovici L, Bologa CG. Lead-like, Drug-like or "Pub-
like": How different are they? J Comput Aided Mol Des. 2007;21: 113–119.
doi:10.1007/s10822-007-9105-3

21. Rankovic Z. CNS Physicochemical Property Space Shaped by a Diverse Set of Molecules with
Experimentally Determined Exposure in the Mouse Brain. J Med Chem. 2017;60: 5943–5954.
doi:10.1021/acs.jmedchem.6b01469

22. McKerrow JH, Lipinski CA. The rule of five should not impede anti-parasitic drug development.
Int J Parasitol Drugs Drug Resist. 2017;7: 248–249. doi:10.1016/j.ijpddr.2017.05.003

23. Sander T, Freyss J, von Korff M, Rufener C. DataWarrior: An Open-Source Program For
Chemistry Aware Data Visualization And Analysis. J Chem Inf Model. 2015;55: 460–473.
doi:10.1021/ci500588j

24. Keiser J, Tritten L, Adelfio R, Vargas M. Effect of combinations of marketed human
anthelmintic drugs against *Trichuris muris in vitro* and *in vivo*. Parasit Vectors. 2012;5: 292.
doi:10.1186/1756-3305-5-292

25. Ho NFH, Geary TG, Barsuhn CL, Sims SM, Thompson DP. Mechanistic studies in the
transcuticular delivery of antiparasitic drugs II: *ex vivo/in vitro* correlation of solute transport by
Ascaris suum. Mol Biochem Parasitol. 1992;52: 1–13. doi:10.1016/0166-6851(92)90031-E

26. Burns AR, Wallace IM, Wildenhain J, Tyers M, Giaever G, Bader GD, et al. A predictive model
for drug bioaccumulation and bioactivity in *Caenorhabditis elegans*. Nat Chem Biol. 2010;6:
549–557. doi:10.1038/nchembio.380

27. Maya C, Ortiz M, Jiménez B. Viability of *Ascaris* and other helminth genera non larval eggs in
different conditions of temperature, lime (pH) and humidity. Water Sci Technol J Int Assoc
Water Pollut Res. 2010;62: 2616–2624. doi:10.2166/wst.2010.535

28. Moser W, Schindler C, Keiser J. Efficacy of recommended drugs against soil transmitted
helminths: systematic review and network meta-analysis. BMJ. 2017;358: j4307.
doi:10.1136/bmj.j4307

29. Tilney LG, Connelly PS, Guild GM, Vranich KA, Artis D. Adaptation of a nematode parasite to
living within the mammalian epithelium. J Exp Zoolog A Comp Exp Biol. 2005;303A: 927–945.
doi:10.1002/jez.a.214

30. Wakelin D. The Development of the Early Larval Stages of *Trichuris muris* in the Albino
Laboratory Mouse. J Helminthol. 1969;43: 427–436. doi:10.1017/S0022149X00004995

31. Vejzagić N, Kringel H, Bruun JM, Roepstorff A, Thamsborg SM, Grossi AB, et al. Temperature
dependent embryonic development of *Trichuris suis* eggs in a medicinal raw material. Vet
Parasitol. 2016;215: 48–57. doi:10.1016/j.vetpar.2015.10.031

32. Nolf LO. Experimental studies on certain factors influencing the development and viability of
the ova of the human *Trichuris* as compared with those of the human *Ascaris*. Am J Epidemiol.
1932;16: 288–322. doi:10.1093/oxfordjournals.aje.a117862

33. Fahmy MA. An investigation on the life cycle of *Trichuris muris*. Parasitology. 1954;44: 50–57.

34. Zhang J, Yang PL, Gray NS. Targeting cancer with small molecule kinase inhibitors. Nat Rev
Cancer. 2009;9: 28. doi:10.1038/nrc2559

35. Wilding B, Klempier N. Newest Developments in the Preparation of Thieno[2,3-*d*]pyrimidines.
Org Prep Proced Int. 2017;49: 183–215. doi:10.1080/00304948.2017.1320513

36. Folkes AJ, Ahmadi K, Alderton WK, Alix S, Baker SJ, Box G, et al. The Identification of 2-(1H-
Indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-
d]pyrimidine (GDC-0941) as a Potent, Selective, Orally Bioavailable Inhibitor of Class I PI3
Kinase for the Treatment of Cancer. J Med Chem. 2008;51: 5522–5532. doi:10.1021/jm800295d

37. González Cabrera D, Le Manach C, Douelle F, Younis Y, Feng T-S, Paquet T, et al. 2,4-
Diaminothienopyrimidines as Orally Active Antimalarial Agents. J Med Chem. 2014;57: 1014–
1022. doi:10.1021/jm401760c

38. Gonzàlez Cabrera D, Douelle F, Le Manach C, Han Z, Paquet T, Taylor D, et al. Structure–
Activity Relationship Studies of Orally Active Antimalarial 2,4-Diamino-thienopyrimidines. J
Med Chem. 2015;58: 7572–7579. doi:10.1021/acs.jmedchem.5b01156

39. Johnston KL, Cook DAN, Berry NG, Hong WD, Clare RH, Goddard M, et al. Identification and
prioritization of novel anti-*Wolbachia* chemotypes from screening a 10,000-compound diversity
library. Sci Adv. 2017;3: eaao1551. doi:10.1126/sciadv.aao1551

Supporting information captions

709   **S1 Figure. Raw data separated by batch for the *in vivo* hatching experiment.** Each point indicates

710   one mouse that has been infected with *T. muris* eggs that had been treated with deionised water plus

711   1% v/v DMSO (control) or deionised water plus 1% v/v DMSO and final concentration 100μM

712   **OX02926** for 14 days. Blue line indicates mean for each treatment group.