Surprise About Sensory Event Timing Drives Cortical Transients in the Beta Frequency Band

T. Meindertsma[1,2,3], N.A. Kloosterman[2,3,4], A.K. Engel[1], E.J. Wagenmakers[2], T.H. Donner[1,2,3]

[1]*Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany;* [2]*Department of Psychology,* [3]*Amsterdam Brain and Cognition, University of Amsterdam, Amsterdam, Netherlands;* [4]*Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Max Planck Institute for Human Development, Berlin, Germany*

Address for correspondence:
Thomas Meindertsma
Dept. of Psychology
University of Amsterdam
Nieuwe Achtergracht 129B, 1018 WS Amsterdam, The Netherlands
Email: t.meindertsma@uva.nl
        t.donner@uke.de

Author contributions:
Conceptualization, T.M., N.A.K., E.J.W. and T.H.D.; Investigation, T.M.; Formal analysis T.M.; Analytic tools, E.J.W. and N.A.K.; Writing - Original draft, T.M. and T.H.D; Writing - Review & Editing, T.M., N.A.K., A.K.E., E.J.W. and T.H.D.; Funding Acquisition, A.K.E. and T.H.D.; Supervision, T.H.D.

Number of pages: 21
Number of figures: 5
Number of tables/multimedia/3D models: 0
Number of words Abstract: 248
Number of words Significance Statement: 119
Number of words Introduction: 499
Number of words Discussion: 1287

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by the Netherlands Organization for Scientific Research (NWO, dossiernummer 406-14-016, to T.H.D. and T.M.); the Amsterdam Brain and Cognition priority program (ABC2014-01, to T.H.D.); the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 604102 (Human Brain Project) (to T.H.D. and A.K.E.); and the German Research Foundation (DFG): Heisenberg Professorship DO 1240/3-1 (to T.H.D.), the Collaborative Research Centers SFB 936 (Projects A2/A3, A7, to A.K.E., T.H.D.) and TRR 169 (Project B1 to A.K.E.). We thank Udo Böhm for methodological advice and all members of the Donner lab for helpful discussion.

Abstract

Learning the statistical structure of the environment is crucial for adaptive behavior. Humans and non-human decision-makers seem to track such structure through a process of probabilistic inference, which enables predictions about behaviorally relevant events. Deviations from such predictions cause surprise, which in turn helps improve inference. Surprise about the timing of behaviorally relevant sensory events drives phasic responses of neuromodulatory brainstem systems, which project to the cerebral cortex. Here, we developed a computational model-based magnetoencephalography (MEG) approach for mapping the resulting cortical transients across space, time, and frequency, in the human brain (N=28, 17 female). We used a Bayesian ideal observer model to learn the statistics of the timing of changes in a simple visual detection task. This model yielded quantitative trial-by-trial estimates of temporal surprise. The model-based surprise variable predicted trial-by-trial variations in reaction time more strongly than the externally observable interval timings alone. Trial-by-trial variations in surprise were negatively correlated with the power of cortical population activity measured with MEG. This surprise-related power suppression occurred transiently around the behavioral response, specifically in the beta frequency band. It peaked in parietal and prefrontal cortices, remote from the motor cortical suppression of beta power related to overt report (button press) of change detection. Our results indicate that surprise about sensory event timing transiently suppresses ongoing beta-band oscillations in association cortex. This transient suppression of frontal beta-band oscillations might reflect an active reset triggered by surprise, and is in line with the idea that beta-oscillations help maintain cognitive sets.

Significance statement

The brain continuously tracks the statistical structure of the environment to anticipate behaviorally relevant events. Deviations from such predictions cause surprise, which in turn drives neural activity in subcortical brain regions that project to the cerebral cortex. We used magnetoencephalography in humans to map out surprise-related modulations of cortical population activity across space, time, and frequency. Surprise was elicited by variable timing of visual stimulus changes requiring a behavioral response. Surprise was quantified by means of an ideal observer model. Surprise predicted behavior as well as a transient suppression of beta frequency-band oscillations in frontal cortical regions. Our results are in line with conceptual accounts that have linked neural oscillations in the beta-band to the maintenance of cognitive sets.

Introduction

Humans and other organisms continuously adapt their behavior to the statistical structure of their environment. This suggests that the brain is equipped with neural machinery for statistical learning, which can interact with the processes driving goal-directed behavior. Of particular importance here is surprise (Dayan and Yu, 2006; O'Reilly et al., 2013), a violation of one's expectation about the next event, which might indicate a sudden change in the environmental structure, and can transiently boost central arousal state, increasing the organism's sensitivity and learning rate (Yu and Dayan, 2005; Nassar et al., 2012).

Expectation, uncertainty, and surprise are intricately related. The precision of expectations scales with uncertainty, that is, the width of the distribution of observed events: high uncertainty precludes forming precise expectations. Violations of expectations cause surprise, the level of which depends on the difference between the expected and actually observed event (often termed prediction error). These intuitions can be formalized within the framework of Bayesian statistics and used to search for neurophysiological correlates (see Materials and Methods: *Bayesian ideal observer model: General approach and rationale*).

One important dimension of the environment is the timing of relevant sensory events (Gibbon et al., 1997; Nobre et al., 2007). Two lines of work have studied the neural basis of temporal expectation effects. One has shown that environments with rhythmic (i.e., precise) temporal structure entrain neural oscillations in the cerebral cortex, the phase of which then modulates sensory cortical responses, perception, and cognition (Lakatos et al., 2008; Schroeder and Lakatos, 2009; Rohenkohl and Nobre, 2011; Rohenkohl et al., 2012; Riecke et al., 2015; van Ede et al., 2017). In these rhythmic changes of the environment, surprise is minimized (once the structure is learned expectations match observations). Consequently, this first line of work has identified neural correlates of temporal expectation, rather than of surprise.

The second line of work has studied neural responses of subcortical, neuromodulatory centers, specifically, dopaminergic centers of the midbrain, to sensory events entailing reward. Because event timing here varied non-periodically from trial to trial as in many natural environments, this work could link phasic neuromodulatory responses to temporal surprise (Hollerman and Schultz, 1998; Fiorillo et al., 2008). Surprise-driven phasic responses might also occur in other neuromodulatory brainstem systems, such the noradrenergic system (Dayan and Yu, 2006). Because brainstem neuromodulatory systems have widespread projections to the cortical networks underlying goal-directed behavior, one would expect changes in cortical population activity elicited by surprise (Bouret and Sara, 2005). However, this second line of work on temporal expectation has focused on surprise-related activity in subcortical systems.

Here, we studied responses to surprise about the timing of sensory events in human cortex. A computational model-based magnetoencephalography (MEG) approach enabled us to map surprise-related cortical transients across space, time, and frequency. We used a Bayesian model that accumulated previously experienced durations of the interval between visual changes into posterior beliefs about the next interval duration. This ideal observer model provided trial-to-trial measures of temporal surprise, which predicted modulations of prefrontal and parietal cortical beta-band dynamics.

Materials & Methods

This paper reports a re-analysis of an MEG data set that has previously been used for a study into decision-related feedback signals in visual cortex (Meindertsma et al., 2017). Here, we focus on those aspects of the experimental design that are most

136 relevant for the issue addressed in the current paper: uncertainty and surprise about
137 the timing of the experimental events specified below. We refer to our previous paper
138 (Meindertsma et al., 2017) for a more detailed description of the visual stimulus and
139 the behavioral task.
140
Participants
Thirty-one volunteers participated in the experiment. Two participants were excluded
due to incomplete data and one participant did not complete the experiment due to
poor quality of simultaneously acquired pupil data. Thus, 28 participants (17 female,
age range 20 - 54 years, mean age 28.3, SD 9.2) were included in the analysis. All
participants had normal or corrected-to-normal vision and no known history of
neurological disorders. The experiment was conducted in accordance with the
Declaration of Helsinki and approved by the local ethics committee of the Hamburg
Medical Association. Each participant gave written informed consent.
150
Stimulus
MEG was measured while subjects viewed the intermittent presentation of a target
(full contrast Gabor patch; diameter: 2°) and reported the on- and offset of the target
(Figure 1A). Gabor targets flickered at 10 Hz (counter-phase) through alternation of
two out-of phase Gabors every 50 ms. This caused steady-state evoked responses
over visual cortex at 10 and 20 Hz (data not shown), distinct in terms of spectral
profile, topography and functional characteristics from the surprise-related
modulations we focused on here. The target was located in either the lower left or
lower right visual field quadrant (eccentricity: 5°, counterbalanced between subjects),
surrounded by a rotating mask (17°x17° grid of black crosses), and superimposed on
a gray background. The mask rotated at a speed of 160°/s. The target was separated
from the mask by a gray "protection zone" subtending about 2° around the target
(Bonneh et al., 2001). Subjects fixated on a fixation mark (red outline, white inside,
0.8° width and length) centered on the mask in the middle of the screen. Stimuli were
presented using the Presentation Software (NeuroBehavioral Systems, Albany, CA,
USA, RRID:SCR_002521). Stimuli were back-projected on a transparent screen
using a Sanyo PLC-XP51 projector with a resolution of 1024x768 pixels at 60 Hz.
Subjects were seated 58 centimeters from the screen in a whole-head
magnetoencephalography (MEG) scanner setup in a dimly lit room.
170

171
Figure 1: Behavioral task. A. Schematic depiction of the stimulus and task. A salient, flickering
target (Gabor patch) temporarily appeared and disappeared on a rotating background. Subjects fixated
on the red fixation mark and reported stimulus changes either by direct button press or silently counting
the disappearances and reporting the total number at the end of the run. **B.** The interval duration
between stimulus changes was randomly drawn from one of three distributions that corresponded to

three hazard rates (left), resulting in distinct distributions of intervals (right, average histogram over
subjects). **C.** Example time courses of target presence (1 = present, 0 = absent) drawn from these
distributions.
180
Behavioral task and experimental design
The subjects' task was to maintain stable fixation and detect the physical offsets and
onsets of the target, the predictability of which fluctuated from trial to trial, and the
mean predictability of which varied systematically across blocks. To this end, the
interval durations between stimulus changes were sampled from three different
distributions in the different blocks. These distributions were computed so as to
produce three predetermined so-called hazard functions, which describe the
probability that an event will occur at a particular time, given that it has not occurred
yet. The hazard function formalizes the expectation of a change and affects human
reaction times in simple detection tasks (Luce, 1986). The hazard function can be
computed as follows:
192
193 $$\lambda_t = \frac{f_t}{1 - F_t},$$ Eq. 1
194
where λ_t is the value of the hazard function at time point t, f_t is the value of
distribution f on time point t, and F_t is the area under the curve of distribution f from
$-\infty$ to time t.
We used the following procedure to construct three 'environments', referred to
as 'Short', 'Long', and 'Flat' below. We first selected three hazard functions that
systematically differed in their level of predictability (Figure 1B, C). We then
computed the actual distributions of intervals by re-arranging Eq. 1 as follows:
202
203 $$f_t = \lambda_t * (1 - F_t) ,$$ Eq. 2
204
The interval durations were then randomly selected from f. Specifically, the temporal
environments were defined as follows:
Short: The hazard function was a narrow Gaussian distribution with a mean of
2 s and a standard deviation of 0.2 s. This resulted in nearly periodic and, thus,
largely predictable intervals between events.
Long: This condition used the same hazard function as the previous
condition, but with a larger mean and standard deviation (6 s and 0.6 s, respectively)
thus rendering event timings less predictable (Fiorillo et al., 2008).
Flat: The hazard function was flat with a mean of 6s, yielding the least
predictable interval durations. The resulting distribution of interval durations, f_t,
therefore, approximated an exponential distribution; characterizing a memory-less
process (i.e. the timing of the next event could not be predicted from previously
encountered intervals, Feller, 1959).
Computational analysis with a Bayesian model (Fig. 2) described below
confirmed that the sampled intervals from these three environments gave rise to
different mean levels of uncertainty and surprise (Fig. 2G,H). The three environments
were presented in separate three-minute blocks.
Within each of the above temporal environments, there were two behavioral
tasks. Both tasks required subjects to monitor the changes of the small visual target.
In one task (called *Detection-button*), they were asked to report those changes
immediately. Specifically, subjects reported target offsets or onsets by pressing a
button with the right index or middle finger, respectively. In the other task (*Detection-*
count), they were asked to count and report the changes at the end of the block.
Subjects silently counted the number of target offsets and reported the total in

response to a 4-AFC question at the end to the block. The two tasks were randomly
selected before each block under the constraint that both would occur equally often.
We here analyzed both task conditions, but only found robust effect for Detection-
button.
All subjects completed a total of 6 blocks of the Short environment, and 16
blocks of the other Long and Flat environments, resulting in about the same number
of trials per environment. Additionally, subjects performed a motion-induced
blindness task and a functional localizer task, which were not relevant for the current
study, but are reported in our previous paper (Meindertsma et al., 2017). All blocks
within an environment were completed in succession; the order of environments was
counter-balanced across subjects.
240
Bayesian ideal observer model: General approach and rationale
We developed an ideal observer model to quantify surprise and uncertainty about the
timing of sensory events (i.e., the target on- and offsets). The model tracked the
evolving predictive distribution of upcoming interval durations; more specifically, it
computes the posterior predictive of unobserved interval durations, conditional on the
observed data, throughout each block of the experiment. We assumed that subjects
tracked the temporal statistics of the task in a similar way, and we used the posterior
predictive distribution as a proxy of the subjects' belief states (i.e., their prediction of
the timing of the next stimulus change).
While we used an ideal observer model that prescribed the optimal inference
for our task, we are agnostic to the precise inference process that was used by our
subjects and we do not claim that subjects used the exact computations used by the
model. Our central assumption was that subjects accumulated observations
throughout each block (i.e., over more than just one or two previous intervals). This
assumption was derived from a substantial body of work on other forms of learning
and evidence accumulation (Sutton and Barto, 1998; Gold and Shadlen, 2007; Glaze
et al., 2015), and it was supported by the findings described in Results. Our model
implemented the normative accumulation strategy by perfectly integrating across the
entire history of the observations (here: of interval durations) and updating internal
representations accordingly. A practical benefit of this approach was that it did not
require fitting of model parameters, for which our current data did not provide
sufficiently strong constraints.
The only free parameter in the model was the level of temporal estimation
noise, which we allowed to scale with the magnitude of the interval duration
according to Weber's law (Gibbon et al., 1997). To this end, we transformed the
discrete values of the observed intervals into Gaussian distributions that were used
to update the model (see next section). The mean of these distributions was equal to
the observed interval t and their standard deviation was equal to the observed
interval t times a Weber's fraction (coefficient of variation, Gibbon et al., 1997). We
simulated the model with 34 Weber's fraction values ranging from 0.001 to 0.5
(0.001, 0.05:0.01:0.35, 0.4, 0.5). We then computed the correlation between the
measured single-trial reaction times (pooled across all subjects) and surprise (see
Bayesian ideal observer model: Implementation), separately for each Weber fraction,
and selected the Weber fraction that maximized this correlation. To this end, we fitted
a second order polynomial to the correlation coefficients as a function of Weber's
fraction and extracted the maximum of the polynomial. This yielded a Weber's
fraction of 0.17 (Figure 2F), which was used for all analyses reported in this paper.
Using model-based surprise from a noise-free version of the model yielded
qualitatively identical results (data now shown).
280

 Bayesian ideal observer model: Implementation
We assumed that the subjects used a model in which the observed intervals have
been generated from a gamma distribution with parameters alpha (shape) and beta
(scale). These parameters were given uninformative prior distributions (Lee and
Wagenmakers, 2013), which were updated by the data to posterior distributions.
Using the interval duration distributions as the observations, we could obtain
the expectations about to-be-observed intervals by generating posterior predictives
(i.e., drawing an alpha-beta pair from the joint posterior distribution and then drawing
a predicted interval from the associated gamma distribution; repeating this process
many times yields a posterior predictive distribution for the to-be-observed interval).
We assumed that the subjects updated their belief state after each observation of a
new interval duration. Likewise, the model was updated after every interval *t* by
computing a new posterior predictive distribution, based on the durations of intervals
1:t and the prior.
We generated a posterior predictive distribution over the to-be-observed
intervals using Gibbs sampling (a Markov chain Monte Carlo, or MCMC, algorithm;
Andrieu et al., 2003) in the software JAGS (Plummer, 2003) and Matlab (version
R2013a, RRID:SCR_001622). We used two Markov chains with different starting
points comprised of 2500 samples per chain with 500 samples burn-in, for a
combined total of 4000 samples. The posterior predictive MCMC samples $Y_{1\ldots4000}$ for
the next interval, *t+1*, were then summarized by a gamma distribution using the
functions 'gamfit' and 'gampdf' in Matlab (Figure 2A,B):
303

$$Y^{(t+1)}_{1\ldots4000} \sim Gamma(\kappa_{t+1}, \theta_{t+1}) \equiv f_{t+1}, \qquad \text{Eq. 3}$$

305
where $Y^{(t+1)}_{1\ldots4000}$ are the MCMC samples, and κ_{t+1} and θ_{t+1} are the parameters of the
gamma distribution f_{t+1}; hence, f_{t+1} is the continuous posterior predictive distribution
for the upcoming interval after having observed the preceding intervals *1...t*.
To be able to relate trial-to-trial uncertainty and surprise to behavior and the
MEG data, we extracted two information theoretic metrics from the time-evolving
posterior predictive distribution f_{t+1} (i.e., belief).
Uncertainty: We quantified trial-to-trial uncertainty about the timing of the
upcoming interval *t+1* as the entropy of the posterior predictive distribution f_{t+1} (i.e.,
the posterior predictive based on intervals *1…t*):
315

$$H_{t+1} = -\int_0^\infty (f_{t+1}(x) * \log f_{t+1}(x))\, dx, \qquad \text{Eq. 4}$$

317
where H_{t+1} is the entropy after intervals *1…t*, and the integral is over all possible
values *x* for the upcoming interval. Entropy depended on the width of f_{t+1}, and thus
uncertainty was higher when predictions of interval durations were less precise
(Figure 2A,C,D). For clarity, in what follows we will use the term entropy when
referring to this uncertainty.
Surprise: For every upcoming interval *t+1*, we computed the surprise about
the corresponding interval duration in terms of the Shannon information conveyed by
the interval duration x_{t+1}, given the posterior predictive distribution f_{t+1}:
326

$$I_{t+1} = -\log f_{t+1}(x_{t+1}), \qquad \text{Eq. 5}$$

328
where I_{t+1} is the information gained by interval *t+1*, given f_{t+1}. Thus, surprise was
defined as the negative log-probability of the upcoming interval *t+1*, given the
intervals that had been presented so far.

We added one further transformation in the computation of surprise. The
surprise measure defined in Eq. 5 quantified the surprise about the upcoming event
timing based on the posterior predictive distribution f_{t+1}, but disregarding the time
elapsed in the current interval. It is unlikely that exactly this distribution translated into
subjects' level of surprise: as time passed and no event occurred in a given interval,
all interval durations shorter than the elapsed time become impossible. Subjects
likely discounted these impossible intervals in their expectation of the timing of the
upcoming event, which should have also affected their level of surprise. In other
words, their internal representation of the posterior predictive distribution changed
dynamically throughout each trial, as a function of elapsed time. We constructed a
time-varying version of the posterior predictive distribution f_{t+1}, which was also
conditioned on the elapsed time on interval t. This version was equal to f_{t+1} for
elapsed time equal to 0 and then increasingly deviated from f_{t+1} as elapsed time grew.
We approximated this time-varying distribution, denoted as f'_{t+1} in the following, by
setting all probabilities in f_{t+1} up to the current time point to zero and renormalizing the
remaining distribution to integrate to 1 (Figure 2B). We then computed surprise
based on this new distribution f'_{t+1} using Eq. 5. The time-variant prior f'_{t+1} converged
to 1 as time passed, and thus surprise approached zero for longer intervals.
350

Regressing computational variables against behavior
We used reaction time (RT) during Detection-button as behavioral readout of the
impact of uncertainty and surprise. Accuracy approached ceiling for all subjects, due
to the high saliency of the target. We computed and compared mean RTs per
environment and stimulus event (target off- and onset).
We also correlated RT to our trial-to-trial estimates of surprise and entropy.
RT was log-transformed so as to normalize the skewed RT distributions. To test if the
model-based surprise fitted the behavioral (RT) data better than a linear combination
of just the two previous interval durations (i.e., a leaky accumulation with strong
leak), we used multiple linear regressions to compare the following two nested
models:
362

363   M1:   *log(RT) ~ Interval$_t$*Env*Event + Interval$_{t-1}$*Env*Event*
364   M2:   *log(RT) ~ Interval$_t$*Env*Event + Interval$_{t-1}$*Env*Event + Surprise*Con*Event,*
365

where *Interval$_t$* and *Interval$_{t-1}$* corresponded to the durations of the two intervals
preceding the visual change (i.e., interval on trial *t* and *t-1*), and *Surprise* was the
computational model-derived metric. Predictors were multiplied by categorical
variables envrionment (*Env*, the three different temporal environments) and *Event*
(target offset or onset). Both variables strongly affected RT (Figure 3A). We fitted
both M1 and M2 and compared the fits per subject using adjusted R^2.
372

MEG data collection
Magnetoencephalography (MEG) data were acquired on a CTF 275 MEG system
(VSM/CTF Systems, Port Coquitlam, British Columbia, Canada) with a sample rate of
1200 Hz. The location of the subjects' head was measured in real-time using three
fiducial markers placed in the both ears and on the nasal bridge to control for
excessive movement. Furthermore, electrooculogram (EOG) and electrocardiogram
(ECG) were recorded to aid artifact rejection. All data were recorded in sets of four
blocks of three minutes duration (or two blocks at the end of an environment set).
381
382
383

MEG data analysis

Preprocessing. The data were analyzed in Matlab (version R2013a, The Mathworks, Natick, MA, USA, RRID:SCR_001622) using the Fieldtrip (Oostenveld et al., 2011, RRID:SCR_004849) toolbox and custom-made software.

Trial extraction. In blocks involving subjects' reports, we extracted trials of variable duration, centered on subjects' button presses, from the 3 min blocks of continuous stimulation. We call this method for trial extraction "response-locked". The following constraints were used to avoid mixing data segments from different percepts when averaging across trials: (i) The maximum trial duration ranged from −1.5 s to 1.5 s relative to report; (ii) when another report occurred within this interval, the trial was terminated 0.5 s from this report; (iii) when two reports succeeded one another within 0.5 s, no trial was defined; (iv) for the analysis of Detection-button blocks, we included only those reports that were preceded by a physical change of the target stimulus within 0.2 to 1 s, thus discarding reports not related to stimulus changes. We used this method for the analyses related to surprise. In an alternative analysis of all Detection blocks, trials were defined in the same way as described above, but now aligned to physical target on- and offsets ("stimulus-locked"). In the Detection-count task, no button responses were given during the block, so stimulus-locked trial extraction was the only option. We used this method for the analysis related to entropy (see Kloosterman et al., 2015b & Meindertsma et al., 2017 for a similar procedure).

Artifact rejection. All epochs that contained artifacts caused by environmental noise, eye-blinks, muscle activity or squid jumps were excluded from further analysis using standard automatic methods included in the Fieldtrip toolbox. Epochs that were marked as containing an artifact were discarded after every artifact detection step. For all artifact detection steps the artifact thresholds were set individually for all subjects. Both of these choices aimed at optimization of artifact exclusion. Line-noise was filtered out by subtracting the 50, 100, 150 and 200 Hz frequency components from the signal.

Time-frequency decomposition. We used a sliding window Fourier transform to compute the time-frequency representation for each sensor and each trial of the MEG data. The sliding window had a length of 200 ms and a time step size of 50 ms, with one Hanning taper (frequency range 5-35 Hz, frequency resolution 5 Hz and frequency step size 1 Hz). The data was baseline corrected for every frequency bin and MEG sensor separately. The baseline was computed by averaging single-trial power over a baseline time window. The baseline time windows ranged from −1.25 to −0.75 s for response-locked and −1 to −0.5 s for stimulus-locked analyses, respectively. The time course of every frequency bin and sensor combination was baseline corrected by subtracting the single-trial baseline power at that frequency and dividing by the mean baseline power across trials within an experimental environment. We used the single-trial baseline power for subtraction to eliminate the effect of slow power fluctuations, because any surprise-related power modulations could only have occurred after the sensory event that elicited surprise. We used the mean baseline for division in order to minimize noise in the single-trial estimates of the single-trial power modulation values. This division was used to compensate for the common decay of power with frequency, which hinder identification of effects at higher frequencies, and to normalize the single-trial modulation values (Siegel and Donner, 2010). It did not systematically alter the association of power modulation values with other variables.

Source reconstruction. We used an adaptive linear spatial filtering method called linear beamforming (Van Veen et al., 1997; Gross et al., 2001) to estimate single-trial modulations of MEG power at the source level. We computed a common

filter for a baseline time window (1 to 0.5 s before response), a 'transient' time
window, and a frequency band of interest (0 to 0.5 s after response, 20 Hz +/- 4 Hz
spectral smoothing, see dashed box in Figure 4A). The transient time window and
frequency band of interest were selected based on cluster-based statistics at the
sensor level (see next section). We used the measured head positions and individual
single-shell volume conductor models, based on individual images from T1-weighted
structural MRI. We computed the power values, in both baseline and transient time
windows, for each trial and source grid point (i.e., voxel) as follows. First, we
projected the sensor-level MEG power values from the time window of interest as
well as from a baseline time window through the common spatial filter. Second, we
converted the estimated power values during the time window of interest into units of
power modulation, again by subtracting and dividing by the corresponding baseline
power values.
449
Correlating single-trial computational variables to MEG power
We correlated the MEG power modulation to our measures of entropy and surprise,
as derived using our model (see *Bayesian ideal observer model: Implementation*)
across trials. Although intricately related (see Introduction), uncertainty and surprise
entailed different computations (see above). A key difference was *when* during the
course of a trial these two quantities where computed. So, we reasoned that neural
correlates of these computational quantities should also differ in their dynamics:
uncertainty about event timing should be reflected in the neural baseline state before
occurrence of the sensory event, whereas surprise should be reflected in a transient
response elicited by that event. Thus, we used different components of the single-
trial MEG power estimates for the analyses of entropy and surprise.
Entropy: We correlated entropy to the MEG power modulation separately in
every MEG sensor and frequency bin. This was done within subject and separately
for the three environments. There are structural differences in entropy and surprise
between these environments (Figure 2G,H), thus pooling over these conditions might
result in inflated correlations that reflect session differences instead of the true
correlation between entropy and MEG power. We reasoned that entropy should
affect baseline or tonic arousal, where high entropy should cause higher arousal. As
our task was continuous, we considered the time window right before the stimulus
change the best reflection of a baseline state. For this reason we averaged the MEG
power over the time period right before a stimulus change (-0.5 to -0.25s with respect
to the target offset or onset) before correlating to entropy.
The results were then averaged over the three environments and transformed
with the Fisher z transformation (Fisher, 1915):
474
$$z = 0.5 \cdot \ln \left(\frac{1+r}{1-r} \right) \hspace{3cm} \text{Eq. 6}$$
476
We used two-tailed permutation tests with a cluster-based correction for
multiple corrections to test the correlation coefficients against zero (Efron and
Tibshirani, 1998; Maris and Oostenveld, 2007).
Surprise: Correlations between surprise and MEG power modulation were
performed using the same method, with the following exceptions. First, we attuned
the analysis in two ways to account for the correlation between surprise and RT
(Figure 2F, 3). Because of this correlation, any post-stimulus correlations between
surprise and MEG power modulation might reflect differences in the timing of the
button press. We performed this analysis response-locked, because these RT
differences are difficult to disentangle from genuine effects of surprise when the
power modulations are time-locked to the stimulus change. Additionally, to account

for confounding effects of RT and the duration of the previous interval, we also
performed a partial correlation analysis between surprise and MEG power
modulation with the interval duration preceding the stimulus change or RT as
covariate. Second, for the correlation between surprise and MEG power modulation
we did not average over a specific time window, but instead performed correlations
separately for every time point, resulting in a 3-dimensional matrix of correlations
(sensor * frequency bin * time point). Consequently, we also performed cluster-based
permutation statistics over these three dimensions. The correlations that survived
cluster correction were visualized by integrating (i.e. computing the area under the
curve) over sensors and frequency bins (for the time course), sensors and time
points (for the frequency spectrum), frequency bins and time points (for the
topography) or just over sensors for the time frequency representation (see Hipp et
al., 2012 for a similar approach).
To assess the robustness of the emerging clusters we performed a cross-
validation analysis using a leave-one-out procedure. To this end, we repeated the
cluster-based permutation statistics on all possible iterations of N-1 subjects, each
time using the resulting cluster as a mask to calculate the average correlation in the
left-out subject, separately for target offset and onset trials. These values were tested
against zero and against each other across subjects using permutation tests (10.000
permutations).
We also computed the correlation between trial-to-trial power modulation
averaged over the whole cluster and log(RT). The resulting correlations were tested
against zero across subjects using a permutation test (10.000 permutations).
The transient modulations of MEG power estimated for each voxel in the
source grid, derived by means of source reconstruction (see *MEG data analysis:*
Source reconstruction), were correlated to the trial-to-trial measure of surprise. This
was done separately within each subject and the resulting correlations averaged over
subjects after Fischer's z-transformation (Eq. 6). For comparison, we also computed
the average modulations of MEG power in the same time window and frequency
band. The resulting maps of correlation or average power modulation were
nonlinearly aligned to a template brain (Montreal Neurological Institute) using the
individual images from structural MRI. To test the similarity of the spatial topography
of the correlation to the average modulation of power, we correlated the two
corresponding source maps per subject and tested the correlation coefficients again
zero on the group level by means of a permutation test (10.000 permutations).
523

Results

Subjects (N=28) performed a simple visual detection task reporting on- and offsets of
a small, but salient target stimulus (Figure 1A). In different blocks, target events were
administered using three different temporal environments (Figure 1B,C) translating
into different overall levels of uncertainty and surprise about the timing of target
events (Figure 2G,H). In order to quantify these two computational variables not only
across conditions, but also across individual trials, we developed a Bayesian belief-
updating model. The model incorporated the evolving beliefs (i.e. the posterior
predictive distributions) of an ideal observer about the temporal intervals between the
sensory events. Beliefs were dynamically updated across trials and even within trials
(for surprise, see *Materials and Methods*). From these time-evolving probability
distributions, we extracted trial-by-trial measures of information-theoretic entropy
(quantifying uncertainty) and surprise, which we related to the behavior and neural
dynamics of our participants.
538

Estimates of entropy and surprise fluctuated across trials, especially in the
early part of each block (Figure 2C-E). The trial averages of both measures within
each block also varied lawfully between the different experimental conditions, scaling
with the predictability of the stimulus changes (Figure 2G,H). Estimates were
smallest for the Short condition, intermediate for the Long condition, and largest for
the Flat condition. As expected, variations in entropy and surprise were weakly
correlated across trials (r=0.13 for Short and Flat, r=0.19 for Long condition, Figure
2G), since both measures were computed from the same probability distribution
(Materials and Methods). Even so, these two variables entailed distinct computations,
possibly by distinct neural circuits. Critically, both computational variables could be
computed at different times during a trial, thus possibly leading to different dynamical
modulations cortical population activity.

578
579
580

Surprise predicts reaction time

The model-derived computational variable surprise predicted subjects' reaction time (RT) in the detection task. Mean RT scaled with the different temporal environments in the same way as surprise and entropy, with the fastest RTs for Short and slowest RTs for the Flat condition (Figure 3A, compare with Figure 2E). RT correlated with surprise also at the single trial level (Figure 3B). We did not find robust correlations to RT for entropy.

We also tested whether model-derived surprise (entailing accumulation of intervals across the entire experimental block) predicted RT over and above a linear combination of only the two previous intervals (entailing, e.g. a leaky accumulation with strong leak). To this end, we used a nested regression model, which quantified the predictive power of a combination of surprise and the previous two intervals in accounting for the influence of temporal environments and target on- or offset, on RT (Materials and Methods). We compared this against a simpler model with only the two previous intervals. Because model-based surprise depended on all previous intervals, the comparison between the above two nested models assessed the impact on reaction time of intervals beyond the second one. We used adjusted R^2 for comparison, which penalized model complexity. This comparison yielded higher adjusted R^2 values for the model including surprise in 22 of 28 subjects (Figure 3C), indicating that surprise predicted RT over and above the duration of the previous two intervals.

Taken together, these results indicate that subjects tracked the temporal structure of the task by accumulating interval distributions at least over more than two intervals, akin to what was prescribed by the ideal observer model. We next searched the whole-brain MEG data for a dynamical neurophysiological signature of this process. To this end, we focused on the trial-to-trial fluctuations of surprise within each of the environments (Short, Long, Flat), which were more pronounced than the differences in mean surprise between environments.

Figure 3: Link between computational variables and behavior. A. Average reaction time (RT) per interval distribution, separate for reports of target offsets and onsets. Bars show average over subjects; black dots depict average per subject. *** p<0.001, ** p<0.01, permutation tests across subjects, 10.000 permutations (differences between target off- and onsets, p-values were p=0, p=0 and p=0.003, for Short, Long and Flat, respectively; between conditions, p=0, p=0 and p= 0.001, for Short-Long, Short-Flat and Long-flat, respectively). **B.** Correlation between trial-to-trial fluctuations in Entropy (left)/Surprise (right) and log-transformed RT. Mean correlation coefficient Entropy: mean r=0.03 (s.d.=0.02), p=0.17 and r=-0.04 (s.d.=0.03), p=0.07 for off- and onset, respectively, difference off-on p=0.47; Surprise: mean r=0.09 (s.d.=0.02), p=0.007 and r=0.15 (s.d.=0.03), p=0 for off and onset, respectively, difference off-on p=0.48. **C.** Model comparisons between partial (including only 2 previous intervals) and full regression models (further including surprise; see main text). Difference in adjusted R^2. Positive values indicate support for full model. Dots depict individual subjects; bars depict the mean, error bars depict SEM.

Widespread cortical beta-band transient driven by surprise
We mapped out the cortical responses to trial-to-trial fluctuations in surprise by
correlating the model-based surprise measures to modulations of MEG power,
around the time of subjects' behavioral responses to sensory events. We did this in
an exhaustive fashion across every time and frequency bin and MEG sensor and
tested for clusters of significant correlations across these three dimensions, while
applying cluster-based multiple comparison correction (*Materials and Methods*). This
approach revealed negative correlations in the beta (~20 Hz) frequency range, as
well as in the lowest frequency bin resolved (5 Hz), indicating that higher surprise
was associated with lower power in these frequency ranges. The peak in this
negative correlation cluster started about 0.2 s before and reached its maximum
about 0.25 s after subjects' report of the stimulus change. This cluster exhibited
several peaks over central, left frontal, and to a lesser extent left parietal cortex
(Figure 4A,C).
For all analyses shown in Figures 4 and 5, we used partial correlations,
controlling for reaction time, and we focused on the Detection-button task that
entailed immediate behavioral report of the change of the visual target (Materials and
Methods). We controlled for reaction time because (i) the data showed that the latter
was affected by surprise (Figure 3), and (ii) motor responses are known to modulate
beta-power around the time of response (Donner et al., 2009). Thus, button-presses
could have potentially influenced the modulation by surprise. We focused on
Detection-button because (i) we could only establish links between surprise and
behavior for this task and (ii) it allowed us to lock neural dynamics more closely to the
conscious registration of the visual change. When performing the correlation analysis
for the Detection-count task (then locked to the physical stimulus change), we did not
obtain any in significant correlation clusters. In two control analyses, we confirmed
that the above results were robust to (i) using the 'raw' correlation between surprise
and MEG power and (ii) controlling for the preceding interval duration. Both analyses
resulted in highly similar clusters of negative correlations (data not shown).
The surprise-related cluster was robust and not driven by outliers, and the
effect was not specific to the type of stimulus event (target on- or offset). We used a
leave-one-out cross-validation procedure to test the robustness of the correlations on
both target on- and offsets (Materials and Methods). We found robust negative
correlations in the left-out subjects (Figure 4D). Furthermore, the correlation was
found for both target offsets and onsets (Figure 4D, mean -0.036, -0.025, SEM 0.010,
0.006, p = 0.006, 0.002, for target offsets and onsets, respectively; difference:
p=0.26, permutation tests, 10.000 permutations).
As expected from previous work on modulations of MEG power around motor
responses (Donner et al., 2009), the overall modulation of MEG power in the time-
frequency window of the surprise-correlation cluster (16-24 Hz, 0–0.5 s from
response, normalized by the baseline 1-0.5 s before response) peaked in bilateral
motor cortex (Figure 4B). But the component of beta-power modulations that
correlated with trial-by-trial surprise showed a different cortical distribution, with
negative correlations that peaked in the central sulcus, extending from motor- to
more frontal cortex, and in left frontal and parietal cortex (compare Figure 4B and
4C). Indeed, there was no similarity between the individual topographies of the
surprise-linked and the overall power modulations (mean correlation across subjects:
r=-0.02, p=0.42). These observations indicate that the report-locked modulation
linked to surprise and of overall power were located in distinct cortical networks.
673

Figure 4: Widespread cortical beta-band transient driven by surprise. A. Exhaustive partial correlation (controlling for RT) between trial-to-trial measures of surprise and MEG power modulation in all sensors, time and frequency bins results in one cluster (cluster-based correction for multiple comparison, p = 0.001, two-sided) of negative correlation. Different panels show different dimensions of the cluster by integrating over the other dimensions; top left: time course, top right: spatial topography, bottom left: time-frequency representation; bottom right: frequency spectrum. B. Source reconstruction of the power modulation in the time window in which surprise-MEG correlation was strongest (dashed box in panel A). C. Source-reconstructed illustration of the correlation between transient modulation and trial-to-trial surprise depicted in panel B. These source maps are not statistically thresholded, but instead serve for comparing the correlation's spatial distribution with the transient power modulation in panel B (average correlation between surprise and power modulation across subjects = 0.02, p=0.42). D. Leave-one-out cross-validation of the cluster found in panel A, separately for target offsets and onsets. Cluster-based permutation was performed on N-1 subjects and the average correlation in the resulting cluster was computed for the remaining subject (black dots); bars show averages over subjects. Correlation values were tested against 0 (permutation test; ** p<0.01; p=0.005 and p=0.002 for target offsets and onsets, respectively, p=0.19 for offsets-onsets difference). E. Correlation between MEG power in the cluster and log(RT) for separate distributions and average RT; permutation tests. All *** p <0.001, all offset-onset differences p>0.05 (lowest was p=0.21).

The surprise-related cluster for target offsets and onsets both exhibited a bimodal in the frequency domain, similar to the pooled analysis (Figure 5; compare to Figure 4A): next to the peak around 20 Hz just after response, an additional peak was evident in the lowest frequency bin resolved (5 Hz). For offsets, the effect was quite sustained in time (-0.25 to 0.5s around response); the topography showed peaks over parietal and occipital cortex and over left frontal cortex (Figure 5A). By contrast, the cluster for target onsets was more confined in time (with a sharp peak ~0.1s after report) and a different topography that peaked over central parietal cortex (Figure 5B). Taken together, our results suggest that perceptual surprise about both target on- and offsets elicited cortical transients in the beta-band. We consider them general dynamical correlates of temporal surprise monitoring. In addition, stimulus changes seem to have recruited additional processes expressed in the very low (<= 5 Hz) frequency range.

Finally, we asked whether the trial-to-trial fluctuations in beta-power modulations also predicted trial-to-trial variations in subjects' (log-transformed) RTs. Here, we used the Pearson correlation values (i.e., without regressing out RT; *Materials and Methods*). Just as surprise, beta-power in the cluster also robustly predicted RT (Figure 4E). These correlations were negative, as expected based on the negative correlation between surprise and MEG-power (Figure 4A). We also compared the strength of this correlation between MEG-power and RT to the strength of the correlation between surprise and RT across, this correlation between correlations was positive, but not significant (r=0.19, p=0.33).

 No robust correlations between MEG baseline power and entropy
We did not find any evidence for a correlation of the raw baseline (-0.5 to 0 s with
respect to stimulus change) MEG power with uncertainty, as measured in entropy.
Correlations between entropy and MEG power spectra in the time window before
stimulus change did not result in any significant (sensor-frequency) clusters that
survived multiple-comparison correction (data not shown). It is likely that this lack of
robust correlation reflected the continuous reduction in trial-to-trial variations of
entropy over the course of each block (Figure 2C).
724

725
726 **Figure 5: Separate correlations for offsets and onsets.** Correlation analysis and cluster-based
727 statistics performed separately for target offsets (**A**) and target onsets (**B**); p=0.001 for both analyses.
728

Discussion

In this study, we comprehensively mapped cortical transients elicited by surprise
about the timing of sensory events. We used a Bayesian updating model to estimate
trial-to-trial variations of surprise and correlated these to subjects' behavior as well as
to neural dynamics, across the cortical surface. The model-derived surprise
estimates predicted across-trial and environment variations in RT. The surprise
estimates also predicted transient suppressions of low-frequency and beta-band
power in a widespread network comprising motor-, prefrontal and parietal cortical
regions, predominantly in the left hemisphere. The model-derived surprise estimates
were more closely related to both behavior and cortical dynamics than the mere trial-
to-trial variations in externally observable interval timings.
The signatures of surprise we uncovered in the beta frequency band were
quite similar around target on- and offset (Figure 5). This stands in sharp contrast to
the opposite beta-band modulation during (illusory or veridical) target
disappearances and reappearances, proposed to reflect a decision-related feedback
signal to in visual cortex (Meindertsma et al., 2017). The beta-band transients
identified here likely reflected a distinct process that did not encode the content of the
perceptual change, but rather the level of surprise about it.
One possibility is that surprise is computed in those fronto-parietal cortical
networks exhibiting the surprise-related modulation of beta-oscillations observed
here. Another possibility is that the surprise-related modulations are inherited from
other regions projecting to those fronto-parietal networks. Indeed, neuromodulatory
brainstem systems are a prominent candidate source. In particular the dopaminergic
and noradrenergic systems are driven by temporal expectation and surprise (Aston-
Jones and Cohen, 2005; Dayan and Yu, 2006; Fiorillo et al., 2008). Further, there is

mounting evidence for a link between neuromodulation and beta-band power in
visual cortex (Belitski et al., 2008; Donner and Siegel, 2011; Safaai et al., 2015;
Zaldivar et al., 2018).
Specifically, phasic responses in dopaminergic nuclei, encode not only
reward, but also the expected timing of reward arrival. The strength of these phasic
neuronal responses inversely scales with predictability of the timing of reward, in line
with encoding surprise about reward arrival, and it also predicted behavioral
anticipation of reward (i.e. licking behavior) in monkeys (Fiorillo et al., 2008). Our
current study complements this previous work, by unraveling the cortex-wide
dynamics elicited by surprising events. Our design did not involve rewards but rather
neutral, yet behaviorally relevant sensory events.
In a previous report based on the same data set as the current one
(Kloosterman et al., 2015a), we showed that mean pupil dilation responses during
the perceptual changes scaled in amplitude across the three environments in line
with mean surprise as shown in the current Figure 2H. Pupil dilation is closely linked
to phasic responses in neuromodulatory brain systems, in particular the
noradrenergic locus coeruleus (Joshi et al., 2016; Reimer et al., 2016; de Gee et al.,
2017). Thus, if the surprise-related modulations of cortical activity observed here
were driven by phasic neuromodulation, one would expect to find correlations
between single-trial pupil responses and surprise (Preuschoff et al., 2011; Nassar et
al., 2012). Due to the sluggish dynamics of the peripheral pupil apparatus (Hoeks
and Levelt, 1993; De Gee et al., 2014), testing for trial-by-trial correlations between
pupil dilations and surprise (or, likewise, between baseline pupil diameter and
uncertainty) in our experiment requires dedicated analysis approaches that tease
apart fluctuating baseline levels and responses evoked by individual events. Using a
general linear model (Hoeks and Levelt, 1993; De Gee et al., 2014), we failed to
obtain reliable single-trial pupil responses and correlations to single-trial surprise
(data not shown). This failure was likely, at least in part, due to the rapid nature of the
current experimental design. Future work should use more widely spaced intervals to
test whether pupil dilations reflect trial-to-trial variations of surprise.
Our current study provides a comprehensive picture of the cortical transients
elicited by surprise, by systematically mapping these transients across the cortical
surface and time-frequency plane. Previous work in humans has also studied neural
correlates of model-derived measures of surprise, although this entailed surprise
about stimulus identity, and not timing. Electrophysiological work found surprise
about cue identity to modulate the P3 component of the EEG event-related potential
as well as motor cortical excitability (Bestmann et al., 2008; Mars et al., 2008).
Functional magnetic resonance imaging work linked surprise about the spatial
location of stimuli to transients in posterior parietal cortex (O'Reilly et al., 2013). An
EEG study dissociated oscillatory neural signatures of surprise and evidence
accumulation (Gould et al., 2012). This latter study also found surprise-related
modulation of beta-band power primarily at frontal and parieto-occipital electrodes,
but the underlying cortical distribution was not estimated. Future studies of surprise in
other domains (e.g. about cue identity) should use a similar approach to assess if
surprise-related cortical transients are domain-general or -specific. Further,
simultaneous EEG and MEG recordings (Schurger et al., 2015) are necessary to
unravel the relationship between surprise-linked modulations of fronto-parietal beta-
band oscillations and of the P3-component.
Another line of work has investigated the functional role of externally
entrained low-frequency oscillations in temporal expectation. For fixed intervals,
alpha phase in sensory cortices was found to be predictive of expected time of target
arrival and lowered the threshold for sensory detection (Lakatos et al., 2008; Cravo et

al., 2011, 2013; Rohenkohl and Nobre, 2011). Alpha oscillations might reflect
rhythmic fluctuations in cortical excitability, entrained by rhythmic sensory input,
which aids stimulus processing and perceptual performance (Schroeder and Lakatos,
2009). The high variability in interval durations (see Figure 1B,C inset) might explain
the lack of alpha-band effects in our study. First, the range of possible durations was
too broad to form predictions that fall within a specific phase of an alpha cycle.
Second, even when oscillatory phase was modulated by temporal expectation in our
task, the trial-to-trial variability would make it difficult to align trials and make these
modulations visible.

It is tempting to relate our results to conceptual accounts of the functional role
of beta-band oscillations in the brain (Engel and Fries, 2010; Spitzer and Haegens,
2017). One account (Engel and Fries, 2010) holds that beta-band oscillations help
maintain the current sensorimotor or cognitive state (termed the 'status quo').
Another account (Spitzer and Haegens, 2017) holds that beta-band oscillations help
activate the currently relevant task sets. In both frameworks, the need for maintaining
the current status quo, or task set, is low when surprise (the violation of expectation,
or probability of change in the environment) is high, in line with our observation of a
suppression of beta-band oscillations under high surprise.

While our current work presents an important first step towards unraveling the
modulation of cortical dynamics by surprise, it is limited in that we only studied
environments with constant statistical structure within each block. Once a posterior
distribution has been learned, there remains no unexpected uncertainty, only
expected uncertainty (Yu and Dayan, 2005). By contrast, the statistical structure of
natural environments is often volatile. Richer experimental designs, that are volatile
and include unmarked changes, allow for probing into richer, presumably hierarchical
dynamics (Sugrue et al., 2004; Nassar et al., 2012; Meyniel et al., 2015). A more
volatile task-environment would also lead to an increase in trial-to-trial variability of
our entropy measure, providing a better-suited context to study the effects of this
type of uncertainty on cortical processing. Our ongoing work aims to push beyond
these limits by using richer environmental statistics that require more complex
inference processes.

To conclude, we here uncovered a novel signature of temporal surprise that
affected an elementary perceptual decision (target detection) and was characterized
by a temporally focal, but spatially widespread, modulation of cortical population
activity. This modulation might be instrumental in translating inferences about the
behaviorally-relevant temporal structure into its consequences for action.
842

References

Andrieu C, de Freitas N, Doucet A, Jordan MI (2003) An Introduction to MCMC for
Machine Learning. Mach Learn 50:5–43.
Aston-Jones G, Cohen JD (2005) An integrative theory of locus coeruleus-
norepinephrine function: adaptive gain and optimal performance. Annu Rev
Neurosci 28:403–450.
Belitski A, Gretton A, Magri C, Murayama Y, Montemurro MA, Logothetis NK, Panzeri
S (2008) Low-frequency local field potentials and spikes in primary visual cortex
convey independent visual information. J Neurosci 28:5696–5709.
Bestmann S, Harrison LM, Blankenburg F, Mars RB, Haggard P, Friston KJ, Rothwell
JC (2008) Influence of uncertainty and surprise on human corticospinal
excitability during preparation for action. Curr Biol 18:775–780.
Bonneh YS, Cooperman A, Sagi D (2001) Motion-induced blindness in normal
observers. Nature 411:798–801.
Bouret S, Sara SJ (2005) Network reset: A simplified overarching theory of locus

coeruleus noradrenaline function. Trends Neurosci 28:574–582.

Burnham KP, Anderson DR (2004) Multimodel inference: Understanding AIC and
BIC in model selection. Sociol Methods Res 33:261–304.

Cravo a. M, Rohenkohl G, Wyart V, Nobre a. C (2011) Endogenous modulation of
low frequency oscillations by temporal expectations. J Neurophysiol 106:2964–
2972.

Cravo AM, Rohenkohl G, Wyart V, Nobre AC (2013) Temporal expectation enhances
contrast sensitivity by phase entrainment of low-frequency oscillations in visual
cortex. J Neurosci 33:4002–4010.

Dayan P, Yu AJ (2006) Phasic norepinephrine: a neural interrupt signal for
unexpected events. Network 17:335–350.

de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH
(2017) Dynamic modulation of decision biases by brainstem arousal systems.
Elife 6:e23232.

De Gee JW, Knapen T, Donner TH (2014) Decision-related pupil dilation reflects
upcoming choice and individual bias. Proc Natl Acad Sci U S A 11:618–625.

Donner TH, Siegel M (2011) A framework for local cortical oscillation patterns.
Trends Cogn Sci 15:191–199.

Donner TH, Siegel M, Fries P, Engel AK (2009) Buildup of choice-predictive activity
in human motor cortex during perceptual decision making. Curr Biol 19:1581–
1585.

Efron B, Tibshirani RJ (1998) An Introduction to the Bootstrap. Boca Raton, FL:
Chapman & Hall/CRC Press.

Engel AK, Fries P (2010) Beta-band oscillations-signalling the status quo? Curr Opin
Neurobiol 20:156–165.

Feller W (1959) An Introduction to Probability Theory and Its Applications (Shewhart
WE, WIlks SS, eds)., Second. Wiley Publication in Statistics.

Fiorillo CD, Newsome WT, Schultz W (2008) The temporal precision of reward
prediction in dopamine neurons. Nat Neurosci 11:966–973.

Fisher RA (1915) Frequency Distribution of the Values of the Correlation Coefficient
in Samples from an Indefinitely Large Population. Biometrika 10:507.

Gibbon J, Malapani C, Dale CL, Gallistel C (1997) Toward a neurobiology of temporal
cognition: advances and challenges. Curr Opin Neurobiol 7:170–184.

Glaze CM, Kable JW, Gold JI (2015) Normative evidence accumulation in
unpredictable environments. Elife 4:1–27.

Gold JI, Shadlen MN (2007) The neural basis of decision making. Annu Rev Neurosci
30:535–574.

Gould IC, Nobre AC, Wyart V, Rushworth MFS (2012) Effects of decision variables
and intraparietal stimulation on sensorimotor oscillatory activity in the human
brain. J Neurosci 32:13805–13818.

Gross J, Kujala J, Hamalainen M, Timmermann L, Schnitzler A, Salmelin R (2001)
Dynamic imaging of coherent sources: Studying neural interactions in the
human brain. Proc Natl Acad Sci U S A 98:694–699.

Hipp JF, Hawellek DJ, Corbetta M, Siegel M, Engel AK (2012) Large-scale cortical
correlation structure of spontaneous oscillatory activity. Nat Neurosci 15:884–
890.

Hoeks B, Levelt WJM (1993) Pupillary dilation as a measure of attention : A
quantitative system analysis. Behav Res Methods, Instruments, Comput 25:16–
26.

Hollerman JR, Schultz W (1998) Dopamine neurons report an error in the temporal
prediction of reward during learning. Nat Neurosci 1:304–309.

Joshi S, Li Y, Kalwani RM, Gold JI (2016) Relationships between Pupil Diameter and

Neuronal Activity in the Locus Coeruleus , Colliculi , and Cingulate Cortex Article
Relationships between Pupil Diameter and Neuronal Activity in the Locus
Coeruleus , Colliculi , and Cingulate Cortex. Neuron 89:1–14.

Kloosterman N a., Meindertsma T, van Loon AM, Lamme V a. F, Bonneh YS, Donner
TH (2015a) Pupil size tracks perceptual content and surprise. Eur J Neurosci
41:1068–1078.

Kloosterman NA, Meindertsma T, Hillebrand A, Van Dijk BW, Lamme VAF, Donner
TH (2015b) Top-down modulation in human visual cortex predicts the stability of
a perceptual illusion. J Neurophysiol 113:1063–1076.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE (2008) Entrainment of
Neuronal Oscillations as a Mechanism of Attentional Selection. Science (80-)
320:110–113.

Lee MD, Wagenmakers E-J (2013) Bayesian Cognitive Modeling: A Practical Course,
1st ed. New York: Cambridge University Press.

Luce RD (1986) Response Times: Their Role in Inferring Elementary Mental
Organization, 1st ed. Oxford University Press, USA;

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-
data. J Neurosci Methods 164:177–190.

Mars RB, Debener S, Gladwin TE, Harrison LM, Haggard P, Rothwell JC, Bestmann
S (2008) Trial-by-trial fluctuations in the event-related electroencephalogram
reflect dynamic changes in the degree of surprise. J Neurosci 28:12539–12545.

Meindertsma T, Kloosterman NA, Nolte G, Engel AK, Donner TH (2017) Multiple
Transient Signals in Human Visual Cortex Associated with an Elementary
Decision. J Neurosci 37:5744–5757.

Meyniel F, Schlunegger D, Dehaene S (2015) The Sense of Confidence during
Probabilistic Learning: A Normative Account. PLOS Comput Biol 11:e1004305.

Nassar MR, Rumsey KM, Wilson RC, Parikh K, Heasly B, Gold JI (2012) Rational
regulation of learning dynamics by pupil-linked arousal systems. Nat Neurosci
15:1040–1046.

Nobre A, Correa a., Coull J (2007) The hazards of time. Curr Opin Neurobiol
17:465–470.

O'Reilly JX, Schüffelgen U, Cuell SF, Behrens TEJ, Mars RB, Rushworth MFS (2013)
Dissociable effects of surprise and model update in parietal and anterior
cingulate cortex. Proc Natl Acad Sci U S A 110:E3660-9.

Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open source
software for advanced analysis of MEG, EEG, and invasive electrophysiological
data. Comput Intell Neurosci 2011:156869.

Plummer M (2003) JAGS: A program for analysis of Bayesian graphical models using
Gibbs sampling. Proc 3rd Int Work Distrib Stat Comput (DSC 2003):20–22.

Preuschoff K, 't Hart BM, Einhäuser W (2011) Pupil Dilation Signals Surprise:
Evidence for Noradrenaline's Role in Decision Making. Front Neurosci 5:115.

Reimer J, Mcginley MJ, Liu Y, Rodenkirch C, Wang Q, Mccormick DA, Tolias AS
(2016) Pupil fluctuations track rapid changes in adrenergic and cholinergic
activity in cortex. Nat Commun 7:1–7.

Riecke L, Sack AT, Schroeder CE (2015) Endogenous Delta/Theta Sound-Brain
Phase Entrainment Accelerates the Buildup of Auditory Streaming. Curr Biol
25:3196–3201.

Rohenkohl G, Cravo AM, Wyart V, Nobre AC (2012) Temporal Expectation Improves
the Quality of Sensory Information. J Neurosci 32:8424–8428.

Rohenkohl G, Nobre a. C (2011) Alpha Oscillations Related to Anticipatory Attention
Follow Temporal Expectations. J Neurosci 31:14076–14084.

Safaai H, Neves R, Eschenko O, Logothetis NK, Panzeri S (2015) Modeling the

effect of locus coeruleus firing on cortical state dynamics and single-trial sensory
processing. Proc Natl Acad Sci 112:201516539.

Schroeder CE, Lakatos P (2009) Low-frequency neuronal oscillations as instruments
of sensory selection. Trends Neurosci 32:9–18.

Schurger A, Sarigiannidis I, Naccache L, Sitt JD, Dehaene S (2015) Cortical activity
is more stable when sensory stimuli are consciously perceived. Proc Natl Acad
Sci 112:E2083–E2092.

Siegel M, Donner TH (2010) Linking Band-Limited Cortical Activity to fMRI and
Behavior. In: Simultaneous EEG and fMRI. Recording, Analysis, and Application
(Ullsperger M, Debener S, eds), pp 1–23. New York: Oxford University Press.

Spitzer B, Haegens S (2017) Beyond the Status Quo: A Role for Beta Oscillations in
Endogenous Content (Re-) Activation. eneuro:ENEURO.0170-17.2017.

Sugrue LP, Corrado GS, Newsome WT (2004) Matching Behavior and the
Representation of Value in the Parietal Cortex. Science (80-) 304:1782–1788.

Sutton RS, Barto AG (1998) Reinforcement Learning : An Introduction, First.
Cambridge: MIT Press.

van Ede F, Niklaus M, Nobre AC (2017) Temporal Expectations Guide Dynamic
Prioritization in Visual Working Memory through Attenuated α Oscillations. J
Neurosci 37:437–445.

Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A (1997) Localization of brain
electrical activity via linearly constrained minimum variance spatial filtering.
IEEE Trans Biomed Eng 44:867–880.

Yu AJ, Dayan P (2005) Uncertainty, neuromodulation, and attention. Neuron 46:681–
692.

Zaldivar D, Goense J, Lowe SC, Logothetis NK, Panzeri S (2018) Dopamine Is
Signaled by Mid-frequency Oscillations and Boosts Output Layers Visual
Information in Visual Cortex. Curr Biol 28:224–235.e5.