A comprehensive model of DNA fragmentation for the preservation of High Molecular Weight DNA

Tomas Klingström[1,*], Erik Bongcam-Rudloff[1] and Olga Vinnere Pettersson[2]

[1] SLU-Global Bioinformatics Centre, Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Uppsala, 756 51, Sweden.
[2] Uppsala Genome Center, Science for Life Laboratory, Dept. of Immunology, Genetics and Pathology, Uppsala University, BMC, Box 815, SE-752 37 UPPSALA

* To whom correspondence should be addressed. Tel: +4618-672126; Fax: None; Email: tomas.klingstrom@slu.se

ABSTRACT

During DNA extraction the DNA molecule undergoes physical and chemical shearing, causing the DNA to fragment into shorter and shorter pieces. Under common laboratory conditions this fragmentation yields DNA fragments of 5-35 kilobases (kb) in length. This fragment length is more than sufficient for DNA sequencing using short-read technologies which generate reads 50-600 bp in length, but insufficient for long-read sequencing and linked reads where fragment lengths of more than 40 kb may be desirable.

This study provides a theoretical framework for quality management to ensure access to high molecular weight DNA in samples. Shearing can be divided into physical and chemical shearing which generate different patterns of fragmentation. Exposure to physical shearing creates a characteristic fragment length where DNA fragments are cut in half by shear stress. This characteristic length can be measured using gel electrophoresis or instruments for DNA fragment analysis. Chemical shearing generates randomly distributed fragment lengths visible as a smear of DNA below the peak fragment length. By measuring the peak of DNA fragment length and the proportion of very short DNA fragments both sources of shearing can be measured using commonly used laboratory techniques, providing a suitable quantification of DNA integrity of DNA for sequencing with long-read technologies.

INTRODUCTION

The structure of the DNA molecule has been known for more than 60 years and it is without a doubt the most intensively studied molecule in our bodies. However, there has been little need to evaluate the mechanisms of DNA fragmentation at length scales beyond a few thousand base pairs (Mulcahy *et al.*, 2016). With the rising importance of long-read sequencing technologies and mapping technologies utilising linked reads (Schadt *et al.*, 2010; Goodwin *et al.*, 2016) it is however evident that a greater understanding of DNA structural integrity and fragmentation must be achieved to support better quality management practices for high-molecular-weight DNA (HMW-DNA) (English *et al.*, 2012; Malentacchi *et al.*, 2015; Goodwin *et al.*, 2016). DNA is generally considered a relatively stable analyte as it is highly resistant to physical shear at the lengths required for PCR and short-read next generation sequencing (NGS) (Morente *et al.*, 2006; Lou *et al.*, 2014; Robb *et al.*, 2014; Ellervik and Vaught, 2015). Prevention of chemical degradation over short periods of time is achieved using routine procedures. Due to this perceived stability, the research on the preservation of HMW-DNA has long been neglected (Anchordoquy and Molina, 2007). Purity, yield and amplification success are seen as sufficient quality parameters when evaluating DNA sample quality making it hard to estimate if DNA in samples is suitable for long-read sequencing (Bustin *et al.*, 2009; Le Page *et al.*, 2013; Mathay *et al.*, 2016; Mulcahy *et al.*, 2016; Office of Biorepositories and Biospecimen Research *et al.*, n.d.).

In this study we create a comprehensive framework of the significant factors defining the fragment length distribution of DNA provided for sequencing. Biobank research and published method papers

provide valuable empirical data on fragmentation observed under common laboratory conditions. Yet, data is too sparse to untangle the potential pre-analytical variables and complex interdependencies necessary to identify the key molecular mechanisms contributing to fragmentation (Robb *et al.*, 2014; Malentacchi *et al.*, 2016a). Therefore, this work takes an approach where experimental data is assessed and theories are generated by a wide variety of fields affected by DNA fragmentation. By merging data from the following fields, we are thereby able to present a model explaining the key contributing factors to DNA fragmentation when preparing biospecimen for high throughput genomics research:

- Early research on the physical properties of DNA conducted in the years after identifying the structure of the DNA helix (Bowman and Davidson, 1972; Adam and Zimm, 1977; Dancis, 1978).

- Investigations into the chemical properties and degradation of DNA in vitro/in vivo (Lindahl, 1993) that ultimately led to the discovery of cellular repair mechanisms of DNA and the Nobel prize for these discoveries.

- Archaeological research investigating the decay of DNA over long periods of time and the decay products generated during the process. Indicating that the initial rate of decay is substantially higher than can be explained by depurination (Sawyer *et al.*, 2012) but that long term fragmentation follow a 1^{st} order reaction where depurination is a key driving factor (Overballe-Petersen *et al.*, 2012) which for genomic DNA is close to what we can expect from depurination at pH 7.5 (Allentoft *et al.*, 2012).

- Forensic research provides a baseline for DNA fragmentation patterns in different tissues under conditions where no stabilization of samples is provided (Bär *et al.*, 1988; Johnson and Ferris, 2002).

- The stability of plasmids, linearized plasmids and genomic DNA for pharmaceutical applications (Evans *et al.*, 2000). In particular, it should be noted that a similar survey to determine pharmacologically relevant factors of DNA degradation has been published (Rossmanith *et al.*, 2011). Under these conditions DNA degradation due to depurination was estimated to be the dominant factor for degradation of samples.

- Polymer science and physical chemistry studying the behaviour of polymers (Vanapalli *et al.*, 2006), and simulating electrostatic interactions *in silico* explaining the influence of folding and protection of DNA under high ionic-strength conditions (Cinque *et al.*, 2010; Jiang *et al.*, 2015).

A large number of potentially relevant chemical mechanisms for degradation have been identified in pharmacological research studying the degradation of DNA-based pharmaceuticals. For the purposes of DNA extraction, using aqueous solutions oxidation and depurination are estimated to be the relevant factors (Pogocki and Schöneich, 2000; Rossmanith *et al.*, 2011). DNA sequencing of archaeological findings support these conclusions as they indicate that even DNA retrieved in conditions far from optimal show a degradation pattern consistent with depurination being the main source of fragmentation (Allentoft *et al.*, 2012). By adjusting for this baseline of degradation other, more situational, factors for degradation such as the influence of specific DNA extraction can be identified and used to generate a minimal viable explanation for DNA fragmentation under laboratory conditions.

A MINIMAL EXPLANATION OF DNA FRAGMENTATION

When tracking DNA fragmentation in a laboratory environment the degradation process of DNA is initiated at the start of the warm ischemia time for tissue collection or blood draw (or equivalent) when collecting fluids (Johnson and Ferris, 2002; Betsou *et al.*, 2010).

Chemical factors

The disruption of the cellular environment triggers a number of processes leading to the degradation of DNA by oxidizing attacks and degradation by nucleases, unless preservation measures are taken (Elmore, 2007; Rossmanith *et al.*, 2011). This process can be halted through freezing or drying of samples. It can also be prevented by the addition of preservatives, in the form of chelating agents which bind metal ions that catalyse the production of hydroxyl radicals as well as Mg^{2+} ions in Mg^{2+}-dependent endonucleases (Lahiri and Schnabel, 1993; Rossmanith *et al.*, 2011). Chemical degradation by these pathways is the immediate threat to DNA integrity and may rapidly produce a smear of short fragments. At the same time, these factors make DNA vulnerable to physical stress by "nicking" (the introduction of single strand breaks in double stranded DNA), but this issue is easily dealt with by the addition of buffers containing chelators, or a combination of EDTA + ethanol, which are highly effective in preventing degradation by oxidizing attacks (Evans *et al.*, 2000). It should, however, be noted that EDTA on its own actually catalyses the formation of hydroxyl radicals and therefore increases degradation rates unless combined with a radical scavenger (Evans *et al.*, 2000).

The other major cause of chemical shearing of DNA is depurination followed by β-elimination which is an acid-catalysed first order reaction (An *et al.*, 2014). As the reaction is driven by protonation it is impossible to completely prevent it in aqueous solutions, and the complete dehydration of DNA requires encapsulation to protect the sample from atmospheric water as the DNA molecule binds water in its minor and major groove (Bonnet *et al.*, 2010). Storing samples at pH 8 (Brunstein, 2015), reducing molecular mobility and temperature, along with the avoidance of primary amines, can reduce the reaction rate to levels sufficiently low to ensure that DNA maintain integrity for most long-read purposes for centuries (An *et al.*, 2014).

Physical stress

Fragmentation due to physical stress has an immediate impact on HMW-DNA. Susceptibility to physical stress is directly dependent on the length and folding of DNA molecules (Anchordoquy and Molina, 2007; Cinque *et al.*, 2010; Lengsfeld *et al.*, 2011), and during extraction DNA is subjected to physical forces generating a peak abundance of 5-35 kb under common laboratory conditions (Malentacchi *et al.*, 2013, 2016b). Laminar flows have for a long time been seen as the main culprit behind DNA fragmentation (Dancis, 1978), but chemists specialising in the study of polymers in turbulent flow have provided convincing evidence indicating that turbulence dominates the influence of laminar flows under most conditions (Vanapalli *et al.*, 2006). These conclusions are further supported by the teams of Anchordoquy (Lengsfeld *et al.*, 2011) and Baigl (Cinque *et al.*, 2010) who have investigated the factors influencing the degradation of DNA for pharmaceutical applications (Anchordoquy and Molina, 2007).

These conclusions on physical shear stress significantly simplify the work to minimise the influence of physical shear in DNA processing, as it means that most DNA fragmentation occurs when the contour length of DNA exceeds the Kolmogorov length of a turbulent flow. The Kolmogorov length scale defines the smallest possible eddies occurring in a system before they dissipate into energy. To understand this connection between quantum mechanics and biology it is necessary to know that all turbulence can be modelled as consisting of small vortices that in turn consist of smaller vortices. At a certain threshold, it is no longer possible to divide vortices into smaller ones, and at the Kolmogorov length scale these distances are so small that the spatial movement of molecules turns into vibrations (*i.e.* heat). Calculations by Vanapalli *et al.* have proven that the pull of a DNA molecule stuck in two vortices at the same time is sufficient to break the strand in half and that this is the primary cause of fragmentation under most hydrodynamic conditions (Vanapalli *et al.*, 2006). During sample processing fragmentation occurs over time until all DNA molecules have been degraded to a length where all fragments are short enough to remain within a single eddy, which is consistent with experimental results with commonly used laboratory equipment, such as vortex mixers and pipettes (Yoo *et al.*,

2011), and results suggest that the breakage rate is a function of the shear rate rather than the shear stress (Bowman and Davidson, 1972; Meacle *et al.*, 2007).

Freezing and long-term storage

Finally, DNA can be fragmented by freezing, thawing, and long-term storage. Repeated freeze-thaw cycles are the most significant source of extensive damage but continued chemical reactions while the samples are frozen also play a role. Researchers frequently assume that nothing happens to DNA inside a frozen sample (Hubel *et al.*, 2014), but in reality both chemical processes and physical movement occur even at low temperatures. Between -39 and -50 °C, the DNA molecule becomes rigid and brittle as it enters a glassy state (Norberg and Nilsson, 1996) while protons in the water remain mobile even at temperatures close -268 °C (Benton *et al.*, 2016). There is a lack of long term studies at -80 °C or –196 °C, and the complex behaviour of water at low temperatures makes it impossible to estimate at what temperatures chemical activity generated by protons ceases to occur. Using the Arrhenius equation to estimate an upper bound of activity we can, however, estimate that a strand break occur at a rate of approximately once every 5 seconds in a 3.2 billion base pair genome at 25 °C and pH 8. Storage in a -80 °C freezer reduces the rate to less than once per 10 000 years (An et al., 2014).

During freezing and thawing two other factors may contribute to the fragmentation of DNA (Anchordoquy *et al.*, 2001; Shao *et al.*, 2012). As a sample cools beneath the freezing temperature of fluids, it will undergo nucleation where water molecules (or other fluids) start assembling into crystals. Solutes are not incorporated into the solid ice crystals, resulting in the sample separating into solid components and unfrozen liquid components with high concentration of solutes. Creating an environment where two sources of DNA shearing has been proposed (Anchordoquy *et al.*, 2001; Brunstein, 2015):

- As ice crystals form, solutes are crowded into gaps with increasingly high concentrations of solutes, thereby drastically increasing the potential for oxidising attacks and protonation of the DNA chain, leading to chemical shearing of the DNA polymer.

- As ice crystals form, DNA polymers are partially embedded within the ice crystals. The mechanical stress could thereby lead to the breakage of DNA molecules with a higher likelihood of breakage occurring in longer polymers.

Based on laboratory experiments, it is evident that chemical shearing is present under certain circumstances (Davis *et al.*, 2000) but can be completely avoided as highly purified samples of short DNA fragments (<2 kb) have remained intact during repeated freeze-thawing in several studies (Ross *et al.*, 1990; Thornton *et al.*, 2005; Integrated DNA Technologies, 2014; Rossmanith *et al.*, 2011). Rossmanith *et al.* could even show that short amplicons (274 bp) remained intact while the same amplicons when located within longer genomic sequences from *Salmonella typhimurium* (4 857 kb) and *Listeria monocytognes* (1 160 kb) were sheared during their freeze-thaw experiments (Rossmanith *et al.*, 2011).

Research in cryopreservation of oocytes, embryos and sperm (Amidi *et al.*, 2016; Kopeika *et al.*, 2015) suggest that shearing of DNA during freeze-thaw can be reduced by the addition of antioxidants, indicating that oxidation of DNA by radicals may, at least *in vivo*, be a significant cause of DNA fragmentation. Such conclusions must however be considered highly tentative as there is a substantial differences in the chemical environment inside the cell compared to the *in vitro* conditions covered by this study. The safeguards necessary to avoid physical shearing by ice crystal formation are also poorly understood. The addition of glycerol (Calcott and Gargett, 1981; Schaudien *et al.*, 2007) is known to protect DNA during freeze-thaw cycles but we have been unable to determine why samples seemingly subjected to physical shearing may converge to lengths anywhere between 2000 bp and

25 000 bp in different studies (Röder *et al.*, 2010; Ross *et al.*, 1990; Schaudien *et al.*, 2007; Shao *et al.*, 2012).

DISCUSSION

Long-read sequencing enables studies of complex genomic events, e.g. structural re-arrangements, copy number variations, repeat expansions, etc. Hence, it is obvious that the long-read technologies, such as PacBio, 10x Genomics and Oxford Nanopore will be increasingly popular. This constitutes a challenge for laboratories and biobanks, as best practice routines for storage of biological samples and DNA extractions need to be adjusted in order to allow delivery of HMW-DNA suitable for long-read sequencing (Malentacchi *et al.*, 2016a). Currently, the Oxford Nanopore MinION, GridION and PromethION are able to sequence reads up to 1 Mbp in length (Oxford Nanopore Technologies, 2017). The optimal results obtained by 10x Chromium systems are obtained if the sample contains a high number of molecules above 50 kbp, and up to a few hundred kilobases, and the PacBio systems can produce reads up to 60-70 kbp (Bickhart et al., 2017; Lok et al., 2017; Paajanen et al., 2017). These numbers will most likely rise due to technology development, and biobanks should be ready to be able to provide researchers with DNA samples of adequate quality and quantity.

Current biobanking practices provide DNA with a high yield and purity when tested, but suffer from a high level of variability in regards to DNA integrity (Malentacchi *et al.*, 2013, 2015). Improved protocols and subtle alterations of protocols can, however, provide significant improvements. The SPIDIA-ring trials show that biobanks are able to rapidly improve their operations when provided with sufficient guidance (Malentacchi *et al.*, 2016a). It is therefore important that an acceptable definition of DNA fragmentation and robust units of measurement are made available to the research community to improve the availability of HMW-DNA for researchers. In this regard the SPIDIA project (Malentacchi *et al.*, 2013) and the Global Genome Biodiversity Network (GGBN) (Mulcahy *et al.*, 2016) have provided pioneering work on the subject on how to objectively evaluate and compare DNA fragmentation in samples using gel-electrophoresis and a *Hin*dIII ladder. Both projects propose a "greater than X kb" approach where the percentage of DNA above a certain "DNA threshold" is measured and used as an indicator of DNA integrity. The SPIDIA project used the 4.36 kb size marker as its X-value and supplemented it with the measure peak intensity to provide an assessment of DNA integrity (Ciniselli *et al.*, 2015). The GGBN on the other hand decided to forego the usage of a peak intensity and instead proposed the X value to be dependent on the intended utilization of samples but with the 9.416 bp size marker as a default as it is substantially longer than standard HTS reads and would be an appropriate minimum for long-read sequencing (Mulcahy *et al.*, 2016).

The influence of physical shear stress and chemical degradation are, however, substantially different from each other. We therefore believe that two separate measures of DNA fragmentation are necessary to help researchers to evaluate the integrity of their DNA in an adequate manner.

The characteristic fragment length (CFL): The most abundant fragment length of the sample. In gel-electrophoresis (both conventional and capillary), this value corresponds to the peak intensity of the signal.

The smear ratio (SR): The proportion of DNA with a fragment length of less than 500 bp. This value can be easily calculated using the imageJ open source software to integrate over the signal intensity curve as described by Mulcahy *et al.* (Mulcahy *et al.*, 2016).

The characteristic fragment length is routinely reported when DNA integrity is assessed using gel-electrophoresis. Based on the conclusions drawn in this paper it can be deemed suitable as a direct but conservative estimate of how physical shear stress limits the DNA fragment length in a sample.

The smear ratio provides an estimate of the proportion of DNA that has been shredded to short fragment lengths due to freeze-thaw shearing or chemical degradation. From a technical perspective the work to produce a "less than X kb" measurement is identical to the "greater than X kb" proposed by

GGBN, but it provides researchers with a more informative measurement. Samples with highly preserved content will have a lopsided distribution around the characteristic fragment length with a long tail of shorter DNA fragments. Changes in the maximum fragment length by physical shear stress will easily be detected by a shift in peak fragment length and a "less than X kb" measurement provides a corresponding estimate of the amount of DNA being sheared to very short fragments due to chemical degradation and/or freeze-thaw.

These two measurements therefore provide laboratories with the means to monitor changes in processing and allow end users to assess the potential of obtaining sufficiently HMW DNA from a resource. For advanced users and quality management it is, however, still attractive to also make the entire fragment distribution plot available (Permenter *et al.*, 2015). Based on the experience of National Genomics Infrastructure - Sweden, the following instrumentation can provide researchers with this type of information: Agilent Bioanalyser (Agilent Technologies), Fragment Analyzer (Advanced Analytical) and Femto Pulse (Advanced Analytical), where two latter systems can process at least 8 samples in parallel providing both the characteristic fragment length and smear ratio in a single analysis (figure 1).

The first megabase plus read was recently announced using the MinION technology from Oxford Nanopore (Oxford Nanopore Technologies, 2017). At such long-read lengths, currently standard methods require modifications and gentle DNA extraction methods such as phenol–chloroform extraction (for reference the Rad003 protocol used by Loman lab to extract DNA for generating the megabase-length read is available at http://lab.loman.net/protocols/). The shearing of DNA caused by the extraction step itself is, at least in high-throughput laboratories such as biobanks, likely to generate fragments with a characteristic length of more than 50 kb, while turbulent forces during other parts of the process restrict the fragment length to 5-35 kb (Shao *et al.*, 2012; Malentacchi *et al.*, 2015). Among commonly used extraction methods, precipitation based DNA extraction generate the highest molecular weight of DNA (Malentacchi *et al.*, 2015) but variations in fragment length (Shao *et al.*, 2012) and other quality parameters mean that kit-by-kit comparisons will be necessary for the selection of suitable best practice procedures for DNA extraction depending on the intended usage of the samples.

CONCLUSIONS

Based on the model presented above, laboratories are advised to focus on three areas if they wish to increase the integrity of extracted DNA samples (table 1). During the retrieval of samples it is important to quickly stabilise the sample by preventing enzymatic degradation and generation of free radicals within the sample. After the cells have been lysed it is important to handle the sample with outmost care to minimise physical shearing (gentle mixing, slow pipetting, using wide-bore pipette tips, etc) and then store DNA at low temperature or in a completely dehydrated state to avoid depurination.

Physical degradation by shear stress is mainly caused by microscale turbulence. The turbulence fragments DNA until fragments reach a characteristic peak length. Physical forces are strongest in the centre of the DNA polymer but nicks and random events may cause strand breaks elsewhere. This generates a characteristic peak length roughly corresponding to half the length of the smallest fragment length prone to degradation in the system. When measured by gel electrophoresis this generates a characteristic pattern with a peak intensity of fluorescence followed by a long tail smear below the peak and a shorter upstream section of fragments that, due to random/chemical effects, reached a length protected from turbulence but still longer than half the critical length.

Freeze-thaw events may contribute significant damage to DNA but is highly dependent on parameters such as polymer size, ion strength, rate of temperature drops, presence of nicks etc. Currently, there is no generalized theory to predict fragmentation but preservation with means such as storage in 50 % glycerol has been shown to prevent shearing. Furthermore, it seems that variations in upstream

protocols have a profound impact on the vulnerability to freeze-thaw cycles as evidenced by in-house testing not identifying any fragmentation of samples exposed to repeated freeze-thaw cycles.

To support further work on maintaining DNA integrity it is important that researchers report relevant data on the integrity of DNA. DNA integrity data is often omitted or reported using inadequate measurements such as electrophoresis with *Hind*III digested λ DNA, which does not provide relevant information above 23 kb. Current literature on maintaining longer DNA for high throughput applications is therefore limited and this study provides a foundation for further work in quality management and process optimisation for the production of HMW-DNA for sequencing.

Figure 1, example of high quality DNA with minimal smear (section B) and a characteristic fragment length of 129 292 bp as measured by the Femtopulse (Advanced Analytical). High quality samples often contain a plateau on the right hand side of the characteristic fragment length, meaning that the CFL should not be interpreted as a measure of the longest fragment length available for sequencing. But rather as a measurement of the turbulent forces restricting the fragment length of the sample. To provide longer reads suitable for long-read sequencing or linked-read technology, the DNA extraction process can be altered to reduce physical shearing. With continuous monitoring it is also possible to quickly any inadvertent changes influencing DNA integrity as chemical shearing an increased smear ratio or a drop in characteristic fragment length is immediately measured using the above described technique.

Table 1, key parameters for identifying causes of DNA fragmentation and how to reduce it.

	Source	Time scale	Mechanism	Threat profile	Preventive measure	Damage pattern
Sample collection	Enzymatic degradation	Immediate	Naturally occuring DNAses attack the DNA polymer	An immediate threat to the integrity of DNA unless enzymes are inhibited or destroyed.	Addition of Proteinase degrading the protein or chelators binding metal ions functioning as co-factors.	DNA fragmentation with specific cleavage sites
	Oxidation	Seconds to years	Metal ions catalysing the production of hydroxyl radicals	An immediate threat to the integrity of DNA unless chelating agents and/or a radical scavenger is added.	Addition of chelators or EDTA + Ethanol	Smearing
Processing	Turbulence	Immediate	Physical shearing	Poorly optimised machinery and careless handling of sample will cause physical shearing of the DNA.	Freezing, dehydration or maintaining a high P/E-value	Characteristic fragment length
	Depurination	Weeks to years	Protonation of purines lead to depurination followed by beta elimination	A slow reaction that may cause issues under acidic conditions or long term storage at room temperature.		Smearing
Storage	Freeze-thaw	Immediate	Physical shearing	The damage mechanisms of freeze-thawing DNA are still poorly understood *in vitro* but damage is proportional to the number of freeze-thaw cycles the sample has undergone. Flash freezing or the addition of a compound allowing samples to vitrify rather than form ice crystals seem effective.	Addition of a protective agent such as glycerol or flash freezing to low temperatures in order to prevent large ice crystals from forming.	Characteristic fragment length
	Freeze-thaw	Immediate	Crowding of solutes leading to hydrolysis and/or oxidation	The damage mechanisms of freeze-thawing DNA are still poorly understood in vitro but damage is proportional to the number of freeze-thaw cycles the sample has undergone. Flash freezing or the addition of a compound allowing samples to vitrify rather than form ice crystals seem effective as solutes are not crowded into the vicinity of the DNA polymer. Minimising the number of freeze-thaw cycles is also an obvious solution.	Maintaining a high purity of extracted DNA and possibly addition of protective agents.	Smearing
	Depurination	Weeks to years	Protonation of purines leading to depurination followed by beta elimination	A slow reaction that may cause issues under acidic conditions or long term storage at high temperature or with insufficient removal of water	Low temperature storage or complete dehydration.	Smearing

ACKNOWLEDGEMENT

The authors would like to thank Dr Jaan Hong for his help with drawing our blood as well as PhD candidate Ida Höijer and Dr Oskar Karlsson for their help in the laboratory that have allowed us to run confirmatory tests.

FUNDING

This work was financed by the BBMRI-LPC and the B3Africa projects. BBMRI-LPC is supported by the European Community's 7th Framework Programme (FP7/20072013) grant agreement no. 313010, B3Africa is supported by the European Union s Horizon 2020 research and innovation programme under grant agreement No 654404. Work performed at Uppsala Genome Center – NGI Uppsala has been funded by RFI – VR and SciLifeLab Sweden.

REFERENCES

Adam, R.E., Zimm, B.H., 1977. Shear degradation of DNA. Nucleic Acids Res. 4, 1513–1538. https://doi.org/10.1093/nar/4.5.1513

Allentoft, M.E., Collins, M., Harker, D., Haile, J., Oskam, C.L., Hale, M.L., Campos, P.F., Samaniego, J.A., Gilbert, M.T.P., Willerslev, E., Zhang, G., Scofield, R.P., Holdaway, R.N., Bunce, M., 2012. The half-life of DNA in bone: measuring decay kinetics in 158 dated fossils. Proc. R. Soc. B Biol. Sci. 279, 4724–4733. https://doi.org/10.1098/rspb.2012.1745

Amidi, F., Pazhohan, A., Shabani Nashtaei, M., Khodarahmian, M., Nekoonam, S., 2016. The role of antioxidants in sperm freezing: a review. Cell Tissue Bank. 17, 745–756. https://doi.org/10.1007/s10561-016-9566-5

An, R., Jia, Y., Wan, B., Zhang, Y., Dong, P., Li, J., Liang, X., 2014. Non-Enzymatic Depurination of Nucleic Acids: Factors and Mechanisms. PLoS ONE 9, e115950. https://doi.org/10.1371/journal.pone.0115950

Anchordoquy, T.J., Allison, S.D., Molina, M. d.C, Girouard, L.G., Carson, T.K., 2001. Physical stabilization of DNA-based therapeutics. Drug Discov. Today 6, 463–470. https://doi.org/10.1016/S1359-6446(01)01739-1

Anchordoquy, T.J., Molina, M.C., 2007. Preservation of DNA. Cell Preserv. Technol. 5, 180–188. https://doi.org/10.1089/cpt.2007.0511

Bär, W., Kratzer, A., Mächler, M., Schmid, W., 1988. Postmortem stability of DNA. Forensic Sci. Int. 39, 59–70. https://doi.org/10.1016/0379-0738(88)90118-1

Benton, O., Sikora, O., Shannon, N., 2016. Classical and quantum theories of proton disorder in hexagonal water ice. Phys. Rev. B 93. https://doi.org/10.1103/PhysRevB.93.125143

Betsou, F., Lehmann, S., Ashton, G., Barnes, M., Benson, E.E., Coppola, D., DeSouza, Y., Eliason, J., Glazer, B., Guadagni, F., Harding, K., Horsfall, D.J., Kleeberger, C., Nanni, U., Prasad, A., Shea, K., Skubitz, A., Somiari, S., Gunter, E., International Society for Biological and Environmental Repositories (ISBER) Working Group on Biospecimen Science, 2010. Standard Preanalytical Coding for Biospecimens: Defining the Sample PREanalytical Code. Cancer Epidemiol. Biomarkers Prev. 19, 1004–1011. https://doi.org/10.1158/1055-9965.EPI-09-1268

Bickhart, D.M., Rosen, B.D., Koren, S., Sayre, B.L., Hastie, A.R., Chan, S., Lee, J., Lam, E.T., Liachko, I., Sullivan, S.T., Burton, J.N., Huson, H.J., Nystrom, J.C., Kelley, C.M., Hutchison, J.L., Zhou, Y., Sun, J., Crisà, A., Ponce de León, F.A., Schwartz, J.C., Hammond, J.A., Waldbieser, G.C., Schroeder, S.G., Liu, G.E., Dunham, M.J., Shendure, J., Sonstegard, T.S., Phillippy, A.M., Van Tassell, C.P., Smith, T.P.L., 2017. Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. Nat. Genet. 49, 643–650. https://doi.org/10.1038/ng.3802

Bonnet, J., Colotte, M., Coudy, D., Couallier, V., Portier, J., Morin, B., Tuffet, S., 2010. Chain and conformation stability of solid-state DNA: implications for room temperature storage. Nucleic Acids Res. 38, 1531–1546. https://doi.org/10.1093/nar/gkp1060

Bowman, R.D., Davidson, N., 1972. Hydrodynamic shear breakage of DNA. Biopolymers 11, 2601–2624. https://doi.org/10.1002/bip.1972.360111217

Brunstein, J., 2015. Freeze-thaw cycles and nucleic acid stability: what's safe for your samples? MLO. Med. Lab. Obs. 47, 44–45.

Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J., Wittwer, C.T., 2009. The MIQE Guidelines:

Minimum Information for Publication of Quantitative Real-Time PCR Experiments. Clin. Chem. 55, 611–622. https://doi.org/10.1373/clinchem.2008.112797

Calcott, P.H., Gargett, A.M., 1981. Mutagenicity of freezing and thawing. FEMS Microbiol. Lett. 10, 151–155. https://doi.org/10.1111/j.1574-6968.1981.tb06227.x

Ciniselli, C.M., Pizzamiglio, S., Malentacchi, F., Gelmini, S., Pazzagli, M., Hartmann, C.C., Ibrahim-Gawel, H., Verderio, P., 2015. Combining qualitative and quantitative imaging evaluation for the assessment of genomic DNA integrity: The SPIDIA experience. Anal. Biochem. 479, 60–62. https://doi.org/10.1016/j.ab.2015.03.023

Cinque, L., Ghomchi, Y., Chen, Y., Bensimon, A., Baigl, D., 2010. Protection of Human Genomic DNA from Mechanical Stress by Reversible Folding Transition. ChemBioChem 11, 340–343. https://doi.org/10.1002/cbic.200900734

Dancis, B.M., 1978. Shear breakage of DNA. Biophys. J. 24, 489–503. https://doi.org/10.1016/S0006-3495(78)85396-X

Davis, D.L., O'Brien, E.P., Bentzley, C.M., 2000. Analysis of the degradation of oligonucleotide strands during the freezing/thawing processes using MALDI-MS. Anal. Chem. 72, 5092–5096.

Ellervik, C., Vaught, J., 2015. Preanalytical Variables Affecting the Integrity of Human Biospecimens in Biobanking. Clin. Chem. 61, 914–934. https://doi.org/10.1373/clinchem.2014.228783

Elmore, S., 2007. Apoptosis: A Review of Programmed Cell Death. Toxicol. Pathol. 35, 495–516. https://doi.org/10.1080/01926230701320337

English, A.C., Richards, S., Han, Y., Wang, M., Vee, V., Qu, J., Qin, X., Muzny, D.M., Reid, J.G., Worley, K.C., Gibbs, R.A., 2012. Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-Read Sequencing Technology. PLoS ONE 7, e47768. https://doi.org/10.1371/journal.pone.0047768

Evans, R.K., Xu, Z., Bohannon, K.E., Wang, B., Bruner, M.W., Volkin, D.B., 2000. Evaluation of Degradation Pathways for Plasmid DNA in Pharmaceutical Formulations via Accelerated Stability Studies. J. Pharm. Sci. 89, 76–87. https://doi.org/10.1002/(SICI)1520-6017(200001)89:1<76::AID-JPS8>3.0.CO;2-U

Goodwin, S., McPherson, J.D., McCombie, W.R., 2016. Coming of age: ten years of next-generation sequencing technologies. Nat. Rev. Genet. 17, 333–351. https://doi.org/10.1038/nrg.2016.49

Hubel, A., Spindler, R., Skubitz, A.P.N., 2014. Storage of Human Biospecimens: Selection of the Optimal Storage Temperature. Biopreservation Biobanking 12, 165–175. https://doi.org/10.1089/bio.2013.0084

Integrated DNA Technologies, 2014. Oligonucleotide Stability Study.

Jiang, Y.-W., Ran, S.-Y., He, L.-L., Wang, X.-H., Zhang, L.-X., 2015. Decondensation behavior of DNA chains induced by multivalent cations at high salt concentrations: Molecular dynamics simulations and experiments. Chin. Phys. B 24, 118701. https://doi.org/10.1088/1674-1056/24/11/118701

Johnson, L.A., Ferris, J.A.J., 2002. Analysis of postmortem DNA degradation by single-cell gel electrophoresis. Forensic Sci. Int. 126, 43–47. https://doi.org/10.1016/S0379-0738(02)00027-0

Kopeika, J., Thornhill, A., Khalaf, Y., 2015. The effect of cryopreservation on the genome of gametes and embryos: principles of cryobiology and critical appraisal of the evidence. Hum. Reprod. Update 21, 209–227. https://doi.org/10.1093/humupd/dmu063

Lahiri, D.K., Schnabel, B., 1993. DNA isolation by a rapid method from human blood samples: Effects of MgCl2, EDTA, storage time, and temperature on DNA yield and quality. Biochem. Genet. 31, 321–328. https://doi.org/10.1007/BF00553174

Le Page, C., Köbel, M., de Ladurantaye, M., Rahimi, K., Madore, J., Babinszky, S., Bachvarov, D.R., Bachvarova, M., Beauchamp, M.-C., Cass, C.E., Chadwick, D., Colleen, C., Damaraju, S., Dufour, J., Gotlieb, W.H., Kalloger, S.E., Portelance, L., McAlpine, J.N., Matte, I., Piché, A., Shaw, P., Roehrl, M.H.A., Vanderhyden, B.C., Watson, P.H., Huntsman, D.G., Provencher, D.M., Mes-Masson, A.-M., 2013. Specimen Quality Evaluation in Canadian Biobanks

Participating in the COEUR Repository. Biopreservation Biobanking 11, 83–93. https://doi.org/10.1089/bio.2012.0044

Lengsfeld, C.S., Munson, L., Lentz, Y.K., Anchordoquy, T.J., 2011. DNA Hydrodynamic Degradation Controlled by Kolomogorov Length Scales in Pipe Flow. J. Pharm. Sci. 100, 3088–3095. https://doi.org/10.1002/jps.22582

Lindahl, T., 1993. Instability and decay of the primary structure of DNA. Nature 362, 709–715. https://doi.org/10.1038/362709a0

Lok, S., Paton, T.A., Wang, Z., Kaur, G., Walker, S., Yuen, R.K.C., Sung, W.W.L., Whitney, J., Buchanan, J.A., Trost, B., Singh, N., Apresto, B., Chen, N., Coole, M., Dawson, T.J., Ho, K., Hu, Z., Pullenayegum, S., Samler, K., Shipstone, A., Tsoi, F., Wang, T., Pereira, S.L., Rostami, P., Ryan, C.A., Tong, A.H.Y., Ng, K., Sundaravadanam, Y., Simpson, J.T., Lim, B.K., Engstrom, M.D., Dutton, C.J., Kerr, K.C.R., Franke, M., Rapley, W., Wintle, R.F., Scherer, S.W., 2017. *De Novo* Genome and Transcriptome Assembly of the Canadian Beaver (*Castor canadensis*). G3amp58 GenesGenomesGenetics 7, 755–773. https://doi.org/10.1534/g3.116.038208

Lou, J.J., Mirsadraei, L., Sanchez, D.E., Wilson, R.W., Shabihkhani, M., Lucey, G.M., Wei, B., Singer, E.J., Mareninov, S., Yong, W.H., 2014. A review of room temperature storage of biospecimen tissue and nucleic acids for anatomic pathology laboratories and biorepositories. Clin. Biochem. 47, 267–273. https://doi.org/10.1016/j.clinbiochem.2013.12.011

Malentacchi, F., Ciniselli, C.M., Pazzagli, M., Verderio, P., Barraud, L., Hartmann, C.C., Pizzamiglio, S., Weisbuch, S., Wyrich, R., Gelmini, S., 2015. Influence of pre-analytical procedures on genomic DNA integrity in blood samples: The SPIDIA experience. Clin. Chim. Acta 440, 205–210. https://doi.org/10.1016/j.cca.2014.12.004

Malentacchi, F., Pazzagli, M., Simi, L., Orlando, C., Wyrich, R., Hartmann, C.C., Verderio, P., Pizzamiglio, S., Ciniselli, C.M., Tichopad, A., Kubista, M., Gelmini, S., 2013. SPIDIA-DNA: An External Quality Assessment for the pre-analytical phase of blood samples used for DNA-based analyses. Clin. Chim. Acta 424, 274–286. https://doi.org/10.1016/j.cca.2013.05.012

Malentacchi, F., Pizzamiglio, S., Ibrahim-Gawel, H., Pazzagli, M., Verderio, P., Ciniselli, C.M., Wyrich, R., Gelmini, S., 2016a. Second SPIDIA-DNA External Quality Assessment (EQA): Influence of pre-analytical phase of blood samples on genomic DNA quality. Clin. Chim. Acta 454, 10–14. https://doi.org/10.1016/j.cca.2015.12.032

Malentacchi, F., Pizzamiglio, S., Ibrahim-Gawel, H., Pazzagli, M., Verderio, P., Ciniselli, C.M., Wyrich, R., Gelmini, S., 2016b. Data and performances evaluation of the SPIDIA-DNA Pan-European External Quality Assessment: 2nd SPIDIA-DNA laboratory report. Data Brief 6, 980–984. https://doi.org/10.1016/j.dib.2016.01.062

Mathay, C., Hamot, G., Henry, E., Mommaerts, K., Thorlaksdottir, A., Trouet, J., Betsou, F., 2016. Method Validation for Extraction of Nucleic Acids from Peripheral Whole Blood. Biopreservation Biobanking 14, 520–529. https://doi.org/10.1089/bio.2016.0011

Meacle, F. j., Zhang, H., Papantoniou, I., Ward, J. m., Titchener-Hooker, N. j., Hoare, M., 2007. Degradation of supercoiled plasmid DNA within a capillary device. Biotechnol. Bioeng. 97, 1148–1157. https://doi.org/10.1002/bit.21275

Morente, M.M., Mager, R., Alonso, S., Pezzella, F., Spatz, A., Knox, K., Kerr, D., Dinjens, W.N.M., Oosterhuis, J.W., Lam, K.H., Oomen, M.H.A., van Damme, B., van de Vijver, M., van Boven, H., Kerjaschki, D., Pammer, J., Lopez-Guerrero, J.A., Llombart Bosch, A., Carbone, A., Gloghini, A., Teodorovic, I., Isabelle, M., Passioukov, A., Lejeune, S., Therasse, P., van Veen, E.-B., Ratcliffe, C., Riegman, P.H.J., 2006. TuBaFrost 2: Standardising tissue collection and quality control procedures for a European virtual frozen tissue bank network. Eur. J. Cancer 42, 2684–2691. https://doi.org/10.1016/j.ejca.2006.04.029

Mulcahy, D.G., Macdonald, K.S., Brady, S.G., Meyer, C., Barker, K.B., Coddington, J., 2016. Greater than X kb: a quantitative assessment of preservation conditions on genomic DNA quality, and a proposed standard for genome-quality DNA. PeerJ 4, e2528. https://doi.org/10.7717/peerj.2528

Norberg, J., Nilsson, L., 1996. Glass transition in DNA from molecular dynamics simulations. Proc. Natl. Acad. Sci. U. S. A. 93, 10173–10176.

Office of Biorepositories and Biospecimen Research, National Cancer Institute, National Institutes of Health, U.S. Department of Health and Human Services, n.d. Summary Report of the NIST-NCI Biospecimen Quality Assessment & Standards Development Workshop.

Overballe-Petersen, S., Orlando, L., Willerslev, E., 2012. Next-generation sequencing offers new insights into DNA degradation. Trends Biotechnol. 30, 364–368. https://doi.org/10.1016/j.tibtech.2012.03.007

Oxford Nanopore Technologies, 2017. World first: continuous DNA sequence of more than a million bases achieved with nanopore sequencing. [WWW Document]. Nanoporetech. URL https://nanoporetech.com/about-us/news/world-first-continuous-dna-sequence-more-million-bases-achieved-nanopore-sequencing?utm_content=64564686&utm_medium=social&utm_source=twitter (accessed 12.19.17).

Paajanen, P., Kettleborough, G., Lopez-Girona, E., Giolai, M., Heavens, D., Baker, D., Lister, A., Wilde, G., Hein, I., Macaulay, I., Bryan, G.J., Clark, M.D., 2017. A critical comparison of technologies for a plant genome sequencing project. bioRxiv 201830. https://doi.org/10.1101/201830

Permenter, J., Ishwar, A., Rounsavall, A., Smith, M., Faske, J., Sailey, C.J., Alfaro, M.P., 2015. Quantitative analysis of genomic DNA degradation in whole blood under various storage conditions for molecular diagnostic testing. Mol. Cell. Probes 29, 449–453. https://doi.org/10.1016/j.mcp.2015.07.002

Pogocki, D., Schöneich, C., 2000. Chemical Stability of Nucleic Acid–Derived Drugs. J. Pharm. Sci. 89, 443–456. https://doi.org/10.1002/(SICI)1520-6017(200004)89:4<443::AID-JPS2>3.0.CO;2-W

Robb, J.A., Gulley, M.L., Fitzgibbons, P.L., Kennedy, M.F., Cosentino, L.M., Washington, K., Dash, R.C., Branton, P.A., Jewell, S.D., Lapham, R.L., 2014. A Call to Standardize Preanalytic Data Elements for Biospecimens. Arch. Pathol. Lab. Med. 138, 526–537. https://doi.org/10.5858/arpa.2013-0250-CP

Röder, B., Fruhwirth, K., Vogl, C., Wagner, M., Rossmanith, P., 2010. Impact of Long-Term Storage on Stability of Standard DNA for Nucleic Acid-Based Methods. J. Clin. Microbiol. 48, 4260–4262. https://doi.org/10.1128/JCM.01230-10

Ross, K.S., Haites, N.E., Kelly, K.F., 1990. Repeated freezing and thawing of peripheral blood and DNA in suspension: effects on DNA yield and integrity. J. Med. Genet. 27, 569–570. https://doi.org/10.1136/jmg.27.9.569

Rossmanith, P., Röder, B., Frühwirth, K., Vogl, C., Wagner, M., 2011. Mechanisms of degradation of DNA standards for calibration function during storage. Appl. Microbiol. Biotechnol. 89, 407–417. https://doi.org/10.1007/s00253-010-2943-2

Sawyer, S., Krause, J., Guschanski, K., Savolainen, V., Pääbo, S., 2012. Temporal Patterns of Nucleotide Misincorporations and DNA Fragmentation in Ancient DNA. PLoS ONE 7, e34131. https://doi.org/10.1371/journal.pone.0034131

Schadt, E.E., Turner, S., Kasarskis, A., 2010. A window into third-generation sequencing. Hum. Mol. Genet. 19, R227–R240. https://doi.org/10.1093/hmg/ddq416

Schaudien, D., Baumgärtner, W., Herden, C., 2007. High Preservation of DNA Standards Diluted in 50% Glycerol: Diagn. Mol. Pathol. 16, 153–157. https://doi.org/10.1097/PDM.0b013e31803c558a

Shao, W., Khin, S., Kopp, W.C., 2012. Characterization of Effect of Repeated Freeze and Thaw Cycles on Stability of Genomic DNA Using Pulsed Field Gel Electrophoresis. Biopreservation Biobanking 10, 4–11. https://doi.org/10.1089/bio.2011.0016

Thornton, M., Gladwin, A., Payne, R., Moore, R., Cresswell, C., McKechnie, D., Kelly, S., March, R., 2005. Automation and validation of DNA-banking systems. Drug Discov. Today 10, 1369–1375. https://doi.org/10.1016/S1359-6446(05)03560-9

Vanapalli, S.A., Ceccio, S.L., Solomon, M.J., 2006. Universal scaling for polymer chain scission in turbulence. Proc. Natl. Acad. Sci. 103, 16660–16665. https://doi.org/10.1073/pnas.0607933103

Yoo, H.-B., Lim, H.-M., Yang, I., Kim, S.-K., Park, S.-R., 2011. Flow cytometric investigation on degradation of macro-DNA by common laboratory manipulations. J. Biophys. Chem. 02, 102–111. https://doi.org/10.4236/jbpc.2011.22013