Building a tumor atlas: integrating single-cell RNA-Seq data with spatial transcriptomics in pancreatic ductal adenocarcinoma

Reuben Moncada[1], Marta Chiodin[1], Joseph C. Devlin[1], Maayan Baron[1], Cristina H. Hajdu[2], Diane Simeone[2], and Itai Yanai[1]

[1] Institute for Computational Medicine, NYU Langone Health, New York

[2] Pancreatic Cancer Center, NYU Langone Health, New York

To understand the architecture of a tissue it is necessary to know both the cell populations and their physical relationships to one another. Single-cell RNA-Seq (scRNA-Seq) has made significant progress towards the unbiased and systematic characterization of the cell populations within a tissue, as well as their cellular states, by studying hundreds and thousands of cells in a single experiment. However, the characterization of the spatial organization of individual cells within a tissue has been more elusive. The recently introduced 'spatial transcriptomics' method (ST) reveals the spatial pattern of gene expression within a tissue section at a resolution of one thousand 100 μm spots, each capturing the transcriptomes of ~10-20 cells. Here, we present an approach for the integration of scRNA-Seq and ST data generated from the same sample of pancreatic cancer tissue. Using markers for cell-types identified by scRNA-Seq, we robustly deconvolved the cell-type composition of each ST spot, to generate a spatial atlas of cell proportions across the tissue. Studying this atlas, we found that distinct spatial localizations accompany each of the three cancer cell populations that we identified. Strikingly, we find that subpopulations defined in the scRNA-Seq data also exhibit spatial segregation in the atlas, suggesting such an atlas may be used to study the functional attributes of subpopulations. Our results provide a framework for creating a tumor atlas by mapping single-cell populations to their spatial region, as well as the inference of cell architecture in any tissue.

INTRODUCTION

Tremendous technological advances have enabled a molecular view of cancer at

unprecedented resolution[1]. Single-cell RNA-sequencing (scRNA-Seq) has emerged as a

powerful tool that provides an unbiased and systematic characterization of the cells present in a

given tissue[2–4]. Indeed, the application of scRNA-Seq to patient tumors has uncovered multiple

cancer subtypes, cellular subpopulations, and has highlighted intercellular cross-talk within the

tumor microenvironment [5–12]. Using a non-malignant cell transcriptome as a reference, large

copy number variations across entire chromosomes can also be inferred from the

transcriptomes of malignant cells[7]. However, due to the necessity of cellular dissociation prior to

sequencing of individual cells, the spatial context for each cell is lost thus limiting insight into the

manner by which they compose a tumor.

Recently, methods have recently been introduced that provide spatially resolved transcriptomic

profiling[13–15] on the basis of a limited set of genes (typically < 20). In order to integrate spatial

information with scRNA-Seq data, these methods are incredibly useful. For example, *in situ*

hybridization (ISH) gene expression atlases[16,17] have made for useful references for cellular

localization. Using the ISH atlas as a guide, these groups were able to accurately map rare

subpopulations in two different organisms using a small subset of genes. However, such atlases

to guide the localization of each cell do not exist for solid tumors which have an unpredictable

tissue architecture and gene expression patterns. Thus, high-throughput and comprehensive

mapping of single-cells onto tissue requires robust integration of multiple methods.

The recently developed Spatial Transcriptomics (ST) method is unique in its potential for

seamless integration with scRNA-Seq data. ST enables spatially resolved transcriptomic

profiling of tissue sections using spatially barcoded oligo-deoxythymidine (oligo-dT) microarrays,

allowing for unbiased mapping of transcripts over entire tissue sections[18] (Figure 1). Stahl *et al.*

first used ST to characterize unique histological features of the olfactory bulb and breast cancer

tissue, distinguishing genes expressed in invasive cancer versus ductal cancer *in situ*[18].

However, as is the case of previously reported spatially resolved transcriptomic tools[13,19,20], a

main limitation of the ST method is its lack of cellular resolution: each spot captures the

transcriptomes of ~10-20 neighboring cells. Thus, in order to extract ST's full potential, it would

be necessary to combine its data with a distinct data modality such as scRNA-Seq.

Here, we present the integration of scRNA-Seq with the ST method. In our method, a single-cell

tumor suspension is generated and processed using the inDrop platform to identify cell-types

present and to infer the cellular identity of each cluster using a set of marker genes. From the

same tumor, tissue sections are also cryosectioned and processed using the ST method to

provide an unbiased map of all expressed transcripts across the tissue section. Because each

ST spot is a mosaic of transcripts from all cells present within the spot, we computationally

deconvolve each spot to precisely estimate cell-type proportions across the tissue using the

scRNA-Seq-identified cell-type markers. Finally, deconvolving each ST spot yields individual,

spatially resolved transcriptomes for each identified cell-type, allowing for comparison of cells

across spatially distinct regions of tissue.

We used this approach to study a tissue from the same pancreatic ductal adenocarcinoma

(PDAC) tumor (Figure 1). We found three distinct pancreatic cancer cell populations – among

five other non-cancer cell-types – in the scRNA-Seq data. Deconvolving the ST data on the

PDAC tissue section, we find that the three cancer cell populations occupy distinct physical

regions. Our analysis demonstrates the plausibility of using two powerful technologies to

construct a comprehensive cellular atlas for any heterogeneous tissue.

RESULTS

Identifying cell populations in pancreatic cancer with single-cell RNA-Seq

Two hours after a tumor was resected from a patient, it cleared pathology and arrived in our lab,

where it was immediately processed for scRNA-seq and ST (see Methods). We processed the

single-cell suspension using the inDrop platform[21], collecting approximately 4,000 cells. After

sequencing, initial analysis, and filtering, 806 transcriptomes remained for analysis with an

average of approximately 4,000 unique molecular identifiers (UMIs) and 1,800 unique genes per

cell (Figure S1).

A tSNE analysis of the 806 cells based upon 615 dynamically expressed genes revealed eight

distinct clusters (Figure 2A). To characterize each cluster, we identified cluster-specific gene

expression (see Methods). Figure 2B shows the expression of a set of markers across the

individual cells that were influential in our cell-type inference. The detected cell-types include

fibroblasts (expressing S100A4[22]), macrophages (expressing of FCGR3A and CD14[23]), tuft cells

(expressing AVIL and TRPM5[24]), CD8 T-cells (expressing CD8A[25]), and red blood cells

(expressing HBB, HBA2, HBA1, and HBD)[26]. We did not identify acinar or ductal cells perhaps

due to our procedure for making a single-cell suspension. However, from our previous analyses

of human and mouse pancreata we already have strong markers for these cell types[27].

We inferred that three of the clusters correspond to cancer populations: (1) a cell population

expressing DPCR1, CLDN4, CEACAM6, CA9, GABRP, and AGR2, all of which are upregulated

in pancreatic cancer cells[28–33] (henceforth, Cancer A), (2) a cell population that is exocrine-like

based on the Moffitt et al. classification based on the expression of REG3A, REG1A, CFTR,

SLC4A4, AQP3, and SPINK1[34–38] (Cancer B), and (3) a cell population that displays high

expression of pancreatic cancer-associated genes including S100P, LAMC2, TM4SF1, GABRP,

and NPM1[32,39–41] (Cancer C) (Figure 2B-C). Based on a recent classification of PDAC

subtypes[42], we infer that both the Cancer A and the Cancer B population are aligned most

closely to the 'pancreatic-progenitor' subtype based on high expression of the transcription

factors that define this class (Figure S2).

Spatial transcriptomics (ST) of pancreatic cancer tissue

We cryosectioned unfixed, frozen tissue sections for ST analysis, from the same pancreatic

cancer tumor sample used to generate the scRNA-Seq data (Figure 2). The sections were

mounted onto a spatially barcoded microarray slide (see Methods). After staining the tissue, the

section was presented to a trained pathologist for histological annotation of distinct tissue

features (Figure 3A-D). We thus defined four regions: (1) high in cancer cells and desmoplasia,

(2) the duct epithelium, (3) normal pancreatic tissue, and (4) inflamed tissue. The slide was then

processed with the ST protocol: involving cDNA synthesis, *in vitro* transcription amplification,

library construction, and sequencing[18]. Analyzing the sequence reads, we demultiplexed the

reads and identified their spatial location within the tissue using the ST spatial-specific barcodes

of the array. We detected approximately 2,000 UMIs and approximately 1,000 unique genes per

spot. Mapping the distribution of UMIs and unique genes over the tissue spots indicates they

are uniformly distributed (Figure 3E).

The ST data allows one to examine for a particular gene across the tissue. Figure 3G shows the

spatial gene expression profiles of four genes. As shown, CRISP3 is localized to the duct

epithelium region of the tissue section, while COL1A2 shows localized expression in the

desmoplasia region. We next asked if the ST spot transcriptomes can be clustered into co-

expressed regions. For this, we performed principal components analysis (PCA) on the 1,339

most dynamically expressed genes across all spots. Figure 3H indicates the scores of the first

three PCs mapped to the tissue showing distinct tissue regions. The regions demarcated by

these PCs confirm the pathologist-annotated sections of the tissue (Fig. 3A). For example, the

PC3-high spots spatially localize to the duct epithelium of the tissue section. The genes

contributing the most (highest loadings) to PCs one through three showed enriched Gene

Ontology terms for cell migration and collagen fibril organization (PC1); cell migration and

collagen induced tyrosine kinase activity (PC2); and digestive processes (PC3).

Deconvolution of spatial transcriptomic data using single-cell transcriptomic data

In our previous work, we studied the transcriptomes of human and mouse pancreata at single-

cell resolution[43]. Using the cell-type specific markers that we identified by scRNA-Seq, we

deconvolved previously reported bulk pancreas transcriptomic data using an algorithm we

named Bseq-SC (Bulk-sequencing single-cell deconvolution analysis). Here, we apply the same

algorithm to the deconvolution of the spatial transcriptome spots to understand the

regionalization of each cell-type (Figure 4A).

In order to test Bseq-SC's accuracy in deconvolving cell-types, we assembled 46 artificial

mixtures of cell-type specific transcriptomes at distinct proportions (Figure S3A). We then

deconvolved the cell-type proportions using Bseq-SC and compared with the true proportions

(Figure S3A). We found R^2 values greater than 0.9 for each of the six examined cell-types,

indicating Bseq-SC's ability to accurately deconvolve cell-proportions (Figure S3B).

A set of marker genes specific to each cell-type was compiled to form a basis matrix for

deconvolution (Figure 4A). Although neither acinar nor ductal cells were detected in our scRNA-

Seq data, we added to our basis matrix the previously identified strong marker genes for these

cell-types (Figure S3)[44]. Deconvolving each spot transcriptome into individual cell-type,

proportions we depicted the fraction of cell-types composing each spot with pie-charts replacing

the ST spots (Figure 4B).

When plotting the relative proportions of each cancer population identified, we found that they

spatially segregate across the tissue (Figure 4C). Interestingly, while the Cancer A cells are

sparsely concentrated across the tissue, the Cancer B cells localized strongly to the duct

epithelium and the Cancer C population strongly localized to the region high in cancer tubules

and desmoplasia.

Figure 4D shows the cell-type proportions for all of the examined spots. As the earlier PC

analysis suggested, the annotated histological features correspond to areas with distinct cell-

type compositions. The region described as high in cancer tubules and desmoplasia was

deconvolved predominantly into the cancer C population and fibroblasts (Figure 4D, light blue

and dark blue, respectively). The duct epithelium region was enriched for Cancer B cells and

ductal cells (Figure 4C, light green). Also as expected, the normal pancreatic tissue was

enriched for acinar cells (Figure 4C, light green). Interestingly, the Cancer A cells were found to

be localized in multiple regions of the tissue (Figure 4D, light green).

Intra-cell population spatial relationships

While the Bseq-SC algorithm deconvolves a given bulk ST spot to its relative cell proportions

(Figure 4), we added a second step to further infer the full transcriptome of each of the

composing cell-types (Figure 5A). Using these fully deconvolved spatial transcriptomes, we

hypothesized that we could detect subpopulations of a particular cell-type across space. We

started with a PCA on the deconvolved Cancer B spot transcriptomes (Figure 5B). To ask

whether there is a correspondence between the Cancer B spot transcriptomes and their spatial

position on the tissue, we annotated each spot with a specific color, using a two dimensional

color map (Figure 5B). Mapping these colors back to the tissue we found that the PCA clusters

of spots co-segregate spatially on the tissue (Figure 5C). In particular, spots located in the

ductal epithelium region are clustered in PC space, while spots in the normal pancreatic area of

the also cluster in PC space (Figure 5B).

We next asked whether the same genes that contribute to subpopulations in the ST

deconvolved spots, also delineate subpopulations amongst the scRNA-Seq clusters. For

several genes we indeed found such co-separation (Figure 5D-F, Table S1-S2). For example

REG1A – a gene implicated in the ductal to acinar transition[45] – is one of the top 30 contributors

to PC1 (Figure 5D). Its expression in the tissue also showed a distribution that is skewed to the

ductal epithelium. Notably, this subpopulation structure is validated by the scRNA-Seq data

where REG1A also shows subpopulation structure (Figure 5F). As another example, APOL1

shows a separation in both the ST data and the scRNA-Seq data (Figure S5) (Figure 5F-I).

Finally Table S1 indicates additional genes with co-separation in both the ST and scRNA-Seq

data for both the Cancer B and Cancer C deconvolved transcriptomes.

DISCUSSION

Here we describe the first application of scRNA-seq (using the inDrop system) and ST on the same tumor sample. By integrating these two orthogonal methods we extend existing analyses for studying spatial transcriptomics data by deconvolving the spot transcriptomes to individual cell-types using the scRNA-Seq-defined cell-type markers (Figure 1). Previous scRNA-Seq studies cluster cells and then annotate each cluster based upon its uniquely expressed genes. Our approach of coupling scRNA-Seq and spatial transcriptomics offers another method for annotating an scRNA-Seq cluster: since there are histological features within the tissue section that can be visually distinguished and annotated, it is possible to match the location of an scRNA-Seq cluster after our deconvolution step with the histological region where they localize to guide cell-type inference (Figure 4).

Here we identify three populations of cancer cells that we distinguish using their gene expression profiles. Past work using integrative gene expression analyses[34,42,46] or virtual microdissections[47] defined subtypes of PDAC that largely overlap between studies. Here, two of the three identified cancer cell populations (Cancer A and Cancer B) appear to align strongly to the 'pancreatic progenitor' subtype previously described based on high expression of the transcription factors characteristic of this subtype (Figure S2)[42,46]. Interestingly, the third cancer cell population (Cancer C) displays a gene expression program that was not defined in past molecular characterizations of pancreatic cancer. This cluster of cancer cells is enriched for the expression of TM4SF1 and CLDN1, genes associated with invasive traits[48–50], and as well as the expression of other genes known to be associated with pancreatic cancer progression (S100P, LAMC2, NPM1)[39–41]. In our analysis we did not identify any acinar or ductal cells, perhaps due to the nature of the cell suspension preparation (see Methods). Although a defining characteristic of pancreatic cancer is a dense desmoplastic reaction – which consists of fibroblasts, stellate cells, endothelial cells, and immune cells[51] – only fibroblasts and immune cells were identified in our scRNA-Seq data (Figure 2). Our inability to fully capture the cells present in the stroma likely reflects the specific preparation of the single-cell suspension necessary for scRNA-Seq, suggesting that perhaps additional steps need to be taken to fully dissociate the stromal and endothelial compartments of the desmoplasia.

Interestingly, when we localize each of the observed cancer populations onto the ST spots based on their deconvolved proportions, only the Cancer C population is spatially restricted to the cancerous and desmoplastic region. When we map the fibroblast population onto the tissue,

we also find them to be strongly restricted to the cancerous and desmoplastic region and not

highly concentrated around the pancreatic tissue. Both pancreatic cancer cells and stromal cells

secrete factors and cooperate to promote the aggressive nature of the disease[51–53]. Because

the fibroblasts identified in this scRNA-Seq dataset localize to the desmoplasia region with the

Cancer C population and not to the fibrotic region surrounding the normal pancreatic tissue, this

particular fibroblast population perhaps represents a population of cells in the stroma that is

acting to promote the growth of these cancer cells.

An advantage of our integration is the mapping of populations onto irregular tissue

architectures. With the fully deconvolved ST data, we identify spatially segregated

subpopulations of the Cancer A and Cancer B populations (Figures 5 and S5). In the case of the

REG1A enriched Cancer cell B subpopulation, this may potentially represent a subset of cells

that have transitioned from an acinar cell phenotype to a ductal cell phenotype based on a

report suggesting the involvement of REG1A in this process[45].

The construction of a tumor atlas has far reaching impact, particularly with regard to the

identification and classification of cell-populations that comprise such a heterogeneous tissue.

The advent of scRNA-Seq has allowed for the identification of cancer subtypes and non-

malignant cell subpopulations[10,54]; the framework for atlas construction described here can aid

in assigning potential functional roles of cellular subtypes based on spatial localization (relative

to the tissue, or relative to the other cells present). Additionally, by applying scRNA-Seq and ST

on the same biological sample as we describe here, rare subpopulations specific to the sample

can be mapped to the same tissue of origin. In the case of tumors for which the precise

composition of different tumor sub-classifications are likely to vary from individual to individual,

the subtype composition and spatial localization can be ascertained for a given patient, and can

perhaps be correlated with patient outcome.

Our approach for integration has a number of limitations. First, the number of spots with the

current implementation of ST is just over 1,000 spots that span a ~6 mm x ~6 mm array.

Second, the resolution is low; each spot is 100 μm in diameter, thus capturing roughly 10-20

cells depending on cell types captured and the thickness of the tissue section (generally ranging

from 8 to 20 μm). Additionally, the distance between the center of two adjacent spots is 200 μm,

therefore the array of ST spots does not cover the entire area of the tissue. At this scale there is

insufficient resolution, for example, to study the architecture of pancreatic islets which have a

diameter similar to that of the ST spots. Despite these limitations we observed interesting

cellular architectures within the tumor and it would be interesting to apply this approach beyond

cancer, for example to organs such as the testes, embryonic stages where the entire organism

can fit on the slide, and dynamic processes such as bacterial infection.

METHODS

Tumor sample handling and dissociation. The pancreatic ductal adenocarcinoma tumor was

delivered in RPMI (Fisher) on ice directly from the operating room to the lab after clearing

pathology (~2 hours). The tumor resection was rinsed in ice cold PBS and cut into ~4-5 mm^3

pieces from which 1 mm thick slices were taken and set aside in ice-cold PBS. The remaining

~3-4 mm^3 pieces were embedded in chilled OCT and snap-frozen in isopentane cooled with

liquid N_2. The 1 mm tissue slices stored in PBS was further minced with scalpels to < 1 mm^3.

Tissue was rinsed from the dish with ice cold PBS and pelleted by centrifuging at 300 x g for 3

minutes at 4 degrees. PBS was aspirated and 5 ml 0.25% pre-warmed trypsin-EDTA with 10

U/µl DNaseI (Roche) was added and put into a 37 degree water bath for 30 minutes with gentle

inversion every 5 minutes. The resulting suspension was filtered through a 100 µm cell strainer

to remove larger chunks of undigested tissue. Enzymatic digestion was quenched with the

addition of FBS to a final concentration of 10%. Cells were pelleted by centrifuging the

suspension at 300 x g for 3 minutes at 4 degrees and washed twice with 5 ml ice-cold PBS.

After a final spin at 300 x g for 3 minutes, the cells were resuspended in PBS to a final

concentration of 10,000 cells/ml. The resulting viability was >95% as shown by trypan blue

exclusion.

inDrop library preparation and scRNA-Seq. From the single-cell suspension, 4000 cells were

encapsulated using the inDrop platform and reverse transcription (RT) reaction was performed

as previously described[43]. The number of PCR cycles performed for final library amplification

ranged from 9-12 cycles. Paired end sequencing was performed on an Illumina NextSeq500

sequencer.

Clustering of single-cell data and marker gene selection. tSNE was applied to the single-cell

RNA-Seq data using only highly expressed (fraction of total TPM above 0.5) and highly variant

genes (Fano-factor above mean-dependent threshold). To define cell-types from the tSNE

analysis we used a density-clustering method, DBscan[55]. This approach revealed 8 cell types,

from which we used the top preferentially expressed genes ($p < 10^{-5}$, Kolmogorov Smirnov test)

to infer the cluster identity.

Spatial Transcriptomics of PDAC tissue. Approximately 3-4 mm^3 sized pieces of the tumor

tissue were embedded in cold OCT and snap frozen in isopentane prior to cryosectioning.

Sections were cut to a thickness of 10 μm and mounted onto the spatially barcoded arrays.

Tissue fixation, staining, imaging, and subsequent Spatial Transcriptomics library preparation

was performed as previously described[56] with the following changes: after RNA amplification by

in vitro transcription (IVT) and subsequent bead clean-up, second RT reaction was performed

using random hexamers, eliminating the need for a primer ligation step[57]. The number of PCR

cycles to amplify the final libraries ranged from 9-13 cycles. Paired end sequencing was

performed on an Illumina NextSeq500 sequencer. FASTQ files were processed using an

adapted version of the celseq2 pipeline to demultiplex the spots to their spatial location.

Statistical deconvolution and cell-type specific expression estimation. In order to calculate

the cell type proportions, we generated a basis matrix of for each cell type using a list of specific

marker genes to generate a signature for each cell type as previously described[43]. This basis

matrix was then used to estimate cell type proportions for each individual ST spot using

CIBERSORT[58]. Additionally, with the full single cell data for over 19,000 genes we calculated

the "cell type specificity" as the likelihood a particular gene would be expressed in each of the

cell types. The specificity was then factored alongside the cell type proportions to estimate the

cell type specific expression of each gene in each spot according to the following

formulas.

$$E_n = \left((B * P_n) * S\right) * \frac{B}{E_{spot}}$$

$$E_{spot} = \Sigma\left((B * P_n) * S\right)$$

Where *n* is the cell type, E_n is the expression of a gene in cell type *n*, *B* is the raw UMI count for

that given gene in a single spot, P_n is the proportion of cell type *n* in that single spot, *S* is the cell

type specificity for the given gene, and E_{spot} is the sum of expression estimates for the gene

over *n* cell types.

FIGURE CAPTIONS

Figure 1. A schematic for the integration of scRNA-Seq and ST. A surgically resected

PDAC tumor sample was split and processed in parallel by scRNA-Seq and ST. scRNA-Seq

was performed using inDrop - a microfluidic implementation of CEL-Seq2 - to produce a gene

expression matrix. After clustering, each cluster is defined as a particular cell-type according to

their transcriptomes. A cryosection of an OCT embedding of the rest of the sample was used for

ST analysis to produce a gene expression matrix where each column is a spot transcriptome.

Integrating the two datasets allows us to deconvolve each spot into its comprising cell-types

Figure 2. Identifying cell-types present in a PDAC tumor sample.

(A) tSNE plot of scRNA-Seq data produced from a part of the PDAC tumor sample studied

here. After sequencing and filtering, 806 cells were analysed (see Methods). The tSNE

analysis is based on 615 dynamically expressed genes (see Methods). Colors indicate

cell-types characterized by marker expression (**B**).

(B) Heatmap indicating the expression of cell-type specific markers used to infer cell-type

identity.

(C) For four clusters, separate tSNE plots are shown with colors indicating the expression of

levels of the noted marker genes. Expression values are scaled by the k-nearest

neighbors smoothed expression described by Wagner et al (bioRxiv 2017).

(D) PCA on the Cancer A and Cancer B populations using 742 and 543 dynamically

expressed genes, respectively.

Figure 3. Pancreatic cancer spatial transcriptomics analysis

(A) Annotated H&E staining of a section of PDAC tumor tissue on the ST slide. The

annotations indicate a region high in cancer cells and desmoplasia (red), normal

pancreatic tissue (blue), normal duct tissue (yellow), and inflammation (green). Note the

spots in the background.

(B) Inset of pancreatic tumor tubules and surrounding desmoplasia. White arrowheads point

to tumor cells organizing around tubules. Black arrowheads show the surrounding

stroma and desmoplasia.

(C) Inset of healthy pancreatic tissue. Arrowheads indicate the acini.

(D) Inset of duct epithelium and inflamed tissue. White arrowheads indicate the pancreatic

ducts and the black arrowheads point to inflammatory cells with smaller nuclei.

(E) Histogram of the number of detected transcripts (left) and genes (right) for each spot.

(F) The maps show the detected transcripts (left) and genes (right) in the indicated spatial

location.

(G) Spatial gene expression profiles for the indicated four genes.

(H) Principal components analysis on the spot transcriptomes. Color in each plot indicates

the score for the PC. Note the regional localization of expression. Below, enriched gene

ontology terms for genes with high loadings for the PC.

Figure 4. Deconvolution of spatial transcriptome data with scRNA-Seq-defined cell-type

markers.

(A) A schematic of ST deconvolution using the Bseq-SC algorithm. Using a set of marker

genes for each cell-type (left), the Bseq-SC algorithm deconvolves bulk transcriptomes

into individual cell-type proportions.

(B) The pie-charts indicate the deconvolved proportions of cell-types in the specified tumor

region.

(C) Heat maps of select cell-type proportions over the ST spots.

(D) Cell-type proportions across the entire studied tissue section.

Figure 5. Subpopulation spatial mapping.

(A) Schematic of the full deconvolution of bulk Spatial Transcriptomics data into individual

cell-type spatial transcriptomes

(B) PCA of the deconvolved Cancer B gene expression. Only spots with a proportion of

Cancer B cells >10% are selected for analysis. Spots are colored by a two-dimensional

color map.

(C) ST spots, colored as in B. Note the clustering of similar color shades.

(D) Same as B, colored according to the deconvolved expression levels of the REG1A gene.

(E) Same as C, colored according to the deconvolved expression levels of the REG1A gene.

(F) Subpopulations in the inDrop data. PCA of Cancer B population (Figure 2D), colored

according to the expression levels of REG1A.

SUPPLEMENTARY FIGURES AND TABLES

Figure S1. inDrop statistics. On the left, histogram of unique transcripts per cell (log10). Right, histogram of unique genes per cell.

Figure S2. Expression of pancreatic progenitor subtype TFs in Cancer A cells and Cancer B cells. Expression of transcription factors defining of the 'pancreatic progenitor' subtype described in (30).

Figure S3. Testing Bseq-SC accuracy.

(A) Comparing known cell-type proportions with the deconvolved proportions for simulated bulk samples. 46 bulk mixtures were assembled from our previous pancreas data[43] by two different strategies. In the first strategy (1-36), we generated mixes comprised of equal combinations of three cell types, two cell types and one mixture of equal amounts of all six cell types. The transcriptome of each cell-type was derived from 150 cells annotated to that cell type. In the second strategy (37-46), 150 cells are randomly selected from the ~7,000 single-cell matrix. Note that the 10 mixtures are nearly identical and reflect biological proportions of the single cell data.

(B) A strong correlation was detected between the deconvolved and true proportions.

Figure S4. Basis matrix of marker genes for deconvolution of ST data.

Figure S5. Subpopulation spatial mapping.

    (A) PCA of the deconvolved Cancer A gene expression. Only spots with a proportion of Cancer A cells >10% are selected for analysis. Spots are colored by a two-dimensional color map.

    (B) ST spots, colored as in A. Note the clustering of similar color shades.

    (C) Same as A, colored according to the deconvolved expression levels of the APOL1 gene.

    (D) Same as B, colored according to the deconvolved expression levels of the APOL1 gene.

    (E) Subpopulations in the inDrop data. PCA of Cancer A population (Figure 2D), colored according to the expression levels of APOL1.

Table S1. Additional genes showing co-separation for the Cancer B and C deconvolved transcriptomes.

Genes co-separating Cancer A	Genes co-separating Cancer B
TFF2	CLDN2
TFF3	TIMP1
DUOX2	TGM2
SERPING1	KRT19
CD74	DDIT4
SOD1	
MUC1	
PRSS8	
MCL1	
PLAT	

References:

1. Offit, K. A decade of discovery in cancer genomics. *Nat. Rev. Clin. Oncol.* **11,** 632–634 (2014).
2. Kolodziejczyk, A. A., Kim, J. K., Svensson, V., Marioni, J. C. & Teichmann, S. A. The Technology and Biology of Single-Cell RNA Sequencing. *Mol. Cell* **58,** 610–620 (2015).
3. Papalexi, E. & Satija, R. Single-cell RNA sequencing to explore immune cell heterogeneity. *Nat. Rev. Immunol.* **18,** 35–45 (2017).
4. Ofengeim, D., Giagtzoglou, N., Huh, D., Zou, C. & Yuan, J. Single-Cell RNA Sequencing: Unraveling the Brain One Cell at a Time. *Trends Mol. Med.* **23,** 563–576 (2017).
5. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science* **344,** 1396–401 (2014).
6. Tirosh, I. *et al.* Single-cell RNA-seq supports a developmental hierarchy in human oligodendroglioma. *Nature* **539,** 309–313 (2016).
7. Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352,** 189–96 (2016).
8. Venteicher, A. S. *et al.* Decoupling genetics, lineages, and microenvironment in IDH-mutant gliomas by single-cell RNA-seq. *Science (80-.).* **355,** eaai8478 (2017).
9. Darmanis, S. *et al.* Single-Cell RNA-Seq Analysis of Infiltrating Neoplastic Cells at the Migrating Front of Human Glioblastoma. *Cell Rep.* **21,** 1399–1410 (2017).
10. Chung, W. *et al.* Single-cell RNA-seq enables comprehensive tumour and immune cell profiling in primary breast cancer. *Nat. Commun.* **8,** 15081 (2017).
11. Zheng, C. *et al.* Landscape of Infiltrating T Cells in Liver Cancer Revealed by Single-Cell Sequencing. *Cell* **169,** 1342–1356.e16 (2017).
12. Horning, A. M. *et al.* Single-cell RNA-seq reveals a subpopulation of prostate cancer cells with enhanced cell cycle-related transcription and attenuated androgen response. *Cancer Res.* canres.1924.2017 (2017). doi:10.1158/0008-5472.CAN-17-1924
13. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* **348,** aaa6090 (2015).
14. Shah, S., Lubeck, E., Zhou, W. & Cai, L. In Situ Transcription Profiling of Single Cells Reveals Spatial Organization of Cells in the Mouse Hippocampus. *Neuron* **92,** 342–357

(2016).

15.   Long, X., Colonell, J., Wong, A. M., Singer, R. H. & Lionnet, T. Quantitative mRNA
imaging throughout the entire Drosophila brain. *Nat. Methods* **14,** 703–706 (2017).

16.   Achim, K. *et al.* High-throughput spatial mapping of single-cell RNA-seq data to tissue of
origin. *Nat. Biotechnol.* **33,** 503–9 (2015).

17.   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat. Biotechnol.* **33,** 495–502 (2015).

18.   Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by
spatial transcriptomics. *Science* **353,** 78–82 (2016).

19.   Junker, J. P. *et al.* Genome-wide RNA Tomography in the zebrafish embryo. *Cell* **159,**
662–75 (2014).

20.   Chen, J. *et al.* Spatial transcriptomic analysis of cryosectioned tissue samples with Geo-
seq. *Nat. Protoc.* **12,** 566–580 (2017).

21.   Klein, A. M. *et al.* Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic
Stem Cells. *Cell* **161,** 1187–1201 (2015).

22.   Strutz, F. *et al.* Identification and characterization of a fibroblast marker: FSP1. *J. Cell*
Biol. **130,** 393–405 (1995).

23.   Gautier, E. L. *et al.* Gene-expression profiles and transcriptional regulatory pathways that
underlie the identity and diversity of mouse tissue macrophages. *Nat. Immunol.* **13,**
1118–28 (2012).

24.   Gerbe, F., Legraverend, C. & Jay, P. The intestinal epithelium tuft cells: specification and
function. *Cell. Mol. Life Sci.* **69,** 2907–17 (2012).

25.   Mahnke, Y. D., Brodie, T. M., Sallusto, F., Roederer, M. & Lugli, E. The who's who of T-
cell differentiation: Human memory T-cell subsets. *Eur. J. Immunol.* **43,** 2797–2809
(2013).

26.   Hardison, R. C. Evolution of hemoglobin and its genes. *Cold Spring Harb. Perspect. Med.*
2, a011627 (2012).

27.   Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas
Reveals Inter- and Intra-cell Population Structure. *Cell Syst.* **3,** (2016).

28.   Yan, J. *et al.* High expression of diffuse panbronchiolitis critical region 1 gene promotes
cell proliferation, migration and invasion in pancreatic ductal adenocarcinoma. *Biochem.*
Biophys. Res. Commun. **495,** 1908–1914 (2018).

29.   Tsutsumi, K. *et al.* Claudin-4 expression predicts survival in pancreatic ductal
adenocarcinoma. *Ann. Surg. Oncol.* **19 Suppl 3,** S491-9 (2012).

30.   Chen, J. *et al.* CEACAM6 induces epithelial-mesenchymal transition and mediates
invasion and metastasis in pancreatic cancer. *Int. J. Oncol.* **43,** 877–85 (2013).

31.   Li, G., Feng, G., Achour, Y., Genin, C. & Tostain, J. MN/CA9 as a novel molecular marker
for the detection of cancer. *Expert Opin. Med. Diagn.* **1,** 91–7 (2007).

32.   Takehara, A. *et al.* Gamma-aminobutyric acid (GABA) stimulates pancreatic cancer
growth through overexpressing GABAA receptor pi subunit. *Cancer Res.* **67,** 9704–12
(2007).

33.   Ferreira, R. M. M. *et al.* Duct- and Acinar-Derived Pancreatic Ductal Adenocarcinomas
Show Distinct Tumor Progression and Marker Expression. *Cell Rep.* **21,** 966–978 (2017).

34.   Collisson, E. A. *et al.* Subtypes of pancreatic ductal adenocarcinoma and their differing
responses to therapy. *Nat. Med.* **17,** 500–3 (2011).

35.   Direito, I., Paulino, J., Vigia, E., Brito, M. A. & Soveral, G. Differential expression of
aquaporin-3 and aquaporin-5 in pancreatic ductal adenocarcinoma. *J. Surg. Oncol.* **115,**
980–996 (2017).

36.   Huang, X., Huang, L. & Shao, M. Aquaporin 3 facilitates tumor growth in pancreatic
cancer by modulating mTOR signaling. *Biochem. Biophys. Res. Commun.* **486,** 1097–
1102 (2017).

37. Li, Q. *et al.* Reg proteins promote acinar-to-ductal metaplasia and act as novel diagnostic
and prognostic markers in pancreatic ductal adenocarcinoma. *Oncotarget* **7,** 77838–
77853 (2016).
38. Liu, X. *et al.* REG3A accelerates pancreatic cancer cell growth under IL-6-associated
inflammatory condition: Involvement of a REG3A-JAK2/STAT3 positive feedback loop.
Cancer Lett. **362,** 45–60 (2015).
39. Arumugam, T., Simeone, D. M., Van Golen, K. & Logsdon, C. D. S100P promotes
pancreatic cancer growth, survival, and invasion. *Clin. Cancer Res.* **11,** 5356–64 (2005).
40. Kosanam, H. *et al.* Laminin, gamma 2 (LAMC2): a promising new putative pancreatic
cancer biomarker identified by proteomic analysis of pancreatic adenocarcinoma tissues.
Mol. Cell. Proteomics **12,** 2820–32 (2013).
41. Zhu, Y. *et al.* NPM1 activates metabolic changes by inhibiting FBP1 while promoting the
tumorigenicity of pancreatic cancer cells. *Oncotarget* **6,** 21443–51 (2015).
42. Bailey, P. *et al.* Genomic analyses identify molecular subtypes of pancreatic cancer.
Nature **531,** 47–52 (2016).
43. Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas
Reveals Inter- and Intra-cell Population Structure. *Cell Syst.* **3,** 346–360.e4 (2016).
44. Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas
Reveals Inter- and Intra-cell Population Structure. *Cell Syst.* (2016).
doi:10.1016/j.cels.2016.08.011
45. Li, Q. *et al.* Reg proteins promote acinar-to-ductal metaplasia and act as novel diagnostic
and prognostic markers in pancreatic ductal adenocarcinoma. *Oncotarget* **7,** 77838–
77853 (2016).
46. Aguirre, A. J., Hruban, R. H. & Raphael, B. J. Integrated Genomic Characterization of
Pancreatic Ductal Adenocarcinoma. *Cancer Cell* **32,** 185–203.e13 (2017).
47. Moffitt, R. A. *et al.* Virtual microdissection identifies distinct tumor- and stroma-specific
subtypes of pancreatic ductal adenocarcinoma. *Nat. Genet.* **47,** 1168–1178 (2015).
48. Cao, J. *et al.* TM4SF1 Regulates Pancreatic Cancer Migration and Invasion
<i>In Vitro</i> and <i>In Vivo</i>
Cell. Physiol. Biochem. **39,** 740–750 (2016).
49. Yang, J. *et al.* TM4SF1 Promotes Metastasis of Pancreatic Cancer via Regulating the
Expression of DDR1. *Sci. Rep.* **7,** 45895 (2017).
50. Kondo, J. *et al.* Claudin-1 expression is induced by tumor necrosis factor-alpha in human
pancreatic cancer cells. *Int. J. Mol. Med.* **22,** 645–9 (2008).
51. Rasheed, Z. A., Matsui, W. & Maitra, A. *Pathology of pancreatic stroma in PDAC.*
Pancreatic Cancer and Tumor Microenvironment (Transworld Research Network, 2012).
52. Sato, N., Maehara, N. & Goggins, M. Gene Expression Profiling of Tumor–Stromal
Interactions between Pancreatic Cancer Cells and Stromal Fibroblasts. *Cancer Res.* **64,**
6950–6956 (2004).
53. Ohuchida, K. *et al.* Radiation to stromal fibroblasts increases invasiveness of pancreatic
cancer cells through tumor-stromal interactions. *Cancer Res.* **64,** 3215–22 (2004).
54. Li, H. *et al.* Reference component analysis of single-cell transcriptomes elucidates
cellular heterogeneity in human colorectal tumors. *Nat. Genet.* **49,** 708–718 (2017).
55. Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. A Density-Based Algorithm for Discovering
Clusters in Large Spatial Databases with Noise.
56. Stahl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by
spatial transcriptomics. *Science (80-.).* **353,** 78–82 (2016).
57. Hashimshony, T. *et al.* CEL-Seq2: Sensitive highly-multiplexed single-cell RNA-Seq.
Genome Biol. **17,** (2016).
58. Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.
Nat. Methods **12,** 453–457 (2015).

569
570

Figure 1

Resected tumor

Dissociation to single-cell suspension

Tissue embedding and cryosectioning

Encapsulation of live cells with inDrop

Mount cryosections onto spatially barcoded microarray

Poly-T stretch
Unique molecular identifer (UMI)
Spatial array barcode (ID)
Illumina 5' adapter
T7 Promoter
Cleavage site

Library preparation and sequencing

Library preparation and sequencing

Single-cell UMI count matrix

	Cell 1	Cell 2	Cell 3	· · ·	Cell n
Gene 1	0	1	0		1
Gene 2	4	0	0		0
Gene 3	2	1	5		0
Gene m	0	1	0		1

Spatial Transcriptomics UMI count matrix

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	12	2	0		4
Gene 2	2	0	8		2
Gene 3	4	3	9		0
Gene i	2	3	0		4

Identification of cell populations

Cell type A
Cell type B
Cell type C
Cell type D
Cell type E
Cell type F
Cell type G
Cell type H

tSNE 2
tSNE 1

Gene expression maps

Gene i expression

Low   High

Deconvolution of Spatial Transcriptomics data using single-cell RNA-seq data

Individual cell-type proportions and transcriptomes across tissue

Cell type A   Cell type E
Cell type B   Cell type F
Cell type C   Cell type G
Cell type D   Cell type H

Figure 2

Figure 3

Figure 4

A

Cell types

Cell-specific marker

z-score

Bseq-SC ⊗

ST 'bulk' spots

=

Cell type A
Cell type B
Cell type C
Cell type D

B

Cancer cells A
Cancer cells B
Cancer cells C
Fibroblasts
Macrophages
Tuft cells
CD8 T-cells
Red blood cells
Acinar cells
Ductal cells

C

Duct epithelium
Cancer and desmoplasia
Pancreatic tissue

Cancer cells A   Cancer cells B   Cancer cells C   Fibroblasts

Deconvolved proportion (%)

0   10   20   30   40   50   60   70

D

Figure 5

A

Spatial Transcriptomics UMI count matrix

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	12	2	0		4
Gene 2	2	0	8		2
Gene 3	4	3	9		0
· · ·					
Gene i	2	3	0		4

Full Deconvolution

Fully deconvolved Cell type D Spatial Transcriptome

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	1	0.75	0		0
Gene 2	1.5	0	0		2
					0

Fully deconvolved Cell type C Spatial Transcriptome

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	0	0.25	0		0
Gene 2	0	0	0		2
					4

Fully deconvolved Cell type B Spatial Transcriptome

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	10	0	0		4
Gene 2	0.5	0	3		2
					4

Fully deconvolved Cell type A Spatial Transcriptome

	Spot 1	Spot 2	Spot 3	· · ·	Spot j
Gene 1	1	0	0		0
Gene 2	0	0	5		0
Gene 3	0.25	3	0		0
· · ·					
Gene i	2	0.5	0		1

Cell-specific marker — Cell type A, Cell type B, Cell type C, Cell type D — z-score

B
Cancer B deconvolved spots

PC1, PC2

C
Cancer B

D
REG1A (ST data)

PC1, PC2

TPM ($\log_{10}$): 1.5, 1, 0.5, 0

E
REG1A

PC1, PC2

TPM ($\log_{10}$): 1.5, 1, 0.5, 0

G
REG1A (scRNA-Seq data)

PC1, PC2

TPM ($\log_{10}$): 2.8, 2.6, 2.4, 2.2, 2, 1.8

Figure S1

Figure S2

Figure S3

Figure S4

B

Figure S5