**Evolution at two time frames: polymorphisms from an ancient singular**

**divergence event fuel contemporary parallel evolution**

Steven M. Van Belleghem[1,2,3,4,*], Carl Vangestel[1,2], Katrien De Wolf[1,2], Zoë De Corte[1], Markus Möst[5], Pasi

Rastas[6], Luc De Meester[7] and Frederik Hendrickx[1,2,*]

[1.] Royal Belgian Institute of Natural Sciences, Brussels, Belgium

[2.] Terrestrial Ecology Unit, Biology Department, Ghent University, Ghent, Belgium

[3.] Department of Biology, Center for Applied Tropical Ecology and Conservation, University of Puerto Rico, Rio

Piedras, Puerto Rico

[4.] Department of Zoology, University of Cambridge, Cambridge, United Kingdom

[5.] Institute of Ecology, University of Innsbruck, Innsbruck, Austria

[6.] Ecological Genetics Research Unit, Department of Biosciences, University of Helsinki, Helsinki, Finland

[7.] Laboratory of Aquatic Ecology, Evolution and Conservation, KU Leuven, 3000 Leuven, Belgium

**\* Corresponding authors:**

Steven M. Van Belleghem (VanBelleghemSteven@hotmail.com)

Frederik Hendrickx (Frederik.Hendrickx@naturalsciences.be)

**Abstract**

When species occur in repeated ecologically distinct habitats across their range, adaptation may proceed surprisingly fast and result in parallel evolution. There is increasing evidence that such cases of rapid parallel evolution are fueled by standing genetic variation, but the origin of this genetic variation remains poorly understood. In *Pogonus chalceus* beetles, short- and long-winged ecotypes have diverged in response to contrasting hydrological regimes and can be repeatedly found along the Atlantic European coast. By analyzing genomic variation across the beetles' distribution, we reveal that genomically widespread short-wing selected alleles evolved during a singular divergence event, estimated at ~0.19 Mya. The ancient and differentially selected alleles are currently polymorphic in all populations across the range, allowing for the fast evolution of one ecotype from a small number of random individuals, as low as 5 to 15, of the populations of the other ecotype. Our results suggest that cases of fast parallel ecological divergence might be the result of evolution at two different time frames: ecological divergence in isolation in the past, followed by repeated selection on the divergently evolved alleles after admixture. We suggest that this mechanism may be common and potentially driven by large-scale environmental changes such as glacial cycles.

**Introduction**

Adaptation to local environmental conditions may lead to the evolution of distinct ecotypes and, ultimately, new species (1, 2). Under prolonged periods of geographical isolation, the absence of gene flow allows populations to accumulate new alleles and build-up genome-wide differences in the frequency of these alleles (3, 4). However, increasing evidence demonstrates that ecological divergence may occur surprisingly fast and even in absence of a physical barrier (5–8). As new beneficial mutations are unlikely to accumulate rapidly, these cases of fast adaptation likely involve selection on genetic variation that was present before divergence took place (9–11). Understanding the origin of and factors that maintain this variation is important as it largely determines the rate and direction of genetic adaptation to rapid environmental change (10, 12, 13).

Populations that have recently and repeatedly adapted to ecologically distinct conditions hold the promise to identify the alleles involved in ecological divergence and to reconstruct their evolutionary history (14–16). Under one scenario, these alleles could have evolved as rare neutral or mildly deleterious alleles within the ancestral population and become repeatedly selected (17). Alternatively, divergently selected genetic variants could initially have evolved and built-up in isolation. Secondary contact between the diverged populations may have then resulted in polymorphisms at these adaptive loci, providing the raw genetic material for repeated and rapid evolution when populations later face similar environmental conditions (18–

20). Hence, rapid and repeated ecological divergence could be the result of evolution at two different time frames. This latter idea is appealing as adaptive alleles have in this case already been subjected to a selective filter (10), which may also explain why examples of rapid adaptive divergence often involve parallel adaptation to the same selection pressure. However, complete evidence for this mechanism is currently lacking and requires that (i) differentially selected alleles at multiple unlinked loci diverged within the same time frame, which precedes the more recent parallel divergence events, (ii) differentially selected alleles evolved in at least partial isolation and (iii) populations of each ecotype are polymorphic at these loci to allow for the fast evolution of the alternative ecotype when the selection pressure changes.

Populations of the wing-polymorphic beetle *Pogonus chalceus* have adapted to two contrasting habitat types across Atlantic-Europe; tidal and seasonal salt marshes (Figure 1a). *Tidal* salt-marshes are inundated on an almost daily basis for at most a few hours and are inhabited by *P. chalceus* individuals that have a relatively small body size, short wings and submergence behavior during inundation. In contrast, salt-marshes that are subject to *seasonal* inundations that last for several months harbor *P. chalceus* individuals with a larger body size, fully developed wings and more frequent dispersal behavior (21–23). Populations of both the short- and long-winged ecotypes can be found along the Atlantic coastal region in Europe and often occur in close proximity and even sympatric mosaics (Figure 1b) (24). Despite evidence that ecological divergence is, at least partly, under genetic control (22, 24, 25), previous research based on microsatellite data also revealed very low neutral genetic differentiation within geographic locations (24). This suggests either a very recent differentiation and/or high levels of ongoing gene flow between these ecotypes. These observations render this a most suitable system to infer the origin of the allelic variants that underlie local adaptation (26).

Here we investigate genomic differentiation in multiple ecologically divergent population pairs of *P. chalceus* and reconstruct the evolutionary history of the alleles underlying ecological divergence. In agreement with a past, singular divergence event, we find sharing of the genealogical pattern at unlinked loci showing signatures of selection. However, the apparent potential of these beetles to rapidly and repeatedly adapt to the different tidal and seasonal hydrological regimes, is likely fueled by the maintenance of relatively high frequencies of the selected alleles in the alternative habitat. These results contribute to our understanding of the mechanisms underlying fast and parallel ecological adaptation and the factors determining the evolutionary potential of populations and species facing changing environments.

## Results

*Wing size distribution*

We sampled individuals in four population pairs inhabiting geographically close tidal and seasonally inundated habitats in Belgium (Be; 48 ind.), France (Fr; 48 ind.), Portugal (Po; 16 ind.) and Spain (Sp; 16 ind.), as well as a tidal marsh population in the UK (Uk; 8 ind.) and a seasonally inundated habitat at the Mediterranean coast of France (Me; 8 ind.) (Figure 1a). Individuals from the seasonally inundated habitats had significantly longer wings and larger body sizes compared to those from the tidally inundated marshes ($F_{1,117} = 1904.4$, $P < 0.0001$ for wing length and $F_{1,117} = 162.29$, $P < 0.0001$ for body size). The degree of divergence in wing length between the two ecotypes varied among the four population pairs ($F_{3,117} = 23.11$, $P < 0.0001$), with highly divergent wing lengths in Sp and Po, and some overlap in wing lengths in Be and Fr. Based on these clear-cut differences in wing length, we refer to the populations sampled in the tidal or seasonally inundated habitats as belonging to the short-winged (S) or long-winged (L) ecotype, respectively.

**Figure 1.** *Pogonus chalceus* **sampling and ecotypic divergence. (a.)** Sampling locations and density plots of the wing size distribution in the sampled populations. Blue indicates tidal habitats with short-winged beetles, red indicates seasonal habitats with long-winged beetles. BeS: Belgium-short, BeL: Belgium-long, FrS: France-short, FrL: France-long, UkS: UK-short, MeL: Mediterranean-long, PoS: Portugal-short, PoL: Portugal-long, Sps: Spain-short, SpL: Spain-long. Large circles represent populations included in the present study, small circles represent populations sampled previously (22, 27). **(b.)** Detail of the Fr population (Guérande) where short- and long-winged ecotypes occur in close proximity (< 20 m). This area consists of historic salt-extraction areas that were created by man approximately one thousand years ago and consists of a network of tidally inundated canals (FrS, partly indicated in blue; short-winged populations) interlaced with seasonally inundated salt extraction ponds (FrL, partly indicated in red; long-winged populations).

*Population structure and genome wide divergence and diversity*

Genetic differentiation ($F_{ST}$) among the 10 different populations, based on a quality filtered set of 15,756

SNPs distributed over 1,142 loci obtained by RAD-tag sequencing, varied considerably and ranged from a

low (BeS vs. UkS: $F_{ST} = 0.052$) to a high degree of differentiation (PoS vs. MeL: $F_{ST} = 0.37$) (Table S1).

Genetic differentiation increased significantly with increasing geographic distance between the populations

($r_S = 0.37$, $P = 0.017$) and was higher when populations belonged to a different ecotype ($r_S = 0.33$, $P = 0.02$)

(Figure 2a). When restricting the SNPs to a 'neutral' set wherein we excluded RAD-tags containing a SNP

with a signature of divergent selection (see *Outlier loci*), there was a stronger effect of geographic distance

on genetic differentiation ($r_S = 0.54$, $P = 0.002$), while the significant ecotype effect disappeared ($r_S = 0.09$,

$P = 0.2$). Principal Coordinate Analysis (PCoA) using all SNP data divided samples largely according to

ecotype along the first PCo axis, whereas the second PCo axis grouped samples according to geographic

location (Figure 2b). Bayesian clustering (28) of individuals based on their genotypes supported 8 and 6

genetically distinct populations ($K$) for the 'total' and 'neutral' SNP set, respectively (Figure 2c; Figure S1).

For the neutral SNP set, individuals from the same population pair (except Sp) clustered together as a single

population, irrespective of their ecotype.

Average nucleotide diversity ($\pi$) at RAD-tags did not differ between ecotypes (GLMM with RAD-tag ID as

random effect: Ecotype effect: $F_{1,\ 1977} = 0.31$, $P = 0.6$), but differed among population pairs (Population

effect: $F_{3,\ 1977} = 11.5$, $P < 0.0001$). The most southern populations had a significantly higher $\pi$ compared to

the northern populations, with Sp having the highest $\pi$ compared to all other populations ($P < 0.001$) and

Po having a higher $\pi$ compared to Fr ($P = 0.012$). The difference in nucleotide diversity among population

pairs was also consistent among ecotypes (Population*Ecotype interaction: $F_{3,\ 1977} = 2.2$, $P = 0.09$).

Despite the apparent close genetic relationship of long- and short-winged ecotypes within each geographic

population pair, we observed substantial heterogeneity in $F_{ST}$ across SNPs (Figure 3, Figure S2). A

substantial number of SNPs showed $F_{ST}$ values that exceeded 0.5 in the ecotype comparisons. For some of

these SNPs, different alleles even reached almost complete fixation in the different ecotypes. This

proportion of SNPs with $F_{ST}$ values higher than 0.5 increased towards the more southern population pairs

(Be: 2.1%, Fr: 4.5%, Po: 6.3% and Sp: 11.1%). In contrast, only very few $F_{ST}$ values exceeded 0.5 when

similar ecotypes were compared from different population pairs (e.g. Be versus Fr; Figure S2). SNPs that

were strongly differentiated in one particular population pair were also significantly more differentiated in

any of the other population pairs ($0.498 < r < 0.66$; $P$ all $< 0.0001$; Figure S2), providing evidence for

extensive sharing of highly differentiated SNPs among population pairs.

**Figure 2. Genetic differentiation among the studied *Pogonus chalceus* populations. (a.)** Relationship between $F_{ST}$ and geographic distance. Yellow and green dots depict $F_{ST}$ comparisons of populations from different or the same ecotype, respectively. Open dots represent $F_{ST}$ comparisons involving population Po. **(b.)** Principal Coordinate Analysis (PCoA) for RAD-tag sequenced samples. **(c.)** Population structure of the ten *P. chalceus* populations based on Bayesian clustering (28). The best supported number of clusters was 8 for all RAD-tags and 6 for neutral RAD-tags (see Figure S1). See Figure 1 for population codes.

*Outlier loci*

We identified a total of 512 (3.2%) SNPs in 109 (15%) paired RAD-tags with stronger differentiation as expected by chance in at least one of the ecotype comparisons (BayeScan (29) with false discovery rate = 0.05). A total of 75 (0.48%) SNPs in 32 (6.3%) paired RAD-tags showed allele frequencies that were strongly associated with the ecotypic divergence across all investigated populations (BayEnv2 (30); $\log_{10}$BF = 4) (see Supplementary material - outlier loci; Figure S3 and S4). RAD-tags were mapped to a genome assembly (see Supplementary material – genome assembly and linkage map), showing that SNPs with a high $F_{ST}$ value clustered into several unlinked regions that were distributed over a large proportion of the genome (Figure 3). These regions of elevated divergence were largely consistent across the different population pairs and are primarily clustered on the first half of LG1, across the full length of LG2 and LG3 and at the center of LG4 (Figure 3). In contrast, no elevated levels of differentiation or outlier loci were observed on LG6 to LG10. Yet, some more subtle differences could be observed as exemplified by the central region of LG5 where high genomic differentiation was only observed for the Fr and Sp population pair, but not in the Be and Po population comparison.

**Figure 3. Genomic divergence ($F_{ST}$) between *Pogonus chalceus* ecotype pairs.** Outlier SNPs within each population pair, as identified by BayeScan (29), are indicated in red. Size of SNPs is proportional to the $\log_{10}BF$ reported by BayEnv2 (30) and indicates the degree of support that allele frequencies are significantly correlated with habitat type across all sampled populations. Markers on LG10 are significantly sex-linked.

*Quantifying standing genetic variation*

We estimated the potential of each ecotype to adapt to the alternative habitat by calculating the frequency of alternative alleles at outlier loci for both ecotypes. For individuals of the long-winged ecotype, on average 10% (SpL) to 42% (BeL) of the outlier loci contained at least one allele selected in the alternative, short-winged, ecotype. A similar pattern was found in the opposite direction, with individuals from short-winged populations containing long-wing selected alleles at 10% (SpS) to 48% (BeS) of the outlier loci. Moreover, random sampling of genomic variation in an increasing number of individuals showed a steep increase in the proportion of outlier loci with at least one allele selected in the alternative habitat (Figure 4). This demonstrates that different individuals from the same population generally carry different alleles that are selected in the alternative habitat at different loci. For example, more than 80% of the full set of 75 outlier loci detected with BayEnv2 contain at least one short-wing selected allele in a random sample of only eight long-winged individuals for the populations BeL, FrL and PoL (Figure 4a). This demonstrates the presence of substantial standing genetic variation in individuals sampled in the seasonally inundated marshes to adapt to tidal marshes. Only for the most southern long-winged populations (SpL and MeL), short-wing selected alleles are present at lower frequencies and these long-winged populations are unlikely to contain the full set of alleles selected in the short-winged ecotype. For FrS, PoS and particularly SpS, long-wing selected alleles accumulated at a much lower rate under random sampling of short-winged individuals (Figure 4b).

**Figure 4. Amount of standing genetic variation.** Accumulation curves of **(a.)** the proportion of outlier loci containing at least one short-wing selected allele in a random sample of *N* long-winged individuals and **(b.)** the proportion of outlier loci containing at least one long-wing selected allele in a random sample of *N* short-winged individuals.

*Sequence variation and phylogenetic reconstruction at outlier loci*

Haplotype networks and trees of the 1.2 kb sequence alignments obtained from RAD-tag loci show that haplotypes selected in short-winged populations generally clustered as a strongly supported monophyletic clade of closely related sequences (clade support level > 0.96; Figure 5a, b and S5). This clustering supports a singular mutational origin of short-wing selected alleles at each of the investigated outlier tags. Short-wing selected alleles appeared to be derived as they most frequently constituted a subclade within those selected in long-winged populations (Figure S5). This is in line with the observation that all other species within the genus *Pogonus* are long-winged (31). The average absolute divergence between the differentially selected haplotypes ($d_{XY} = 0.011 \pm 0.0014$) was about 1.65 times higher compared to the average divergence between two randomly chosen haplotypes at these loci ($\pi_{tot, outliers} = 0.0067 \pm 0.00097$, *t*-test: $P < 0.0001$) and highlights a deep divergence between the alleles selected in short- and long-winged populations. Dating the divergence time between the differentially selected alleles using BEAST (32) and the divergence from *P. littoralis* as a calibration point (0.62 Mya (27)), pointed towards highly comparable divergence times among outlier loci (Figure 5b and S5). The divergence time between long- and short-wing selected alleles ranged between 0.12 Mya and 0.28 Mya, with an average of 0.189 Mya ± 0.09 Mya, and suggests that the divergence of short-winged alleles from the long-winged alleles took place during the Late Pleistocene.

**Figure 5. Haplotype structure and diversity at divergently selected loci.** (a.) Haplotype networks of
RAD-tags containing outlier SNPs and at least 10 variable sites (BayEnv2; log10BF > 4) at the different
linkage groups. Haplotypes selected in short-winged populations are depicted in blue, haplotypes selected
in long-winged populations are depicted in red. (b.) Estimated divergence time (Mya) between alleles
selected in short-winged (blue) versus long-winged (red) populations. The tree represents the general
phylogenetic relationship between short- and long-wing selected alleles and the estimated divergence point.
(c.) Relationship between $F_{ST}$ and Tajima's $D$ and (d.) absolute nucleotide divergence, $d_{XY}$, scaled relative
to the divergence from the outgroup species *Pogonus littoralis* in the four population pairs. (e.) Comparison
of nucleotide diversity (π) and Tajima's $D$ at neutral loci of long-winged (L) and short-winged (S)
populations (left) and between haplotypes at outlier RAD-tags selected in L or S populations (right).

In agreement with divergent selection, sequences from more strongly differentiated RAD-tags had a
significantly higher Tajima's $D$ (i.e. increased heterozygosity; $F_{ST}$: $F = 57.36$, $P < 0.0001$; Figure 5c) and
absolute nucleotide divergence between the ecotypes normalized by the divergence from the outgroup $P.$
$littoralis$ ($= d_{XY}/ d_{XY, P. littoralis}$, see Methods for details; $F_{ST}$: $F = 83.9$, $P < 0.0001$; Figure 5d). The significant
relation between $F_{ST}$ and absolute divergence (normalized $d_{XY}$) further supports that the observed
heterogeneity in genomic divergence between the ecotypes is the result of divergent selection of ancient
alleles embedded within a genome that is homogenized between the ecotypes, rather than selection on
recently obtained new mutations (33, 34). Furthermore, a reduced recombination rate was observed between
haplotypes that are divergently selected between long- and short-winged populations ($r^2 = 0.140$) compared
to the recombination rate observed within long-winged haplotypes ($r^2 = 0.184$).

We further observed that nucleotide diversity ($\pi$) of haplotypes selected in the short-winged ecotype was
strongly reduced and tended to be nearly seven times lower ($\pi_S = 0.0009 \pm 0.0003$) compared to those
selected in the long-winged ecotype ($\pi_L = 0.0062 \pm 0.0003$) (GLMM with tagID as random effect: Ecotype
effect: $F_{1, 66} = 25.12$, $P < 0.0001$; Figure 5e). This difference was consistent among the four populations
(Ecotype*Population interaction: $F_{3, 64} = 0.87$; $P = 0.45$). In contrast, nucleotide diversity at neutral RAD-
tags was comparable between both ecotypes (GLMM with RAD-tag as random effect: Ecotype effect: $F_{1,}$
$_{599} = 0.98$, $P < 0.3$; Figure 5e). The nucleotide diversity of the haplotypes selected in long-winged
populations was also comparable to average nucleotide diversity observed at non-outlier loci ($\pi_{tot, neutral RAD-}$
$_{tags} = 0.0057 \pm 0.0003$), showing that only haplotypes selected in the short-winged ecotype have this reduced
nucleotide diversity (Figure 5e). Similarly, Tajima's $D$ of haplotypes selected in the short-winged ecotype
was significantly lower compared to those selected in the long-winged ecotype ($F_{1,27} = 11.7$; $P = 0.002$;
Figure 5e) and suggests a recent spread of short-wing selected alleles along the Atlantic European coast.
241
242

**Discussion**
Understanding the genomic basis of repeated and fast ecological adaptation provides unique insights into
the process of evolutionary diversification (14, 15, 35). While evidence is accumulating that cases of
repeated adaptation are largely driven by selection on standing genetic variation (36), the evolutionary origin
of the selected alleles remains currently poorly understood (20, 37). In *P. chalceus,* several unique
observations help to disentangle the complex history of fast and parallel ecological divergence and the origin
of the adaptive alleles that underlie it. These observations include: (i) Diverged ecotypes within a geographic
region share genomic variation at neutral loci but are characterized by extensive genomic islands of
divergence, widely distributed across the genome. (ii) Genomic regions of elevated divergence are highly
consistent across the four population pairs. (iii) Haplotypes of alleles selected in short-winged populations

cluster monophyletically and are derived. (iv) The pattern of divergence across short-wing selected alleles
and divergence time estimates are comparable among loci. (v) Short-wing associated haplotypes show
strongly reduced diversity, negative Tajima's *D* and reduced recombination with the long-wing associated
haplotypes. (vi) A relatively small subsample of individuals from long-winged populations harbor the
majority of alleles selected in short-winged populations, and vice versa. In summary, these observations
imply that the ecotypes diverged *in situ* within each geographic region, but that the adaptive alleles that
constitute the short-winged ecotype have a singular evolutionary origin and likely evolved within an ancient,
more isolated, subpopulation. The time at which short-wing selected alleles diverged from those selected in
long-winged populations was estimated to be about 0.19 Mya. Subsequent admixture between the long- and
short-winged subpopulation may then have resulted in a genetically polymorphic population from which
the short-winged ecotype re-evolved more recently and repeatedly along the European Atlantic coast by the
concerted selection of these previously diverged alleles.

Despite the ancient divergence of the adaptive alleles, the contemporary repeated distribution of the *P.*
*chalceus* ecotypes likely evolved from highly admixed populations. The low amount of neutral genetic
differentiation should, however, not be interpreted as conclusive evidence for a repeated *in situ* divergence,
because it also may be the result of neutral gene flow between both ecotypes at secondary contact (37–39).
Nonetheless, several additional observations support that the current distribution of the ecotypes involves
the repeated *in situ* evolution of both ecotypes. First, quantifying the amount of short-wing selected alleles
present in long-winged populations revealed that sufficient genetic variation is present in a founding
population of between 5 and 15 long-winged individuals to evolve a short-winged ecotype. Thus, genetic
constraints for the evolution of the short-winged ecotype out of long-winged individuals, and vice versa,
appear to be surprisingly low. Both weak selection as well as high rates of ongoing gene flow between the
ecotypes could explain the relatively high frequency of alleles that are selected in the alternative ecotype
(40). Second, short-winged individuals are unable to disperse by flight between the currently highly isolated
salt-marsh areas. The fragmented distribution of tidal salt marshes along the Atlantic coast renders it
therefore unlikely that they were colonized by short-winged individuals based on terrestrial dispersal alone,
in particular because the species strongly avoids unsuitable habitat patches (31). Third, we previously put
forward a behavioral mechanism that may explain the spatial sorting of these ecotypes into their respective
habitats and which may induce *in situ* divergence of genetically distinct ecotypes (23). Adding to this, the
initial evolution of co-adapted alleles at multiple physically unlinked loci in geographic isolation may have
facilitated the repeated ecological divergence under gene flow (41).

Major geographic expansions and contractions of the tidal and seasonal habitat types and the *P. chalceus*
species distribution have likely occurred since the initial divergence of the short-winged ecotype. We
estimated the evolution of the short-winged ecotype to have occurred during the Mid to Late Pleistocene.
Since then, Europe has been subject to at least one interglacial (0.130 – 0.115 Mya) and one glacial (0.115
– 0.012 Mya) period. These major climatic changes fragmented the Euro-Atlantic coastline, potentially
creating opportunities for the initial evolution of the short-winged ecotype in the partially isolated large
coastal floodplains that extended, for instance, around the North-Sea basin (42). Suggested by the low
nucleotide diversity among the short-wing selected alleles, secondary contact between the short- and long-
winged ecotypes likely took place with small refuge populations of the short-winged ecotype that persisted
during the last glacial period. These refuge populations were likely located more southward relative to the
current species distribution, as it is deemed unlikely that the species persisted at the current northern latitudes
of its distribution during the last glacial maximum (31). In addition, the lower degree of overall genetic
differentiation between the ecotypes in the more northern population pairs Be and Fr, together with their
higher frequencies of alleles selected in the alternative habitat, less profound phenotypic differentiation and
lower overall genetic diversity ($\pi$) are all consistent with a northward expansion of a highly admixed and
dispersive long-winged population and more recent ecotypic divergence. Potentially, this expansion may
have further facilitated the maintenance of deleterious short-wing selected alleles in the expanding long-
winged population (43), which then became repeatedly selected in the emergent tidal (short-winged)
habitats.
305
The two-step process of initial divergence in an ancient more isolated population and subsequent admixture
putatively also applies to other examples of fast and repeated ecological divergence. Repeated ecological
divergence at the same loci has been reported in some iconic examples of parallel evolution, such as
stickleback, cichlid fishes and *Heliconius* butterflies (44–48). These loci have in many cases been assigned
to shared ancient polymorphisms that were present in the population before the evolution of the currently
observed divergent populations. Moreover, many of these loci are sometimes identified and are unlinked
throughout the genome, such as in fruit flies, *Timema* walking sticks and *Littorina* sea snails (7, 49, 50).
The genetic signature of the evolution of the *P. chalceus* ecotypes shows strong analogies to these well-
studied cases of repeated adaptation. In cichlid fishes, moreover, it has been extensively argued that
divergence in isolation and subsequent admixture may have provided the genetic material for the incredibly
diverse and recent adaptive radiations of cichlid fish (11, 51). Untangling the evolutionary history of the
alleles involved in these and other cases will help in better understanding the processes that drive parallel
divergence as well as fast responses to environmental change

**Conclusion**

The initial evolution of co-adapted alleles at multiple physically unlinked loci is facilitated in geographic isolation (3). Subsequent admixture of gene pools may then enrich the adaptive genetic variation and allow for subsequent fast and repeated adaptation. In agreement to this, we demonstrated for the *P. chalceus* populations that the alleles required to adapt to the alternative environment can be maintained in the source population. These loci are expected to be maladaptive within the source population and it is likely both the temporal and spatial repetition of this divergence, combined with relatively high levels of gene flow and range expansion, that maintain these allele frequencies. Glacial cycles, in particular, can be expected to have played an important role in this process as episodes of fission and fusion of the different ecotypes generates strong opportunities for both the evolution of adaptive genetic variants as well as admixture (42). The proposed mechanism illustrates that the distinction between *in-situ* divergence followed by secondary contact is less clear-cut as generally assumed and that both processes can be involved at different time frames. An important implication is that this mechanism might reconcile different views on the geography of ecological divergence in which adaptive divergence between closely related populations is either interpreted as primary divergence, and thus the onset of speciation (2), or the result of secondary introgression after initial ecological divergence in allopatry (37–39).

**Methods**

*Sampling*

Diverged population pairs of *P. chalceus* were collected from both tidal and seasonal salt marshes extending

nearly the entire species range (Figure 1) (31). We sampled four geographically isolated population pairs

(separated between approximately 450 km and 900 km) of a tidal and seasonally flooded inland population

each. Wing and elytral sizes were measured by means of a calibrated ocular with a stereomicroscope. We

further conducted RAD-seq genotyping on two specimens of the long-winged outgroup species *P. littoralis*,

which were sampled in the Axion Delta, Thessaloniki, Greece (Table S2).

*RAD-tag sequencing*

DNA was extracted using the DNA extraction NucleoSpin® Tissue kit (Macherey-Nagel GmBH). Extracted

genomic DNA was normalized to a concentration of 7.14 ng/µl and processed into RAD libraries according

to Etter et al. (2011), using the restriction enzyme SbfI-HF (NEB) (52). Final enrichment was based on 16

PCR cycles. A total of nine RAD libraries including 16 individuals each and, hence, a total of 144

individuals were sequenced paired-end for 100 cycles (i.e. 100 bp) in a single lane on an Illumina HiSeq2000

platform according to manufacturer's instructions. The outgroup *P. littoralis* specimens were sequenced

separately. The raw data was demultiplexed to recover individual samples from the Illumina libraries using

the *process_radtags* module in Stacks v1.20 software (53). Reads were quality filtered when they contained

15 bp windows of mean Phred scores lower than 10. PCR duplicates were identified as almost (i.e. allowing

for sequencing errors) identical reverse read sequences and removed, using a custom Perl script (54).

*Genome assembly*

Total DNA was extracted from individuals captured in the canal habitat of the salt marshes in the Guérande

region (France), using the DNA extraction NucleoSpin® Tissue kit (Macherey-Nagel GmBH). Illumina

paired-end (100 bp) and mate-paired (49 bp) libraries were constructed with insert sizes of 200 bp, 500 bp,

800 bp, 2 kb and 5 kb and sequenced on an Illumina HiSeq2000 system according to the manufacturer's

protocol (Illumina Inc.). Adapter contamination in reads was deleted using Cutadapt v1.4 (55) and reads

that did not have a matching pair after adaptor filtering were removed. Reads were corrected for sequencing

error with SOAPec v2.02 (56), using a *k*-mer size of 17 and a low frequency cutoff of consecutive *k*-mer of

3. Sequencing of the 200 bp, 500bp, 800 bp, 2 kb and 5 kb insert libraries resulted in a total of ~57.7 Gb of

sequencing data, of which 56.6 Gb was retained after data cleaning (Table S3). Reads were assembled using

SOAPdenovo2 (56) using a *k*-mer parameter of 47, which was selected for producing the largest contig and

scaffold N50 size after testing a range of *k*-mer settings between 19 and 71. The short insert libraries were

used for both contig building and scaffolding. The long insert libraries were only used for scaffolding.

SOAPdenovo GapCloser v1.12 tool (56) was used with default settings to close gaps emerging during

scaffolding. We used DeconSeq v0.4.3 (57) to identify and remove possible human, bacterial and viral

contamination in the assembly (Table S4). Completeness of the assembled genome was assessed by

comparing the assembly with a dataset of highly conserved core genes that occur in a wide range of

eukaryotes using the CEGMA pipeline v2.5 (58).

*Linkage map*

To position the genomic scaffolds into linkage groups, we constructed a linkage map by genotyping parents

and offspring (RAD-seq) from four families (Table S5). For the parental generation, we used lab-bred

individuals (F0) whose parents originated from the French population, to ensure that they had not been

mated in the field. A total of 72 F1 offspring ($n$ = 23, 14, 23 and 12 offspring from each family) were raised

till adulthood and subsequently genotyped, together with their parents. To maximize the number of scaffolds

comprising a marker, RAD-tag sequencing was based on a *Pst*I-HF (NEB) digest (6bp recognition site)

instead of the SbfI-HF (NEB) digest (8bp recognition site) of the population genomic analysis. Final

enrichment was based on 16 PCR cycles. Illumina HiSeq sequencing resulted in a total of 237 M paired-

end reads, of which 113 M remained after quality filtering and removal of PCR duplicates. Reads were

mapped to the draft reference genome with BWA-mem (59) using default settings. Linkage map

reconstruction was performed with LepMAP2 (60). LepMAP2 reconstructs linkage maps based on a large

number of markers and accounts for lack of recombination in males due to achiasmatic meiosis, which is

suggested in *P. chalceus* and male Caraboidea in general (61) (see Supplementary material – genome

assembly and linkage map).

*Population genomic analysis*

Quality and clone filtered paired-end reads of the 144 field captured individuals were mapped to a draft

reference genome with BWA-mem (59). Indel realignment, SNP and indel calling was performed with

GATK's UnifiedGenotyper tool (62). Paired-end sequencing of the approximately 200 to 600 bp RAD tag

fragments adjacent to symmetric SbfI restriction sites allowed us to obtain sequence information of 1,200

bp fragments (paired RADtag) around each restriction site. Hence, after SNP calling we retained all sites

within 1,200 bp windows around each SbfI recognition site in the genome, totaling 732,884 bp of sequence.

Haplotype phasing was subsequently first performed with GATK 'read-backed phasing' (62), while the

remaining unphased sequences were phased with Beagle v4.1 (63). A reliable SNP set was then obtained

by filtering out SNPs with minimum genotype quality lower than 20, minimum average depth lower than

10 and a minor allele frequency less than 0.01. This resulted in 27,757 SNPs, distributed over 712 paired

RADtags, with an average individual depth of 62.9 ($\pm$ 51 std), of which 15,756 were present in at least 80%

of the individuals.

*Analysis of population structure*

Pairwise $F_{ST}$-statistics (64) across RAD-tags were calculated for all pairwise population comparisons using

Genepop v4.5.1 (65). Principal Coordinate Analysis was performed using *adegenet* in R (66). To minimize

dependence due to physical linkage among SNPs, we randomly selected one single SNP per paired RAD-

tag. We also constructed a 'neutral' subset by excluding SNPs located on scaffolds showing signatures of

divergent selection (20.5% of all SNPs). As a criterion, we excluded scaffolds containing a SNP with a

$\log_{10}(BF) > 3$ as determined by BayEnv (67). The Pearson correlation between genetic divergence and either

geographic distance between the populations or ecotype (coded as $1 = $ different ecotype and $0 = $ identical

ecotype) was assessed by a Mantel test in the vegan v2.2-1 package in R v3.1.3 (68). Based on these two

datasets, we used the Bayesian clustering algorithm implemented in STRUCTURE v2.3.4. (28) to assign

individuals into $K$ clusters based on their multilocus genotype. We applied an admixture model with three

independent runs for each $K = $ 2-10, 100,000 MCMC repetitions with a burn-in of 30,000, correlated allele

frequencies among populations and no prior information on population origin. Default settings were used

for the prior parameters. The best supported number of clusters ($K$) was determined from the increase in the

natural logarithm of the likelihood of the data for different numbers of assumed populations.

*Outlier loci detection*

Support for loci showing significantly higher degrees of differentiation was first detected with BayeScan2.1

(29) within each population pair (Be, Fr, Po and Sp). BayeScan assumes that divergence at each locus

between populations is the result of population specific divergence from an ancestral population as well as

a locus specific effect. The prior odds of the neutral model was set to 10. Twenty pilot runs, 5,000 iterations

each, were set to optimize proposal distributions and final runs were performed for 50,000 iterations,

outputting every tenth iteration, and a burn-in of 50,000 iterations. Detection of outlier loci is particularly

vulnerable to false positives (69). To account for this, we applied a false discovery rate (FDR) correction of

0.05, meaning that the expected proportion of false positives is 5% (29).

To test for the presence of SNPs whose alleles are directionally selected in the two habitats across all

populations, we used the approach implemented in BayEnv2 (30, 67). This method identifies SNPs whose

allele frequencies are strongly correlated with an environmental variable given the overall covariance in

allele frequencies among populations. The covariance in allele frequencies, which represents the null model

against which the effect, $\beta$, of an environmental variable on the allele frequencies of each SNP is tested, was

estimated based on all SNPs present in at least 80% of the individuals. This covariance matrix was strongly

correlated with the $F_{ST}$ matrix (Mantel-test: $r_S$= 0.87), indicating that it accurately reflects the genetic

structuring of the populations. For each SNP, the posterior probability of a null model assuming no effect

of the environment ($\beta = 0$) is compared against the alternative model which includes the effect of the

environmental variable. As environmental variable, we assigned tidal habitats (BeS, FrS, PoS, SpS and UkS)

the value -1 and seasonal inundated habitats (BeL, FrL, PoL, SpL and MeL) the value 1. The degree of

support that variation at a SNP covaries with the habitat wherein the population was sampled is then given

by the Bayes Factor (BF), the ratio of the posterior probabilities of the alternative versus the null model. For

both the estimation of the covariance structure and the environmental effect, a total of 100,000 iterations

was specified.

*Reconstructing the evolutionary history of outlier loci*

To gain insight into the evolutionary history of the alleles differentiating the two ecotypes, we reconstructed

haplotypes of the 1200 bp long paired RAD-tag loci for each individual. Sites with a read depth lower than

10 or a genotype quality lower than 20 were treated as missing. Haplotypes could be reconstructed for 627

paired RAD-tags with on average 671 bp genotyped in at least 75 % of the individuals. We constructed split

networks with the NeighbourNet algorithm using SplitsTree4 (70) for all RAD-tags that contained an outlier

SNP with a Bayes Factor (BF) support level larger than 3 based on the BayEnv2 analysis. Haplotypes were

subsequently split in two groups according to base composition at the outlier SNP with the highest support

and visualized on the networks.

We further calculated for all RAD-tags the following haplotype statistics with the EggLib v2.1.10 Python

library (71): haplotype based $F_{ST}$ (72), average pairwise difference ($d_{XY}$) between both ecotypes, total

nucleotide diversity ($\pi_{tot}$), nucleotide diversity within the long- and short-winged ecotype ($\pi_L$ and $\pi_S$,

respectively) and Tajima's *D*. Comparison of measures of $d_{XY}$ ($\approx 2\mu t + \theta_{Anc}$) and $\pi$ ($\approx 4N\mu$) between RAD-

tags depend, besides the average coalescence time between haplotypes, also on the mutation rate ($\mu$) of the

RAD-tag. As we are primarily interested in comparing values of these statistics among RAD-tags

independent of their mutation rate, we normalized these values by the average number of nucleotide

differences between *P. chalceus* and the outgroup species *P. littoralis* (73). More specifically, we first

calculated $d_{XY}$ between haplotypes of *P. chalceus* and *P. littoralis* ($d_{xy,\,littoralis}$), and divided both $\pi_{tot}$ and $d_{XY}$

by this value.

We estimated the divergence time between haplotypes selected in short- and long-winged populations with

BEAST 1.7.1 (32). The analysis was restricted to outlier RAD-tags (15 in total) that are also present in the

outgroup species *P. littoralis* and that contained on average at least 10 segregating sites among the sequences

of *P. chalceus*. This latter criterium was implemented to ensure a sufficiently high substitution rate for

reliable time calibration. The tree was calibrated using the divergence from *P. littoralis*, estimated at 0.62 ±

0.8MY, as calibration point (27). We assumed a GTR substitution model, a strict clock model and standard

coalescent tree prior. Analyses were run by default for 10 million generations of which the first 2 million

generations were treated as burn-in and discarded for the calculation of posterior probability estimates.

**Data accessibility**

476    Raw sequencing reads are available in the NCBI Short Read Archive under BioProject PRJNA381601. The

477    genome assembly, ordered using the linkage map, is available under accession NEEE00000000. Reads of

478    genome assembly: SAMN06684244-SAMN06684249; RAD-seq data of *Pogonus chalceus*:

479    SAMN06691389-SAMN06691532; RAD-seq data of *Pogonus littoralis*: SAMN06691533-

480    SAMN06691534; RAD-seq data for linkage map construction: SAMN06806679- SAMN06806758.

**Acknowledgments**

This study could not have been performed without access to *P. chalceus* collection of the late Konjev

Desender. We are grateful to Alexandre Ramos, Viki Vandomme and Lut Van Nieuwenhuyse for help in

collecting the samples, Jonas Van Belleghem for measuring wings and Simon Martin for valuable

bioinformatics help, Karim Gharbi (Edinburgh Genomics Center) for extensive support in the preparation

of the RAD-seq libraries and Chris Jiggins for valuable comments on the manuscript. This work was

supported by funding received from the FWO-Flanders (PhD grant to SVB) and the Belgian Science Policy

(MO/36/ 025, BR/121/PI/GENESORT and BR/175/PI/PARAWINGS to FH) and was partly conducted

within the framework of the Interuniversity Attraction Poles program IAP (SPEEDY) – Belgian Science

Policy. The genomic analyses were carried out using the STEVIN Supercomputer Infrastructure at Ghent

University, funded by Ghent University, the Flemish Supercomputer Center (VSC), the Hercules

Foundation and the Flemish Government – Department EWI.

**Author contributions**

498    SMVB and FH designed the research and analyzed the data. SMVB, CV, KDW and FH performed the

499    research. SMVB, FH, CV, ZDC, MM and LDM wrote the paper. PR contributed analytical tools.

**References**

1.    Schluter D (2009) Evidence for ecological speciation and its alternative. *Science* 323:737–741.

2.    Nosil P (2012) *Ecological Speciation* (Oxford University Press, Oxford, UK).

3.    Coyne JA, Orr HA (2004) *Speciation* (Sinauer Associates, Sunderland, MA, USA).

4.    Feder JL, Flaxman SM, Egan SP, Comeault A a., Nosil P (2013) Geographic mode of speciation and genomic
divergence. *Annu Rev Ecol Evol Syst* 44:73–97.

5.    Arnegard ME, et al. (2014) Genetics of ecological divergence during speciation. *Nature* 511(7509):307–311.

6.    Richardson JL, Urban MC, Bolnick DI, Skelly DK (2014) Microgeographic adaptation and the spatial scale of
evolution. *Trends Ecol Evol* 29(3):165–176.

7.    Riesch R, et al. (2017) Transitions between phases of genomic differentiation during stick-insect speciation.
*Nat Ecol Evol* 1:82.

8.    Hendry AP (2016) *Eco-evolutionary dynamics* (Princeton University Press, Princeton, US).

9.    Colosimo PF, et al. (2005) Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin
alleles. *Science* 307:1928–1933.

10.    Barrett RDH, Schluter D (2008) Adaptation from standing genetic variation. *Trends Ecol Evol* 23(1):38–44.

11.    Meier JI, et al. (2017) Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat Commun* 8:14363.

12.    Barton NH, Keightley PD (2002) Understanding quantitative genetic variation. *Nat Rev Genet* 3(1):11–21.

13.    Savolainen O, Lascoux M, Merilä J (2013) Ecological genomics of local adaptation. *Nat Rev Genet*
14(11):807–820.

14.    Arendt J, Reznick D (2008) Convergence and parallelism reconsidered: what have we learned about the
genetics of adaptation? *Trends Ecol Evol* 23(1):26–32.

15.    Elmer KR, Meyer A (2011) Adaptation in the age of ecological genomics: insights from parallelism and
convergence. *Trends Ecol Evol* 26(6):298–306.

16.    Rosenblum EB, Parent CE, Brandt EE (2014) The molecular basis of phenotypic convergence. *Annu Rev Ecol*
*Evol Syst* 45(1):203–226.

17.    Orr AH, Betancourt AJ (2001) Haldane's seive and adaptation from the standing genetic variation. *Genetics*
157(1977):875–884.

18.    Feder JL, et al. (2003) Allopatric genetic origins for sympatric host-plant shifts and race formation in
*Rhagoletis*. *Proc Natl Acad Sci* 100(18):10314–10319.

19.    Schluter D, Conte GL (2009) Genetics and ecological speciation. *Proc Natl Acad Sci U S A* 106(1):9955–9962.

20.    Welch JJ, Jiggins CD (2014) Standing and flowing: The complex origins of adaptive variation. *Mol Ecol*
23(16):3935–3937.

21.    Dhuyvetter H, Gaublomme E, Desender K (2004) Genetic differentiation and local adaptation in the salt-marsh
beetle *Pogonus chalceus*: a comparison between allozyme and microsatellite loci. *Mol Ecol* 13(5):1065–1074.

22.    Van Belleghem SM, Hendrickx F (2014) A tight association in two genetically unlinked dispersal related traits
in sympatric and allopatric salt marsh beetle populations. *Genetica* 142:1–9.

23.    Van Belleghem SM, De Wolf K, Hendrickx F (2016) Behavioral adaptations imply a direct link between
ecological specialization and reproductive isolation in a sympatrically diverging ground beetle. *Evolution*
70(8):1904–1912.

24.    Dhuyvetter H, Hendrickx F, Gaublomme E, Desender K (2007) Differentiation between two salt marsh beetle
ecotypes: evidence for ongoing speciation. *Evolution* 61(1):184–193.

25.    Desender K (1989) Heritability of wing development and body size in a carabid beetle, *Pogonus chalceus*
Marsham, and its evolutionary significance. *Oecologia* 78:513–520.

26.    Raeymaekers JAM, Backeljau T (2015) Recurrent adaptation in a low-dispersal trait. *Mol Ecol* 24(4):699–701.

27.    Van Belleghem SM, Roelofs D, Hendrickx F (2015) Evolutionary history of a dispersal-associated locus across
sympatric and allopatric divergent populations of a wing-polymorphic beetle across Atlantic Europe. *Mol Ecol*
24:890–908.

28.    Falush D, Stephens M, Pritchard JK (2007) Inference of population structure using multilocus genotype data:
dominant markers and null alleles. *Mol Ecol Notes* 7(4):574–578.

29.    Foll M, Gaggiotti O (2008) A genome-scan method to identify selected loci appropriate for both dominant and
codominant markers: a Bayesian perspective. *Genetics* 180(2):977–93.

30.    Günther T, Coop G (2013) Robust identification of local adaptation from allele frequencies. *Genetics*
195(1):205–220.

31.    Turin H (2000) *De Nederlandse loopkevers: verspreiding en ecologie* (KNNV Uitgeverij).

32.    Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol*

7:214.

33. Cruickshank TE, Hahn MW (2014) Reanalysis suggests that genomic islands of speciation are due to reduced
diversity, not reduced gene flow. *Mol Ecol* 23(13):3133–3157.

34. Han F, et al. (2017) Gene flow, ancient polymorphism, and ecological adaptation shape the genomic landscape
of divergence among Darwin's finches. *Genome Res*:1–12.

35. Stern DL (2013) The genetic causes of convergent evolution. *Nat Rev Genet* 14(11):751–764.

36. Conte GL, Arnegard ME, Peichel CL, Schluter D (2012) The probability of genetic parallelism and
convergence in natural populations. *Proc R Soc B* 279(1749):5039–5047.

37. Faria R, et al. (2014) Advances in Ecological Speciation: an integrative approach. *Mol Ecol* 23(3):513–521.

38. Bierne N, Gagnaire PA, David P (2013) The geography of introgression in a patchy environment and the thorn
in the side of ecological speciation. *Curr Zool* 59(1):72–86.

39. Foote AD (2017) Sympatric speciation in the genomic era. *Trends Ecol Evol* 2327:1–11.

40. Stuart YE, et al. (2017) Contrasting effects of environment and genetics generate a continuum of parallel
evolution. *Nat Ecol Evol* 1(158):1–7.

41. Fry JD (2003) Multilocus models of sympatric speciation: Buch versus Rice versus Felsenstein. *Evolution*
57(8):1735–1746.

42. Busschers FS, Kasse C, Balen RTV, Vandenberghe J, Cohen KM (2007) Late Pleistocene evolution of the
Rhine-Meuse system in the southern North Sea basin: imprints of climate change, sea-level oscillation and
glacio-isostacy. *Quat Sci Rev* 26:3216–3248.

43. Travis JMJ, et al. (2007) Deleterious mutations can surf to high densities on the wave front of an expanding
population. *Mol Biol Evol* 24(10):2334–2343.

44. Hohenlohe P a, Phillips PC, Cresko W a (2010) Using population genomics to detect selection in natural
populations: key concepts and methodological considerations. *Int J Plant Sci* 171(9):1059–1071.

45. Jones FC, et al. (2012) The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*
484(7392):55–61.

46. Brawand D, et al. (2014) The genomic substrate for adaptive radiation in African cichlid fish. *Nature* 513:375–
381.

47. Colosimo PF, et al. (2005) Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin
alleles. *Science* 307:1928–1933.

48. Van Belleghem SM, et al. (2017) Complex modular architecture around a simple toolkit of wing pattern genes.
*Nat Ecol Evol* 1(3):52.

49. Michel AP, et al. (2010) Widespread genomic divergence during sympatric speciation. *Proc Natl Acad Sci U*
*S A* 107(21):9724–9729.

50. Ravinet M, et al. (2016) Shared and nonshared genomic divergence in parallel ecotypes of *Littorina saxatilis*
at a local scale. *Mol Ecol* 25:287–305.

51. Meier JI, Sousa VC, Marques DA, Selz OM (2017) Demographic modelling with whole-genome data reveals
parallel origin of similar *Pundamilia* cichlid species after hybridization. *Mol Ecol* 26:123–141.

52. Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA (2011) SNP discovery and genotyping for
evolutionary genetics using RAD sequencing. *Methods Mol Biol* 772(2):1–19.

53. Catchen J, Hohenlohe P a, Bassham S, Amores A, Cresko W a (2013) Stacks: an analysis tool set for population
genomics. *Mol Ecol* 22(11):3124–3140.

54. Kerth C (2012) Scripts for RAD. https://github.com/claudiuskerth/scripts_for_RAD.

55. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet*
17:10–12.

56. Luo R, et al. (2012) SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler.
*Gigascience* 1(1):18.

57. Schmieder R, Edwards R (2011) Fast identification and removal of sequence contamination from genomic and
metagenomic datasets. *PLoS One* 6(3):e17288.

58. Parra G, Bradnam K, Korf I (2007) CEGMA: a pipeline to accurately annotate core genes in eukaryotic
genomes. *Bioinformatics* 23(9):1061–1067.

59. Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
*arXiv*:1303.3997v1.

60. Rastas P, Calboli FCF, Guo B, Shikano T, Merilä J (2015) Construction of ultradense linkage maps with Lep-
MAP2: Stickleback F2 recombinant crosses as an example. *Genome Biol Evol* 8(1):78–93.

61. Serrano J (1981) Male achiasmatic meiosis in Caraboidea (Coleoptera, Adephaga). *Genetica* 57:131–137.

62. McKenna A, et al. (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-

| 611 | | generation DNA sequencing data. *Genome Res* 20(9):1297–303. |
| 612 | 63. | Browning SR, Browning BL (2011) Haplotype phasing: existing methods and new developments. *Nat Rev* |
| 613 | | *Genet* 12(10):703–714. |
| 614 | 64. | Weir BS, Cockerham CC (1984) Estimating F-Statistics for the analysis of population structure. *Evolution* |
| 615 | | 38(6):1358–1370. |
| 616 | 65. | Rousset F (2008) Genepop'007: a complete re-implementation of the genepop software for Windows and |
| 617 | | Linux. *Mol Ecol Resour* 8(1):103–106. |
| 618 | 66. | Jombart T, Ahmed I (2011) *adegenet 1.3-1*: new tools for the analysis of genome-wide SNP data. |
| 619 | | *Bioinformatics* 27(21):3070–3071. |
| 620 | 67. | Coop G, Witonsky D, Di Rienzo A, Pritchard JK (2010) Using environmental correlations to identify loci |
| 621 | | underlying local adaptation. *Genetics* 185(4):1411–1423. |
| 622 | 68. | Oksanen J (2007) Multivariate analyses of ecological communities in R: vegan tutorial. 39. |
| 623 | 69. | Lotterhos KE, Whitlock MC (2015) The relative power of genome scans to detect local adaptation depends on |
| 624 | | sampling design and statistical method. *Mol Ecol* 24:1031–1046. |
| 625 | 70. | Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol* |
| 626 | | 23(2):254–67. |
| 627 | 71. | De Mita S, Siol M (2012) EggLib: processing, analysis and simulation tools for population genetics and |
| 628 | | genomics. *BMC Genet* 13(1):27. |
| 629 | 72. | Hudson RR, Slatkin M, Maddison WP (1992) Estimation of levels of gene flow from DNA sequence data. |
| 630 | | *Genetics* 132:583–589. |
| 631 | 73. | Patterson N, Richter DJ, Gnerre S, Lander ES, Reich D (2006) Genetic evidence for complex speciation of |
| 632 | | humans and chimpanzees. *Nature* 441(June):1103–1108. |
| 633 | | |