Running head

PHYLODYNAMIC MODEL ADEQUACY

Title

Phylodynamic model adequacy using posterior predictive simulations

Authors

Sebastian Duchene[1]*, Remco Bouckaert[2,3], David A. Duchene[4], Tanja Stadler[5,6], Alexei J.

Drummond[2]

[1]Department of Biochemistry and Molecular Biology, Bio21 Molecular Science and Biotechnology Institute,

University of Melbourne, Australia.

[2]Centre for Computational Evolution, University of Auckland, Auckland, New Zealand.

[3]Max Planck Institute for the Science of Human History, Jena, Germany.

[4]School of Life and Environmental Sciences, University of Sydney, Sydney, Australia.

[5]Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland.

[6]Swiss Institute of Bioinformatics, Lausanne, Switzerland.

* To whom correspondence should be addressed:

Sebastian Duchene

Email: sebastian.duchene@unimelb.edu.au

Abstract

Rapidly evolving pathogens, such as viruses and bacteria, accumulate genetic change at a

similar timescale over which their epidemiological processes occur, such that it is possible

to make inferences about their infectious spread using phylogenetic time-trees. For this

purpose it is necessary to choose a phylodynamic model. However, the resulting inferences

are contingent on whether the model adequately describes key features of the data. Model

adequacy methods allow formal rejection of a model if it cannot generate the main features

of the data. We present TreeModelAdequacy (TMA), a package for the popular BEAST2

software, that allows assessing the adequacy of phylodynamic models. We illustrate its

utility by analysing phylogenetic trees from two viral outbreaks of Ebola and H1N1

influenza. The main features of the Ebola data were adequately described by the coalescent

exponential-growth model, whereas the H1N1 influenza data was best described by the

birth-death SIR model.

Keywords

Model adequacy, Bayesian phylogenetics, Phylodynamics, Viral evolution, Posterior

predictive simulation, BEAST2

Phylogenetic trees depict the evolutionary relationships between groups of organisms. In

the context of infectious diseases, pathogen genetic data can be used to infer such trees. By

assuming a substitution model and including independent information about time, one can

calibrate the molecular clock to obtain time-trees, where the branch lengths correspond to

units of time, and internal nodes of the tree represent the timing of divergence events. For

rapidly evolving viruses and bacteria it is possible to use the sampling times as time-

calibrations (Drummond et al. 2003; Rieux and Balloux 2016). In these organisms, genetic

change and epidemiological or ecological processes occur over a similar timescale. Thus,

time-trees can be informative about epidemiological dynamics, a field of research known as

phylodynamics (Holmes et al. 1993; Grenfell et al. 2004; Kühnert et al. 2011; Volz et al.

2013).

Phylodynamic models describe the distribution of node times, branch lengths, and

sampling times. In full Bayesian phylogenetic analyses, the tree, parameters for the

molecular clock, the substitution model and the phylodynamic model can be estimated

simultaneously using molecular data. In this Bayesian framework, the phylodynamic model

is effectively a 'tree prior' (e.g. du Plessis and Stadler 2015). The simplest phylodynamic

models are the coalescent exponential-growth (CE) and the coalescent constant-size (CC).

These two models have very different expectations about the shape of phylogenetic trees,

with exponentially growing populations tending to produce trees with longer external

branches than those evolving under constant population sizes (O'Meara, 2012; Volz et al.,

2013). Alternative phylodynamic models are the birth-death models, which include a

parameter to describe the sampling rate, so they have an expectation on the number of

taxa and their distribution over time (Stadler 2010; Stadler et al. 2012).

Choosing an appropriate model is important to draw reliable inferences from parameters of

interest. For instance, the CE and the constant birth-death (BD) models can estimate the

basic reproductive number, R_o (Frost & Volz, 2010; Stadler et al., 2012; Volz et al., 2013),

which is defined as the average number of new cases that a single case will generate over

the course of its infection in a fully susceptible population (Anderson and May 1979, 1992).

Failing to account for complex epidemiological dynamics can bias the estimate of this key

parameter (Stadler et al. 2014; Alkhamis et al. 2016; Ratmann et al. 2016). A Bayesian

approach to selecting a phylodynamic model is to estimate marginal likelihoods for a pool

of models and selecting that with the highest marginal likelihood (Baele et al. 2012; Baele et

al. 2016), but it is also possible to obtain weighted averages of parameter estimates based

on the support for each model (e.g. Baele, Li, Drummond, Suchard, & Lemey, 2013; R.

Bouckaert & Drummond, 2017; Huelsenbeck, Larget, & Alfaro, 2004; Li & Drummond,

2012).

Model adequacy in phylogenetics

Model selection methods only allow a relative comparison of a set of models, but they

cannot determine whether any of the models in question could have generated key features

of the data at hand (i.e. absolute model fit). Such information however is key to avoid

unreliable inferences from a model and to improve our understanding of the biological

processes that produced the data. Absolute model fit can be assessed via model adequacy

methods, where a model is considered 'adequate' if it is capable of generating the main

features of the empirical data. Consequently, model adequacy allows the user to formally

reject a model or to identify aspects of the data that are poorly described, instead of

ranking it with respect to other models, as is the case with model testing (e.g. Goldman

1993; Bollback 2002; Ripplinger and Sullivan 2010; Brown 2014).

Model adequacy is typically conducted by fitting a model to the empirical data, and

generating synthetic data from the model in question, a procedure that is similar to a

parametric bootstrap (Goldman 1993). The adequacy of the model is determined depending

on whether the synthetic data are similar to the empirical data, according to a descriptive

test statistic (Gelman and Shalizi 2013; Gelman et al. 2014). The test statistics should

summarise key aspects of the data or a combination of the data and parameter estimates

(Gelman et al. 1996). Examples of test statistics that have been used to assess the

substitution model include the multinomial likelihood or a measure of compositional

homogeneity (Goldman 1993; Huelsenbeck et al. 2001; Foster 2004). The joint clock model

and tree prior key can be assessed using the expected number of substitutions in individual

branch lengths of the tree as test statistics (Duchêne et al. 2015). For DNA barcoding the

number of OTUs and multinomial likelihood have been shown to be effective test statistics

(Barley and Thomson 2016). Phylodynamic and diversification models are fitted to

phylogenetic trees and their parameters depend on the distribution of nodes, such that

some useful test statistics include the ratio of external to internal branch lengths, the tree

height, and measures of phylogenetic tree imbalance (Revell et al. 2005, 2008; Drummond

and Suchard 2008; Höhna et al. 2015). Clearly, designing test statistics is not trivial, but they

should attempt to explicitly test some of the assumptions of the model. For example, the

CE and CC models have different expectations of the ratio of external to internal branch

lengths, such that this may be a useful test statistic.

Bayesian model adequacy

Bayesian model adequacy consists of a posterior predictive framework (Rubin 1981, 1984;

Bollback 2002; Brown 2014a, 2014b; Lewis et al. 2014; Höhna et al. 2017). The posterior

distribution of the model in question is approximated given the empirical data, for example

using Markov chain Monte Carlo (MCMC). Samples from the MCMC are drawn to simulate

data sets under the model used for the empirical analysis. For example, the posterior

distribution of the growth rate and population size parameters of the CE model can be

sampled to simulate phylogenetic trees. Such simulations are known as posterior predictive

simulations. Test statistics are then calculated for every posterior predictive simulation (i.e.

for every simulated tree) to generate a distribution of values according to the model. A

posterior predictive probability, similar to the frequentist p-value, can be calculated by

determining where the value of the test statistic for the empirical data (i.e. the empirical

phylogenetic tree) falls with respect to the posterior predictive distribution (Gelman et al.

2014). Following Gelman et al. (2014) we refer to the posterior predictive probability as p_B to

differentiate it from the frequentist p-value. A useful guideline to determine whether the

model is adequate is to determine whether a test statistic within the 95% credible interval

(Bollback 2002; Brown 2014a). This approach is sometimes conservative, particularly when

test statistics do not follow a Gaussian distribution, and other methods of calculating

posterior predictive probabilities are also possible (Gelman et al. 2014; Höhna et al. 2017).

Combining multiple test statistics leads to multiple testing, which can be addressed by

using a multivariate p_B values (Drummond and Suchard 2008). However, Gelman et al.

(2014) suggest considering each test statistically separately to assess individual aspects of

the model and the data, which is the approach taken here.

Bayesian model adequacy is similar to Approximate Bayesian Computation (ABC)

techniques in that both methods use test statistics from simulated data. The aim of ABC is

to approximate the posterior by comparing test statistics from simulations from the prior

and the empirical data (Csilléry et al. 2010; Ratmann et al. 2012; Poon 2015), which

sometimes leads to biases for model testing (Robert et al. 2011). In contrast, in model

adequacy the simulations are generated from the posterior distribution and they are not

used to approximate the posterior. In spite of these differences, test statistics developed for

ABC can be useful to assess model adequacy.

"TreeModelAdequacy" package in BEAST2

We implemented a computational framework to assess the adequacy of phylodynamic

models as a package for BEAST2 (Bouckaert et al., 2014). Analyses as outlined in Fig 1 are

easy to set up through BEAUti, the graphical user interface for BEAST, to generate an xml

file with the tree, the model, and test statistics. Our package, TreeModelAdequacy (TMA),

takes a tree with branch lengths proportional to time. The tree can be a summary tree from

BEAST2, or estimated using a different method.

We fit phylodynamic models available in BEAST2 by approximating the posterior

distribution of the parameters of the model using MCMC. To generate the posterior

predictive simulations, we draw random samples from the posterior for the parameters

from the MCMC after removing the burn-in phase, and we simulate phylogenetic trees

using stochastic simulations and master equations using MASTER (Vaughan and

Drummond 2013) or the coalescent simulator in BEAST2. The last step consists in

calculating test statistics for the empirical data and for the posterior predictive simulations,

which depends on the TreeStat2 package (available at

http://github.com/alexeid/TreeStat2). The user can select a large number of test statistics

(Supplementary material). At the end of the analysis, p_B values and quantiles for the

posterior predictive distribution are shown, but they can also be visualised in Tracer

(available at: http://beast.bio.ed.ac.uk/tracer) or using an R script included in the package.

At present, the range of phylodynamic models that can be assessed includes the CC and CE

coalescent models, the constant BD with serial sampling (Stadler et al. 2012), and the birth-

death susceptible-infected-recovered model (BDSIR; Kühnert et al. 2014).

Our implementation allows parallelisation of the tree simulation step, which can increase

computational speed when the simulation conditions require extensive calculations. This

step can also be conducted independently on a computer cluster. Our standalone

application TreeModelAdequacyAnalyser can also compute test statistics and p_B values,

even for trees generated in a different program than BEAST2, given that the posterior

predictive trees are provided. TMA is open source and freely available under a LGPL licence.

It can be downloaded from BEAUTi2 (part of BEAST2), and the documentation and

example files are available at: http://github.com/sebastianduchene/tree_model_adequacy.

To verify our implementation, we conducted a simple simulation experiment. We simulated

100 trees under each of the four phylodynamic models (CC, CE, BD, and BDSIR) using

BEAST2 and MASTER and analysed them with the matching model. We assessed their

adequacy according to nine test statistics (Supplementary material). The parameters for our

simulations were based on analyses of 72 whole genome sequences of Ebola virus (Gire et

al. 2014). We found that the p_B values for all test statistics were between 0.025 and 0.975 for

about 95% of each set of simulations, indicating that our implementation is correct

(Supplementary material).

Phylodynamic model adequacy in empirical virus data

West African Ebola virus

We obtained a phylogenetic tree inferred in a previous study from 72 Ebola virus whole

genome samples collected during the 2013-2016 epidemic (Gire et al. 2014). The samples

were collected from May to July 2014 in Sierra Leone. These data have been used in

previous studies to estimate epidemiological parameters, with estimates of R_o ranging from

1.5 to 2.5, depending on the phylodynamic model (Stadler et al., 2014; Volz & Pond, 2014).

An important consideration about our analysis is that we assume that the tree topology and

divergence time estimates are sufficiently accurate and that the data are informative.

We inferred phylodynamic parameters for the Ebola virus tree using four models; CC, CE,

the BD, and the BDSIR (Kühnert et al. 2014). The CE, BD, and BDSIR models can estimate R_o

if information about the sampling process (for the birth-death models) or present number

of infected individuals (for the coalescent) is available. In this case, we assumed that the

sampling proportion was 0.7 (Gire et al. 2014) by fixing this parameter in the BD and BDSIR

models. For the remaining parameters of these two models, we used the same prior

distributions as in a previous analysis of these data (Stadler et al. 2014). For the CE model

we used a Laplace distribution and a $1/x$ distribution as priors for the growth rate and for the

effective population size, respectively. We ran an MCMC of 10^7 steps and generated 1,000

posterior predictive simulations, and we computed four test statistics.

To compare the different models, we calculated p_B for four test statistics; the tree height,

the slope ratio of a lineages-through-time (LTT) plot, the ratio of external to internal branch

lengths, and the Colless index of phylogenetic imbalance (Fig. 2; Supplementary material).

Importantly, the slope ratio of the LTT plot has been found to be informative for inferring

epidemiological parameters using ABC (Saulnier et al. 2017).

In the CC model the p_B was < 0.05 for all test statistics, with the exception of the tree height

at 0.06 (Fig. 2 and Supplementary material Fig. S1). The CE and BD models described these

data better, with most p_B values between 0.11 and 0.56. The p_p value for the Colless index in

the CE model was the lowest for both of these models, at 0.04. The BDSIR model had

overall low p_B values, from to 0.01 to 0.18, with the lowest values found for the ratio of

external to internal branch lengths (0.03) and for the slope ratio of the LTT plot (0.01). The

fact that most p_B values for the CE and BD models frequently fall near the centre of the

posterior predictive distributions is consistent with the rapid spread that Ebola virus was

undergoing at the time the sequences were collected.

For the CE, BD, and BDISIR models we can estimate R_o and the infectious period, $1/\delta$. The R_o

median estimates were: 1.6 (95% credible interval (CI): 1.12 – 2.2) for the BD, 1.21 (95% CI:

1.1 – 1.5), for the CE, and 1.59 (95% CI: 1.22 – 1.94) for the BDSIR. Estimates for the

infectious period in calendar days were: 5.46 (95% CI: 4.14 – 7.24) for the BD, 2.90 (95% CI:

1.75 – 5.50) for the CE, and 5.21 (95% CI: 4.24 – 7.02) for the BDSIR. The estimates from

these models were very similar and overlapped with those from previous studies (Stadler et

al. 2014). Although the BDSIR did not capture some of the main features of these data, this

model is similar to the BD when the number of susceptible individuals is very large, which

probably explains the overlap in R_o estimates between these models.

2009 H1N1 Influenza

We obtained a phylogenetic tree from a previous study (Hedge et al. 2013),which was

estimated from 328 whole genome samples from the 2009 H1N1 Influenza pandemic. The

samples were collected from April to December 2009, such that they encompass a large

portion of the duration of the pandemic. We used a similar method as for the Ebola virus

phylogenetic tree to fit the four phylodynamic models (CC, CE, BD, and BDSIR). However,

instead of fixing the sampling proportion we used an informative prior distribution of the

infectious period via the *becomeUnifectiousRate* parameter, with a normal distribution of

mean 85 and standard deviation of 15 (corresponding to an infectious period of about 4.45

days).

The CC and CE models had p_B values of 0.00 for all four test statistics, such that they did not

adequately describe any of these aspects of the tree. The BD model had p_B values of 0.53

and 0.09 for the tree height and the slope ratio of the LTT plot, and of 0.00 for the ratio of

external to internal branch lengths (Fig. 2 and Supplementary material Fig. S2). In contrast,

the BDSIR model overall described the H1N1 tree better overall than the other three

models, with p_B values of between 0.07 and 0.44. This result is consistent with the sampling

time of the data, which includes the start of the pandemic and the decline in the number of

infections towards the end of the year. We calculated an R_o of mean 3.01 (95% CI: 2.5 – 3.7)

at the start of the pandemic in January that declined to $R_o < 1$ around June, when infectious

spread was lower. This estimate is similar to those made in previous studies based on

census data (Forsberg White et al. 2009), but in the higher range of those based on the CE

model for samples collected in early stages of the pandemic (e.g. Hedge et al. 2013). For

comparison, the R_o estimate from the BD model, which appeared inadequate, was

substantially lower, with a mean of 1.02 (95% CI: 1.00 – 1.03).

Conclusion

Model adequacy methods are useful to understand the biological processes that generate

the data, such as the evolutionary branching process. For example, our approach reveals

that in July of 2014 the West African Ebola outbreak was still growing exponentially and

that the 2009 H1N1 influenza virus pandemic had evidence of a depletion of susceptible

individuals in December. In some cases, identifying models that indadequately describe key

aspects of the data may improve estimates of parameters of interest, such as R_o in our H1N1

influenza analyses. One consideration of our approach is that it requires an accurate

estimate of a single phylogenetic tree. Clearly, the phylogenetic tree should be inferred

using informative sequence data and the sensitivity to the prior should be carefully

examined (Ritchie et al. 2016; Boskova et al. 2018; Möller et al. 2018), which is also the case

for any Bayesian analysis. For example, a tree estimated from uninformative sequence data

will be driven by the prior and will necessarily appear to be adequately described by the

matching model, potentially leading to increased rates of type 2 errors. A limitation of our

method is that it does not account for phylogenetic uncertainty, which can be addressed by

comparing sets of trees from the posterior with those from the posterior predictive

distribution. However, this approach will require the development of new test statistics and

model assessment criteria. Model adequacy in phylogenetics will benefit from further

development of methods to assess more sophisticated phylodynamic models, such as those

that account for population structure (Kühnert et al. 2016; Müller et al. 2017a, 2017b; Volz

and Siveroni 2018), and techniques to improve the interpretation of p_B values for test

statistics that are not normally distributed, such at the Colless index. Model adequacy

software, such as TMA, will be key to address these questions.

Acknowledgements

The authors thank Timothy Vaughan for valuable discussions. SD was supported by a

McKenzie fellowship and a Dyason grant from the University of Melbourne. AJD was

supported by a Rutherford fellowship

(http://www.royalsociety.org.nz/programmes/funds/rutherford-discovery/) from the Royal

Society of New Zealand. TS was supported in part by the European Research Council under

the Seventh Framework Programme of the European Commission (PhyPD: grant

agreement number 335529).

Conflict of interest

306 The authors gave final approval to the manuscript and have no conflict of interest to

307 declare.

Author contributions

310 SD, RB, and AJD wrote the computer code. SD analysed the data. TS, SD, and AJD designed

311 the experiments. SD wrote the manuscript with input from all the authors.

Figure Legends

Fig 1. Posterior predictive simulation framework implemented in the TreeModelAdequacy

package. Step (1) consists in Bayesian analysis in BEAST under model M to estimate the

posterior distribution of parameters η, shown with the arrow and posterior density in

orange. In step (2) samples from the posterior are drawn to simulate phylogenetic trees,

known as posterior predictive simulations, using MASTER as shown by the green arrow. In

step (3) the posterior predictive simulations are analysed in TreeStat to generate the

posterior predictive distribution of test statistic T_a, shown by the blue arrow and probability

density. Finally, T_a is also computed for the tree from the empirical data using TreeStat,

shown by the red arrow, to calculate a posterior predictive probability (p_B). Test statistics

and p_B values can also be computed for trees generated in other programs using

TreeModelAdequacyAnalyser, given that the tree from the empirical data and the posterior

predictive simulations are provided.

Fig 2. Model adequacy results for the two empirical data sets, West African Ebola and 2009

H1N1 influenza. The histograms show the distribution of two test statistics, the tree height

and the slope ratio of the lineages-through-time (LTT) plot, for the posterior predictive

simulations. The black points also show the distribution of the test statistics, and they have

been jittered along the y-axis to improve visualisation. The red lines denote the value for

the tree estimated from the empirical data. The posterior predictive p-value, p_B, is shown

for each test statistic. A feature of the empirical phylogenetic tree described by a test

statistic, such as the tree height, is considered adequately described by the model if the

empirical value falls within the 95% quantile range of the posterior predictive distribution,

such that the $p_B > 0.05$. Two more test statistics were computed, the ratio of external to

internal branch lengths and the Colless index, which are shown in Supplementary material

Fig. S1. Note that for the H1N1 influenza analyses, the values are shown in a $\log_{10}$ scale.

Supplementary figure captions

Fig S1. Model adequacy results for Ebola virus data with four test statistics. The colours and

symbols match those in Fig 2.

Fig S2. Model adequacy results for H1N1 influenza data with four test statistics. The colours

and symbols match those in Fig 2. Note that the values are shown in $\log_{10}$ scale.

References

Alkhamis M.A., Perez A.M., Murtaugh M.P., Wang X., Morrison R.B. 2016. Applications of

Bayesian phylodynamic methods in a recent US porcine reproductive and respiratory

syndrome virus outbreak. Front. Microbiol. 7:67.

Anderson R.M., May R.M. 1979. Population biology of infectious diseases: Part I. Nature.

280:361.

Anderson R.M., May R.M. 1992. Infectious diseases of humans: dynamics and control.

Oxford university press.

Baele G., Lemey P., Bedford T., Rambaut A., Suchard M.A., Alekseyenko A. V. 2012.

Improving the accuracy of demographic and molecular clock model comparison while

accommodating phylogenetic uncertainty. Mol. Biol. Evol. 29:2157–2167.

Baele G., Lemey P., Suchard M.A. 2016. Genealogical working distributions for Bayesian

model testing with phylogenetic uncertainty. Syst. Biol. 65:250–264.

Baele G., Li W.L.S., Drummond A.J., Suchard M.A., Lemey P. 2013. Accurate model

selection of relaxed molecular clocks in bayesian phylogenetics. Mol. Biol. Evol.

30:239–243.

Barley A.J., Thomson R.C. 2016. Assessing the performance of DNA barcoding using

posterior predictive simulations. Mol. Ecol. 25:1944–1957.

Bollback J.P. 2002. Bayesian model adequacy and choice in phylogenetics. Mol. Biol. Evol.

19:1171–1180.

Boskova V., Stadler T., Magnus C. 2018. The influence of phylodynamic model

specifications on parameter estimates of the Zika virus epidemic. Virus Evol. 4:vex044.

Bouckaert R., Drummond A.J. 2017. bModelTest: Bayesian phylogenetic site model

averaging and model comparison. BMC Evol. Biol. 17:42.

Bouckaert R., Heled J., Kühnert D., Vaughan T., Wu C.-H., Xie D., Suchard M.A., Rambaut

A., Drummond A.J. 2014. BEAST 2: a software platform for Bayesian evolutionary

analysis. PLOS Comput. Biol. 10:e1003537.

Brown J.M. 2014a. Detection of implausible phylogenetic inferences using posterior

predictive assessment of model fit. Syst. Biol. 63:334–348.

Brown J.M. 2014b. Predictive approaches to assessing the fit of evolutionary models. Syst.

Biol. 63:289–292.

Csilléry K., Blum M.G.B., Gaggiotti O.E., François O. 2010. Approximate Bayesian

computation (ABC) in practice. Trends Ecol. Evol. 25:410–418.

Drummond A.J., Pybus O.G., Rambaut A., Forsberg R., Rodrigo A.G. 2003. Measurably

evolving populations. Trends Ecol. Evol. 18:481–488.

Drummond A.J., Suchard M.A. 2008. Fully Bayesian tests of neutrality using genealogical

summary statistics. BMC Genet. 9:68.

Duchêne D.A., Duchêne S., Holmes E.C., Ho S.Y.W. 2015. Evaluating the adequacy of

molecular clock models using posterior predictive simulations. Mol. Biol. Evol.

32:2896–2995.

Forsberg White L., Wallinga J., Finelli L., Reed C., Riley S., Lipsitch M., Pagano M. 2009.

Estimation of the reproductive number and the serial interval in early phase of the

2009 influenza A/H1N1 pandemic in the USA. Influenza Other Respi. Viruses. 3:267–

276.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst. Biol. 53:485–495.

Frost S.D.W., Volz E.M. 2010. Viral phylodynamics and the search for an "effective number

of infections." Philos. Trans. R. Soc. London B Biol. Sci. 365:1879–1890.

Gelman A., Carlin J.B., Stern H.S., Dunson D.B., Vehtari A., Rubin D.B. 2014. Model

checking. Bayesian data analysis. Boca Raton, Florida: CRC press Boca Raton, FL. p.

141–163.

Gelman A., Meng X.-L., Stern H. 1996. Posterior predictive assessment of model fitness via

realized discrepancies. Stat. Sin.:733–760.

Gelman A., Shalizi C.R. 2013. Philosophy and the practice of Bayesian statistics. Br. J. Math.

Stat. Psychol. 66:8–38.

Gire S.K., Goba A., Andersen K.G., Sealfon R.S.G., Park D.J., Kanneh L., Jalloh S., Momoh

M., Fullah M., Dudas G. 2014. Genomic surveillance elucidates Ebola virus origin and

transmission during the 2014 outbreak. Science (80-.). 345:1369–1372.

Goldman N. 1993. Simple diagnostic statistical tests of models for DNA substitution. J. Mol.

Evol. 37:650–661.

Grenfell B.T., Pybus O.G., Gog J.R., Wood J.L.N., Daly J.M., Mumford J.A., Holmes E.C.

2004. Unifying the epidemiological and evolutionary dynamics of pathogens. Science.

303:327–332.

Hedge J., Lycett S.J., Rambaut A. 2013. Real-time characterization of the molecular

epidemiology of an influenza pandemic. Biol. Lett. 9:20130331.

Höhna S., Coghill L.M., Mount G.G., Thomson R.C., Brown J.M. 2017. P3: Phylogenetic

Posterior Prediction in RevBayes. Mol. Biol. Evol.

Höhna S., May M.R., Moore B.R. 2015. TESS: an R package for efficiently simulating

phylogenetic trees and performing Bayesian inference of lineage diversification rates.

Bioinformatics. 32:789–791.

Holmes E.C., Zhang L.Q., Simmonds P., Rogers A.S., Brown A.J.L. 1993. Molecular

investigation of human immunodeficiency virus (HIV) infection in a patient of an HIV-

infected surgeon. J. Infect. Dis. 167:1411–1414.

Huelsenbeck J.P., Larget B., Alfaro M.E. 2004. Bayesian phylogenetic model selection using

reversible jump Markov chain Monte Carlo. Mol. Biol. Evol. 21:1123–1133.

Huelsenbeck J.P., Ronquist F., Nielsen R., Bollback J.P. 2001. Bayesian inference of

phylogeny and its impact on evolutionary biology. Science (80-.). 294:2310–2314.

Kühnert D., Stadler T., Vaughan T.G., Drummond A.J. 2014. Simultaneous reconstruction of

evolutionary history and epidemiological dynamics from viral sequences with the

birth–death SIR model. J. R. Soc. Interface. 11:20131106.

Kühnert D., Stadler T., Vaughan T.G., Drummond A.J. 2016. Phylodynamics with migration:

A computational framework to quantify population structure from genomic data. Mol.

Biol. Evol. 33:2102–2116.

Kühnert D., Wu C.H., Drummond A.J. 2011. Phylogenetic and epidemic modeling of rapidly

evolving infectious diseases. Infect. Genet. Evol. 11:1825–1841.

Lewis P.O., Xie W., Chen M.-H., Fan Y., Kuo L. 2014. Posterior predictive Bayesian

phylogenetic model selection. Syst. Biol. 63:309–321.

Li W.L.S., Drummond A.J. 2012. Model averaging and Bayes factor calculation of relaxed

molecular clocks in Bayesian phylogenetics. Mol. Biol. Evol. 29:751–761.

Möller S., du Plessis L., Stadler T. 2018. Impact of the tree prior on estimating clock rates

during epidemic outbreaks. Proc. Natl. Acad. Sci. USA. 115:4200–4205.

Müller N.F., Rasmussen D.A., Stadler T. 2017a. MASCOT: Parameter and state inference

under the marginal structured coalescent approximation. bioRxiv.:188516.

Müller N.F., Rasmussen D.A., Stadler T. 2017b. The Structured Coalescent and Its

Approximations. Mol. Biol. Evol. 34:2970–2981.

O'Meara B.C. 2012. Evolutionary inferences from phylogenies: a review of methods. Annu.

Rev. Ecol. Evol. Syst. 43:267–285.

du Plessis L., Stadler T. 2015. Getting to the root of epidemic spread with phylodynamic

analysis of genomic data. Trends Microbiol. 23:383–386.

Poon A.F.Y. 2015. Phylodynamic inference with kernel ABC and its application to HIV

epidemiology. Mol. Biol. Evol. 32:2483–2495.

Ratmann O., Donker G., Meijer A., Fraser C., Koelle K. 2012. Phylodynamic inference and

model assessment with approximate bayesian computation: influenza as a case study.

PLoS Comput. Biol. 8:e1002835.

Ratmann O., Hodcroft E.B., Pickles M., Cori A., Hall M., Lycett S., Colijn C., Dearlove B.,

Didelot X., Frost S. 2016. Phylogenetic Tools for Generalized HIV-1 Epidemics:

Findings from the PANGEA-HIV Methods Comparison. Mol. Biol. Evol. 34:185–203.

Revell L.J., Harmon L.J., Collar D.C. 2008. Phylogenetic signal, evolutionary process, and

rate. Syst. Biol. 57:591–601.

Revell L.J., Harmon L.J., Glor R.E. 2005. Under-parameterized model of sequence evolution

leads to bias in the estimation of diversification rates from molecular phylogenies.

Syst. Biol. 54:973–983.

Rieux A., Balloux F. 2016. Inferences from tip-calibrated phylogenies: a review and a

practical guide. Mol. Ecol. 25:1911–1924.

Ripplinger J., Sullivan J. 2010. Assessment of substitution model adequacy using frequentist

and Bayesian methods. Mol. Biol. Evol. 27:2790–2803.

Ritchie A.M., Lo N., Ho S.Y.W. 2016. The impact of the tree prior on molecular dating of

data sets containing a mixture of inter-and intraspecies sampling. Syst. Biol. 66:413–

425.

Robert C.P., Cornuet J.-M., Marin J.-M., Pillai N.S. 2011. Lack of confidence in approximate

Bayesian computation model choice. Proc. Natl. Acad. Sci. USA. 108:15112–15117.

Rubin D.B. 1981. Estimation in parallel randomized experiments. J. Educ. Stat. 6:377–401.

Rubin D.B. 1984. Bayesianly justifiable and relevant frequency calculations for the applied

statistician. Ann. Stat. 12:1151–1172.

Saulnier E., Alizon S., Gascuel O. 2017. Assessing the accuracy of Approximate Bayesian

Computation approaches to infer epidemiological parameters from phylogenies.

PLOS Comput. Biol. 13:e1005416.

Stadler T. 2010. Sampling-through-time in birth-death trees. J. Theor. Biol. 167:696–404.

Stadler T., Kouyos R., von Wyl V., Yerly S., Böni J., Bürgisser P., Klimkait T., Joos B., Rieder

P., Xie D. 2012. Estimating the basic reproductive number from viral sequence data.

Mol. Biol. Evol. 29:347–357.

Stadler T., Kühnert D., Rasmussen D.A., du Plessis L. 2014. Insights into the early epidemic

spread of Ebola in Sierra Leone provided by viral sequence data. PLoS Curr. 6.

Vaughan T.G., Drummond A.J. 2013. A stochastic simulator of birth–death master

equations with application to phylodynamics. Mol. Biol. Evol. 30:1480–1493.

Volz E., Pond S. 2014. Phylodynamic analysis of Ebola virus in the 2014 Sierra Leone

epidemic. PLoS Curr. 6.

Volz E., Siveroni I. 2018. Bayesian phylodynamic inference with complex models.

bioRxiv.:268052.

Volz E.M., Koelle K., Bedford T. 2013. Viral phylodynamics. PLOS Comput. Biol. 9:e1002947.

Empirical data (tree)

(1) BEAST

Bayesian analysis
(model M, with parameters η)

Posterior distribution
P(η|tree,M)

Posterior
probability

η

(2) MASTER

Sample P(η|tree,M) to
simulate trees

Posterior predictive
simulations

TreeStat
(3)

(3) TreeStat

T_a

	West African Ebola (2014)		**H1N1 Influenza (December 2009)**	
Constant-size coalescent	$p_B = 0.06$	$p_B = 0.01$	$p_B = 0.00$	$p_B = 0.00$
Exp. growth coalescent	$p_B = 0.27$	$p_B = 0.55$	$p_B = 0.00$	$p_B = 0.00$
Constant birth-death	$p_B = 0.56$	$p_B = 0.21$	$p_B = 0.53$	$p_B = 0.09$
Birth death SIR	$p_B = 0.18$	$p_B = 0.01$	$p_B = 0.47$	$p_B = 0.07$
	Tree height (years)	Slope ratio of LTT plot	Tree height (years)	Slope ratio of LTT plot