Title

Assessment of an organ-specific *de novo* transcriptome

of the nematode trap-crop, *Solanum sisymbriifolium.*

Authors/Affiliations

Alexander Q Wixom[a], N Carol Casavant[a], Joseph C Kuhl[a], Fangming Xiao[a], Louise-Marie

Dandurand[b], and Allan B Caplan[a*]

[a]Department of Plant Sciences, University of Idaho, Moscow, ID 83844-2333

[b]Department of Entomology, Plant Pathology, and Nematology, University of Idaho, Moscow, ID

83844-2329

Contact

*Corresponding author e-mail: acaplan@uidaho.edu

Running Title

Running title: A *de novo S. sisymbriifolium* transcriptome

Abstract

Solanum sisymbriifolium, also known as "Litchi Tomato" or "Sticky Nightshade," is an undomesticated and poorly researched plant related to potato and tomato. Unlike the latter species, *S. sisymbriifolium* induces eggs of the cyst nematode, *Globodera pallida*, to hatch and migrate into its roots, but then arrests further nematode maturation. In order to provide researchers with a partial blueprint of its genetic make-up so that the mechanism of this response might be identified, we used single molecule real time (SMRT) sequencing to compile a high quality *de novo* transcriptome of 41,189 unigenes drawn from individually sequenced bud, root, stem, and leaf RNA populations. Functional annotation and BUSCO analysis showed that this transcriptome was surprisingly complete, even though it represented genes expressed at a single time point. By sequencing the 4 organ libraries separately, we found we could get a reliable snapshot of transcript distributions in each organ. A divergent site analysis of the merged transcriptome indicated that this species might have undergone a recent genome duplication and re-diploidization. Further analysis indicated that the plant then retained a disproportionate number of genes associated with photosynthesis and amino acid metabolism in comparison to genes with characteristics of R-proteins or involved in secondary metabolism. The former processes may have given *S. sisymbriifolium* a bigger competitive advantage than the latter did.

Introduction

Solanum sisymbriifolium (SSI), otherwise known as "litchi tomato", "morelle de Balbis", or "sticky nightshade", is an undomesticated relative of potato and tomato. For more than a decade, SSI has been investigated as a trap-crop (a plant that attracts nematodes but kills them before they can reproduce) for nematodes such as Globodera pallida that normally parasitize potatoes and tomatoes (Timmermans, 2005; Dandurand and Knudsen, 2016). It is also a potential source of anti-protozoan (Meyre-Silva et al., 2013) and anti-molluscan (Bagalwa et al., 2010) metabolites. If the genetic basis for these protective processes could be identified, it might be possible to transfer these traits, either through cross-breeding or through modern transgenic technologies, from this weed to its domesticated relatives. However, while the genomes of potato and tomato have been studied extensively, spiny solanums, like SSI, have not (Yang et al., 2014). Only 54 SSI nucleotide sequences have been submitted to NCBI as of 2016. This ignorance about the biology and genetics of the spiny solanums could be masking a wealth of genetic resources that could be used to protect agriculturally important crops.

Most bioinformatic analyses of a species begin with the assembly and annotation of a complete genome. Once assembled, these data can be searched for genes encoding a particular protein or RNA sequence. For those working on a species that has not been studied extensively in the past, and which is only being studied now in order to conduct a limited number of experiments, whole genome sequencing can be more expensive and time consuming than can be justified. In these circumstances, alternative methods using sequencing technologies that are generally referred to as next-generation sequencing (NGS), have allowed researchers to by-pass whole genome sequencing in favor of generating a smaller database, one depleted of the silent regions of the genome and of genes that are not contributing to the phenotypes of interest. Most commonly, this is done using Illumina or 454 platforms that generate 10's and 100's of millions of short reads from cDNA copies of all of the mRNAs expressed during a given moment of time. Once obtained, these sequences can then be merged in silico into full length protein coding sequences. However, this de novo transcriptome can sometimes prove problematic. Short reads derived from highly conserved coding domains and repetitively organized genes can potentially be aligned and joined into chimeric assemblies that cannot be verified or removed because there is no independently

sequenced genome available to serve as an extended template or scaffold to ensure that the
merged sequences are indeed co-linear (Yang and Smith, 2013). A recent technical
improvement, Pacific Biosciences' single-molecule real-time (SMRT) "sequencing by
synthesis" strategy, has become sufficiently accurate and attainably priced to be utilized by
small research groups. The benefit of using SMRT sequencing is that it produces vastly
longer reads than previous methodologies, although with lower coverage (Eid *et al.*, 2009).
The longer reads allow researchers to establish a transcriptome consisting of nearly complete
open reading frames free of the kinds of errors possible when sequences must be assembled
in silico from short reads (Ocwieja *et al.*, 2012; Zhang *et al.*, 2014).

The specific goal of the current project was to establish a four organ (bud, leaf, stem
and root) *de novo* transcriptome of SSI. In doing so, we wanted to ensure that the final
sequences were high-quality and consisted of genes that were biologically relevant and not
artifacts of some *in silico* assembly process. This transcriptome will provide a reference
library to be used in future RNA-seq experiments to identify genes for nematode and other
pathogen resistances in SSI.

Results

Establishing a SMRT sequenced transcriptome

Before any sequencing was attempted, the genome size of SSI was estimated using flow cytometry (Supplemental Figure 1). This showed that the genome mass of SSI was approximately 4.73 pg per 2C, or 2,315 mega-base pairs per 1C. By comparison, Arumuganathan and Earle, 1991, using the same technology estimated that the tomato genome massed between 1.88 to 2.07 pg per 2C while tetraploid potato massed between 3.31 to 3.86 pg per 2C. Thus, these initial measurements gave SSI a genome size greater than tetraploid potato. Despite their unusual length (Paul and Banerjee, 2015), SSI has 24 chromosomes, like diploid potatoes and many other Solanaceae (data not shown). Due to the size of this genome, and our interest in generating a database of protein-coding genes, we elected to sequence the SSI transcriptome rather than its genome.

Generating an SSI transcriptome was done using Single-Molecule Real Time (SMRT) sequencing by PacBio Sciences (Eid *et al.*, 2009) that does not need to assemble short reads into one contiguous sequence. Rather than producing only short reads, SMRT technology can provide reads up to 60,000 bp along a single molecule of DNA. This allows the capture of entire genes with one read rather than chunking it into many small bits that have to be assembled later. This sequencing strategy gives higher coverage than Sanger-based reactions like those performed on Applied Biosystems™ gene analysis instruments, and longer reads than Illumina or Roche 454.

The Iso-Seq pipeline classifies the sequences as either full-length non-chimeric (FLNC), or non-full length reads. Full length reads are those containing both 5' and 3' adapters, in addition to the poly(A) tail. The reads containing these parts in the expected order, i.e. 5' adapter–poly(A)–3' adapter, with no additional copies of these parts, are classified as non- chimeric. The FLNC reads in the present dataset were corrected with the non-full length reads using Iterative Cluster for Error correction and the Pacific Biosciences Quiver algorithm (https://github.com/PacificBiosciences/SMRT-Analysis/wiki/ConsensusTools-v2.3.0-Documentation).

In an attempt to improve our ability to detect differences in the suites of genes expressed in different parts of the plant, we generated cDNA from 4 organs; leaves, stems,

roots, and unopened flower buds. We then independently carried out SMRT sequencing of all 4 samples. Finally, all corrected FLNC reads were merged *in silico* and redundancy was removed using CD-HIT-EST (Li and Godzik, 2006).

This SMRT sequencing strategy created 231,712 total corrected FLNC sequences (Table 1) using the aforementioned pipeline. These sequences had a GC content of 41.2%. CD-HIT-EST was then used to reduce the redundant sequences to sets with 100% identity. This lowered the number of sequences to 139,611 with a GC content of 41.0%. The GC content continued to decrease as the identity was reduced using CD-HIT-EST. At 80% identity, there were 32,315 sequences, with an estimated GC content of 39.7%. The decrease in GC content could be due to the methodology of CD-HIT-EST that retains the longest sequence during the reduction process. Because of this, reads that spanned untranslated regions of a transcript were expected to be favored over those only consisting of coding regions.

In the end, we chose to work with a final SMRT dataset that had been reduced to 90% identity and consisted of a set of 41,189 sequences with a GC content of 39.9%. We judged that this estimate of the number of transcripts present in the 4 organs would most likely err on the high side, yet still retain most splice variants within a gene, as well as many paralogs and single nucleotide polymorphisms between alleles of this obligate outbreeder.

Evidence based Quality Control of the SMRT Transcriptome

We performed an internal quality check by sequencing 45 randomly chosen clones from a cDNA library using Sanger Dye Deoxy technology (ABI 3730, Applied Biosystems). Bowtie2 (Langmead and Salzberg, 2012) was then used to find the most likely equivalent of each clone in our SSI transcriptome. A manual comparison of the Sanger-sequenced clone and the assembled transcript was done using DNA Strider (Marck, 1988). Firstly, all 45 cDNAs were found in the SSI transcriptome (Figure 1A). Secondly, only two of these SMRT-derived sequences appeared to be chimeric (Figure 1B), and based on the length of non-homologous stretches, could have been transcribed from different members of the same gene family rather than been created by misassembly. During our analysis of the SSI transcriptome, we did find entries that consisted of inverted repeats of entire gene sequences. These inverted repeats likely occurred during the preparation of the cDNA library prior to

sequencing rather than during sequencing or subsequent computational processing as can be

found in Illumina or 454 assemblies (Loman *et al.*, 2012; Luo *et al.*, 2012).

When the SSI transcriptome was analyzed using Mercator (Lohse *et al.*, 2014), it

contained 38.6% unannotated sequences. This was markedly fewer than the percent

unannotated sequences of either potato (50.8%) or tomato (46.4%) transcriptomes processed

in the same way via Mercator (Supplemental Figure 4). Other than that, the binned profile

of SSI was very similar to the published transcriptomes (Supplemental Figure 3) of these

plants. This led us to believe that our transcriptome was at least of comparable quality with

the working transcriptomes of these two better studied species.

PfamScan (Finn *et al.*, 2008) was also used to annotate the domains of the

transcriptome. This program uses HMMer (Eddy, 1998) domain annotations, and used in

combination with protocols established by Sarris *et al.*, 2016, allowed for the annotation of

domains found in the amino acid sequences translated from the assembled transcriptome.

This annotated 84.7% of the transcriptome with at least one recognizable domain. There

were fewer unannotated sequences in the SSI transcriptome than in the STU and SLY

transcriptomes (Supplemental Table 1). The reduced number of unannotated sequences

found in the SMRT transcriptome might reflect the fact that this set had undergone a

conservative reduction to 90% identity. Alternatively, the reduced number of unannotated

SSI sequences could have resulted from the fact that we had only sampled the four most

frequently studied organs of a "normally" growing plant, that is, plants manifesting a

physiological state which has been extensively studied in numerous species, while the STU

and SLY transcriptomes were compiled from plants sampled over a much broader range of

life-history stages and growth conditions ranging from fruit and tuber development to

exposure to biotic and abiotic stresses where the functions of many genes are still under

investigation.

To further test the quality and completeness of our transcriptome, BUSCO

benchmarking (Simão *et al.*, 2015) was performed. The BUSCO database was established to

allow researchers to assess the completeness of new genomes or transcriptomes based on the

detection of a set of universal, single-copy orthologs. We found 93% intact BUSCO

archetypes (889 genes) in the SSI SMRT transcriptome, 30.2% (289) of these were found in

multiple copies, while an additional 2.2% (21) of the BUSCO archetypes were present in

fragments, and 4.8% (46) were missing entirely (Table 2). These numbers representing genes

expressed during a single growth condition of SSI, were only 4.7 percentages different from

the numbers of BUSCO archetypes found in the entire SMRT sequenced genome of *A.*

thaliana.

SSI has the same number of chromosomes as most other Solanaceae, and does not

appear to be polyploid (data not shown), yet the BUSCO analysis showed that SSI had more

duplicate copy archetypes than diploid and tetraploid potato. This high number of similar

sequences could point to the fact that our transcriptome has not been reduced far enough, or

could be one line of evidence that SSI has undergone extensive genome duplication or

hybridization in the past. This latter hypothesis was evaluated by divergent gene analysis as

has been done with plants such as wheat (Krasileva *et al.*, 2013). When the program

Freebayes (Garrison and Marth, 2012) was run using a defined diploid setting, it output

information stating there were genes that had more than 2 alleles or paralogues. We redid the

analysis using defined triploid and tetraploid settings and found that even after merging

sequences with more than 90% identity using CD-HIT-EST, the SSI transcriptome contained

1,348 genes with 3 distinguishable alleles or paralogues and furthermore, 44 genes with 4

distinguishable copies (Table 3). It was noteworthy that no gene had more than 4 alleles or

paralogues. A simple explanation for these multiple gene variants, that would be consistent

with the BUSCO analysis, was that SSI underwent a genome duplication followed by

diploidization in the past and that over time, some of the duplicated loci acquired additional

mutations while other loci were lost. To determine if this proposed duplication was restricted

to one chromosome, or one chromosomal arm, the 44 genes with 4 alleles were mapped onto

SLY chromosomes (Supplemental Table 2). There were "4–allele" genes found on 11 of 12

SLY chromosomes which indicated, assuming that genes dispersed in tomato were not linked

when the two species diverged, that SSI has undergone a full genome duplication rather than

a segmental duplication within one chromosome.

Since SSI is not as well-known as other Solaneacae, we employed OrthoMCL v2.0.9

(Li *et al.*, 2003) to illustrate some of the common features its gene complement showed with

those of other plants. Protein sequences from our SSI transcriptome (translated using the

program ESTScan (Iseli *et al.*, 1999)), and protein sequences from tomato (SLY), potato

(STU), eggplant (SME), *Arabidopsis thaliana* (ATH), papaya (CPA), grapes (VVI), peaches

(PPE), black cottonwood (PTR), oranges (CSI), alfalfa (MTR), maize (ZMA) and rice (OSA)

were merged into 45,234 orthologous groups (gene families). In this set, 6097 orthologous

groups were shared by all 13 species (Figure 2), an overlap well within the range of previous

studies (Yang *et al.*, 2014). Each species had many additional groups that were not shown in

this diagram because they were not shared with all members of this set of plants.

Interestingly, even closely related species like SSI, STU, SLY, and SME had hundreds of

groups not found in each other. When the annotations of the SSI unique set were compared

to the full transcriptome, several functional groups showed a disproportionate increase. It is

possible that these disproportionately expanded sets, that included photosynthetic genes, and

genes for amino acid and vitamin metabolism (Supplemental Figure 5), diverged so much

more than groups such as those for cell wall composition, and hormone and secondary

metabolism, because expansion of the former traits gave SSI a competitive edge over other

species in their habitat. Overall, though, there were fewer groups of genes unique to SSI

than unique to STU and SME. As noted previously, this could merely reflect the fact that our

data came from a single-point snapshot of only 4 organs and so would have lacked those

transcripts specifically expressed during fruit and seed set, germination, senescence, abiotic

stress, pathogen attacks, and numerous other stages of a plant's lifecycle.

Using highly conserved orthologous genes, i.e. subunits of Rubisco, provisional

phylogenies were created for nuclear-encoded and chloroplast-encoded genes using the

aforementioned species (data not shown). In doing so, we concluded that nuclear SSI was

most closely related to eggplant, which has been noted previously (Särkinen *et al.*, 2015),

while chloroplast SSI was more closely related to tomato. This dichotomy has also been seen

by others (Miz *et al.*, 2008) and interpreted to indicate that SSI had undergone an ancient

hybridization and afterwards retained the chloroplast genome from one parent, and much of

the nuclear genome from another. However, many more SSI genes will have to be compared

with the genes of many more South American plants to confirm that this hybridization

occurred.

Building a snapshot of organ-associated gene expression

Since we had maintained separate cDNA pools from individual organs, it was

possible to backtrack each sequence within the final transcriptome to obtain a provisional

profile of gene expression throughout the plant (Figure 3A). This analysis showed that there were 8019 sequences expressed solely in buds, 4957 solely in roots, 5349 solely in leaves, 4198 solely in stems, and 7212 sequences expressed in all tissues. That left 11,538 sequences that were expressed in more than one organ but not in all 4.

This backtracking allowed us to construct an expression snapshot that showed how different genes were being expressed at the time the organs were harvested. Using several in-house Python scripts, we recorded the number of reads for genes that had common annotations for several different physiological processes.

A set of light-harvesting complex genes (LHC-I) were predictably found in aerial organs with few exceptions (Figure 3B), demonstrating that the backtracking program could extract biologically useful information about sequences with specified characteristics from the merged transcriptome. In order to determine if this kind of analysis of SMRT sequences could categorize the expression of very different sets of genes, we constructed an inter-organ expression profile of genes that encoded both a leucine-rich repeat (LRR) domain together and a nucleotide binding (NB-ARC) domain (Figure 3C), a pairing frequently found in pathogen resistance genes (R-genes). This profile of R-gene prevalence in SSI, potato, and tomato indicated that there was a reduction of these genes in the SMRT transcriptome compared to the other two species (Table 4). Three of these potential R-genes were then assayed by semi-quantitative PCR (primers found in Supplemental Table 3) and quantified using a sample of cDNA from the same pool that had been sequenced, and a sample from an independently-prepared, unsequenced cDNA pool (Figure 4). In order to assess whether a SMRT data set could be a reliable indicator of gene expression, both the *in silico* and PCR measurements of gene expression were normalized in kind to an actin sequence (Ssi032526). The physical measurements of expression of two of the three genes matched the expression snapshot extremely well, but the third gene (Ssi038051) was more abundant in stems and buds than expected based on its SMRT expression snapshot. This confirms that whole transcriptome snapshots can provide a provisional picture of organ differences in gene expression, but further shows that the expression of each gene of interest needs to be verified biologically, most usefully by multiple independent tests.

Discussion

The creation of a *de novo* transcriptome necessitates massive amounts of follow-up

analyses, both *in silico* and biologically, to estimate its reliability.

We initially employed both Illumina and 454 sequencing (data not shown) in order

to compensate for the different kinds of errors to which each method was prone (Luo *et al.*,

2012). Screening this assembly with genes randomly selected from an SSI cDNA library

revealed that 20% of these genes failed to match any of the assembled sequences in this

database (Supplemental Figure 2A), and of those that matched, 40% appeared to be chimeric

(Supplemental Figure 2B). In contrast, all of these cDNAs were found in our SMRT

sequenced transcriptome and few were patently chimeric (Figure 1).

A number of factors are known to exasperate misassembly including the presence of

large gene families and of repeatedly occurring kmers in the dataset (Moreton *et al.*, 2015).

Even though we did not sequence the SSI genome, we found 4 lines of evidence indicating

that it might be complex enough to pre-dispose our transcriptome to these kinds of assembly

mistakes. First, the nuclear DNA content of SSI was larger than most diploid Solanaceae,

roughly the same size as a tetraploid potato (Supplemental Figure 1). Second, divergent gene

analysis indicated that the SSI transcriptome was unusually complex and contained 3 and 4

distinguishable alleles for many genes (Table 3). Third, there were only 67 putative R-genes,

that is, genes containing a nucleotide binding domain (NB-ARC) and a leucine-rich repeat

domain (LRR), in the SMRT sequenced dataset compared to the 309 in STU, and 137 in

STU (Table 4). Finally, an unusually high percentage of the BUSCO gene set were present

in multiple copies in SSI even though our transcriptome could only consist of a portion of

all the genes that are likely to be encoded in its DNA (Table 2). One model consistent with

these 4 facts was that SSI had, sometime in the past, undergone a partial or complete genome

duplication. Over time, as diploidy was re-established, some of the duplicated alleles or

paralogues diverged, while others were lost. Nevertheless, enough of the expanded gene

families remained to confound the alignment programs that tried to differentiate between

their members. While these kinds of errors might be correctable with the use of other

assembly programs, we chose, instead to create an assembly-independent transcriptome

using SMRT technology.

At the moment, SMRT technology does not provide the sequence coverage or depth

that can be obtained with Illumina or 454 sequencing. In order to increase our chances of

sampling uncommon organ-specific transcripts, we prepared independent cDNA pools from

4 organs of the plant. Using an in-house script

(https://github.com/AlexWixom/Transcriptome_scripts), we were able to increase the value

of the final library by generating expression snapshots for genes of interest in each organ.

These expression snapshots are no substitute for a more thorough RNA-seq study, but they

do provide a preliminary assessment of a plant's biology at the time of harvest. Using these

snapshots, we recognized different patterns of expression of individual LHC-1 genes (Figure

3B) within the photosynthetic parts of the plant. We also saw that 2 of the 3 R-genes re-

examined by PCR showed the same expression pattern in two independent RNA and cDNA

preparations as found in the transcriptome itself (Figure 4). Thus, in the absence of RNA-

seq studies or experimental evidence for the role of a specific locus, this kind of library

assembly could be used to direct researchers to the subset of R-genes most likely responsible

for the resistance in a given organ.

R-genes coding for recognition proteins are commonly perceived as sentinels that are

awaiting activation by molecules introduced during infection (Jones and Dangl, 2006).

Therefore, the reduced number of R-genes found in SSI (67), compared to both potato (309),

or tomato (137) was unexpected. This discrepancy could be explained in any one of several

ways. First, SSI might be using proteins with novel domain structures in place of classic R-

genes. Second, sequencing depth might simply have been inadequate to capture all R-genes

that were actually being expressed at low levels. Finally, SSI could be relying on rapidly

inducing transcription of R-genes after an infection has occurred. Any one of these

hypotheses is worthy of continuing analysis.

With this transcriptome as an example, we have established a protocol that opens the

door to further genetic mining of previously uncharacterized species. The completeness of

our database indicates that *de novo* transcriptomes not only provide an economical and time-

saving way to study a new species, but can also provide expression data that could not be

gleaned from a genomic sequence.

Materials and Methods

Plant and Culture Conditions

S. sisymbriifolium (SSI) seeds obtained from C. Brown (USDA-ARS, Prosser WA) were germinated in soil. Nodes from a single plant were sterilized for 20 min using 10% NaClO with 0.05% Tween20. Plant material was then washed 3x with sterile distilled H_2O and put into 120 mL baby food jars containing standard Murashige and Skoog salts, pH 5.6, 3% sucrose, 0.7% agar, 100 µg mL^{-1} myo-inositol, 2.0 µg mL^{-1} glycine, 1 µg mL^{-1} thiamine, 0.5 µg mL^{-1} pyridoxine, and 0.5 µg mL^{-1} nicotinic acid. A single plant was chosen as the progenitor of all of the plants used in this study. All of its descendants were maintained at 25˚C in 16 h light, and subcultured vegetatively every 4 wk. Over the course of the project, rooted clones with at least 4-6 leaves were put into 2 L of hydroponic medium (Yoshida *et al.*, 1976), referred to here as Fake Field. Each container was diffusely aerated through an aquarium stone, maintained at constant volume by the addition of distilled water, and emptied and refilled with fresh hydroponic medium every 7 d. Hydroponic containers were maintained at 22˚C, 16 h light with an irradiance level of 0.0006 W m^{-2}. Illumination was provided by GE Lighting Fluorescent lamps (13781, F96T12/CW/1500). After a 2 wk lag-time, plants began producing 1-3 new leaves each wk, and flowered continuously afterwards. All experiments were performed on plants that had not been infected or wounded in any way previously.

RNA extraction

RNA was extracted from SSI bud, stem, leaf, and root and infected root organs adapted from the protocol in Casavant *et al.*, 2017. Adaptions included use of a coffee grinder to homogenize tissue with the addition of dry ice to maintain RNA integrity.

Genome size estimation by Flow Cytometry

Healthy green leaf tissues were collected from SSI plantlets growing *in vitro*. Roughly 1 cm^2 (0.01g or less) of leaf was chopped in 1mL ice cold LB01 buffer for 1.5-2 min (Doležel *et al.*, 1989). The LB01 buffer contained 50µg mL^{-1} RNase stock and 50µg mL^{-1} propidium iodide (25% PI stock in DMSO) per mL of LB01. Each sample was chopped

with a fresh razor blade in a clean Pyrex petri plate. The finely chopped suspension was then

filtered through a 50μm nylon mesh filter (Partec 04-0042-2317). This filtered suspension

was kept in the dark at 4˚C for between 15-90 min before it was analyzed.

Genome size estimations were made using a BD FACSARIA Flow Cytometer

(IBEST Imaging Core, University of Idaho, Moscow, ID, USA). A green laser at 488nm was

used to excite the propidium iodide stained cells and was then collected in the PE-A channel.

Thresholds for PE-A were set at 1,000 and FSC at 500. The voltages were set so the major

peak (2C) of the SSI samples were near 50,000 on the linear scale. Four suspensions were

made from separate donor plants once a day for three consecutive days. Two replicates of

two external standards were also used daily in addition to the 4 SSI samples. External

standards included *Solanum lycopersicum cv.* Stupicke polni tyckove rane (2C= 1.96 pg

DNA) and *Glycine max* cv. Polanka (2C= 2.50 pg DNA) (Doležel *et al.*, 1992; Doležel *et al.*,

1994) which were chosen because their genome sizes were in the expected range of SSI. One

repetition of internal standards was run using tomato and soybean. DNA content was

estimated using the equation described by Doležel *et al.*, 2007.

Library Preparation for Iso-Seq

SMRT library preparation and sequencing were performed by the National Center for

Genome Resources (Santa Fe, New Mexico). The Iso-Seq libraries for four organs, root,

stem, leaf and bud, were prepared for Isoform Sequencing (Iso-Seq) using the Clontech

SMARTer PCR cDNA Synthesis Kit and the BluePippin Size Selection System protocol as

described by Pacific Biosciences (https://goo.gl/ij71Hh) with the following modifications.

For cDNA conversion, 3 μg of total RNA was put into each Clonetech SMARTer reaction.

From the PCR optimization procedure specified in the protocol, it was determined that 14

cycles of PCR would be sufficient for amplification of each organ's cDNA. Amplification

was followed by size selection on each sample to obtain three size bins (0.5-2 kb, 1.5-3 kb

and 2.5-6 kb) using the Blue Pippin (Sage Science, Beverly, Massachusetts) instrument. The

amplified and size selected cDNA products were made into SMRTbell Template libraries per

the Isoform Sequencing protocol referenced above. Libraries were prepared for sequencing

by annealing a sequencing primer (component of the SMRTbell Template Prep Kit 1.0) and

then binding polymerase to this primer-annealed template. The polymerase-bound template

was bound to MagBeads (P/N 100-125-900) (https://goo.gl/wdZErU) and sequencing was

performed on a PacBio RS II instrument. 12 v3 SMRTcells were run for the root tissues, 14

for the leaf tissues, 9 for the stem tissues, and 12 for the bud tissues for a total of 47

SMRTcells (Pacific Biosciences, P/N 100-171-800). The libraries from each organ were

separately sequenced using P6C4 polymerase and chemistry and 240-minute movie times

(Pacific Biosciences, P/N 100-372-700, P/N 100-356-200).

Single Molecule Real Time Sequencing

All SMRT cells for a given organ were run through the Iso-Seq pipeline included in

the SMRT Analysis software package. First, reads of insert (ROIs, previously known as

circular consensus sequences or CCS) were generated using the minimum filtering

requirement of 0 or greater passes of the insert and a minimum read quality of 75. This

allowed for the high yields going into subsequent steps, while providing high accuracy

consensus sequences where possible. The pipeline then classified the ROI in terms of full-

length, nonchimeric and non-full length reads. This was done by identifying the 5' and 3'

adapters used in the library preparation as well as the poly(A) tail. Only reads that contained

all three in the expected arrangement and did not contain any additional copies of the adapter

sequence within the DNA fragment were classified as full-length non-chimeric copies.

Finally, all full-length non-chimeric reads were run through the Iterative Clustering for Error

correction algorithm then further corrected by the Pacific Biosciences Quiver algorithm

(https://github.com/PacificBiosciences/cDNA_primer/wiki/Understanding-PacBio-

transcriptome-data). Once the Iso-Seq pipeline result was available for each organ, the results

were combined into a single data set and redundant sequences were removed using CD-HIT-

EST (Li and Godzik, 2006).

Illumina Sequencing for divergent gene analysis

Extracted total RNA from each previously stated organ was sent to Eurofins

Genomics (Ebersberg, Germany) for library preparation and sequencing. Prior to library

preparation, quality control (QC) was performed on individual tubes of RNA and equal

aliquots of each preparation were blended into one pool. The Illumina library cDNA was

prepared using randomly-primed first and second strand synthesis, followed by gel sizing

and PCR amplification. The library was then physically normalized and found to have insert

sizes of 250-450 bp.

Illumina sequencing was performed on a MiSeq v3 2x300. The read sequences were

clipped using Trimmomatic, version 0.32 (Bolger *et al.*, 2014), and bases with a Phred score

< 20 were removed. Trimmed reads shorter than 150 bp were removed; this step could

remove none, one, or both mates of a read-pair. Digital normalization was applied to the

Illumina reads in order to reduce redundant information present in these large datasets. A

coverage cutoff of 30 and a kmer size of 20 decreased the data to 28% of the initial

31,310,146 reads. BWA (Li and Durbin, 2009) was used to map read pairs to the *Solanum*

tuberosum cultivar Desiree chloroplast genome (GenBank accession DQ38616.2). Only

unmapped reads were retained.

Annotation of Sequences

Mercator sequence annotation was performed using the TAIR, PPAP, KOG, CDD,

IPR, BLAST CUTOFF of 80, and ANNOTATE options (Lohse *et al.*, 2014).

Annotating Protein Domains of Translated Sequences

PfamScan (Finn *et al.*, 2009) was run on the SSI transcriptomes following protocols

set forth by Sarris *et al.*, 2016.

Biological Quality Check of *in silico* Sequences

45 clones from a cDNA library (Express Genomics, Average insert size=1 kb,

Vector= pExpress 1) were randomly selected and sequenced via Sanger Dye-Deoxy DNA

Sequencing (ABI 3730). These sequences were then aligned to the transcriptomes using

Bowtie2 (Langmead and Salzberg, 2012) set for local alignment and best hit only. These

aligned sequences were then manually compared for possible chimeric features.

Evolutionary Comparison of SSI to 13 Other Species

The evolutionary clustering and comparison protocols were adapted from those set

out in Yang *et al.*, 2014. See Supplemental Table 4 for species used and online download

sources.

Divergent Gene Analysis to determine Ploidy

Phasing of the SMRT transcriptome was completed using unassembled Illumina sequences adapted from protocols established by Krasileva *et al.*, 2013, with the addition of an in-house Python script to quantify single-nucleotide polymorphisms present per sequence (https://github.com/AlexWixom/Transcriptome_scripts/freePloidy.py).

Creation of Expression Snapshots Using only SMRT Sequences

In-house Python scripts were used to backtrack final transcriptome sequences to each organ using CD-HIT-EST cluster files (https://github.com/AlexWixom/Transcriptome_scripts).

Expression Snapshot Validation

Sequence specific oligonucleotides were designed for several genes that were then used to obtain semi-quantitative PCR expression snapshots on the same cDNA used to obtain our SSI transcriptome (referred to as the "Sequenced" sample), as well as on a second cDNA pool prepared from RNA collected from independently grown plants (referred to as the "Unsequenced" sample). PCR fragment bands were quantified with a local background subtracted and normalized to actin (following the procedure established by Casavant *et al.*, 2017). The primers for these genes can be found in Supplemental Table 3 with the proposed gene description.

Accession Numbers

The SMRT sequenced transcriptome has been deposited at DDBJ/EMBL/GenBank under the accession GGFC00000000. The version described in this paper is the first version, GGFC01000000.

Data availability

File S1 contains detailed descriptions of all supplemental files. Sequence data are available at GenBank and the accession numbers are GGFC00000000. In-house code used to generate data can be found at https://github.com/AlexWixom/Transcriptome_scripts.

Author Contributions

NCC prepared many of the RNA samples, and all of the PCR analyses. AQW performed all remaining research, statistical analysis, and figure assemblies. AQW and ABC contributed to the experimental design. AQW wrote the manuscript with assistance from ABC. JCK, FX, and L-MD provided equipment and facilities, bacterial clones and plant material, and assisted in data interpretation. All authors have read and approved the manuscript.

Acknowledgments

The authors would like to thank Dr. R. Tripepi for providing us with vegetatively propagated *S. sisymbriifolium*. We are also extremely grateful to the National Center for Genomics Research for preparation and sequencing of SMRT libraries. This research was funded by the Idaho Potato Commission, the Northwest Potato Research Consortium, and USDA-APHIS Farm Bill 10201, 10007.

Arumuganathan, K. and Earle, E. (1991). Nuclear dna content of some important plant
species. *Plant molecular biology reporter*, 9(3):208–218.

Bagalwa, J.-J. M., Voutquenne-Nazabadioko, L., Sayagh, C., and Bashwira, A. S. (2010).
Evaluation of the biological activity of the molluscicidal fraction of solanum sisymbriifolium
against non target organisms. *Fitoterapia*, 81(7):767–771.

Bolger, A., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for illumina
sequence data. *Bioinformatics (Oxford, England)*, 30(15):2114–2120.

Casavant, N. C., Kuhl, J. C., Xiao, F., Caplan, A. B., and Dandurand, L.-M. (2017).
Assessment of globodera pallida rna extracted from solanum roots. *Journal of nematology*,
49(1):12.

Dandurand, L.-M. and Knudsen, G. (2016). Effect of the trap crop *Solanum sisymbriifolium*
and two biocontrol fungi on reproduction of the potato cyst nematode, *Globodera pallida*.
Annals of Applied Biology, 169(2):180–189.

Doležel, J., Binarová, P., and Lcretti, S. (1989). Analysis of nuclear dna content in plant cells
by flow cytometry. *Biologia plantarum*, 31(2):113–120.

Doležel, J., Doleželová, M., and Novák, F. (1994). Flow cytometric estimation of nuclear
dna amount in diploid bananas (musa acuminata andm. balbisiana). *Biologia plantarum*,
36(3):351–357.

Doležel, J., Greilhuber, J., and Suda, J. (2007). Estimation of nuclear dna content in plants
using flow cytometry. *Nature protocols*, 2(9):2233–2244.

Doležel, J., Sgorbati, S., and Lucretti, S. (1992). Comparison of three dna fluorochromes for
flow cytometric estimation of nuclear dna content in plants. *Physiologia plantarum*,
85(4):625–631.

Eddy, S. R. (1998). Profile hidden markov models. *Bioinformatics*, 14(9):755.

Eid, J., Fehr, A., Gray, J., Luong, K., Lyle, J., Otto, G., Peluso, P., Rank, D., Baybayan, P.,
Bettman, B., *et al.* (2009). Real-time dna sequencing from single polymerase molecules.
Science, 323(5910):133–138.

Finn, R. D., Mistry, J., Tate, J., Coggill, P., Heger, A., Pollington, J. E., Gavin, O. L.,
Gunasekaran, P., Ceric, G., Forslund, K., *et al.* (2009). The pfam protein families database.
Nucleic acids research, 38(suppl_1):D211–D222.

Finn, R. D., Tate, J., Mistry, J., Coggill, P. C., Sammut, S. J., Hotz, H.-R., Ceric, G.,

Forslund, K., Eddy, S. R., Sonnhammer, E. L., *et al.* (2008). The pfam protein families database. *Nucleic acids research*, 36(suppl 1):D281–D288.

Garrison, E. and Marth, G. (2012). Haplotype-based variant detection from short-read sequencing. https://arxiv.org/abs/1207.3907v2.

Iseli, C., Jongeneel, C. V., and Bucher, P. (1999). Estscan: a program for detecting, evaluating, and reconstructing potential coding regions in est sequences. In *ISMB*, volume 99, pages 138–148.

Jones, J. D. and Dangl, J. L. (2006). The plant immune system. *Nature*, 444(7117):323–329.

Krasileva, K. V., Buffalo, V., Bailey, P., Pearce, S., Ayling, S., Tabbita, F., Soria, M., Wang, S., Akhunov, E., Uauy, C., *et al.* (2013). Separating homeologs by phasing in the tetraploid wheat transcriptome. *Genome Biology*, 14(6):R66.

Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with bowtie 2. *Nature methods*, 9(4):357–359.

Li, H. and Durbin, R. (2009). Fast and accurate short read alignment with burrows–wheeler transform. *Bioinformatics*, 25(14):1754–1760.

Li, L., Stoeckert, C. J., and Roos, D. S. (2003). Orthomcl: identification of ortholog groups for eukaryotic genomes. *Genome research*, 13(9):2178–2189.

Li, W. and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13):1658–1659.

Lohse, M., Nagel, A., Herter, T., May, P., Schroda, M., Zrenner, R., Tohge, T., Fernie, A. R., Stitt, M., and Usadel, B. (2014). Mercator: a fast and simple web server for genome scale functional annotation of plant sequence data. *Plant, cell & environment*, 37(5):1250–1258.

Loman, N. J., Misra, R. V., Dallman, T. J., Constantinidou, C., Gharbia, S. E., Wain, J., and Pallen, M. J. (2012). Performance comparison of benchtop high-throughput sequencing platforms. *Nature biotechnology*, 30(5):434–439.

Luo, C., Tsementzi, D., Kyrpides, N., Read, T., and Konstantinidis, K. T. (2012). Direct comparisons of illumina vs. roche 454 sequencing technologies on the same microbial community dna sample. *PloS one*, 7(2):e30087.

Marck, C. (1988). 'dna strider': a 'c'program for the fast analysis of dna and protein sequences on the apple macintosh family of computers. *Nucleic acids research*, 16(5):1829–1836.

Meyre-Silva, C., Niero, R., Bolda Mariano, L. N., Gomes do Nascimento, F., Vicente Farias,
I., Gazoni, V. F., dos Santos Silva, B., Giménez, A., Gutierrez-Yapu, D., Salamanca, E., *et al.*
(2013). Evaluation of antileishmanial activity of selected brazilian plants and identification
of the active principles. *Evidence-Based Complementary and Alternative Medicine*, 2013.

Miz, R. B., Mentz, L. A., and Souza-Chies, T. T. (2008). Overview of the phylogenetic
relationships of some southern brazilian species from section torva and related sections of
"spiny solanum"(solanum subgenus leptostemonum, solanaceae). *Genetica*, 132(2):143–
158.

Moreton, J., Izquierdo, A., and Emes, R. D. (2015). Assembly, assessment, and availability
of de novo generated eukaryotic transcriptomes. *Frontiers in genetics*, 6.

Ocwieja, K. E., Sherrill-Mix, S., Mukherjee, R., Custers-Allen, R., David, P., Brown, M.,
Wang, S., Link, D. R., Olson, J., Travers, K., *et al.* (2012). Dynamic regulation of hiv-1 mrna
populations analyzed by single-molecule enrichment and long-read sequencing. *Nucleic*
acids research, 40(20):10345–10355.

Paul, A. and Banerjee, N. (2015). Phylogenetic relationship of some species of *SOLANUM*
based on morphological, biochemical and cytological parameters. *Indian Journal of*
Fundamental and Applied Life Sciences, 5(3):51–56.

Särkinen, T., Barboza, G. E., and Knapp, S. (2015). True black nightshades: Phylogeny and
delimitation of the morelloid clade of solanum. *Taxon*, 64(5):945–958.

Sarris, P. F., Cevik, V., Dagdas, G., Jones, J. D., and Krasileva, K. V. (2016). Comparative
analysis of plant immune receptor architectures uncovers host proteins likely targeted by
pathogens. *BMC biology*, 14(1):8.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M.
(2015). Busco: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics*, 31(19):3210.

Timmermans, B. G. (2005). *Solanum sisymbriifolium (Lam.): a trap crop for potato cyst*
nematodes.

Yang, X., Cheng, Y.-F., Deng, C., Ma, Y., Wang, Z.-W., Chen, X.-H., and Xue, L.-B. (2014).
Comparative transcriptome analysis of eggplant (solanum melongena l.) and turkey berry
(solanum torvum sw.): phylogenomics and disease resistance analysis. *BMC genomics*,
15(1):412.

Yang, Y. and Smith, S. A. (2013). Optimizing de novo assembly of short-read rna-seq data

for phylogenomics. *BMC genomics*, 14(1):328.

Yoshida, S., Douglas, A. F., James, H. C., and Kwanchai, A. G. (1976). Laboratory manual

for physiological studies of rice. *Manila: IRRI*, 56:69–77.

Zhang, W., Ciclitira, P., and Messing, J. (2014). Pacbio sequencing of gene families-a case

study with wheat gluten genes. *Gene*, 533(2):541–546.

Figure Legends

Figure 1: Correspondence between 45 randomly selected Sanger-sequenced SSI cDNAs and the SMRT transcriptome. A) Bowtie2 was used to determine the presence of 45 cDNA clones in the SMRT SSI transcriptome. B) Alignment of matched sequences to SMRT SSI transcriptome was performed using DNA Strider and manually evaluated as either equivalent or chimeric. All Sanger sequenced clones were found in the SMRT dataset and a small percentage were found to be chimeric. See Supplemental Figure 2 for corresponding analysis of Illumina sequenced, Velvet/Oases transcriptome.

Figure 2: Shared and restricted orthologous genes among 13 species. All species shown here shared 6067 core orthologs. Each petal shows the number of gene groups unique to each species. Not shown are groups shared by only 2–12 species. *Solanum sisymbriifolium*, SSI; *Solanum tuberosum*, STU; *Solanum lycopersicum*, SLY; *Solanum melongena*, SME; *Arabidopsis thaliana*, ATH; *Carica papaya*, CPA; *Vitis vinifera*, VVI; *Prunus persica*, PPE; *Populus trichocarpa*, PTR; *Citrus sinensis*, CSI; *Medicago truncatula*, MTR; *Zea mays*, ZMA; and *Oryza sativa*, OSA. See Supplemental Figure 5 for gene ontology bins for the SSI unique groups, and Supplemental Table 4 for sources of datasets.

Figure 3: Final SMRT transcriptome sequences were backtracked through the de-redundification process to the organ sub-transcriptomes. A) Flower plot of genes expressed in all, or in only one, organ. Each petal shows the number of genes only expressed in one organ. In the center are the number of genes expressed in all 4 organs. Not shown are genes expressed in 2 or 3 organs. B) Green-tissue specificity of sequences annotated as genes involved in the light harvesting complex-I pathway via Mercator. C) Sequences annotated as putative resistance genes because they contained nucleotide binding (NB-ARC) and leucine rich repeat (LRR) domains showed varied expression patterns. As shown on the scales on the right of (B) and (C), the darker the color, the more times the sequence was found in that organ.

Figure 4: Comparison of expression of 3 putative R-gene sequences in the SMRT

database to semi-quantitative PCR from 2 cDNA preparations. A) The expression of

three genes with LRR and NB-ARC domains characteristic of the R-genes and an actin

isoform is shown in the heat map at the left and compared on the right to semi-quantitative

PCR of those same genes in two independently prepared cDNA pools, one from the pool

used to generate the transcriptome (Sequenced) and one prepared independently, and not

used to make the transcriptome (Unsequenced). B) The expression of each PCR product from

each pool was quantified and then normalized to the expression of an actin isoform

(Ssi032526). Data (biological replicates, n=2) are represented as mean ± STD. See

Supplemental Table 3 for primers used.

Tables

Table 1: Summary of the SSI transcriptome derived using SMRT technology. Organ sub-transcriptomes were sequenced and combined from 33,170 root, 99,924 bud, 50,825 leaf, and 47,793 stem reads. See Supplemental Figure 3 and Supplemental Figure 4 for gene ontology bins of this transcriptome.

SMRT Assembly		SSI
Total raw reads		231,712
Read lengths		300-7883
Total raw reads size (bp)		362,086,346
GC content		41.17
	Number	139,611
	Total length	237,865,670
Transcripts	N50	2,050
	Max length	7,883
	GC content	40.97
	Number	41,189
	Total length	74,642,518
Unigenes	N50	2,158
	Max length	7,883
	GC content	39.90

642 **Table 2: BUSCO assessment for completeness of 3 transcriptomes and one genome.** The SSI

643 transcriptome appears to be nearly complete, but contains a disproportion number of duplicated

644 sequences. See Supplemental Table 4 for sources of datasets.

	Transcriptome			Genome
	SSI (%)	Tomato (%)	Potato (%)	ATH (%)
Complete BUSCOs	93	96.2	86.7	97.7
Complete Single-copy BUSCOs	62.8	94.2	64.1	N/A
Complete Duplicated BUSCOs	30.2	2.0	22.6	N/A
Fragmented BUSCOs	2.0	0.8	4.7	0.6
Missing BUSCOs	4.1	3.0	8.6	1.7

Table 3: Divergent gene assessment of allele and/or paralog number in the SSI transcriptome.

648   4-allele genes were mapped to tomato chromosomes, see Supplemental Table 2.

Allele or Paralog	# of genes
Homozygous	17,773
2	22,098
3	1,358
4	44

651 **Table 4: R-gene profile of potato (STU), tomato (SLY), and the SSI transcriptome.** The SSI

652 database had fewer assigned R genes (based on the presence of nucleotide-binding domains and

653 leucine-rich repeats within the same open reading frame) than either SLY or STU genomes. Refer to

654 Supplemental Table 1 for full domain annotation statistics.

	R-genes
SSI	67
STU	309
SLY	137

657 **Figures**

(A)   (B)

Figure 1: Correspondence between 45 randomly selected Sanger-sequenced SSI cDNAs and

the SMRT transcriptome. A) Bowtie2 was used to determine the presence of 45 cDNA clones in

the SMRT SSI transcriptome. B) Alignment of matched sequences to SMRT SSI transcriptome was

performed using DNA Strider and manually evaluated as either equivalent or chimeric. All Sanger

sequenced clones were found in the SMRT dataset and a small percentage were found to be chimeric.

See Supplemental Figure 2 for corresponding Illumina sequenced, Velvet/Oases assembled

transcriptome.

Figure 2: Shared and restricted orthologous genes among 13 species. All species shown here shared 6067 core orthologs. Each petal shows the number of gene groups unique to each species. Not shown are groups shared by only 2–12 species. *Solanum sisymbriifolium*, SSI; *Solanum tuberosum*, STU; *Solanum lycopersicum*, SLY; *Solanum melongena*, SME; *Arabidopsis thaliana*, ATH; *Carica papaya*, CPA; *Vitis vinifera*, VVI; *Prunus persica*, PPE; *Populus trichocarpa*, PTR; *Citrus sinensis*, CSI; *Medicago truncatula*, MTR; *Zea mays*, ZMA; and *Oryza sativa*, OSA. See Supplemental Figure 5 for gene ontology bins for the SSI unique groups, and Supplemental Table 4 for sources of datasets.

Figure 3: Final SMRT transcriptome sequences were backtracked through the de-redundification process to the organ sub-transcriptomes. A) Flower plot of genes expressed in all, or in only one, organ. Each petal shows the number of genes only expressed in one organ. In the center are the number of genes expressed in all 4 organs. Not shown are genes expressed in 2 or 3 organs. B) Green-tissue specificity of sequences annotated as genes involved in the light harvesting complex-I pathway via Mercator. C) Sequences annotated as putative resistance genes because they contained a nucleotide binding domain (NB-ARC) and leucine rich repeat (LRR) domains show varied expression patterns. As shown on the scales on the right of (B) and (C), the darker the color, the more times the sequence was found in that organ.

(A)

SSI sequence ID	A. thaliana homologue	Leaf	Root	Stem	Bud
Ssi035516	ADR1-L1 (at4g33300) NB-ARC	0.64 ± 0.22	0.99 ± 0.13	0.20 ± 0.02	0.39 ± 0.00
Ssi038051	domain-containing disease resistance protein (at1g50180)	0.88 ± 0.40	0.43 ± 0.24	0.69 ± 0.06	0.64 ± 0.13
Ssi036337	RPM1 (at3g07040)	0.22 ± 0.02	1.56 ± 0.56	0.19 ± 0.05	0.36 ± 0.10

(B)

Figure 4: Comparison of expression of 3 putative R-gene sequences in the SMRT database to semi-quantitative PCR from 2 cDNA preparations. A) The expression of three genes with LRR and NB-ARC domains characteristic of the R-genes and an actin isoform is shown in the heat map at the left and compared on the right to semi-quantitative PCR of those same genes in two independently prepared cDNA pools, one from the pool used to generate the transcriptome (Sequenced) and one prepared independently, and not used to make the transcriptome (Unsequenced). B) The expression of each PCR product from each pool was quantified and then normalized to the expression of an actin isoform (Ssi032526). Data (biological replicates, n=2) are represented as mean ± STD. See Supplemental Table 3 for primers used.