Kinetics of winter deacclimation in response to temperature determines dormancy status in

grapevines (*Vitis* spp.)

Alisson P. Kovaleski[1], Bruce I. Reisch[1], and Jason P. Londo[1,2*]

[1] School of Integrative Plant Science – Horticulture Section, Cornell University-New York State

Agricultural Experiment Station, 630 W. North Street, Geneva, New York, USA.

[2] United States Department of Agriculture, Agricultural Research Service, Grape Genetics

Research Unit, 630 W. North Street, Geneva, New York, USA.

[*] Corresponding author

APK – e-mail: ap874@cornell.edu

BIR – e-mail: bruce.reisch@cornell.edu

JPL – e-mail: jason.londo@ars.usda.gov; phone: (315) 787-2463

Submission: 31 Jan 2018

Tables: 1

Figures: 4

Word count: 5276

Supplementary Tables: 1

Title

Kinetics of winter deacclimation in response to temperature determines dormancy status in

grapevines (*Vitis* spp.)

Running title

Kinetics of deacclimation resolves dormancy status

Highlight

Deacclimation rates in *Vitis* species appear to follow enzyme kinetics; and using deacclimation

rates, dormancy transitions can be studied in a quantitative manner.

Abstract

Bud dormancy and cold hardiness are critical adaptations for surviving winter cold

stress for temperate perennial plant species, with shifting temperature-based responses during

the winter. The objective of this study was to uncover the relationship between dormancy

transition (chilling requirement) and temperature on the loss of cold hardiness. Dormant

cuttings of European grapevine cultivars (*Vitis vinifera*; 'Cabernet Franc', 'Cabernet

Sauvignon', 'Riesling', and 'Sauvignon blanc') and wild grapevine species (*V. aestivalis, V.

amurensis*, and *V. riparia)* were examined to determine the relationship between chilling

requirement and temperature on rate of deacclimation (k_{deacc}). Differential thermal analysis was

used to determine k_{deacc} using mean low temperature exotherms. Effect of chill was evaluated

as the deacclimation potential (Ψ_{deacc}), which was the change in k_{deacc} due to chill

accumulation. Results indicate that Ψ_{deacc} varies dependent on dormancy state, following a

logarithmic response to chill accumulation. The effect of temperature on k_{deacc} was exponential

at low and logarithmic at high temperatures. The combination of Ψ_{deacc} and k_{deacc} resulted in

good prediction of deacclimation, demonstrating that k_{deacc} can be used as a quantitative

determinant of dormancy transition. This information can be used to refine models predicting

effects of climate change on dormancy and cold hardiness in grapevine.

Key words

Bud, cold hardiness, deacclimation, dormancy, grapevines, rates, supercooling, *Vitis*.

Abbreviations

DTA – Differential thermal analysis

HTE – High temperature exotherm

LTE – Low temperature exotherm

k_{deacc} – Deacclimation rate

Ψ_{deacc} – Deacclimation potential

Chill – Chill accumulation

t – time

T – temperature

Introduction

Due to their general stationary habit, plants have evolved many different coping mechanisms to survive stressful environments such as drought and temperature. With regard to winter in temperate climates, annual plants typically survive as dehydrated seeds while perennial plants develop dormancy and cold hardiness. Dormancy is the temporary cessation of visible growth in meristem containing structures, such as buds, and is divided in three types: para-, endo-, and ecodormancy (Lang *et al.*, 1987). Paradormancy is the suspension due to physiological factors within the plant but outside of the dormant structure, such as the suppression of lateral growth by the apical meristem (i.e. apical dominance). Paradormant buds transition into endodormancy as daylength and temperatures decrease, a state suspension of growth due to unknown endogenous factors within the dormant structure, preventing growth during times when environmental conditions fluctuate between conducive and inhibitory (Horvath *et al.*, 2003). During endodormancy, it is generally understood that exposure to low, non-freezing temperatures is necessary (chilling requirement) to transition to the third dormancy state, ecodormancy. Tissues maintain a dormant state during ecodormancy due to unsuitable environmental conditions.

Concomitantly with the onset of dormancy, plants from areas where temperatures drop below freezing develop cold hardiness through different mechanisms. Some plants, or plant

tissues, tolerate intracellular ice formation, in which cells are extremely dehydrated by the

formation of extracellular ice, and the remaining water is bound to proteins (Burke *et al.*, 1976)

likely solidifying in a glass-like state. Other plants present a freezing resistance mechanism,

where water supercools in the intracellular spaces (Burke *et al.*, 1976). Supercooling is a

process through which water can remain in liquid state below its equilibrium freezing point,

down to a maximum of ~ -40 °C (Bigg, 1953). These plants are typical of mid-latitude

temperate climates, where temperatures do not drop below -40 °C during the winter. In buds

that present supercooling ability, such as those of grapevines (*Vitis* spp.), cold hardiness may

be measured through differential thermal analysis (DTA). This method uses thermoelectric

modules that record voltage changes associated with the release of heat due to phase change

of water into ice (Mills *et al.*, 2006). In grapevine buds, two peaks can be identified using DTA:

a high temperature exotherm (HTE), typically around – 5 °C; and a low temperature exotherm

(LTE), which varies depending on the conditions experienced by the bud. The HTE represents

the freezing of extracellular water (Andrews *et al.*, 1984), which is normal to the process of

resisting cold. The LTE, however, is the freezing of intracellular water, and it happens when

the supercooling point has been exceeded (Mills *et al.*, 2006). The LTE represents the lethal

temperature for a given bud, is correlated with manual assessments of bud death (Wolf and

Cook, 1994), and mean LTE has been adopted as a consistent measurement of bud cold

hardiness.

A general U-shaped pattern of changing cold hardiness is followed during the winter

through three different stages: acclimation, maintenance, and deacclimation. This pattern is

thought to be primarily driven by air temperature (Ferguson *et al.*, 2011, 2014; Londo and

Kovaleski, 2017), although other climatic aspects may also be important (Antivilo *et al.*, 2017).

Acclimation appears to occur primarily during endodormancy, while deacclimation is enhanced

during ecodormancy (Ferguson *et al.*, 2011, 2014; Londo and Kovaleski, 2017). Depending on

the climate and genotype, the transition between endo- and ecodormancy may occur toward

the end of the acclimation stage, or during the maintenance stage of cold hardiness. This

transition is typically evaluated by collecting dormant buds from the field and placing them in

growth permissive conditions, followed by monitoring the time needed to reach budbreak

(Weinbaum *et al.*, 1989; Lloyd and Firth, 1990; Cook and Jacobs, 2000; Fan *et al.*, 2010;

Zhang and Taylor, 2011). The chilling requirement has been met when 50% budbreak occurs

within a time threshold {e.g. 3 weeks for peaches [*Prunus persica* (L.) Batsch; Lloyd and Firth,

1990]; 4 weeks for grapevines (Londo and Johnson, 2014)}. This method represents a

subjective threshold for the change in endo- to ecodormancy transition. As a result,

determining the molecular and metabolic cues and the consequences of this transition, such

as budbreak, are very difficult to determine and poorly understood (Penfield, 2008).

As the dynamics of acclimation and deacclimation change with dormancy state

(Ferguson *et al.*, 2011, 2014), and dormancy state is determined by accumulation of chilling

hours, a future climate may create issues for both cold hardiness and budbreak phenology.

Average global temperatures are increasing (Walsh *et al.*, 2014), although episodes of acute

cold weather in the northern hemisphere are likely to increase as well through more frequent

arctic oscillations (Kolstad *et al.*, 2010). In addition to direct effects to cold hardiness,

increasing temperatures will enhance chill accumulation in higher latitudes, while lower

latitudes will experience a decrease (Luedeling *et al.*, 2011). Changes in chill accumulation will

modify the dynamics of dormancy transitions both in deciduous tree crops and other woody

perennials, causing excessive responsiveness to warm spells or erratic budbreak and growth.

This will ultimately require a shift towards higher latitudes and elevations of tree crops and

assisted migration of forest species as they lag behind their optimal climate niche (Gray and

Hamann, 2013). Therefore, it is important to understand how both temperature and dormancy

status affect the loss of cold hardiness. The objective of this study was to better predict and

understand the relationship between chill accumulation, the dormancy transition, and the loss

of cold hardiness in wild and cultivated grapevines.

Materials and methods

Buds of four different species were collected multiple times during the dormant seasons

of 2014/15, 2015/16, and 2016/17. *V. vinifera* L. 'Cabernet Franc', 'Cabernet Sauvignon',

'Riesling', and 'Sauvignon blanc' were collected from local vineyards (42.705N, 76.973W; and

42.845N, 77.004W), while *V. aestivalis* Michx. (two clones: PI483138, PI483143), *V.*

amurensis Rupr. (three clones: PI588632, PI588635, PI588641), and *V. riparia* Michx. (four

clones: PI588275, PI588562, PI588653, PI588711) were collected from the USDA Plant

Genetic Resources Unit in Geneva, New York. For *V. vinifera*, buds from nodes 3 to 20 from

the base of a cane were used, while for the other species, due to constraints regarding the

number of clones, buds beyond position 20 were used as in Londo and Kovaleski (2017).

Hourly weather data from the closest Network for Environment and Weather

Applications station (NEWA; http://www.newa.cornell.edu/) was used to compute chill

accumulation using the "North Carolina" model (Shaltout and Unrath, 1983). The start date for

each season was chosen based on when chill started to consistently accumulate instead of

being negated, as determined by the NC model. These dates were 11 Sept 2014, 19 Sept

2015, and 24 Sept 2016.

Upon collection, canes were prepared into single- or two-node cuttings and placed

submerging the basipetal cut surface in cups of water. The cups were placed into growth

chambers at 2, 4, 7, 8, 10, 11, 22, or 30 °C. Not every genotype or temperature was used at all

collection points, but this information is available in Supplementary Table S1. Differential

thermal analysis (DTA) was performed to determine the cold hardiness of individual buds as

their low temperature exotherm [LTE; see Mills *et al.* (2006) for details]. Briefly, buds are

excised from the cane and placed on a thermoelectric module in a plate. These plates are

placed in a programmable freezer with a cooling rate of -4 °C/hour. Changes in voltage due to

heat release in the freezing of water are measured by the thermoelectric module and recorded

using a Keithley data logger (Tektronix, Beaverton, OR) attached to a computer. Between 4

and 8 buds were used at any time point to determine mean LTE, and the intervals between

measurement days varied between chill accumulation level and temperature treatment, with

details provided in Supplemental Table 1. Because lower rates of deacclimation were expected

in lower temperatures and lower chill accumulations, lower temperature treatments were

typically surveyed with wider separation between time points compared with high temperature

treatments (Table S1).

Statistics

Based on sample size across temperatures and across sampled years, data sets for *V.*

vinifera were analyzed separately for each cultivar, while data for accessions of *V. aestivalis*,

V. amurensis and *V. riparia* were combined at the species level. Analyses were separated into

the effect of chill accumulation and the effect of temperature on rate of deacclimation. The

datasets used for each analysis are specified in Supplementary Table S1.

Effect of Chill Accumulation. In order to assess the effect of chill accumulation in the rate of

deacclimation (k_{deacc}), or the deacclimation potential (Ψ_{deacc}), individual rates were calculated

using linear regression in R (ver. 3.3.0, R Foundation for Statistical Computing) for each

temperature and chill accumulation as factors. Although every temperature within a chill

accumulation had the same data for day 0 (field collection), temperatures were still allowed to

have different intercepts in order to reduce the effect of day 0 on k_{deacc}. From this regression

model, data points that had a studentized residual ≥2.5 were considered outliers and removed

from the dataset, and the model was re-fit. The k_{deacc} at each chill accumulation were then

transformed to percentage for 4, 7, 10, 11, and 22 °C, standardizing to their k_{deacc} at highest

chill accumulation (either 1440 – 7 and 11 °C – or 1580 chill units – 4, 10, and 22 °C). The

Ψ_{deacc} was estimated as a logistic regression. Initial estimation used the drc library, but final

estimation used the nls() function with the port algorithm, following the equation: $\Psi_{deacc} (\%) =$

$1/{1 + e^{[b \times (\ln Chill - c)]}}$, where *b* and *c* are the estimated parameters, and *Chill* is the chill

accumulation at the time of sample collection. The parameter *c* is the inflection point of the

logistic and *b* is the slope associated with the logistic regression.

Effect of Temperature. Initially, to study the effect of temperature, k_{deacc} obtained from multiple

linear regression (temperature and chill accumulation as factors; Supplementary Table S1) in

lower chill accumulations were re-scaled by dividing by the respective Ψ_{deacc} (e.g., because the

Ψ_{deacc} was estimated to be at ~30% at 860 chill units for 'Riesling', the rates obtained from that

data set were divided by 0.30). Since the effect of chill accumulation was estimated to be ~ 0

% for 360 chill units, these were excluded from the dataset for estimation of temperature effect.

Arrhenius plots were created to evaluate temperature responses and data were then divided

into two data sets, low (2 – 11 °C) and high temperatures (10 – 30 °C), and temperature

effects on deacclimation rates were evaluated separately. The rationale for doing this is

discussed below.

Instead of using the initially calculated rates derived from the above linear models, the

complete data sets were used with the effect of temperature as a continuous variable. The

values for the different chill accumulation points were normalized to that of full chill by

multiplying the Ψ_{deacc} at any given chill accumulation, effectively removing the effect of chill by

mathematically reducing the time for deacclimation. Lower chill accumulations were thus

included in the calculations instead of using only those where chill requirements were initially

assumed to be fulfilled (1440 and 1580 chill accumulations).

For low temperatures, the effect was assumed to be an exponential, and therefore the

rates were calculated as $k_{\text{deacc}} = m \times e^{(n \times T)}$, where *m* and *n* are the parameters estimated,

and *T* is the temperature in °C. For high temperatures, a logarithmic curve was used: $k_{\text{deacc}} =$

$q \times \ln(T - r)$, where *q* and *r* are the parameters estimated, and *T* is the temperature in °C. For

both low and high temperatures, parameters were estimated using nls(), with the port

algorithm.

The fits of all non-linear curves (effect of chill accumulation, low and high temperatures)

were tested using Effron's pseudo-R^2 with the Rsq function in the soilphysics library, which is

calculated using the sum of squares of ordinary residuals from the models. The final model for

loss of cold hardiness from an initial point is then $\Delta_{LTE}(T, Chill) = k_{\text{deacc}} \times \Psi_{\text{deacc}} \times t$, where

the change in LTE (Δ_{LTE}) is a function of the temperature (*T*) it was exposed to, how much chill

accumulation (*Chill*) had passed before exposed to deacclimation temperatures, and the time

(*t*) of exposure. A linear model was used to evaluate the accuracy of the predictions using the

complete data set. For this, instead of temperatures, $k_{\text{deacc}} = \begin{cases} m \times e^{(n \times T)}, T < 12\,°C \\ q \times \ln(T - r), T \geq 12\,°C \end{cases}$ was

used, and instead of accumulated chill, the Ψ_{deacc} was used. From this model ($\Delta_{LTE} = \beta \times k_{\text{deacc}}$

$\times \Psi_{\text{deacc}} \times$ time), considering estimations were appropriate, the β associated with the estimated

k_{deacc} and Ψ_{deacc} was expected to be 1.

Results

After removing outliers using multiple linear regression, data sets kept ≥ 90% of the

observations for all genotypes, with the exception of *V. amurensis*, which had 19.5% of

observations removed (Table 1). Linear behavior appears to describe well the effects of

temperature and chill accumulation in the deacclimation of grapevine buds (Figure 1). A

logistic regression appeared to be a good fit for the Ψ_{deacc} (Figure 2), with pseudo-R^2 values

greater than 0.87 for all genotypes except for 'Cabernet Sauvignon' and *V. amurensis* (0.65

and 0.79, respectively; Table 1). The parameter *e* is the natural logarithm of the inflection

point, and there was little difference in this parameter for these genotypes (e.g., 6.77 to 6.88,

for 'Sauvignon blanc' and 'Riesling', respectively, which is equivalent to 872 to 972 chill units).

Scaled deacclimation rates based on Ψ_{deacc} and corresponding Arrhenius plot (Figure 3)

demonstrated a clear discontinuity between low and high temperatures. A deceleration of the

increase of k_{deacc} as temperature increased was observed, which justifies the different

behaviors used (exponential and logarithmic for low and high temperatures, respectively). The

estimation of temperature effects for low and high temperature intersect between 10 and 11 °C

because of the uncertainty in the exact temperature where there is a change in behaviors. This

is partially due to the lack of data points between 11 and 22 °C, although at 10 and 11 °C the

calculated rates through either model would be very similar. All the *V. vinifera* cultivars had

similar deacclimation rates at low temperatures, but 'Riesling' and 'Cabernet Franc' had higher

deacclimation rates at high temperatures than 'Cabernet Sauvignon' and 'Sauvignon blanc'

(Table 1, Figure 4). *V. riparia* and *V. aestivalis* had similar rates to those of *V. vinifera* at low

temperatures, but *V. riparia* had estimated rates similar to *V. amurensis* in higher

temperatures. *V. aestivalis* was similar to the faster *V. vinifera* ('Cabernet Franc' and

'Riesling'). *V. amurensis* showed the highest deacclimation rates at both low and high

temperatures.

Linear models showed a good fit for all genotypes ($R^2 \geq 0.80$). *V. vinifera* cultivars,

which were analyzed separately had higher R^2 values. The better fit for V. *vinifera* cultivars is

also seen in their β coefficients, which are very close to 1, although *V. aestivalis* and *V. riparia*

had βs of 0.99 and 0.96 respectively. *V. amurensis* had the lowest β, at 0.83.

Discussion

Chilling accumulation, dormancy state, and cold hardiness are complex traits driven by

the interaction of physiological and climactic attributes. The consistency of winter temperatures

is expected to be a major aspect impacting the sustainability and survival of perennial species

in a future climate. This study was conducted to gain a better method for predicting dormancy

transition and chilling requirement in grapevine, as well as to understand how deacclimation

processes (loss of cold hardiness) are impacted by dormancy state.

This paper provides a quantitative assessment of the transition from endo- to
ecodormancy, as evaluated by the Ψ_{deacc}. Unlike the typical dormancy assessment that uses
time to budbreak as a measurement of the transition (Londo and Johnson, 2014), using the
deacclimation rates does not require subjective assumptions currently used to determine
chilling fulfillment; such as having 50% budbreak occur within 28 days. Budbreak occurs only
after the freeze mechanisms associated with cold hardiness are lost (Ferguson *et al.*, 2014;
Salazar-Gutiérrez *et al.*, 2014). Thus, the amount of time needed to reach 50% budbreak in
forcing assays is a combination of the time needed to fully deacclimate, and time required for
deacclimated buds to reach a visible change in phenological stage. This study leverages the
ability to assess loss of hardiness through decreasing supercooling ability as a way to
quantitatively assess rates of deacclimation and its relation to changes in chilling exposure.
Mid-winter warming events are of major concern for temperate crops, yet natural areas
are also affected. In native trees within natural ranges, little to no damage occurs during the
autumn because the freeze resistance readily withstands the minimum temperatures they
usually encounter (Vitasse *et al.*, 2014). However, damage can be substantial in late freeze
events during the spring or unusual warm events in early spring as the potential for
deacclimation increases exponentially (Vitasse *et al.*, 2014). For example, a late spring freeze
in 2007 in North America caused damage in native forests, delaying and reducing canopy
establishment by 16-34 days (Gu *et al.*, 2008). Bokhorst *et al.* (2009) reported that dwarf shrub
vegetation in a sub-Artic region where midwinter temperatures rose to ~7 °C for 2 weeks
followed by further cold temperatures resulted in an 87% reduction in growth during the
summer compared to neighboring undamaged areas. Cannell and Smith (1983) show results
from different authors for forest species in which the time to budbreak is reduced in an
exponential manner. This is similar to our assessment of Ψ_{deacc}, where the upper portion of the
logistic is like the exponential decrease in time to budbreak. The logistic behavior of Ψ_{deacc} in
relation to chill accumulation explains the continuous increase risk of damage of mid-winter
warm spells the later they occur until 100% Ψ_{deacc} is achieved. This is in contrast with the
model developed by Ferguson *et al.* (2011; 2014), where a hard limit for the dormancy
boundary is used to model grape cold hardiness. It is likely that, similar to the behavior
described in our study, forest species also have increased deacclimation rates with higher chill
accumulation, seen as a decrease in the time to budbreak. Additionally, it is important to note

that while canopies may be able to recover from midwinter bud kill events, damage to the

reproductive buds in the case of fruit and nut crops will reduce yields in the season following

damage.

Our assessment of Ψ_{deacc} leading to increases in k_{deacc} agrees with the general

observation that warm temperatures in the early winter are negligible for bud growth (Cannell

and Willett, 1975) and that increasing chill accumulation reduces the heat requirements for

budbreak in fruit species (Couvillon and Erez, 1985; Citadin *et al.*, 2001), and forest species

(Hunter and Lechowicz, 1992; Dantec *et al.*, 2014), considering that loss of hardiness is

required for budbreak to occur (Ferguson *et al.*, 2014). Even at full chilling (100% Ψ_{deacc}),

differences in the k_{deacc} between species and cultivars may be responsible for differences in

early phenological development in the spring, but it is important to recognize there is an

interaction between both. Planted in the same environment, Taulavuori *et al.* (2004) observed

that deacclimation occurs faster in mountain birch [*Betula pubescens* ssp. *czerepanovii* (Orl.)

Hämet-Ahti] ecotypes with origin in areas of low chill accumulation. If the mechanisms are

similar to those described here, what may have been observed in their study was not different

rates of deacclimation, but plants that were at different stages in their deacclimation potential,

or a combination of both. Climate change appears to speed the loss of cold hardiness and

promote earlier leaf-out, although the response is not linear due to interactions with chilling and

photoperiod (Vitasse *et al.*, 2014). In our case, the predicted increase in chill accumulation

(Luedeling *et al.*, 2011) will result in higher Ψ_{deacc} and higher risk of damage should midwinter

warm spells increase in a future climate (Vitasse *et al.*, 2014; Walsh *et al.*, 2014). However,

earlier budbreak may not lead to higher frequency of damage due to frost events if consistently

higher spring temperatures (Chmielewski *et al.*, 2018) accompany future climate changes.

Pagter and Arora (2013) suggest that deacclimation rates are not linear, and that a lag

phase may occur during initial exposure to warm temperatures. The initial lag phase might

explain the differences in deacclimation rates measured in Cragin *et al.* (2017) compared to

those reported in this study, as they only evaluated deacclimation for 4 days. This may also

suggest why they had differences in rates across the years, although this may also be an

effect of different chill accumulation between the two years in their study. Ferguson *et al.*

(2011; 2014) used asymptotic bounds added through a logistic component, deviating from a

strictly linear rate. In our data, although in some cases an initial curvature of the data points

can be observed (e.g., Figure 1, 10 °C – 860 chill units), linear rates were strong descriptors of

the deacclimation rates as observed in the fits of our calculated rates of deacclimation and in

the early multiple linear regression models with temperature and chill accumulation as factors

(R^2 > 0.83, p < 0.001 for all cultivars; Table S1), similar to Wolf and Cook (1992), also looking

at deacclimation rates for grapevines – although they only used 1 temperature. We also

stopped measuring LTEs when those were close to -5 °C, where we had difficulties reading

LTE peaks due to the merge with the much larger high temperature exotherms, and therefore

any curvature in the higher portions of the deacclimation curve would not be observed. Such

curvature is observed in Arora *et al.* (2004) and Kalberer *et al.* (2007), although hardiness loss

stopped at a lower temperature, which was speculated to be due to lack of chill.

Ferguson *et al.* (2011; 2014) used a linear response of the rates of deacclimation in

relation to temperature for 21 *V. vinifera* and two *V. labruscana* Bailey cultivars, with a different

rate and base temperature for endo- and ecodormant buds. In our experiments, deacclimation

occurred in temperatures below those established as the threshold for deacclimation by

Ferguson *et al.* (2014). This suggests that although it may produce a good fit for field data, the

current model may be over-simplifying the response to temperature from a physiological

standpoint. Takeuchi and Kasuga (2017) have also seen deacclimation through electrolyte

leakage of bark and xylem of Japanese white birch [*Betula platyphylla* Sukatchiev var. *japonica*

(Miq.)] at low temperatures, including high below-freezing temperatures (-5 to 0°C). In our

study the temperature dependency of k_{deacc} appears to follow simple enzymatic kinetics,

affected by both the catalytic rate (Arrhenius' law) in lower temperatures, and the denaturation

of enzymes in higher temperatures. This behavior suggests that an active temperature

sensing, as defined by Ruelland and Zachowski (2010), is not necessary for deacclimation.

Although deacclimation is likely controlled by a complex metabolic chain or cycle, the rates of

any given process as a whole are defined by the slowest enzyme – which may change

depending on the temperature (Ruelland and Zachowski, 2010). Additionally, Farquhar and

Richards (1984) proposed that whole plant processes should be analyzed as simple chemical

processes in respect to temperature. Therefore, further comparisons are made in this

discussion with activities of single enzymes and whole metabolic cycles without distinction.

It is notable in the deacclimation process measured in this study that rates are departing

from the exponential increase due to temperature in low range (estimated as ≥ 12 °C) as seen

in the Arrhenius plot (Fig. 3B). This would suggest that some limitation occurs at higher

temperature to the process as a whole: either substrate limitations or enzyme denaturation

(Truhlar and Kohen, 2001). It is important to note, however, that there is no drop in the k_{deacc}

from 22 to 30 °C, but a reduction on the increase of k_{deacc} as compared to from 2 to 10 °C (Fig.

3). This differs from other processes in grapevines, such as root respiration, which has an

exponential response to temperature up to 32 °C for short-term exposure (Huang *et al.*, 2005).

This is likely an adaptation of the enzymes involved in the process of deacclimation,

considering low temperatures in early spring when deacclimation occurs. Rigano *et al.* (2006)

demonstrated similar kinetics when comparing the activity of methyl viologen-dependent nitrate

reductase extracts from psychrophilic and mesophilic algae. Arrhenius plots of the

psychrophilic alga had an inflection point at 15 °C, whereas the mesophilic alga had an

inflection point at 25 °C. We made no attempt in our study to find different inflection points for

each species, as we lacked data in the 12-21 °C range. However, future studies could be

conducted to examine if deacclimation rate is adaptive in widely distributed *Vitis* species as

well. The activity of some enzymes, such as catalases and peroxidases, appears to be

correlated with release of dormancy (Farokhzad *et al.*, 2018) and subsequent budbreak (Pérez

and Burgos, 2004). It may be interesting to measure the activity of these enzymes at different

temperatures to evaluate whether they have a similar response to temperature as that

evaluated in the deacclimation here. Other enzymes involved in the synthesis and degradation

of plant hormones might also be involved, considering that growth inhibition by abscisic acid or

promotion by gibberellins and cytokinins appear to interplay in the control of dormancy

(Horvath *et al.*, 2003).

In a review of photosynthesis responses to temperature, Berry and Björkman (1980)

showed that the phase of exponential increase in photosynthesis rate can end at temperatures

as low as 10 °C and as high as 40 °C, and that both the rates and the optimum temperatures

are also affected by the temperature regime in which plants are grown. For deacclimation rates

in grapevine buds, Cragin *et al.* (2017) suggested an effect of the field LTE (or the intercept)

on the deacclimation rate, which would indicate an effect of the environment. We did not

attempt to use field LTE as a factor in our calculations; however, we expect that it would not

greatly impact k_{deacc}: both 860 and 1580 chill units had similar field LTEs, but very different

k_{deacc} (Fig. 1). We believe that most of the environmental effect on the k_{deacc} is well described

by the chill accumulation through the Ψ_{deacc}.

Cultivar or accession differences are seen within species in cold hardiness (Arora *et al.*,

2004; Ferguson *et al.*, 2011, 2014; Salazar-Gutiérrez *et al.*, 2014, 2016; Szalay *et al.*, 2017),

although deacclimation rates may or may not differ (Arora *et al.*, 2004; Ferguson *et al.*, 2011,

2014; Szalay *et al.*, 2017). The effect of genotypes within species in k_{deacc} is suggested by the

better fit of our curves to *V. vinifera* cultivars as compared to the other species, where

parameters were estimated by combining different accessions in the wild material. However,

our dataset did not allow us to properly analyze each genotype of the wild accessions

separately. It has been suggested that there is no correlation among species between the

maximum cold hardiness or climate of origin and the rate of deacclimation (Arora *et al.*, 2004;

Kalberer *et al.*, 2006). Vitasse *et al.* (2014), however, suggests that the rate of deacclimation is

more related to temperature fluctuations in the area where species evolved, diminishing the

potential for deacclimation in locations where frequent temperature fluctuations occur in the

winter. In our case, *V. riparia* and *V. amurensis*, two species from colder climates as compared

to *V. vinifera*, had higher k_{deacc} at all (*V. amurensis*) or at moderate to high temperatures (*V.*

riparia) than *V. vinifera*. *V. aestivalis*, however, did not appear to differ from the *V. vinifera*

cultivars, in agreement with the assessment by Kalberer *et al.* (2006). Kalberer *et al.* (2006)

also suggested that field and controlled studies can result in different responses to

temperature. In our study, however, we observed higher rates of deacclimation in *V. amurensis*

and *V. riparia* compared to *V. aestivalis*, which is comparable and follows the same behavior

as the "responsiveness" described by Londo and Kovaleski (2017) using field observations.

Cannell and Smith (1983) speculate that average daily temperature is a good descriptor

for timing of budbreak, and that daily amplitude and photoperiod may be ignored. Similarly,

Antivilo *et al.* (2017) saw no or little differences in deacclimation due to low and high

temperature amplitudes, respectively. Although this was not tested, it appears that a linear

approximation may be used for temperature effects on deacclimation, especially in small

temperature amplitudes. Therefore, using daily average temperatures may be appropriate for

field data. However, long-term averages could result in larger error in assessments, and

physiological-based studies should use more precise descriptions of temperature effects such

as those reported here.

It is possible that the accuracy of models for prediction of budbreak may be increased if
they are attached to predictions of cold hardiness and deacclimation kinetics. Rubio *et al.*
(2016) reported both LTE and time to 50% budbreak for *V. vinifera* 'Thompson Seedless',
which appear to be positively correlated in the acclimation period. Although the authors did not
report chill accumulation, it is clear that cold hardiness increases (more negative LTEs) and
days to budbreak decreases once daily mean temperatures below 15 °C start consistently
occurring (May – Southern hemisphere). In their case the time to budbreak appears to reach
the limit of endodormancy [50% budbreak within 28 days as per Londo and Johnson (2014)] in
early June in Chile, but continue dropping until their last evaluation in August. This is in
accordance with our assessment that as chill accumulates, there is an increase in Ψ_{deacc},
leading to potentially faster budbreak. Because of this relationship between Ψ_{deacc} (through
chill accumulation) and k_{deacc} leading to budbreak, we hypothesize that studies that identify
earlier or faster budbreak due to the use of hydrogen cyanamide (Pérez and Burgos, 2004;
Pérez and Lira, 2005) would likely find faster rates of deacclimation as well.
Specific studies linking environmental factors to understand plant phenology in order to
evaluate impacts of climate change are needed (Cleland *et al.*, 2007; Chmielewski *et al.*,
2018). In this study, we demonstrate an objective method for determining the dormancy
transition that could prove useful in other perennial species that utilize supercooling for cold
hardiness [e.g. peach, cherry (*Prunus* spp. L.), azalea (*Rhododendron* spp. L.), larch (*Larix*
kaempferi Sarg.)] in order to understand how they might respond to the future shifts in chill
accumulation and increase in average temperature as a consequence of climate change.
Although measurements of cold hardiness may be different for each species, it appears that
this phenotype can aid in understanding dormancy transitions and spring phenology of plants.
It is clear that simple assessments of deacclimation rates simply using "endo-" and
"ecodormant" material is not appropriate, as there is a continuum of response in deacclimation
rates as chill is accumulated. Additionally, although we made no attempt to evaluate different
chill accumulation models, it is possible that using deacclimation rates may be a more
appropriate way of defining chilling temperatures, considering the more quantitative nature of
this analysis.
459
460   **Supplementary data**

Table S1. Information associated with each deacclimation experiment for each genotype used. Linear models were fit per genotype using chill accumulation and temperature as factors.

Acknowledgements

The work reported here was partially supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, through the Northeast Sustainable Agriculture Research and Education program under subaward number GNE16-130; and by CAPES, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil. The authors would like to acknowledge and thank the following people for contributing to the field collection and processing of dormant bud tissues used in this study: Kathleen Deys, Bill Srmack, John Keeton, Bob Martens, and Matheus Baseggio. The authors also thank Dr. Tim Martinson, Cornell University, for use of freezer facilities, and Anthony Road Wine Co. and Ravines Wine Cellars for access to *V. vinifera* samples.

References

Andrews PK, Sandidge III CR, Toyama TK. 1984. Deep supercooling of dormant and deacclimating *Vitis* buds. American Journal of Enology and Viticulture **35,** 175-177.

Antivilo FG, Paz RC, Keller M, Borgo R, Tognetti J, Juñent FR. 2017. Macro- and microclimate conditions may alter grapevine deacclimation: variation in thermal amplitude in two contrasting wine regions from North and South America. International Journal of Biometeorology **61,** 2033-2045.

Arora R, Rowland LJ, Ogden EL, Dhanaraj AL, Marian CO, Ehlenfeldt MK, Vinyard B. 2004. Dehardening kinetics, bud development, and dehydrin metabolism in blueberry cultivars during deacclimation at constant, warm temperatures. Journal of the American Society for Horticultural Sciences **129,** 667-674.

Berry J, Björkman O. 1980. Photosynthetic response and adaptation to temperature in higher plants. Annual Review of Plant Physiology **31,** 491-543.

Bigg EK. 1953. The supercooling of water. Proceedings of the Physical Society Section B **66,** 688-694.

Bokhorst SF, Bjerke JW, Tømmervik H, Callaghan TV, Phoenix GK. 2009. Winter warming events damage sub-Arctic vegetation: consistent evidence from an experimental manipulation and a natural event. Journal of Ecology **97,** 1408-1415.

Burke MJ, Gusta LV, Quamme HA, Weiser CJ, Li PH. 1976. Freezing and injury in plants. Annual Review of Plant Physiology **27,** 507-528.

Cannell MGR, Smith RI. 1983. Thermal time, chill days and prediction of budburst in *Picea sitchensis*. Journal of Applied Ecology **20,** 951-963.

Cannell MGR, Willett SC. 1975. Rates and times at which needles are initiated in buds of different provenances of *Pinus contorta* and *Picea sitchensis* in Scotland. Canadian Journal of Forest Research **5,** 367-380.

Chmielewski F, Götz K, Weber KC, Moryson S. 2018. Climate change and spring frost damages for sweet cherries in Germany. International Journal of Biometeorology **62,** 217-228.

Citadin I, Raseira MCB, Herter FG, Silva JB. 2001. Heat requirement for blooming and leafing in peach. HortScience **36,** 305-307.

Cleland EE, Chuine I, Menzel A, Mooney HA, Schwartz MD. 2007. Shifting plant phenology in response to global change. Trends in Ecology & Evolution **22,** 357-365.

Cook C, Jacobs G. 2000. Progression of apple (*Malus × domestica* Borkh.) bud dormancy in two mild winter climates. The Journal of Horticultural Science and Biotechnology **75,** 233-236.

Couvillon GA, Erez A. 1985. Influence of prolonged exposure to chilling temperatures on bud break and heat requirement for bloom of several fruit species. Journal of the American Society for Horticultural Sciences **110,** 47-50.

Cragin J, Serpe M, Keller M, Shellie K. 2017. Dormancy and cold hardiness transitions in winegrape cultivars Chardonnay and Cabernet Sauvignon. American Journal of Enology and Viticulture **68,** 195-202.

Dantec CF, Vitasse Y, Bonhomme M, Louvet J, Kremer A, Delzon S. 2014. Chilling and heat requirements for leaf unfolding in European beech and sessile oak populations at the southern limit of their distribution range. International Journal of Biometeorology **58,** 1853-1864.

Fan S, Bielenberg DG, Zhebentyayeva TN, Reighard GL, Okie WR, Holland D, Abbott AG. 2010. Mapping quantitative trait loci associated with chilling requirement, heat requirement and bloom date in peach (*Prunus persica*). New Phytologist **185,** 917-930.

Farokhzad A, Nobakht S, Alahveran A, Sarkhosh A, Mohseniazar M. 2018. Biochemical changes in terminal buds of three different walnut (*Juglans regia* L.) genotypes during dormancy break. Biochemical Systematics and Ecology **76,** 52-57.

Farquhar GD, Richards RA. 1984. Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. Functional Plant Biology **11,** 539-552.

Ferguson JC, Tarara JM, Mills LJ, Grove GG, Keller M. 2011. Dynamic thermal time model of cold hardiness for dormant grapevine buds. Annals of Botany **107,** 389-396.

Ferguson JC, Moyer MM, Mills LJ, Hoogenboom G, Keller M. 2014. Modeling dormant bud cold hardiness and budbreak in twenty-three Vitis genotypes reveals variation by region of origin. American Journal of Enology and Viticulture **65,** 59-71.

Gray LK, Hamann A. 2013. Tracking suitable habitat for tree populations under climate change in western North America. Climatic Change **117,** 289-303.

Gu L, Hanson PJ, Post WM, Kaiser DP, Yang B, Nemani R, Pallardy SG, Meyers T. 2008. The 2007 Eastern US spring freeze: increased cold damage in a warming world? BioScience **58,** 253-262.

Horvath DP, Anderson JV, Chao WS, Foley ME. 2003. Knowing when to grow: signals regulating bud dormancy. Trends in Plant Science **8,** 534-540.

Huang X, Lakso AN, Eissenstat DM. 2005. Interactive effects of soil temperature and moisture on Concord grape root respiration. Journal of Experimental Botany **56,** 2651-2660.

Hunter AF, Lechowicz MJ. 1992. Predicting the timing of budburst in temperate trees. Journal of Applied Ecology **29,** 597-604.

Kalberer SR, Wisniewski M, Arora R. 2006. Deacclimation and reacclimation of cold-hardy plants: Current understanding and emerging concepts. Plant Science **171,** 3-16.

Kalberer SR, Leyva-Estrada N, Krebs SL, Arora R. 2007. Frost dehardening and rehardening of floral buds of deciduous azaleas are influenced by genotypic biogeography. Environmental and Experimental Botany **59,** 264-275.

Kolstad EW, Breiteig T, Scaife AA. 2010. The association between stratospheric weak polar vortex events and cold air outbreaks in the Northern Hemisphere. Quarterly Journal of the Royal Meteorological Society **136,** 886-893.

Lang GA, Early JD, Martin GC, Darnell RL. 1987. Endo-, para-, and ecodormancy: physiological terminology and classification for dormancy research. HortScience **22,** 371-377.

Lloyd J, Firth D. 1990. Effect of defoliation time on depth of dormancy and bloom time for low-chill peaches. HortScience **25,** 1575-1578.

Londo JP, Johnson LM. 2014. Variation in the chilling requirement and budburst rate of wild *Vitis* species. Environmental and Experimental Botany **106,** 138-147.

Londo JP, Kovaleski AP. 2017. Characterization of wild North American grapevine cold hardiness using differential thermal analysis. American Journal of Enology and Viticulture **68,** 203-212.

Luedeling E, Girvetz EH, Semenov MA, Brown PH. 2011. Climate change affects winter chill for temperate fruit and nut trees. PLoS ONE https://doi.org/10.1371/journal.pone.0020155

Mills LJ, Ferguson JC, Keller M. 2006. Cold-hardiness evaluation of grapevine buds and cane tissues. American Journal of Enology and Viticulture **57,** 194-200.

Pagter M, Arora R. 2013. Winter survival and deacclimation of perennials under warming climate: physiological perspectives. Physiologia Plantarum **147,** 75-87.

Penfield S. 2008. Temperature perception and signal transduction in plants. New Phytologist **179,** 615-628.

Pérez FJ, Burgos B. 2004. Alterations in the pattern of peroxidase isoenzymes and transient increases in its activity and in H_2O_2 levels take place during the dormancy cycle of grapevine buds: the effect of hydrogen cyanamide. Plant Growth Regulation **43,** 213-220.

Pérez FJ, Lira W. 2005. Possible role of catalase in post-dormancy budbreak in grapevines. Journal of Plant Physiology **162,** 301-308.

Rigano VDM, Vona V, Lobosco O, Carillo P, Lunn JE, Carfagna S, Esposito S, Caiazzo M, Rigano C. 2006. Temperature dependence of nitrate reductase in the psycrophilic unicellular alga *Koliella antarctica* and the mesophilic alga *Chlorella sorokiniana*. Plant, Cell and Environment **29,** 1400-1409.

Rubio S, Dantas D, Bressan-Smith R, Pérez FJ. 2016. Relationship between endodormancy and cold hardiness in grapevine buds. Journal of Plant Growth Regulation **35,** 266-275.

Ruelland E, Zachowski A. 2010. How plants sense temperature. Environmental and Experimental Botany **69,** 225-232.

Salazar-Gutiérrez MR, Chaves B, Anothai J, Whiting M, Hoogenboom G. 2014. Variation in cold hardiness of sweet cherry flower buds through different phenological stages. Scientia Horticulturae **172,** 161-167.

Salazar-Gutiérrez MR, Chaves B, Hoogenboom G. 2016. Freezing tolerance of apple flower buds. Scientia Horticulturae **198,** 344-351.

Shaltout AD, Unrath CR. 1983. Rest completion prediction model for 'Starkrimson Delicious' apples. Journal of the American Society for Horticultural Sciences **108,** 957-961.

Szalay L, Molnár A, Kovács S. 2017. Frost hardiness of flower buds of three plum (*Prunus domestica* L.) cultivars. Scientia Horticulturae **214,** 228-232.

Takeuchi M, Kasuga J. 2017. Bark cells and xylem cells in Japanese white birch twigs initiate deacclimation at different temperatures. Cryobiology https://doi.org/10.1016/j.cryobiol.2017.11.007

Taulavuori KMJ, Tulavuori EB, Skre O, Nilsen J, Igeland B, Laine KM. 2004. Dehardening of mountain birch (*Betula pubescens* ssp. *czerepanovii*) ecotypes at elevated winter temperatures. New Phytologist **162,** 427-436.

Truhlar DG, Kohen A. 2001. Convex Arrhenius plots and their interpretation. Proceedings of the National Academy of Sciences **98,** 848-851.

Vitasse Y, Lenz A, Körner C. 2014. The interaction between freezing tolerance and phenology in temperate deciduous trees. Frontiers in Plant Science doi: 10.3389/fpls.2014.00541

Walsh J, Wuebbles D, Hayhoe K, *et al.* 2014. Chapter 2: Our Changing Climate. In: Melillo JM, Richmond T, Yohe GW, eds. *Climate Change Impacts in the United States: The Third National Climate Assessment* U.S. Global Change Research Program doi: 10.7930/J0KW5CXT

Weinbaum SA, Polito VS, Muraoka TT. 1989. Assessment of rest completion and its relationship to appearance of tetrads in anthers of 'Nonpareil' almond. Scientia Horticulturae **38,** 69-76.

Wolf TK, Cook MK. 1992. Seasonal deacclimation patterns of three grape cultivars at constant, warm temperature. American Journal of Enology and Viticulture **43,** 171-179.

Wolf TK, Cook MK. 1994. Cold hardiness of dormant buds of grape cultivars: comparison of thermal analysis and field survival. HortScience **29,** 1453-1455.

Zhang J, Taylor C. 2011. The dynamic model provides the best description of the chill process on 'Sirora' pistachio trees in Australia. HortScience **46,** 420-425.

Tables

Table 1. Parameter estimates and model fits for deacclimation potential, low and high temperature deacclimation rates, and linear relationship between LTE and the deacclimation rate × deacclimation potential for 7 *Vitis* genotypes.

Figures

Figure 1. Deacclimation of *V. vinifera* 'Riesling' buds at three temperatures (4, 10, and 22 °C) collected from the field at three different chill accumulations (360, 860, 1580). *Buds at 360 chill accumulation were deacclimated at 11 °C.

Figure 2. Deacclimation potential of *V. vinifera* 'Riesling' buds collected at different chill accumulations in two temperatures. Deacclimation rate at 1580 chill units was used as reference (100 %). The relationship between deacclimation potential and accumulated chill was described by the equation: $\Psi_{deacc} (\%) = {}^{1}/{}_{1\ +\ e^{[-7.23\ \times\ (\ln chill\ -6.88)]}}$, with a pseudo-$R^2$ = 0.87.

Figure 3. Effect of temperature on deacclimation rates of *V. vinifera* 'Riesling'. A) Scaled deacclimation rates (based on deacclimation potential) at three different chill accumulations in response to temperature; B) Arrhenius plot of scaled deacclimation rates. Tick marks in both panels are equivalent and mirrored (e.g. 0 °C = 3.66 × 1000 K^{-1}).

Figure 4. Deacclimation rates of *Vitis* buds as a function of temperature. A) Deacclimation rates at >90 % deacclimation potential for three different *Vitis* genotypes: *V. riparia* (closed circle – full line), *V. vinifera* 'Riesling' (open circles – dashed line), and *V. vinifera* 'Cabernet Sauvignon' (closed squares – dotted line); B) Deacclimation rates of *V. vinifera* 'Riesling' at different chill accumulations as a function of temperature: 1580 (closed circle – full line), 1030 (open circle – dashed line), 860 (closed square – dotted line), 360 (open square – dashdotted line).

Table 1

Species	Cultivar	Deacclimation potential[a]			
		b	e	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	-9.47	6.79	Relative	0.93
	'Cab. Sauvignon'	-8.27	6.80	Relative	0.65
	'Riesling'	-7.23	6.88	Relative	0.87
	'Sauvignon blanc'	-43.66	6.77	Relative	0.93
V. aestivalis		-22.31	6.86	Relative and x	0.97
V. amurensis		-12.94	6.85	Relative	0.79
V. riparia		-9.79	6.82	Relative	0.96
		Low temperature[b]			
		m	n	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	0.0600	0.253	Relative	0.89
	'Cab. Sauvignon'	0.0620	0.218	Relative	0.87
	'Riesling'	0.0675	0.236	Relative	0.92
	'Sauvignon blanc'	0.0542	0.235	Relative and x	0.84
V. aestivalis		0.0965	0.187	Relative and x	0.80
V. amurensis		0.1988	0.211	Relative	0.75
V. riparia		0.0754	0.259	Relative	0.81
		High temperature[c]			
		q	r	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	0.762	7.52	Relative	0.92
	'Cab. Sauvignon'	0.593	7.73	Relative	0.84
	'Riesling'	0.732	7.39	Relative	0.88
	'Sauvignon blanc'	0.513	6.66	Relative	0.83
V. aestivalis		0.830	7.91	Relative and x	0.80
V. amurensis		0.913	3.34	Relative	0.80
V. riparia		0.977	7.32	Relative	0.84
		Linear model[d]			
		β	R^2	N	n (%)
V. vinifera	'Cab. Franc'	0.962	0.88	853	0.96
	'Cab. Sauvignon'	0.997	0.84	1160	0.93
	'Riesling'	1.018	0.90	1260	0.93
	'Sauvignon blanc'	1.002	0.86	745	0.90
V. aestivalis		0.963	0.80	700	0.92
V. amurensis		0.828	0.83	730	0.81
V. riparia		0.987	0.82	1499	0.93

[a] $\Psi_{deacc}\,(\%) = \dfrac{1}{1 + e^{[b \times (\ln Chill - c)]}}$, where *Chill* = chill accumulation

[b] $k_{\mathrm{deacc}} = m \times e^{(n \times T)}$, where *T* = temperature

[c] $k_{\mathrm{deacc}} = q \times \ln(T - r)$, where *T* = temperature

[d] $\Delta_{LTE}(T, Chill) = \beta \times k_{\mathrm{deacc}} \times \Psi_{deacc} \times t$, where *t* = time