1. Title page

(i) Title and running title

Kinetics of winter deacclimation in response to temperature determines dormancy

status and explains budbreak in different *Vitis* species

Deacclimation kinetics resolve dormancy status

(ii) Authors

Alisson P. Kovaleski[1], Bruce I. Reisch[1], and Jason P. Londo[1,2*]

(iii) Author affiliations

[1] School of Integrative Plant Science – Horticulture Section, Cornell University-New

York State Agricultural Experiment Station, 630 W. North Street, Geneva, New York,

USA.

[2] United States Department of Agriculture, Agricultural Research Service, Grape

Genetics Research Unit, 630 W. North Street, Geneva, New York, USA.

[*] Corresponding authors

JPL – e-mail: jason.londo@ars.usda.gov; phone: (315) 787-2463

APK – e-mail: apkovaleski@gmail.com

2. Abstract

Bud dormancy and cold hardiness are critical adaptations for surviving winter cold stress for temperate perennial plant species, with shifting temperature-based responses during the winter. The objective of this study was to uncover the relationship between dormancy transition (chilling requirement) and temperature on the loss of cold hardiness and budbreak. Dormant cuttings of *Vitis vinifera*, *V. aestivalis*, *V. amurensis*, and *V. riparia* were examined to determine the relationship between chilling requirement and temperature on rate of deacclimation (k_{deacc}). Differential thermal analysis was used to determine k_{deacc} using mean low temperature exotherms. Effect of chill was evaluated as the deacclimation potential (Ψ_{deacc}), which was the change in k_{deacc} due to chill accumulation. Budbreak was also evaluated in fully chilled buds at different temperatures. Results indicate that Ψ_{deacc} varies dependent on dormancy state, following a logarithmic response to chill accumulation. The effect of temperature on k_{deacc} was exponential at low and logarithmic at high temperatures. The combination of Ψ_{deacc} and k_{deacc} resulted in good prediction of deacclimation. Budbreak phenology was also explained by differences in k_{deacc}. Deacclimation rates can be used as a quantitative determinant of dormancy transition and budbreak, and to refine models predicting effects of climate change.

3. Keyword index

Bud, cold hardiness, deacclimation, dormancy, grapevines, rates, supercooling, *Vitis*.

4. Introduction

Due to their general stationary habit, plants have evolved many different coping mechanisms to survive stressful environments such as drought and temperature. With regard to winter in temperate climates, annual plants typically survive as dehydrated seeds while perennial plants develop dormancy and cold hardiness. Dormancy is the temporary cessation of visible growth in meristem containing structures, such as buds, and is divided in three types: para-, endo-, and ecodormancy (Lang, Early, Martin, & Darnell, 1987). Paradormancy is the suspension due to physiological factors within the plant but outside of the dormant structure, such as the suppression of lateral growth by the apical meristem (i.e. apical dominance). Paradormant buds transition into endodormancy as daylength and temperatures decrease, a state suspension of growth due to unknown endogenous factors within the dormant structure, preventing growth during times when environmental conditions fluctuate between conducive and inhibitory (Horvath, Anderson, Chao, & Foley, 2003). During endodormancy, it is generally understood that exposure to low, non-freezing temperatures is necessary (chilling requirement) to transition to the third dormancy state, ecodormancy. Tissues maintain a dormant state during ecodormancy due to unsuitable environmental conditions.

Concomitantly with the onset of dormancy, plants from areas where temperatures drop below freezing develop cold hardiness through different mechanisms. Some plants, or plant tissues, tolerate intracellular ice formation, in which cells are extremely dehydrated by the formation of extracellular ice, and the remaining water is bound to proteins (Burke, Gusta, Quamme, Weiser, & Li, 1976) likely solidifying in a glass-like state. Other plants present a freezing resistance mechanism, where

water supercools in the intracellular spaces (Burke et al., 1976). Supercooling is a

process through which water can remain in liquid state below its equilibrium freezing

point, down to a maximum of ~ -40 °C (Bigg, 1953). These plants are typical of mid-

latitude temperate climates, where temperatures do not drop below -40 °C during the

winter. In buds that present supercooling ability, such as those of grapevines (*Vitis*

spp.), cold hardiness may be measured through differential thermal analysis (DTA). This

method uses thermoelectric modules that record voltage changes associated with the

release of heat due to phase change of water into ice (Mills, Ferguson, & Keller, 2006).

In grapevine buds, two peaks can be identified using DTA: a high temperature exotherm

(HTE), typically around – 5 °C; and a low temperature exotherm (LTE), which varies

depending on the conditions experienced by the bud. The HTE represents the freezing

of extracellular water (Andrews, Sandidge, & Toyama, 1984), which is normal to the

process of resisting cold. The LTE, however, is the freezing of intracellular water, and it

happens when the supercooling point has been exceeded (Mills et al., 2006). The LTE

represents the lethal temperature for a given bud, is correlated with manual

assessments of bud death (Wolf & Cook, 1994), and mean LTE has been adopted as a

consistent measurement of bud cold hardiness.

A general U-shaped pattern of changing cold hardiness is followed during the

winter through three different stages: acclimation, maintenance, and deacclimation. This

pattern is thought to be primarily driven by air temperature (Ferguson, Tarara, Mills,

Grove & Keller, 2011; Ferguson, Moyer, Mills, Hoogenboom, & Keller**,** 2014; Londo &

Kovaleski, 2017), although other climatic aspects may also be important (Antivilo et al.,

2017). Acclimation appears to occur primarily during endodormancy, while

deacclimation is enhanced during ecodormancy (Ferguson et al., 2011, 2014; Londo &

Kovaleski, 2017). Depending on the climate and genotype, the transition between endo-

and ecodormancy may occur toward the end of the acclimation stage, or during the

maintenance stage of cold hardiness. This transition is typically evaluated by collecting

dormant buds from the field and placing them in growth permissive conditions, followed

by monitoring the time needed to reach budbreak (Weinbaum, Polito, & Muraoka, 1989;

Lloyd & Firth, 1990; Cook & Jacobs, 2000; Fan et al., 2010; Zhang & Taylor, 2011). The

chilling requirement has been met when 50% budbreak occurs within a time threshold

{e.g. 3 weeks for peaches [*Prunus persica* (L.) Batsch; Lloyd & Firth, 1990]; 4 weeks for

grapevines (Londo & Johnson, 2014)}. This method represents a subjective threshold

for the change in endo- to ecodormancy transition. As a result, determining the

molecular and metabolic cues and the consequences of this transition, such as

budbreak, are very difficult to determine and poorly understood (Penfield, 2008).

As the dynamics of acclimation and deacclimation change with dormancy state

(Ferguson et al., 2011, 2014), and dormancy state is determined by accumulation of

chilling hours, a future climate may create issues for both cold hardiness and budbreak

phenology. Average global temperatures are increasing (Walsh et al., 2014), although

episodes of acute cold weather in the northern hemisphere are likely to increase as well

through more frequent arctic oscillations (Kolstad, Breiteig, & Scaife, 2010). In addition

to direct effects to cold hardiness, increasing temperatures will enhance chill

accumulation in higher latitudes, while lower latitudes will experience a decrease

(Luedeling, Girvetz, Semenov, & Brown, 2011). Changes in chill accumulation will

modify the dynamics of dormancy transitions both in deciduous tree crops and other

woody perennials, causing excessive responsiveness to warm spells or erratic

budbreak and growth. This will ultimately require a shift towards higher latitudes and

elevations of tree crops and assisted migration of forest species as they lag behind their

optimal climate niche (Gray & Hamann, 2013). Therefore, it is important to understand

how both temperature and dormancy status affect the loss of cold hardiness. The

objective of this study was to better predict and understand the relationship between

chill accumulation, the dormancy transition, and the loss of cold hardiness in wild and

cultivated grapevines.

5. Materials and methods

Buds of four different species were collected multiple times during the dormant

seasons of 2014/15, 2015/16, and 2016/17. *V. vinifera* L. 'Cabernet Franc', 'Cabernet

Sauvignon', 'Riesling', and 'Sauvignon blanc' were collected from local vineyards

(42.705N, 76.973W; and 42.845N, 77.004W), while *V. aestivalis* Michx. (two clones:

PI483138, PI483143), *V. amurensis* Rupr. (three clones: PI588632, PI588635,

PI588641), and *V. riparia* Michx. (four clones: PI588275, PI588562, PI588653,

PI588711) were collected from the USDA Plant Genetic Resources Unit in Geneva,

New York. For *V. vinifera*, buds from nodes 3 to 20 from the base of a cane were used,

while for the other species, due to constraints regarding the number of clones, buds

beyond position 20 were used as in Londo & Kovaleski (2017).

Hourly weather data from the closest Network for Environment and Weather

Applications station (NEWA; http://www.newa.cornell.edu/) was used to compute chill

accumulation using the "North Carolina" model (Shaltout & Unrath, 1983). The start date

for each season was chosen based on when chill started to consistently accumulate

instead of being negated, as determined by the NC model. These dates were 11 Sept

2014, 19 Sept 2015, and 24 Sept 2016.

Upon collection, canes were prepared into single- or two-node cuttings and

placed submerging the basipetal cut surface in cups of water. The cups were placed

into growth chambers at 2, 4, 7, 8, 10, 11, 22, or 30 °C. Not every genotype or

temperature was used at all collection points, but this information is available in

Supplementary Table S1. Differential thermal analysis (DTA) was performed to

determine the cold hardiness of individual buds as their low temperature exotherm [LTE;

see Mills et al. (2006) for details]. Briefly, buds are excised from the cane and placed on

a thermoelectric module in a plate. These plates are placed in a programmable freezer

with a cooling rate of -4 °C hour^{-1}. Changes in voltage due to heat release in the

freezing of water are measured by the thermoelectric module and recorded using a

Keithley data logger (Tektronix, Beaverton, OR) attached to a computer. Between 4 and

8 buds were used at any time point to determine mean LTE, and the intervals between

measurement days varied between chill accumulation level and temperature treatment,

with details provided in Supplemental Table 1. Because lower rates of deacclimation

were expected in lower temperatures and lower chill accumulations, lower temperature

treatments were typically surveyed with wider separation between time points compared

with high temperature treatments (Table S1).

In addition, an experiment was designed to compare temperature effects on fully

chilled grapevines (1440 chill accumulation). Buds held at constant temperatures (2, 4,

7, 11, 22 °C) had their budbreak recorded following the modified E-L scale (Dry &

Coombe, 2004) for all *V. vinifera* and *V. riparia*. For this, 5 buds were randomly selected

and E-L number recorded in the same sampling interval as for LTE (Table S1) until

buds were past stage 3 or bud material was exhausted.

Statistics

Based on sample size across temperatures and across sampled years, data sets

for *V. vinifera* were analyzed separately for each cultivar, while data for accessions of *V.*

aestivalis, *V. amurensis* and *V. riparia* were combined at the species level. Analyses

were separated into the effect of chill accumulation and the effect of temperature on rate

of deacclimation. The datasets used for each analysis are specified in Supplementary

Table S1.

<u>*Effect of Chill Accumulation.*</u> In order to assess the effect of chill accumulation in the

rate of deacclimation (k_{deacc}), or the deacclimation potential (Ψ_{deacc}), individual rates

were calculated using linear regression in R (ver. 3.3.0, R Foundation for Statistical

Computing) for each temperature and chill accumulation as factors. Although every

temperature within a chill accumulation had the same data for day 0 (field collection),

temperatures were still allowed to have different intercepts in order to reduce the effect

of day 0 on k_{deacc}. From this regression model, data points that had a studentized

residual ≥2.5 were considered outliers and removed from the dataset, and the model

was re-fit. The k_{deacc} at each chill accumulation were then transformed to percentage for

4, 7, 10, 11, and 22 °C, standardizing to their k_{deacc} at highest chill accumulation (either

1440 – 7 and 11 °C – or 1580 chill units – 4, 10, and 22 °C). The Ψ_{deacc} was estimated

as a logistic regression. Initial estimation used the drc library, but final estimation used

the nls() function with the port algorithm, following the equation:

$\Psi_{deacc} (\%) = {}^{1}/_{1\ +\ e^{[b\ \times\ (\ln Chill\ -c)]}}$, where *b* and *c* are the estimated parameters, and

Chill is the chill accumulation at the time of sample collection. The parameter *c* is the

inflection point of the logistic and *b* is the slope associated with the logistic regression.

Effect of Temperature. Initially, to study the effect of temperature, k_{deacc} obtained from

multiple linear regression (temperature and chill accumulation as factors;

Supplementary Table S1) in lower chill accumulations were re-scaled by dividing by the

respective Ψ_{deacc} (e.g., because the Ψ_{deacc} was estimated to be at ~30% at 860 chill

units for 'Riesling', the rates obtained from that data set were divided by 0.30). Since the

effect of chill accumulation was estimated to be ~ 0 % for 360 chill units, these were

excluded from the dataset for estimation of temperature effect. Arrhenius plots were

created to evaluate temperature responses and data were then divided into two data

sets, low (2 – 11 °C) and high temperatures (10 – 30 °C), and temperature effects on

deacclimation rates were evaluated separately. The rationale for doing this is discussed

below.

Instead of using the initially calculated rates derived from the above linear

models, the complete data sets were used with the effect of temperature as a

continuous variable. The values for the different chill accumulation points were

normalized to that of full chill by multiplying the Ψ_{deacc} at any given chill accumulation,

effectively removing the effect of chill by mathematically reducing the time for

deacclimation. Lower chill accumulations were thus included in the calculations instead

of using only those where chill requirements were initially assumed to be fulfilled (1440

and 1580 chill accumulations).

For low temperatures, the effect was assumed to be an exponential, and

therefore the rates were calculated as $k_{\text{deacc}} = m \times e^{(n \times T)}$, where m and n are the

parameters estimated, and T is the temperature in °C. For high temperatures, a

logarithmic curve was used: $k_{\text{deacc}} = q \times \ln(T - r)$, where q and r are the parameters

estimated, and T is the temperature in °C. For both low and high temperatures,

parameters were estimated using nls(), with the port algorithm.

Fitness of Effects of Chill and Temperature. The fits of all non-linear curves (effect of

chill accumulation, low and high temperatures) were tested using Effron's pseudo-R^2

with the Rsq function in the soilphysics library, which is calculated using the sum of

squares of ordinary residuals from the models. The final model for loss of cold

hardiness from an initial point is then $\Delta_{LTE}(T, Chill) = k_{\text{deacc}} \times \Psi_{\text{deacc}} \times t$, where the

change in LTE (Δ_{LTE}) is a function of the temperature (T) it was exposed to, how much

chill accumulation (*Chill*) had passed before exposed to deacclimation temperatures,

and the time (t) of exposure. A linear model was used to evaluate the accuracy of the

predictions using the complete data set. For this, instead of temperatures, $k_{\text{deacc}} =$

$\begin{cases} m \times e^{(n \times T)}, T < 12\,°\text{C} \\ q \times \ln(T - r), T \geq 12\,°\text{C} \end{cases}$ was used, and instead of accumulated chill, the Ψ_{deacc} was

used. From this model ($\Delta_{\text{LTE}} = \beta \times k_{\text{deacc}} \times \Psi_{\text{deacc}} \times time$), considering estimations were

appropriate, the β associated with the estimated k_{deacc} and Ψ_{deacc} was expected to be 1.

Budbreak. In order to evaluate whether the same temperature effects governing

deacclimation and loss of cold hardiness (measured as LTE) also controlled differences

in the temporal rate of budbreak, a linear model was used. For this, forward selection

was conducted with a reduced model containing an intercept only, and a full model

containing all possible interactions of growing degree-days [GDD = k_{deacc} × t (day)],

temperature, species, and genotype as explanatory variables for bud growth stage. The

forward selection was corrected based on the Bayesian information criterion (BIC).

6. Results

After removing outliers using multiple linear regression, data sets kept ≥ 90% of

the observations for all genotypes, with the exception of *V. amurensis*, which had 19.5%

of observations removed (Table 1). Linear behavior appears to describe well the effects

of temperature and chill accumulation in the deacclimation of grapevine buds (Figure 1).

A logistic regression appeared to be a good fit for the Ψ_{deacc} (Figure 2), with pseudo-R^2

values greater than 0.87 for all genotypes except for 'Cabernet Sauvignon' and *V.*

amurensis (0.65 and 0.79, respectively; Table 1). The parameter *e* is the natural

logarithm of the inflection point, and there was little difference in this parameter for

these genotypes (e.g., 6.77 to 6.88, for 'Sauvignon blanc' and 'Riesling', respectively,

which is equivalent to 872 to 972 chill units).

Scaled deacclimation rates based on Ψ_{deacc} and corresponding Arrhenius plot

(Figure 3) demonstrated a clear discontinuity between low and high temperatures. A

deceleration of the increase of k_{deacc} as temperature increased was observed, which

justifies the different behaviors used (exponential and logarithmic for low and high

temperatures, respectively). The estimation of temperature effects for low and high

temperature intersect between 10 and 11 °C because of the uncertainty in the exact

temperature where there is a change in behaviors. This is partially due to the lack of

data points between 11 and 22 °C, although at 10 and 11 °C the calculated rates

through either model would be very similar. All the *V. vinifera* cultivars had similar

deacclimation rates at low temperatures, but 'Riesling' and 'Cabernet Franc' had higher

deacclimation rates at high temperatures than 'Cabernet Sauvignon' and 'Sauvignon

blanc' (Table 1, Figure 4). *V. riparia* and *V. aestivalis* had similar rates to those of *V.*

vinifera at low temperatures, but *V. riparia* had estimated rates similar to *V. amurensis*

in higher temperatures. *V. aestivalis* was similar to the faster *V. vinifera* ('Cabernet

Franc' and 'Riesling'). *V. amurensis* showed the highest deacclimation rates at both low

and high temperatures.

Linear models showed a good fit for all genotypes ($R^2 \geq 0.80$). *V. vinifera*

cultivars, which were analyzed separately had higher R^2 values. The better fit for V.

vinifera cultivars is also seen in their β coefficients, which are very close to 1, although

V. aestivalis and *V. riparia* had βs of 0.99 and 0.96 respectively. *V. amurensis* had the

lowest β, at 0.83.

The final model in the multiple linear regression for budbreak stage had a

P<0.001 and adjusted-R^2 = 0.66 using an intercept and GDD × species interaction

($Budbreak\ stage\ (species) = a + b \times GDD$). The effect of k_{deacc}, i.e. the effect of

temperature, on budbreak can be seen in figure 5. When LTE and budbreak are

regarded in respect to time, a clear separation was observed between temperatures

(Figure 5A, C, E, G). When GDDs are used, accounting for differences in k_{deacc}, LTE

and budbreak stages for all temperatures are "stacked" (Figure 5B, D, F, H). Within the

same temperature, budbreak occurs at different times for different genotypes (Figure

6A), but is grouped within species when using GDDs (Figure 6B).

7. Discussion

Chilling accumulation, dormancy state, cold hardiness and budbreak are complex

traits driven by the interaction of physiological and climactic attributes. The consistency

of winter temperatures is expected to be a major aspect impacting the sustainability and

survival of perennial species in a future climate. This study was conducted to gain a

better method for predicting dormancy transition and chilling requirement in grapevine,

as well as to understand how deacclimation processes (loss of cold hardiness) and

resulting budbreak are impacted by dormancy state.

This paper provides a quantitative assessment of the transition from endo- to

ecodormancy, as evaluated by the Ψ_{deacc}. Unlike the typical dormancy assessment that

uses time to budbreak as a measurement of the transition (Londo & Johnson, 2014),

using the deacclimation rates does not require subjective assumptions currently used to

determine chilling fulfillment; such as having 50% budbreak occur within 28 days.

Budbreak occurs only after the freeze mechanisms associated with cold hardiness are

lost (Figure 5, 6; Ferguson et al., 2014; Salazar-Gutiérrez, Chaves, Anothai, Whiting, &

Hoogenboom, 2014). Thus, the amount of time needed to reach 50% budbreak in

forcing assays is a combination of the time needed to fully deacclimate, and time

required for deacclimated buds to reach a visible change in phenological stage. This

study leverages the ability to assess loss of hardiness through decreasing supercooling

ability as a way to quantitatively assess rates of deacclimation and its relation to

changes in chilling exposure.

Mid-winter warming events are of major concern for temperate crops, yet natural

areas are also affected. In native trees within natural ranges, little to no damage occurs

during the autumn because the freeze resistance readily withstands the minimum

temperatures they usually encounter (Vitasse, Lenz, & Körner, 2014). However,

damage can be substantial in late freeze events during the spring or unusual warm

events in early spring as the potential for deacclimation increases exponentially (Vitasse

et al., 2014). For example, a late spring freeze in 2007 in North America caused

damage in native forests, delaying and reducing canopy establishment by 16-34 days

(Gu et al., 2008). Bokhorst, Bjerke, Tømmervik, Callaghan, & Phoenix (2009) reported

that dwarf shrub vegetation in a sub-Artic region where midwinter temperatures rose to

~7 °C for 2 weeks followed by further cold temperatures resulted in an 87% reduction in

growth during the summer compared to neighboring undamaged areas. Cannell &

Smith (1983) show results from different authors for forest species in which the time to

budbreak is reduced in an exponential manner. This is similar to our assessment of

Ψ_{deacc}, where the upper portion of the logistic is like the exponential decrease in time to

budbreak. The logistic behavior of Ψ_{deacc} in relation to chill accumulation explains the

continuous increase risk of damage of mid-winter warm spells the later they occur until

100% Ψ_{deacc} is achieved. This is in contrast with the model developed by Ferguson et al.

(2011; 2014), where a hard limit for the dormancy boundary is used to model grape cold

hardiness. It is likely that, similar to the behavior described in our study, forest species

also have increased deacclimation rates with higher chill accumulation, seen as a

decrease in the time to budbreak. Additionally, it is important to note that while canopies

may be able to recover from midwinter bud kill events, damage to the reproductive buds

in the case of fruit and nut crops will reduce yields in the season following damage.

Our assessment of Ψ_{deacc} leading to increases in k_{deacc} agrees with the general

observation that warm temperatures in the early winter are negligible for bud growth

(Cannell & Willett, 1975) and that increasing chill accumulation reduces the heat

requirements for budbreak in fruit species (Couvillon & Erez, 1985; Citadin, Raseira,

Herter, & Silva, 2001), and forest species (Hunter & Lechowicz, 1992; Dantec et al.,

2014), considering that loss of hardiness is required for budbreak to occur (Figure 5, 6;

Ferguson et al., 2014). Analyzing bud phenology at approximately full chilling (1440

Chill accumulation ≥ 95% Ψ_{deacc} for all genotypes), we have seen that k_{deacc} is

responsible for differences in early phenological development in the spring between

genotypes within *V. vinifera*, but species still had different phenology with regard to

GDD. Planted in the same environment, Taulavuori et al. (2004) observed that

deacclimation occurs faster in mountain birch [*Betula pubescens* ssp. *czerepanovii*

(Orl.) Hämet-Ahti] ecotypes with origin in areas of low chill accumulation. If the

mechanisms are similar to those described here, what may have been observed in their

study was not different rates of deacclimation, but plants that were at different stages in

their deacclimation potential, or a combination of both. Climate change appears to

speed the loss of cold hardiness and promote earlier leaf-out, although the response is

not linear due to interactions with chilling and photoperiod (Vitasse et al., 2014). In our

case, the predicted increase in chill accumulation (Luedeling et al., 2011) will result in

higher Ψ_{deacc} and higher risk of damage should midwinter warm spells increase in a

future climate (Vitasse et al., 2014; Walsh et al., 2014). However, earlier budbreak may

not lead to higher frequency of damage due to frost events if consistently higher spring

temperatures (Chmielewski, Götz, Weber, & Moryson, 2018) accompany future climate

changes.

Pagter & Arora (2013) suggest that deacclimation rates are not linear, and that a

lag phase may occur during initial exposure to warm temperatures. The initial lag phase

might explain the differences in deacclimation rates measured in Cragin, Serpe, Keller,

& Shellie (2017) compared to those reported in this study, as they only evaluated

deacclimation for 4 days. This may also suggest why they had differences in rates

across the years, although this may also be an effect of different chill accumulation

between the two years in their study. Ferguson et al. (2011; 2014) used asymptotic

bounds added through a logistic component, deviating from a strictly linear rate. In our

data, although in some cases an initial curvature of the data points can be observed

(e.g., Figure 1, 10 °C – 860 chill units), linear rates were strong descriptors of the

deacclimation rates as observed in the fits of our calculated rates of deacclimation and

in the early multiple linear regression models with temperature and chill accumulation as

factors ($R^2 > 0.83$, $p < 0.001$ for all cultivars; Table S1), similar to Wolf & Cook (1992),

also looking at deacclimation rates for grapevines – although they only used 1

temperature. We also stopped measuring LTEs when those were close to -5 °C, where

we had difficulties reading LTE peaks due to the merge with the much larger high

temperature exotherms, and therefore any curvature in the higher portions of the

deacclimation curve would not be observed. Such curvature is observed in Arora et al.

(2004) and Kalberer, Leyva-Estrada, Krebs, & Arora (2007), although hardiness loss

stopped at a lower temperature, which was speculated to be due to lack of chill.

Ferguson et al. (2011; 2014) used a linear response of the rates of deacclimation

in relation to temperature for 21 *V. vinifera* and two *V. labruscana* Bailey cultivars, with a

different rate and base temperature for endo- and ecodormant buds. In our

experiments, deacclimation occurred in temperatures below those established as the

threshold for deacclimation by Ferguson et al. (2014). This suggests that although it

may produce a good fit for field data, the current model may be over-simplifying the

response to temperature from a physiological standpoint. Takeuchi & Kasuga (2017)

have also seen deacclimation through electrolyte leakage of bark and xylem of

Japanese white birch [*Betula platyphylla* Sukatchiev var. *japonica* (Miq.)] at low

temperatures, including high below-freezing temperatures (-5 to 0°C). In our study the

temperature dependency of k_{deacc} appears to follow simple enzymatic kinetics, affected

by both the catalytic rate (Arrhenius' law) in lower temperatures, and the denaturation of

enzymes in higher temperatures. This behavior suggests that an active temperature

sensing, as defined by Ruelland & Zachowski (2010), is not necessary for

deacclimation. Although deacclimation is likely controlled by a complex metabolic chain

or cycle, the rates of any given process as a whole are defined by the slowest enzyme –

which may change depending on the temperature (Ruelland & Zachowski, 2010).

Additionally, Farquhar & Richards (1984) proposed that whole plant processes should

be analyzed as simple chemical processes in respect to temperature. Therefore, further

comparisons are made in this discussion with activities of single enzymes and whole

metabolic cycles without distinction.

It is notable in the deacclimation process measured in this study that rates are

departing from the exponential increase due to temperature in low range (estimated as

≥ 12 °C) as seen in the Arrhenius plot (Fig. 3B). This would suggest that some limitation

occurs at higher temperature to the process as a whole: either substrate limitations or

enzyme denaturation (Truhlar & Kohen, 2001). It is important to note, however, that

there is no drop in the k_{deacc} from 22 to 30 °C, but a reduction on the increase of k_{deacc} as

compared to from 2 to 10 °C (Fig. 3). This differs from other processes in grapevines,

such as root respiration, which has an exponential response to temperature up to 32 °C

for short-term exposure (Huang, Lakso, & Eissenstat, 2005). This is likely an adaptation

of the enzymes involved in the process of deacclimation, considering low temperatures

in early spring when deacclimation occurs. Rigano et al. (2006) demonstrated similar

kinetics when comparing the activity of methyl viologen-dependent nitrate reductase

extracts from psychrophilic and mesophilic algae. Arrhenius plots of the psychrophilic

alga had an inflection point at 15 °C, whereas the mesophilic alga had an inflection point

at 25 °C. We made no attempt in our study to find different inflection points for each

species, as we lacked data in the 12-21 °C range. However, future studies could be

conducted to examine if deacclimation rate is adaptive in widely distributed *Vitis* species

as well. The activity of some enzymes, such as catalases and peroxidases, appears to

be correlated with release of dormancy (Farokhzad, Nobakht, Alahveran, Sarkhosh, &

Mohseniazar, 2018) and subsequent budbreak (Pérez & Burgos, 2004). It may be

interesting to measure the activity of these enzymes at different temperatures to

evaluate whether they have a similar response to temperature as that evaluated in the

deacclimation here. Other enzymes involved in the synthesis and degradation of plant

hormones might also be involved, considering that growth inhibition by abscisic acid or

promotion by gibberellins and cytokinins appear to interplay in the control of dormancy

(Horvath et al., 2003).

In a review of photosynthesis responses to temperature, Berry & Björkman

(1980) showed that the phase of exponential increase in photosynthesis rate can end at

temperatures as low as 10 °C and as high as 40 °C, and that both the rates and the

optimum temperatures are also affected by the temperature regime in which plants are

grown. For deacclimation rates in grapevine buds, Cragin et al. (2017) suggested an

effect of the field LTE (or the intercept) on the deacclimation rate, which would indicate

an effect of the environment. We did not attempt to use field LTE as a factor in our

calculations; however, we expect that it would not greatly impact k_{deacc}: both 860 and

1580 chill units had similar field LTEs, but very different k_{deacc} (Fig. 1). We believe that

most of the environmental effect on the k_{deacc} is well described by the chill accumulation

through the Ψ_{deacc}.

Cultivar or accession differences are seen within species in cold hardiness (Arora

et al., 2004; Ferguson et al., 2011, 2014; Salazar-Gutiérrez et al., 2014; Salazar-

Gutiérrez, Chaves, & Hoogenboom, 2016; Szalay, Molnár, & Kovács, 2017), although

deacclimation rates may or may not differ (Arora et al., 2004; Ferguson et al., 2011,

2014; Szalay et al., 2017). The effect of genotypes within species in k_{deacc} is suggested

by the better fit of our curves to *V. vinifera* cultivars as compared to the other species,

where parameters were estimated by combining different accessions in the wild

material. However, our dataset did not allow us to properly analyze each genotype of

the wild accessions separately. It has been suggested that there is no correlation

among species between the maximum cold hardiness or climate of origin and the rate of

deacclimation (Arora et al., 2004; Kalberer, Wisniewski, & Arora, 2006). Vitasse et al.

(2014), however, suggests that the rate of deacclimation is more related to temperature

fluctuations in the area where species evolved, diminishing the potential for

deacclimation in locations where frequent temperature fluctuations occur in the winter.

In our case, *V. riparia* and *V. amurensis*, two species from colder climates as compared

to *V. vinifera*, had higher k_{deacc} at all (*V. amurensis*) or at moderate to high temperatures

(*V. riparia*) than *V. vinifera*. *V. aestivalis*, however, did not appear to differ from the *V.*

vinifera cultivars, in agreement with the assessment by Kalberer et al. (2006). Kalberer

et al. (2006) also suggested that field and controlled studies can result in different

responses to temperature. In our study, however, we observed higher rates of

deacclimation in *V. amurensis* and *V. riparia* compared to *V. aestivalis*, which is

comparable and follows the same behavior as the "responsiveness" described by Londo

& Kovaleski (2017) using field observations.

Cannell & Smith (1983) speculate that average daily temperature is a good

descriptor for timing of budbreak, and that daily amplitude and photoperiod may be

ignored. Similarly, Antivilo et al. (2017) saw no or little differences in deacclimation due

to low and high temperature amplitudes, respectively. Although this was not tested, it

appears that a linear approximation may be used for temperature effects on

deacclimation, especially in small temperature amplitudes. Therefore, using daily

average temperatures may be appropriate for field data. However, long-term averages

could result in larger error in assessments, and physiological-based studies should use

more precise descriptions of temperature effects such as those reported here.

It is possible that the accuracy of models for prediction of budbreak may be increased if

they are attached to predictions of cold hardiness and deacclimation kinetics. Rubio,

Dantas, Bressan-Smith, & Pérez (2016) reported both LTE and time to 50% budbreak

for *V. vinifera* 'Thompson Seedless', which appear to be positively correlated in the

acclimation period. Although the authors did not report chill accumulation, it is clear that

cold hardiness increases (more negative LTEs) and days to budbreak decreases once

daily mean temperatures below 15 °C start consistently occurring (May – Southern

hemisphere). In their case the time to budbreak appears to reach the limit of

endodormancy [50% budbreak within 28 days as per Londo & Johnson (2014)] in early

June in Chile, but continue dropping until their last evaluation in August. This is in

accordance with our assessment that as chill accumulates, there is an increase in

Ψ_{deacc}, leading to potentially faster budbreak. Because budbreak occurs following loss of

hardiness, and because of the relationship between Ψ_{deacc} (through chill accumulation)

and k_{deacc} leading to budbreak, we hypothesize that studies that identify earlier or faster

budbreak due to the use of hydrogen cyanamide (Pérez & Burgos, 2004; Pérez & Lira,

2005) would likely find faster rates of deacclimation as well.

Bud phenology timing appears to follow the same temperature response as

deacclimation within species. Therefore, it is likely that loss of cold hardiness may

actually be early stages of growth within the bud, and that the differences in bud

development in regard to GDDs (Figure 6B) between species are due to different bud

morphology. However, there have been no studies comparing early morphological

development in buds of different *Vitis* species. In *V. vinifera*, however, Andreini, Viti, &

Scalabrelli (2009) showed a difference in thermal time requirements between different

genotypes grouped in late, intermediate and early budbreak. They used the same

temperature response for all cultivars, however. This is similar to the assessment of

budbreak by Londo & Johson (2014), in which 'Riesling' and 'Cab. Sauvignon' had

different times to budbreak within the same chill accumulation. In our case, it is clear

that if the same effect of temperature was used between cultivars, budbreak would

demand different thermal time accumulations for 'Riesling' (early budbreak) and 'Cab.

Sauvignon' (late budbreak; Figure 6A). However, when we account for the difference in

the efficiency of the use of temperature (e.g., $k_{\text{deacc}_{Riesling}} > k_{\text{deacc}_{Cab.Sauvignon}}$ at any

given temperature), both *V. vinifera* genotypes have the same GDD requirements for

budbreak (Figure 6B). Therefore, forcing assays using budbreak phenology imply less

efficient deacclimators, such as 'Cabernet Sauvignon', require higher chill accumulation.

Our results indicate that high-chill requirement phenotypes may actually be a

combination of higher chill requirement and low temperature efficiency (low k_{deacc}).

Specific studies linking environmental factors to understand plant phenology in

order to evaluate impacts of climate change are needed (Cleland, Chuine, Menzel,

Mooney, & Schwartz, 2007; Chmielewski et al., 2018). In this study, we demonstrate an

objective method for determining the dormancy transition that could prove useful in

other perennial species that utilize supercooling for cold hardiness [e.g. peach, cherry

(*Prunus* spp. L.), azalea (*Rhododendron* spp. L.), larch (*Larix kaempferi* Sarg.)] in order

to understand how they might respond to the future shifts in chill accumulation and

increase in average temperature as a consequence of climate change. Although

measurements of cold hardiness may be different for each species, it appears that this

phenotype can aid in understanding dormancy transitions and spring phenology of

plants, as we have demonstrated that early bud phenology and deacclimation to cold

are linked and therefore results based on budbreak alone are confounded by dynamics

of the loss of hardiness. It is clear that simple assessments of deacclimation rates

simply using "endo-" and "ecodormant" material is not appropriate, as there is a

continuum of response in deacclimation rates as chill is accumulated. Additionally,

although we made no attempt to evaluate different chill accumulation models, it is

possible that using deacclimation rates may be a more appropriate way of defining

chilling temperatures, considering the more quantitative nature of this analysis.

8. Acknowledgements

The work reported here was partially supported by the National Institute of Food and

Agriculture, U.S. Department of Agriculture, through the Northeast Sustainable

Agriculture Research and Education program under subaward number GNE16-130; and

by CAPES, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil.

The authors would like to acknowledge and thank the following people for contributing

to the field collection and processing of dormant bud tissues used in this study:

Kathleen Deys, Bill Srmack, John Keeton, Bob Martens, and Matheus Baseggio. The

authors also thank Dr. Tim Martinson, Cornell University, for use of freezer facilities, and

Anthony Road Wine Co. and Ravines Wine Cellars for access to *V. vinifera* samples.

The authors have no conflict of interest to declare.

9. References

Andrews P.K., Sandidge III C.R. & Toyama T.K. (1984). Deep supercooling of

dormant and deacclimating *Vitis* buds. *American Journal of Enology and*

Viticulture 35, 175-177.

Andreini L., Viti R. & Scalabrelli G. (2009). Study on the morphological evolution

of budbreak in *Vitis vinifera* L. *Vitis* 4, 153-158.

Antivilo F.G., Paz R.C., Keller M., Borgo R., Tognetti J. & Juñent F.R. (2017).

Macro- and microclimate conditions may alter grapevine deacclimation: variation

in thermal amplitude in two contrasting wine regions from North and South

America. *International Journal of Biometeorology* 61, 2033-2045.

Arora R., Rowland L.J., Ogden E.L., Dhanaraj A.L., Marian C.O., Ehlenfeldt M.K. &

Vinyard B. (2004). Dehardening kinetics, bud development, and dehydrin

metabolism in blueberry cultivars during deacclimation at constant, warm

temperatures. *Journal of the American Society for Horticultural Sciences* 129,

667-674.

Berry J. & Björkman O. (1980). Photosynthetic response and adaptation to

temperature in higher plants. *Annual Review of Plant Physiology* 31, 491-543.

Bigg E.K. (1953). The supercooling of water. *Proceedings of the Physical Society*

Section B 66, 688-694.

Bokhorst S.F., Bjerke J.W., Tømmervik H., Callaghan T.V. & Phoenix G.K. (2009).

Winter warming events damage sub-Arctic vegetation: consistent evidence from

an experimental manipulation and a natural event. *Journal of Ecology* 97, 1408-

1415.

Burke M.J., Gusta L.V., Quamme H.A., Weiser C.J. & Li P.H. (1976). Freezing and

injury in plants. *Annual Review of Plant Physiology* 27, 507-528.

Cannell M.G.R. & Smith R.I. (1983). Thermal time, chill days and prediction of

budburst in *Picea sitchensis*. *Journal of Applied Ecology* 20, 951-963.

Cannell M.G.R. & Willett S.C. (1975). Rates and times at which needles are

initiated in buds of different provenances of *Pinus contorta* and *Picea sitchensis*

in Scotland. *Canadian Journal of Forest Research* 5, 367-380.

Chmielewski F., Götz K., Weber K.C. & Moryson S. (2018). Climate change and

spring frost damages for sweet cherries in Germany. *International Journal of*

Biometeorology 62, 217-228.

Citadin I., Raseira M.C.B., Herter F.G. & Silva J.B. (2001). Heat requirement for

blooming and leafing in peach. *HortScience* 36, 305-307.

Cleland E.E., Chuine I., Menzel A., Mooney H.A. & Schwartz M.D. (2007). Shifting

plant phenology in response to global change. *Trends in Ecology & Evolution* 22,

357-365.

Cook C. & Jacobs G. (2000). Progression of apple (*Malus × domestica* Borkh.) bud

dormancy in two mild winter climates. *The Journal of Horticultural Science and*

Biotechnology 75, 233-236.

Couvillon G.A. & Erez A. (1985). Influence of prolonged exposure to chilling

temperatures on bud break and heat requirement for bloom of several fruit

species. *Journal of the American Society for Horticultural Sciences* 110, 47-50.

Cragin J., Serpe M., Keller M. & Shellie K. (2017). Dormancy and cold hardiness

transitions in winegrape cultivars Chardonnay and Cabernet Sauvignon.

American Journal of Enology and Viticulture 68, 195-202.

Dantec C.F., Vitasse Y., Bonhomme M., Louvet J., Kremer A. & Delzon S. (2014).

Chilling and heat requirements for leaf unfolding in European beech and sessile

oak populations at the southern limit of their distribution range. *International*

Journal of Biometeorology 58, 1853-1864.

Dry P. & Coombe B. (2004). Grapevine growth stages-The modified EL system.

Viticulture 1-Resources.

Fan S., Bielenberg D.G., Zhebentyayeva T.N., Reighard G.L., Okie W.R., Holland D.

& Abbott A.G. (2010). Mapping quantitative trait loci associated with chilling

requirement, heat requirement and bloom date in peach (*Prunus persica*). *New*

***Phytologist* 185, 917-930.**

Farokhzad A., Nobakht S., Alahveran A., Sarkhosh A. & Mohseniazar M. (2018).

Biochemical changes in terminal buds of three different walnut (*Juglans regia* L.)

genotypes during dormancy break. *Biochemical Systematics and Ecology* 76, 52-

57.

Farquhar G.D. & Richards R.A. (1984). Isotopic composition of plant carbon

correlates with water-use efficiency of wheat genotypes. *Functional Plant Biology*

11, 539-552.

Ferguson J.C., Tarara J.M., Mills L.J., Grove G.G. & Keller M. (2011). Dynamic

thermal time model of cold hardiness for dormant grapevine buds. *Annals of*

***Botany* 107, 389-396.**

Ferguson J.C., Moyer M.M., Mills L.J., Hoogenboom G. & Keller M. (2014).

Modeling dormant bud cold hardiness and budbreak in twenty-three Vitis

genotypes reveals variation by region of origin. *American Journal of Enology and*

Viticulture 65, 59-71.

Gray L.K. & Hamann A. (2013). Tracking suitable habitat for tree populations

under climate change in western North America. *Climatic Change* 117, 289-303.

Gu L., Hanson P.J., Post W.M., Kaiser D.P., Yang B., Nemani R., …, Meyers T.

(2008). The 2007 Eastern US spring freeze: increased cold damage in a warming

world? *BioScience* 58, 253-262.

Horvath D.P., Anderson J.V., Chao W.S. & Foley M.E. (2003). Knowing when to

grow: signals regulating bud dormancy. *Trends in Plant Science* 8, 534-540.

Huang X., Lakso A.N. & Eissenstat D.M. (2005). Interactive effects of soil

temperature and moisture on Concord grape root respiration. *Journal of*

Experimental Botany 56, 2651-2660.

Hunter A.F. & Lechowicz M.J. (1992). Predicting the timing of budburst in

temperate trees. *Journal of Applied Ecology* 29, 597-604.

639 **Kalberer S.R., Wisniewski M. & Arora R. (2006). Deacclimation and reacclimation**

640 **of cold-hardy plants: Current understanding and emerging concepts.** *Plant*

641 *Science* **171, 3-16.**

643 **Kalberer S.R., Leyva-Estrada N., Krebs S.L. & Arora R. (2007). Frost dehardening**

644 **and rehardening of floral buds of deciduous azaleas are influenced by genotypic**

645 **biogeography.** *Environmental and Experimental Botany* **59, 264-275.**

647 **Kolstad E.W., Breiteig T. & Scaife A.A. (2010). The association between**

648 **stratospheric weak polar vortex events and cold air outbreaks in the Northern**

649 **Hemisphere.** *Quarterly Journal of the Royal Meteorological Society* **136, 886-893.**

651 **Lang G.A., Early J.D., Martin G.C. & Darnell R.L. (1987). Endo-, para-, and**

652 **ecodormancy: physiological terminology and classification for dormancy**

653 **research.** *HortScience* **22, 371-377.**

655 **Lloyd J. & Firth D. (1990). Effect of defoliation time on depth of dormancy and**

656 **bloom time for low-chill peaches.** *HortScience* **25, 1575-1578.**

658 **Londo J.P. & Johnson L.M. (2014). Variation in the chilling requirement and**

659 **budburst rate of wild** *Vitis* **species.** *Environmental and Experimental Botany* **106,**

660 **138-147.**

Londo J.P. & Kovaleski A.P. (2017). Characterization of wild North American

grapevine cold hardiness using differential thermal analysis. *American Journal of*

***Enology and Viticulture* 68, 203-212.**

Luedeling E., Girvetz E.H., Semenov M.A. & Brown P.H. (2011). Climate change

affects winter chill for temperate fruit and nut trees. *PLoS ONE*

https://doi.org/10.1371/journal.pone.0020155

Mills L.J., Ferguson J.C. & Keller M. (2006). Cold-hardiness evaluation of

grapevine buds and cane tissues. *American Journal of Enology and Viticulture*

57, 194-200.

Pagter M. & Arora R. (2013). Winter survival and deacclimation of perennials

under warming climate: physiological perspectives. *Physiologia Plantarum* 147,

75-87.

Penfield S. (2008). Temperature perception and signal transduction in plants. *New*

***Phytologist* 179, 615-628.**

Pérez F.J. & Burgos B. (2004). Alterations in the pattern of peroxidase

isoenzymes and transient increases in its activity and in H_2O_2 levels take place

during the dormancy cycle of grapevine buds: the effect of hydrogen cyanamide.

***Plant Growth Regulation* 43, 213-220.**

Pérez F.J. & Lira W. (2005). Possible role of catalase in post-dormancy budbreak

in grapevines. *Journal of Plant Physiology* 162, 301-308.

Rigano V.D.M., Vona V., Lobosco O., Carillo P., Lunn J.E., Carfagna S., …, Rigano

C. (2006). Temperature dependence of nitrate reductase in the psycrophilic

unicellular alga *Koliella antarctica* and the mesophilic alga *Chlorella sorokiniana*.

***Plant, Cell and Environment* 29, 1400-1409.**

Rubio S., Dantas D., Bressan-Smith R. & Pérez F.J. (2016). Relationship between

endodormancy and cold hardiness in grapevine buds. *Journal of Plant Growth*

***Regulation* 35, 266-275.**

Ruelland E. & Zachowski A. (2010). How plants sense temperature. *Environmental*

***and Experimental Botany* 69, 225-232.**

Salazar-Gutiérrez M.R., Chaves B., Anothai J., Whiting M. & Hoogenboom G.

(2014). Variation in cold hardiness of sweet cherry flower buds through different

phenological stages. *Scientia Horticulturae* 172, 161-167.

Salazar-Gutiérrez M.R., Chaves B. & Hoogenboom G. (2016). Freezing tolerance of

apple flower buds. *Scientia Horticulturae* 198, 344-351.

Shaltout A.D. & Unrath C.R. (1983). Rest completion prediction model for

'Starkrimson Delicious' apples. *Journal of the American Society for Horticultural*

***Sciences* 108, 957-961.**

Szalay L., Molnár A. & Kovács S. (2017). Frost hardiness of flower buds of three

plum (*Prunus domestica* L.) cultivars. *Scientia Horticulturae* 214, 228-232.

Takeuchi M. & Kasuga J. (2017). Bark cells and xylem cells in Japanese white

birch twigs initiate deacclimation at different temperatures. *Cryobiology*

https://doi.org/10.1016/j.cryobiol.2017.11.007

Taulavuori K.M.J., Taulavuori E.B., Skre O., Nilsen J., Igeland B. & Laine K.M.

(2004). Dehardening of mountain birch (*Betula pubescens* ssp. *czerepanovii*)

ecotypes at elevated winter temperatures. *New Phytologist* 162, 427-436.

Truhlar D.G. & Kohen A. (2001). Convex Arrhenius plots and their interpretation.

***Proceedings of the National Academy of Sciences* 98, 848-851.**

Vitasse Y., Lenz A. & Körner C. (2014). The interaction between freezing tolerance

and phenology in temperate deciduous trees. *Frontiers in Plant Science*

doi: 10.3389/fpls.2014.00541

Walsh J., Wuebbles D., Hayhoe K., Kossin J., Kunken K., Stephens G., …,

Somerville R. (2014). Chapter 2: Our Changing Climate. In *Climate Change*

***Impacts in the United States: The Third National Climate Assessment* (eds J.M.**

Melillo, T. Richmond & G.W. Yohe), pp. 19-67. U.S. Global Change Research

Program doi: 10.7930/J0KW5CXT

Weinbaum S.A., Polito V.S. & Muraoka T.T. (1989). Assessment of rest completion

and its relationship to appearance of tetrads in anthers of 'Nonpareil' almond.

***Scientia Horticulturae* 38, 69-76.**

Wolf T.K. & Cook M.K. (1992). Seasonal deacclimation patterns of three grape

cultivars at constant, warm temperature. *American Journal of Enology and*

***Viticulture* 43, 171-179.**

Wolf T.K. & Cook M.K. (1994). Cold hardiness of dormant buds of grape cultivars:

comparison of thermal analysis and field survival. *HortScience* 29, 1453-1455.

Zhang J. & Taylor C. (2011). The dynamic model provides the best description of

the chill process on 'Sirora' pistachio trees in Australia. *HortScience* 46, 420-425.

10. Tables

Table 1. Parameter estimates and model fits for deacclimation potential, low and high

temperature deacclimation rates, and linear relationship between LTE and the

deacclimation rate × deacclimation potential for 7 *Vitis* genotypes.

Species	Cultivar	Deacclimation potential[a]			
		b	e	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	-9.47	6.79	Relative	0.93
	'Cab. Sauvignon'	-8.27	6.80	Relative	0.65
	'Riesling'	-7.23	6.88	Relative	0.87
	'Sauvignon blanc'	-43.66	6.77	Relative	0.93
V. aestivalis		-22.31	6.86	Relative and X	0.97
V. amurensis		-12.94	6.85	Relative	0.79
V. riparia		-9.79	6.82	Relative	0.96
		Low temperature[b]			
		m	n	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	0.0600	0.253	Relative	0.89
	'Cab. Sauvignon'	0.0620	0.218	Relative	0.87
	'Riesling'	0.0675	0.236	Relative	0.92
	'Sauvignon blanc'	0.0542	0.235	Relative and X	0.84
V. aestivalis		0.0965	0.187	Relative and X	0.80
V. amurensis		0.1988	0.211	Relative	0.75
V. riparia		0.0754	0.259	Relative	0.81
		High temperature[c]			
		q	r	Convergence	Pseudo-R^2
V. vinifera	'Cab. Franc'	0.762	7.52	Relative	0.92
	'Cab. Sauvignon'	0.593	7.73	Relative	0.84
	'Riesling'	0.732	7.39	Relative	0.88
	'Sauvignon blanc'	0.513	6.66	Relative	0.83
V. aestivalis		0.830	7.91	Relative and X	0.80
V. amurensis		0.913	3.34	Relative	0.80
V. riparia		0.977	7.32	Relative	0.84
		Linear model[d]			
		β	R^2	N	n (%)
V. vinifera	'Cab. Franc'	0.962	0.88	853	0.96
	'Cab. Sauvignon'	0.997	0.84	1160	0.93
	'Riesling'	1.018	0.90	1260	0.93

'Sauvignon blanc'	1.002	0.86	745	0.90
V. aestivalis	0.963	0.80	700	0.92
V. amurensis	0.828	0.83	730	0.81
V. riparia	0.987	0.82	1499	0.93

754   [a] $\Psi_{deacc}\ (\%) = 1/{1 + e^{[b \times (\ln Chill - c)]}}$, where *Chill* = chill accumulation

755   [b] $k_{\text{deacc}} = m \times e^{(n \times T)}$, where *T* = temperature

756   [c] $k_{\text{deacc}} = q \times \ln(T - r)$, where *T* = temperature

757   [d] $\Delta_{LTE}(T, Chill) = \beta \times k_{\text{deacc}} \times \Psi_{\text{deacc}} \times t$, where *t* = time

 11. Figures legends

Figure 1. Deacclimation of *V. vinifera* 'Riesling' buds at three temperatures (4, 10, and

22 °C) collected from the field at three different chill accumulations (360, 860, 1580).

*Buds at 360 chill accumulation were deacclimated at 11 °C.

Figure 2. Deacclimation potential of *V. vinifera* 'Riesling' buds collected at different chill

accumulations in two temperatures. Deacclimation rate at 1580 chill units was used as

reference (100 %). The relationship between deacclimation potential and accumulated

chill was described by the equation: $\Psi_{deacc}\ (\%) = {}^{1}\!/\!{}_{1\ +\ e^{[-7.23\ \times\ (\ln Chill\ -6.88)]}}$, with a

pseudo-R^2 = 0.87.

Figure 3. Effect of temperature on deacclimation rates of *V. vinifera* 'Riesling'. A) Scaled

deacclimation rates (based on deacclimation potential) at three different chill

accumulations in response to temperature; B) Arrhenius plot of scaled deacclimation

rates. Tick marks in both panels are equivalent and mirrored (e.g. 0 °C = 3.66 × 1000 K⁻

¹).

Figure 4. Deacclimation rates of *Vitis* buds as a function of temperature. A)

Deacclimation rates at >90 % deacclimation potential for three different *Vitis* genotypes:

V. riparia (closed circle – full line), *V. vinifera* 'Riesling' (open circles – dashed line), and

V. vinifera 'Cabernet Sauvignon' (closed squares – dotted line); B) Deacclimation rates

of *V. vinifera* 'Riesling' at different chill accumulations as a function of temperature:

1580 (closed circle – full line), 1030 (open circle – dashed line), 860 (closed square –

dotted line), 360 (open square – dashdotted line).

Figure 5. Deacclimation and bud phenology of *V. riparia* and *V. vinifera* 'Riesling' in

relation to time (A, C, E, G) and thermal time in GDD (B, D, F, H). Bud phenology as the

"E-L number" (Dry and Coombe, 2004) is shown for 5 temperatures in *V. riparia* and 3

temperatures in 'Riesling', as no bud development was recorded within 90 days at 2 and

4 °C for 'Riesling'. GDDs were calculated based on k_{deacc} for each genotype (see Table

1) at any given temperature multiplied by time in days.

Figure 6. Deacclimation and bud phenology of *V. riparia* (closed circle – full line), *V.*

vinifera 'Riesling' (open circle – dashed line), and *V. vinifera* 'Cabernet Sauvignon'

(closed square – dotted line) at 11 °C in relation to time (A, C) and thermal time in GDD

(B, D).

Supporting information

Table S1. Information associated with each deacclimation experiment for each

genotype used. Linear models were fit per genotype using chill accumulation and

temperature as factors.

A