1

Escherichia coli β-clamp slows down DNA polymerase I dependent nick translation while

accelerating ligation

Amit Bhardwaj[1], Debarghya Ghose[1], Krishan Gopal Thakur[1] and Dipak Dutta[1*]

CSIR-Institute of Microbial Technology, Chandigarh, India[1]

*Corresponding author

E-mail: dutta@imtech.res.in (DD)

Abstract

The nick translation property of DNA polymerase I (Pol I) ensures the maturation of Okazaki fragments by removing primer RNAs and facilitating ligation. However, prolonged nick translation traversing downstream DNA is an energy wasting futile process, as Pol I simultaneously polymerizes and depolymerizes at the nick sites utilizing energy-rich dNTPs. Using an *in vitro* assay system, we demonstrate that the β-clamp of the *Escherichia coli* replisome strongly inhibits nick translation on the DNA substrate. To do so, β-clamp inhibits the strand displacement activity of Pol I by interfering with the interaction between the finger subdomain of Pol I and the downstream primer-template junction. Conversely, β-clamp stimulates the 5' exonuclease property of Pol I to cleave single nucleotides or shorter oligonucleotide flaps. This single nucleotide flap removal at high frequency increases the probability of ligation between the upstream and downstream DNA strands at an early phase, terminating nick translation. Besides β-clamp-mediated ligation helps DNA ligase to seal the nick promptly during the maturation of Okazaki fragments.

Introduction

In replicating *E. coli* cells, DNA polymerase III (Pol III) holoenzyme is the major replicative DNA polymerase [1]. Three Pol III core complexes (each containing $\alpha\epsilon\theta$ polypeptides) interact with the clamp loader complex ($\gamma_1\tau_2\delta\delta'\chi\psi$ polypeptides) and β-sliding clamps at the primer-template junctions of a replication fork [2-8]. The dimeric β-clamp is a doughnut-shaped molecule that accommodates double-stranded (ds) DNA at the central hole [2]. When β is free in solution, ATP-bound form of clamp loader complex drives β-clamp opening and binding to the single-stranded (ss) DNA region of a primer-template junction [9-

12]. Such a clamp loader interaction with the DNA triggers ATP hydrolysis and results in β-clamp loading on the primer-template junction. Subsequently, the clamp loader affinity to the β-clamp and DNA decreases rapidly that allows clamp loader to dissociate [13,14]. Immediately after being loaded onto the DNA, the sliding clamp interacts with Pol III core at the replication fork [15-17]. The β-clamp also has affinity for all other DNA polymerases in *E. coli* [18-25]. These interactions dramatically increase the processivity of the DNA polymerases [18-25]. Furthermore, interactions of β-clamp with DNA ligase and MutS repair proteins have also been reported [19]. Therefore, apart from acting as a processivity factor for the replicating DNA polymerases, the β-clamp may have some other putative roles in the post-replication DNA repair processes.

Synthesis of the leading strand DNA presumably requires a few priming events [26-30], while frequent priming events at the lagging strand generate discrete Okazaki fragments of 1-2kb in length [31,32]. Maturation of the Okazaki fragments eventually produces an intact lagging strand primarily by the nick translating DNA polymerase I (Pol I) and ligase functions. Pol I is a multifunctional protein with a large and a small domains connected by a flexible linker [33,34]. During nick translation, the large domain of Pol I (also known as Klenow fragment) polymerizes from the 3' end at the nick and simultaneously displaces 5' end nucleotides from the template to create variable lengths of oligonucleotide flaps at the nick site [33-37]. Concomitantly, the small domain of Pol I excises the flap by its flap endonuclease activity [33-37]. The strand displacement and flap endonuclease activities are together responsible for the 5' to 3' exonuclease (hereinafter 5' exonuclease) function of Pol I. The Klenow fragment of Pol I also exhibits 3'to 5' exonuclease (hereinafter 3' exonuclease) function that is responsible for the proofreading activity [33].

After synthesizing an Okazaki fragment, the replisome dissociates from the β-clamp. As a result, the "left over" β-clamp remains attached at the 3'-side of the ssDNA gap or nick

[38-41]. Consistently, β-clamp has been shown to dynamically accumulate behind

progressing replication fork in *E. coli* and *Bacillus subtilis* [42,43]. Presumably, β-clamp

subsequently helps in repairing the gap or nick by interacting with Pol I, and DNA ligase

[19,44,45]. Although a growing body of *in vitro* studies in the past have been done to

illuminate the different functions of Pol I, as described above [33-37,46], none of them

addressed the dynamics and biochemistry of these functions in the presence of β-clamp.

Here, we designed *in vitro* experiments to assess the influence of β-clamp on the functions of

Pol I and DNA ligase in the nick translation process. Our approach unraveled a hitherto

unknown mechanism of clamp-assisted DNA gap repair in *E. coli*.

Results

β-clamp inhibits Pol I-mediated nick translation

To probe the effects of β-clamp on nick translation process, we assembled a template

annealing three different oligonucleotides (67, 19 and 28 bases long), as described in

Materials and Methods section. Thus, as depicted in the Fig 1A, we created a template that

has a primed template junction with a 3' recessed end of 19 bases long primer and 20

nucleotides single-stranded gap, which suitably acts as a "placeholder" to loaded-β blocking

its free sliding [12]. The efficiency and accuracy of primer annealing were almost 100%, as

judged by native PAGE (S1 Fig). We loaded β-clamp on the template with the help of clamp

loader and ATP, and initiated the polymerization reaction by adding Pol I and dNTPs. The

pattern of radioactive oligonucleotide bands in the denaturing gel revealed that Pol I filled the

20 nucleotides DNA gap by extending the radiolabeled 19 bases upstream oligonucleotide

and encountered the 28 bases downstream RNA oligonucleotide to initiate nick translation

(Fig 1B). Interestingly, β-clamp slightly stimulated pausing of the nick translating Pol I

scattered between 39th to 67th positions while traversing 28 bases downstream RNA

oligonucleotide (compare the dotted lines in lane 2 and lane 6 of Fig 1B), indicating that β-

clamp could moderately reduce the speed of nick translation. On calculating the rate of nick

translation, we show that the effect of β-clamp on the Pol I dependent nick translation

traversing RNA substrate was nominal (S2 Fig). However, when we used the template with

the 28 bases downstream DNA oligonucleotide, the β-clamp dramatically increased the pause

of nick translating Pol I, as judged by the bulk migration of the nick sites at 30 and 60

seconds time span (Fig 1C; zones II and I).

Fig 1. β-clamp inhibits Pol I-mediated nick translation. (A) The template used in the assay

was prepared by assembling 67 bases, 5'-phosphorylated 28 bases and 5'-radiolabelled

(asterisk) 19 bases oligonucleotides. (**B)** A representative urea denaturing PAGE and the rate

calculation therein (see also S2 Fig) indicate that the presence of β-clamp nominally affected

nick translation traversing 28 bases downstream RNA. (**C)** The urea-denaturing gel shows

that Pol I efficiently translated the nick, degrading 28 bases downstream DNA, but the

presence of β-clamp dramatically reduced the speed of nick translation. The presence of

ligase allowed early ligation when β-clamp slowed down Pol I-mediated nick translation. (**D)**

The urea PAGE represents that the 3'-exo⁻ Pol I-mediated nick translation was slowed down

in the presence of template-loaded β-clamp, in a manner similar to the nick translation with

β-clamp and Pol I combinations, as shown in panel C. Moreover, the presence of template-

loaded β-clamp increases the ligation efficiency during the 3'-exo⁻ Pol I-mediated nick

translation. (**E)** The urea denaturing gel represents that the 5'-exo⁻ Pol I intensely paused at

the 39th nucleotide and downstream positions, affecting the ligation process. The presence of

template-loaded β-clamp moderately enhances these pauses, but facilitates ligation to some

extent. The # indicates the position of truncated by-products that originated from the 67-

nucleotide long product by the 3' exonuclease function of Pol I and 5'-exo⁻ Pol I (panel B, C, E). This truncated product is missing in the assay with 3'-exo⁻ Pol I (panel D). All the experiments were performed at least three times. The values represent mean ± standard deviation (sd).

We conducted the control experiments to show that the clamp loader could efficiently load β-clamp on the above-mentioned specific template types (see the detail in S1 text). For this, we used biotinylated templates or histidine-tag version of β-clamp to show that after a clamp loading reaction, the template or the β-clamp could pull-down β-clamp or template, respectively, with 100% occupancy, while clamp loader proteins were washed away (S3 Fig). This observation suggests that the clamp loading was efficient that allows a stable β-clamp interaction with the DNA containing the primer-template junction. Further, to show that the observed effect β-clamp on nick translation is not an artifact arisen by an interference of clamp loader proteins, we performed clamp-loading reaction and washed the streptavidin-bound biotinylated template to remove the clamp loader proteins. Next, we added Pol I and dNTPs to show that the template-bound β could inhibit nick translation similar to the unwashed control reaction containing clamp loader proteins (S4Fig). This data suggests that the template-loaded β-clamp, but not the free clamp loader proteins, in the reaction mixture interfere with the nick translation by Pol I. These observations are consistent with the original literature, which have demonstrated that β-clamp does not slide freely from a template that has a primer-template junction, and the affinity of clamp loader proteins to β-clamp declines rapidly after latter is loaded on the template [9,47].

We incorporated point mutations in *polA* (the Pol I encoding gene) by site directed mutagenesis to disrupt the 3' and 5' exonuclease activities. The purified 3'-exonuclease-free

(3'-exo⁻) and 5'-exonuclease-free (5'-exo⁻) Pol I were then used to dissect the role of 3' and

5' exonuclease activities of Pol I in the nick translation process. The nick translation

mediated by the 3'-exo⁻ Pol I (Fig 1D) appeared to be faster than the Pol I-mediated nick

translation (compare the intensity of paused bands and 67-nucleotide product between lane 2

of Fig 1C and Fig 1D respectively). To further elaborate this observation, in a separate

experiment, we determined that the rate of nick translation mediated by 3' exo- Pol is faster

than the Pol I-mediated nick translation (S5 Fig). However, in the presence of the template-

loaded β-clamp, 3'-exo⁻ Pol I-mediated nick translation (Fig 1D) was found to be broadly

similar to the Pol I-mediated nick translation patterns (compare lanes 7 and 8 between Fig 1C

and Fig 1D). This observation indicates that the 3'-exonuclease activity of Pol I has little

impact on the speed of nick translation in the presence of template-loaded β-clamp. On the

other hand, the 5'-exo⁻ Pol I exhibited intense pauses at the 39th nucleotide and few bases

downstream positions (dotted zone at lanes 2 and 3; Fig 1E). The template-loaded β-clamp

modestly enhances the intensity of 5'-exo⁻ Pol I pauses (dotted zone at lanes 7 and 8; Fig 1E).

These observations indicate that the elimination of 3'-exonuclease and 5'-exonuclease

activities impair the coordination in the strand displacement synthesis process, leading to

increased and decreased speed of nick translation, respectively, by Pol I.

β-clamp inhibits strand displacement function of Pol I

The Klenow fragment of Pol I exhibits polymerization, 3'exonuclease and 5' to 3'

strand displacement activity [33,34]. To determine if the slow nick translation by clamp-

bound Pol I through the DNA substrate was the result of impaired strand displacement, we

replaced *E. coli* Pol I with the exonuclease-free Klenow (exo⁻ Klenow), which lacks both 3'

and 5' exonuclease functions, in the assay. The exo⁻ Klenow exhibited a trail of scattered

pause complexes, as judged by the band intensities between the 39th to 67th nucleotide

positions (dotted zone at lanes 2 and 3; Fig 2A). Remarkably, the presence of template-loaded

β-clamp further intensified the pausing of exo⁻ Klenow, mostly at the 39th nucleotide position

(zone III; Fig 2A) where the nick translation begins.

Fig 2. β-clamp inhibits strand displacement activity of exo⁻ Klenow and exo⁻ Pol I

(**A**) The urea PAGE shows that exo⁻ Klenow exhibits a slower speed of strand displacement.

Presence of the template-loaded β-clamp stalls the exo⁻ Klenow pauses mostly at the 39th

nucleotide position. The exo⁻ Klenow-mediated strand displacement coupled with ligation

represents that the DNA ligase fails to ligate nicks. However, the presence of template-loaded

β-clamp increases the ligation frequency. (**B**) exo⁻ Pol I exhibits similar functional

consequences on the strand displacement and ligation as observed in case of exo⁻ Klenow

(panel A). The values representing mean ± sd are calculated from three different experiments.

The choice of exo⁻ Klenow of Pol I, in which the entire small domain containing 5'-

exonuclease activity is missing could be non-ideal, because the observed effects might be due

to a conformational change rather than stemming from the loss of 5'-exonuclease activity. To

test this issue, we generated a mutated version of Pol I, which encodes 3'- and 5'-

exonuclease-free Pol I (exo⁻ Pol I) but retains both of the domains. Using the exo⁻ Pol I, we

show that the pattern of paused complexes (dotted zones at the lanes 2 and 3; Fig 2B) was

comparable to the paused complexes observed with the exo⁻ Klenow (Fig 2A) in both the

absence and presence of the template loaded β-clamp. This data indicates that the elimination

of 3' and 5'-exonuclease activities, but not the deletion of small domain, induce the observed

paused complexes in the absence or presence of the template loaded β-clamp. As visually

observed, exo⁻ pol I was apparently more efficient in producing 67 nucleotides long product

than exo⁻ Klenow in the absence of template loaded β-clamp (Fig 2A and 2B).

β-clamp promotes ligation at an early stage of nick-translation

Apart from Pol I, DNA ligase also interacts with the β-clamp [19]. Therefore, the polymerization and flap endonuclease activities of Pol I, which fills the ssDNA gap to produce a nick and continuously generates new 5' ends during nick translation, respectively, may facilitate ligation between the upstream and downstream DNA strands. To probe this working hypothesis, we performed an assay coupled with *E. coli* DNA ligase. We observed that ligase along with Pol I, both in the presence or absence of the template-loaded β-clamp, produced more intense 67 bases long final product than Pol I alone produced in 30 seconds of reaction span (Fig 1C). This observation indicates that ligation by the ligase action during ongoing nick translation quickly contributes creating a fraction of the 67 bases long products. Therefore, almost all of the scattered nick translation products between the 39th and 67th nucleotide positions (marked by the dotted line at lane 2 of Fig 1C), i.e. which were in midway of nick translation, participated in the ligation process to generate 67 bases long product in 30 seconds (lane 4 of Fig 1C). We roughly estimated this contribution of ligase function during nick translation, introducing a new parameter, we termed as "relative ligation frequency" (RLF), that has maximum and minimum values 1 and 0, respectively, by definition (see the Methods section for detail). Thus, RLF is a normalized value or factor that accounts the intensity difference of the 67 bases long product originated in the presence and absence of ligase. Using the formula, RLF is estimated to be 0.84 and 0.96 in the absence and the presence of β-clamp, respectively (Fig 1C) with a statistically insignificant p-value (0.07, T-test) suggesting that β-clamp did not influence the RLF. However, unlike the absence of β-clamp, where the nick translation products were widely scattered (dotted line at lane 2 of Fig 1C), the template-loaded β-clamp restricted nick translation only a few nucleotide downstream of the starting 39th nucleotide position in 30 seconds time span (zone I of lane 7

of Fig 1C). Therefore, the ligase apparently worked at that early nick-point to generate 67

nucleotides long product (lane 9 of Fig 1C). Thus, we infer that by slowing down nick

translation, β-clamp also favors ligation at an early stage of nick translation.

We also performed assays to evaluate the RLF while 3'-exo⁻ and 5'-exo⁻ Pol I

enzymes take part in the nick translation. The RLF was determined to be less when 3'-exo⁻

Pol I was used in place of Pol I for nick translation assay (Fig 1C and 1D). This data

indicates that a faster nick translation by the 3'-exo⁻ Pol I than the Pol I (S5 Fig) inhibits

ligase action during nick translation. However, the presence of template-loaded β-clamp,

which substantially reduced the speed of 3'-exo⁻ Pol I (Fig 1D), increased the RLF to an

extent comparable to the values observed in the assay done with template-loaded β-clamp

and Pol I (Fig 1C). In contrast, using the 5'-exo⁻Pol I in the assay blocked the ligation

process. This data suggests that in the absence of the 5'-exonuclease function, the 3'-

exonuclease and polymerizing centers of Pol I could become hyperactive and thereby blocks

access of ligase to the 3' end of DNA.   However, the presence of template-loaded β-clamp

with the 5'-exo⁻-Pol I facilitated the ligation incrementally with the duration of reactions

(lanes 9 and 10, Fig 1E). These observations indicate that, β-clamp presumably slows down

passage through the nick and stimulate 3'-exonuclease function to backtrack Pol I at the nick

site facilitating ligation process.

Further, to confirm that β-clamp promotes quick ligation at an early stage, we

assessed the influence of β-clamp on the efficiency of ligation when exo⁻ Klenow was

employed for the gap filling and strand displacement processes (Fig 2A). The absence of β-

clamp severely impaired ligation, as the RLF was determined to be very small (Fig 2A). This

data suggests that the strand displacement activity of exo⁻ Klenow pushed the 5' end of

downstream DNA away from the polymerizing 3' end of the upstream DNA strand (lane 2 of

Fig 2A), rendering ligation almost impossible (lane 4 of Fig 2A). On the contrary, the

presence of template-loaded β-clamp greatly facilitated the ligation (Fig 2A). The ligation

efficiencies were similar when full-length exo$^-$ Pol I was used instead of exo$^-$ Klenow, both in

the absence and presence of β-clamp (compare Fig 2A and 2B).These data suggest that by

inhibiting the strand displacement activity, β-clamp extends the duration of the pause of exo$^-$

Klenow or exo$^-$ Pol I at the 39th nucleotides nick position (lanes 7 and 8 of Fig 2A and 2B).

Thus, the β-clamp-mediated weak strand displacement activity of Pol I facilitates DNA ligase

to seal the nick efficiently at an early stage (lane 9 of Fig 2A and 2B).

To visually distinguish between the nick translation and nick translation-coupled

ligation products, we prepared a modified template replacing 28 bases oligonucleotide with a

38 bases oligonucleotide. Thus, the newly assembled template was identical to the template

shown in Fig 1A except for a 10 nucleotide 3'-overhang (S6A Fig). Therefore in a nick

translation coupled ligation reaction the full-length product would be 67 bases long if nick

translation works alone; while coupled ligation would generate a 77 nucleotide long product.

The data presented further emboldened our observation regarding the effect of β-clamp on

ligation. The experiments performed with this template and its limitations are elaborated in

the S1 text.

In a control experiment, we conducted nick translation-coupled ligation experiment

tethering the biotinylated template with the streptavidin beads. After clamp loading, the

streptavidin beads were washed to remove the free clamp and clamp loader proteins, as

mentioned above. Next, we performed the reaction adding 3'-exo- Pol I, ligase, NAD$^+$ and

dNTPs to show only the template-loaded β-clamp, but not free β-clamp or clamp loader

complex β-clamp in our above assays, could modulate the ligation function (S7 Fig).

Loss of DNA contact with the clamp-bound Pol I explains slow nick translation

The finger domain of Pol I interacts with the downstream duplex DNA of the nick site and thereby takes part in displacing the nucleotides of the downstream strand as an oligonucleotide flap [35]. We have depicted this interaction in Fig 3A constructing a model based on the available structures (see Materials and Methods section). Therefore, mutations in the conserved residues of this domain, mainly F771 residue (Fig 3A), inactivate the strand displacement activity of Klenow [30] in a manner similar to the strand displacement activity of β-clamp-bound exo⁻ Klenow and exo⁻ Pol I (Fig 2). Therefore, we hypothesized that β-clamp could allosterically modulate the ternary complex to prevent the interaction between the Klenow and the 5' nucleotides of the downstream DNA strand. To probe this possibility, we slightly modified the template depicted in Fig 1A, for the assay. Here, we radiolabeled the 28 bases long downstream oligonucleotide instead of the 19 bases long upstream oligonucleotide. Additionally, the 28 bases long oligonucleotide had a bromodeoxyuridine (BrdU) base at the 2^{nd} position. BrdU in the DNA can exhibit a photochemical cross-linking at near UV light on encountering the adjacent amino acids of a binding protein, as described [46]. Using this template, we performed the exo⁻ Klenow-mediated strand displacement assay under UV light. In the absence of β-clamp, a small fraction of BrdU containing oligonucleotide formed the cross-linked adduct with exo⁻ Klenow during strand displacement reaction (lane 2 of Fig 3B), suggesting that the exo⁻ Klenow makes contact with the 28 bases oligonucleotide during strand displacement activity. However, the presence of template-loaded β-clamp abolished the cross-linked adduct, suggesting that β-clamp blocked the contact between the exo⁻ Klenow and the 28 bases oligonucleotide (lane 6 of Fig 3B). Free β-clamp and ATP did not affect the formation of the cross-linked adduct (lanes 3 and 4 of Fig

3B, respectively), suggesting that only template-loaded β-clamp has the said effect.

Therefore, the absence of the cross-linked adducts in the presence of β-clamp indicates that

the latter affects the interaction between the exo⁻ Klenow and the downstream DNA (Fig 3A),

affecting strand displacement function during nick translation (Fig 2).

Fig 3. **β-clamp inhibits the contact between the 5'end of nicked DNA and Pol-I.**

(**A**) Structural model of *E. coli* Klenow /DNA complex shows that finger domain of Klenow

makes a contact with downstream nick site. The model shows that the conserved F771 (in

blue surface representation), which participates in the strand displacement, is positioned

between the downstream nicked DNA strands. (**B**) A fraction of BrdU base containing

radiolabelled 28 bases oligonucleotide shows a gel shift under the near UV light, suggesting

that the oligonucleotide makes cross-linked adduct with Klenow during strand displacement

(lane 2). The presence of β-clamp and γ-complex in the absence of ATP (lane 3), or ATP

alone (lane 4) also exhibited similar crosslinking adduct. However, β-clamp, γ-complex, and

ATP together suppressed the formation of crosslinking product (lane 6), suggesting that

loading of the β-clamp on the template blocks the crosslinking. * Represents a minor

contaminated band with the custom synthesized 28 bases oligonucleotide.

Moreover, it is noteworthy that unlike static interaction studies [46], the cross-linking

assay was essentially performed here in a dynamic reaction (strand displacement) process. As

a result, BrdU and the adjacent aminoacyl residue(s) of the finger subdomain of Pol I made a

contact in a short time frame (in seconds) under UV light, leading to a poor level of cross-

linked adducts (Fig 3B).

β-clamp stimulates 5' exonuclease activity of Pol I

The flaps of various lengths generated transiently by Pol I are eventually excised by the flap endonuclease activity of the small domain of Pol I, releasing 1-12nt long products [37]. We show that the flap endonuclease activity of small domain of Pol I in the presence of template-loaded-β-clamp is a crucial step to determine the efficiency of DNA ligase function (Fig 2). Further, to check the effect of template-loaded β-clamp on the flap endonuclease activity of Pol I, we prepared a template where the 5' end of the 28 bases downstream oligonucleotide was radiolabeled (Fig 4A). We observe that the β-clamp modestly increased the rate of mononucleotide release from the 5' end of the 28 bases oligonucleotide by Pol I (Fig 4C and S8A Fig). In another case, we used a 5'-end radiolabeled 38 bases downstream oligonucleotide, of which the first 10 bases formed a flap structure at the 5' end (Fig 4B). The template-loaded β-clamp enhanced the rate of cleavage of the flap by Pol I to 2.5 folds (Fig 4D and S8B Fig).). These data suggest that the template-loaded β-clamp greatly stimulates the flap endonuclease activity of Pol I. We hypothesize that this increased 5' exonuclease activity of the clamp-bound Pol I for mononucleotides or shorter DNA flaps frequently generate 5' ends that helps in early and quick ligation, as shown in Fig 1C.

Fig 4. β-clamp increases the flap endonuclease activity of Pol I

(**A**) and (**B**) show the templates assembled using the various lengths of oligonucleotides. Asterisks (*) indicate the radiolabelled 5'-end of the oligonucleotides. (**C**) A representative denaturing gel indicates that the presence of β-clamp modestly increased the efficiency of single nucleotide cleavage at different time points. Also see S8A Fig for the fraction cleavage rate calculation. (**D**) A representative denaturing gel shows that β-clamp greatly enhanced the efficiency of 10 nucleotide long flap removal. Also see S8B Fig for fraction cleavage rate

338 calculation. All the experiments were done at least three times. The values represent mean ±

339 sd.

Discussion

In brief, we demonstrate that the presence of the template-loaded β-clamp inhibits

contact between the downstream strand and the exo⁻ Klenow (Fig 3), impairing the active

strand displacement activity of exo⁻ Klenow (Fig 2A). This loss of contact ultimately slows

down nick translation by Pol I traversing downstream DNA (Fig 1C). Conversely, β-clamp

barely affects the rate of nick translation traversing RNA substrate (Fig 1B and S2 Fig). Since

DNA:RNA hybrids are thermodynamically less stable than DNA:DNA hybrids [48,49], we

propose that a spontaneous bubbling at the downstream edge of a nick would be greater in the

DNA:RNA hybrid. This phenomenon would allow displacement of the strand to be less

dependent on the active participation of finger domain, making nick translation easier at

DNA:RNA hybrids in the presence of β-clamp (Fig 1B). Such spontaneous bubbling at the

ends of DNA:DNA would be lesser due to its higher thermodynamic stability, causing the

nick translation to slow down (Fig 1C). Thus, we hypothesize that β-clamp prevents faster

nick translation traversing DNA, but not at the RNA primer zone of an Okazaki fragment.

Furthermore, we show that β-clamp enhances the 5' exonuclease activity of Pol I (Fig

4). To elucidate the possible mechanism by which β-clamp influences the 5' exonuclease

activity of Pol I, we first performed molecular docking between β-clamp and Klenow

structures. The model displaying the highest stability under molecular dynamics simulation

was chosen (S9 Fig). In addition, the modeling of Klenow/β-clamp/DNA ternary complex

supports the feasibility of this Klenow/β-clamp interaction as DNA can follow an overall

straight path with minimal strain (S9 Fig). Next, we modeled *E. coli* small domain and

attached it to the N-terminal portion of Klenow with a flexible linker to get a final model

(S10 Fig). We inferred from the model that the interaction between β-clamp and Pol I

occludes the steric conformational space of small domain and pushes it more towards the flap

DNA sites. This hypothesis explains the increased flap endonuclease activity of Pol I bound

to β-clamp (Figs 3C and D).

Our study also unravels how β-clamp helps in the ligation process to repair nicked

DNA. First, β-clamp increases the ligation efficiency by interacting directly with the ligase

(Fig 1C) [19]. Secondly, by slowing down nick translation (Fig 1C) and increasing the 5'

exonuclease activity of Pol I (Fig 4), β-clamp promotes early ligation (Figs 1C and 2). The

RLF was dramatically reduced when we used 3'-exo⁻ Pol I, 5'-exo⁻ Pol I, exo⁻ Pol I, or exo⁻

Klenow versions of Pol I (Figs. 1D, 1E, and 2), suggesting that the loss of concerted

functions of exonucleases and polymerization affected the Okazaki fragment maturation.

Interestingly, β-clamp was able to slow down strand displacement activities of the mutated

polymerases, thereby restoring the ligation function to different extents. Thus, β-clamp serves

as the common interacting platform for replication and repair proteins to maintain overall

DNA metabolism and genome integrity by avoiding prolonged polymerization and

depolymerization steps, and quickly terminating nick translation, as depicted in the schematic

diagram (Fig 5). Therefore, these hitherto unknown aspects of β-clamp function, along with

its previously known role as a processivity factor for Pol I and other DNA polymerases [18-

25] could have immense evolutionary significance in the efficient maintenance of genome

integrity of prokaryotic organisms under challenging environmental milieu.

Fig 5. The model showing influence of β-clamp in gap repair. (A) The model shows that

Pol I removes downstream oligonucleotide by an active nick translation. (red flap).The

upcoming ligase molecule may ligate the polymerizing upstream (green) and the

depolymerising downstream (red) strands occasionally. (**B)** The clamp bound Pol I has been

shown at the nick site. Passive strand displacement for shorter stretch of nucleotides is shown

(short red flap). We hypothesize that efficient flap removal of clamp-bound Pol I frequently

creates new nick sites. As a result, upcoming ligase molecule may have plenty of time to

ligate polymerizing upstream (green) and depolymerising downstream (red) DNA strands at

an early stage.

Materials and Methods

Proteins and reagents used in the assay

PCR of the genes (*dnaN*, *dnaX*, *holA* and *holB*) encoding β-clamp and minimal clamp

loader complex (γ,δ and δ') [50] were performed. Since dnaX gene can produce full length τ

or truncated γ subunits, we performed PCR only for the coding region that encompasses γ

subunit region [51].The PCR products were cloned at the *Nde*I and *Eco*RI sites of pET21a (+)

vector and expressed as tag-less native proteins. The overexpressed proteins were purified

using standard protocols [16,52]. Pol I, exo$^-$ Klenow, *E. coli* DNA ligase, and ultrapure

nucleotides were purchased from New England Biolabs (NEB). Furthermore, to get full

length hexa-histidine-tagged Pol I, PCR amplified *E. coli polA* gene was cloned at the NdeI

and HindIII sites of pET28a (+) vector. Site directed mutagenesis was performed on the

pET28a-*polA* vector using Q5 Site-Directed Mutagenesis Kit from NEB to get 3'-exo$^-$ Pol I

($D^{355}A$, $E^{357}A$), 5'-exo$^-$ Pol I ($D^{13}N$, $D^{138}N$), and exo$^-$ Pol I ($D^{13}N$, $D^{138}N$, $D^{355}A$, $E^{357}A$)

[45,53]. The hexa-histidine-tagged Pol I and its mutated versions were overexpressed and

were purified by nickel-NTA affinity chromatography. The proteins were concentrated,

quantified and stored at -20°C in storage buffer (20 mM Tris-HCl, pH 7.5; 500mM NaCl,

1mM DTT and 50% glycerol). Nicotinamide adenine dinucleotide (NAD$^+$) was purchased

from Sigma-Aldrich. The hexa-histidine-tagged β-clamp was purified by cloning and

overexpressing *dnaN* gene as described in S1 text. The PAGE-purified oligonucleotides

(IDT, Inc., USA) used to prepare templates for the assay were, 67 bases (GGATCCCACAC

TCATTAAAATTAGTCGCTAATGCATTCTAAAGCATTCGCAACGAGAAGATAGA

GG), 19 bases (CCTCTATCTTCTCGTTGC G), 28 bases (GCGACTAATTTTAATGAGTG

TGGGATCC and GCGACUAAUUUUAAUGAGUGUGGGAUCC), and 38 bases (AGAA

CTTGACGCGACTAATTTTAATGAGT GTGGGATCC) in lengths. The 5'-end of 28 bases

primer was phosphorylated. The biotinylated template made for the control experiments are

described in Supporting information.

Assay conditions

5'-end radiolabeling of the desired oligonucleotides were performed using T4

polynucleotide kinase (NEB) and ^{32}P-γATP. The radiolabeled oligonucleotides were purified

using BioGel P-6 columns (Bio-Rad, Inc.) to remove the salts and free radioactivity.

Templates were prepared as described previously [35,54] using combinations of short

oligonucleotides, one of which was 5'-end radiolabeled, as mentioned in the results section.

Briefly, 50nM of each oligonucleotide were mixed with 100 µl annealing buffer (20mM Tris-

Acetate, pH 7.9, 50mM potassium acetate and 10mM magnesium Acetate). The mixture was

heated at 95°C for 2 minutes followed by slow cooling to 37°C to allow annealing of the

oligonucleotides into duplex DNA. The formation of duplex was almost complete, as

observed in a native PAGE (S1 Fig). To allow clamp loader complex to load β-clamp at the

primer template junction, 30 nM β-clamp was incubated with 20nM of annealed template and

30nM clamp loader complex and 1mM ATP in the reaction buffer (20mM Tris-acetate, pH

7.9, 50mM potassium Acetate, 10mM magnesium Acetate, 100µg/ml BSA) at 37°C for 2

minutes. Next, 10μM dNTPs and different versions of Pol I, Klenow and DNA ligase (each at

30nM concentration) were added in the reaction wherever required. For ligation assays, the

reaction mixture was also supplemented with 30μM nicotinamide adenine dinucleotide

(NAD$^+$) to catalyze the ligation process. The final reaction products were separated by urea

denaturing PAGE and were exposed to the phosphor-imager screen. A Fujifilm-9000

phosphor-image scanner was used to visualize the bands. Counts of the bands, or the desired

stretches of the lanes, were recorded for further analyses. A minimum of three experiments

were performed for each reaction to get the average counts and standard deviations. The

relative ligation frequency (RLF), i.e. fraction of nick translating products that take part in

ligation at 30 seconds time-point, was calculated, as defined below. If, X= average counts of

the 67-nucleotide long products generated in the presence of ligase, Y= average counts of the

67-nucleotide long products generated in the absence of ligase, Z = average counts of the

products between 39th and 67th nucleotide positions, then **RLF = (X-Y)/Z**. Nick translation,

single nucleotide and 10 nucleotides cleavage rates were calculated from the initial slopes of

the fitted curves, as described in the S1 text.

The BrdU cross-linking assay

The assay was designed with modifications to the protocol mentioned [46]. Briefly,

the reaction was performed at 37°C under 310 nm of UV light at 2 cm distance for indicated

times to allow cross-linking to occur between the bromodeoxyuridine (BrdU) labeled 28

bases oligonucleotide and nick translating Klenow in the presence or absence of β-clamp.

Since the nick translation reaction was observed to reach completion within a minute (Fig 1),

we allowed crosslinking for one minute. The reaction mixture was loaded on a denaturing gel

to visualize the band shift.

Structural modeling of *E. coli* Klenow/Nicked DNA complex

The structural modeling of Klenow/nicked DNA complex was done based on the crystal

structure of *Geobacillus stearothermophilus.* Klenow complexed with 9 base pairs of duplex

DNA (PDB ID 1L3S). The additional upstream and downstream DNA segments were built

using COOT (See S1 text).

Acknowledgements

464   Authors are grateful to Dr. Jayanta Mukherjee, Bose Institute, India, Dr. Gursharan

465 Kaur, IMTECH, India and Dr. James Boroweic, NYUMC, USA, for reading the manuscript

466 and providing critical suggestions on the manuscript.

References

1. Gefter ML, Hirota Y, Kornberg T, Wechsler JA, Barnoux C. Analysis of DNA

polymerases II and 3 in mutants of *Escherichia coli* thermosensitive for DNA synthesis.

Proc Natl Acad Sci USA. 1971; 68: 3150-3153.

2. Kong XP, Onrust R, O'Donnell M, Kuriyan J. Three-dimensional structure of the beta

subunit of *E. coli* DNA polymerase III holoenzyme: a sliding DNA clamp. Cell 1992; 69:

425-437.

3. Pritchard AE, Dallmann HG, Glover BP, McHenry CS. A novel assembly mechanism for

the DNA polymerase III holoenzyme DnaX complex: association of $\delta\delta'$ with $DnaX_4$

forms $DnaX_3\delta\delta'$. EMBO J. 2000;19: 6536-6545.

4. Lewis JS, Jergic S, Dixon NE. The *E. coli* DNA Replication Fork. Enzymes 2016;39: 31-

88.

5. Beattie TR, Reyes-Lamothe R. A replisome's journey through the bacterial chromosome.
Front Microbiol. 2015; 6: 562.

6. O'Donnell M, Langston L, Stillman B. Principles and concepts of DNA replication in
bacteria, archaea and eukarya. Cold Spring Harb Perspect. 2013; 5: a010108.

7. McInerney P, Johnson A, Katz F, O'Donnell, M. Characterization of a triple DNA
polymerase replisome. Mol Cell 2007;27: 527–538.

8. Reyes-Lamothe R, Sherratt DJ, Leake MC. Stoichiometry and architecture of active DNA
replication machinery in *Escherichia coli*. Science 2010;328: 498–501.

9. Naktinis V, Onrust R, Fang L, O'Donnell M. Assembly of a chromosomal replication
machine: two DNA polymerases, a clamp loader, and sliding clamps in one holoenzyme
particle. II. Intermediate complex between the clamp loader and its clamp. J Biol Chem.
1995;270: 13358–13365.

10. Hingorani MM, and O'Donnell M. ATP binding to the *Escherichia coli* clamp loader
powers opening of the ring-shaped clamp of DNA polymerase III holoenzyme. J Biol
Chem. 1998;273: 24550–24563.

11. Turner J, Hingorani MM, Kelman Z, O'Donnell M. The internal workings of a DNA
polymerase clamp-loading machine. EMBO J. 1999;18: 771–783.

12. Georgescu RE, Kim SS, Yurieva O, Kuriyan J, Kong XP, O'Donnell M. Structure of a
sliding clamp on DNA. Cell 2008;132: 43-54.

13. Bertram JG, Bloom LB, Turner J, O'Donnell M, Beechem JM, Goodman MF. Pre-steady
state analysis of the assembly of wild type and mutant circular clamps of *Escherichia coli*
DNA polymerase III onto DNA. J Biol Chem. 1998;273: 24564–24574.

14. Hingorani MM, Bloom LB, Goodman MF, and O'Donnell, M. Division of labor–
sequential ATP hydrolysis drives assembly of a DNA polymerase sliding clamp around
DNA. EMBO J. 1999;18: 5131–5144.

15. Johnson A, O'Donnell M. Cellular DNA replicases: components and dynamics at the

replication fork. Annu Rev Biochem. 2005;74: 283-315.

16. Kornberg A, Baker T. DNA Replication. Vol. 19. W.H. Freeman & Co. 1992.

17. Stukenberg PT, Studwell-Vaughan PS, O'Donnell M. Mechanism of the sliding beta-

clamp of DNA polymerase III holoenzyme. J Biol Chem. 1991;266: 11328-11334.

18. Indiani C, McInerney P, Georgescu R, Goodman MF, O'Donnell M. A sliding-clamp

toolbelt binds high- and low-fidelity DNA polymerases simultaneously. Mol Cell

2005;19: 805-815.

19. Lopez de Saro FJ, O'Donnell M. Interaction of the β sliding clamp with MutS, ligase, and

DNA polymerase I. Proc Natl Acad Sci USA 2001;98: 8376-8380.

20. Burgers PM, Kornberg A, Sakakibara Y. The dnaN gene codes for the beta subunit of

DNA Polymerase III holoenzyme of *Escherichia coli*. Proc Natl Acad Sci USA 1981;78:

5391-5395.

21. Wagner J, Fujii S, Gruz P, Nohmi T, Fuchs RP. The β clamp targets DNA polymerase IV

to DNA and strongly increases its processivity. EMBO Rep. 2000;1: 484-488.

22. Sutton MD, Opperman T, Walker G C. The *Escherichia coli* SOS mutagenesis proteins

UmuD and UmuD' interact physically with the replicative DNA polymerase. Proc Natl

Acad Sci USA. 1999;96: 12373-12378.

23. Bonner CA, Stukenberg PT, Rajagopalan M, Eritja R, O'Donnell M, McEntee K, Echols

H, Goodman MF. 1992. Processive DNA synthesis by DNA polymerase II mediated by

DNA polymerase III accessory proteins. J Biol Chem. 1999;267: 11431-11438.

24. Hughes Jr AJ, Bryan SK, Chen H, Moses RE, McHenry CS. *Escherichia coli* DNA

polymerase II is stimulated by DNA polymerase III holoenzyme auxiliary subunits. J Biol

Chem. 1991;266: 4568-4573.

25. Sutton MD. Coordinating DNA polymerase traffic during high and low fidelity synthesis.

Biochim Biophys Acta. 2010;1804: 1167-1179.

26. Wang TC. Discontinuous or semi-discontinuous DNA replication in *Escherichia coli*?

Bioessays 2005;27:, 633–636.

27. Heller RC and Marians KJ. Replication fork reactivation downstream of a blocked

nascent leading strand. Nature 2006;439: 557-562.

28. Langston LD and O'Donnell M. DNA replication: keep moving and don't mind the gap.

Mol Cell 2006;23: 155-160.

29. Pomerantz and O'Donnell. The replisome uses mRNA as a primer after colliding with

RNA polymerase. Nature 2008;456: 762–766.

30. Dutta D, Shatalin K, Epshtein V, Gottesman ME, Nudler E. Linking RNA Polymerase

Backtracking to Genomic Instability. Cell 2011;146: 533-543.

31. Ogawa T, Okazaki T. Discontinuous DNA replication. Annu Rev Biochem. 1980;49:

421-457.

32. Tougu K, Marians KJ. The interaction between helicase and primase sets the replication

fork clock. J Biol Chem. 1996;271: 21398-21405.

33. Lovett ST. The DNA Exonucleases of *Escherichia coli*. *EcoSal Plus* 2011;4:

doi:10.1128/ecosalplus.4.4.7.

34. Patel PH, Suzuki M, Adman E, Shinkai A, Loeb LA. Prokaryotic DNA polymerase I:

evolution, structure, and "base flipping" mechanism for nucleotide selection. J Mol Biol.

2001;308: 823-837.

35. Singh K, Srivastava A, Patel SS, Modak MJ. Participation of the fingers subdomain of

Escherichia coli DNA polymerase I in the strand displacement synthesis of DNA. J Biol

Chem. 2007;282: 10594-10604.

36. Srivastava A, Singh K, Modak MJ.. Phe 771 of *Escherichia coli* DNA polymerase I

(Klenow fragment) is the major site for the interaction with the template overhang and the

stabilization of the pre-polymerase ternary complex. Biochem. 2003;42: 3645-3654.

37. Lundquist RC, Olivera BM. Transient generation of displaced single-stranded DNA

during nick translation. Cell 1982;3:, 53-60.

38. Langston LD, O'Donnell M. DNA replication: keep moving and don't mind the gap. Mol

Cell 2006;23: 155-160.

39. Leu FP, Georgescu R, O'Donnell M. Mechanism of the *E. coli* tau processivity switch

during lagging-strand synthesis. Mol Cell 2003;11: 315-327.

40. Stukenberg PT, Turner J, O'Donnell M. An explanation for lagging strand replication:

polymerase hopping among DNA sliding clamps. Cell 1994;78: 877-887.

41. Yang J, Nelson SW, Benkovic SJ. The control mechanism for lagging strand polymerase

recycling during bacteriophage T4 DNA replication. Mol Cell 2006;21: 153-164.

42. Moolman MC, Krishnan ST, Kerssemakers JW, van den Berg A, Tulinski P, Depken M,

et al. Slow unloading leads to DNA-bound β2-sliding clamp accumulation in live

Escherichia coli cells. Nat Commun. 2014;5: 5820

43. Su'etsugu M, Errington J. The replicase sliding clamp dynamically accumulates behind

progressing replication forks in Bacillus subtilis cells. Mol Cell 2011;41: 720-732.

44. Vaisman A, McDonald JP, Noll S, Huston D, Loeb G, Goodman MF,et al. Investigating

the mechanisms of ribonucleotide excision repair in *Escherichia coli*. Mutat Res.

2014;761: 21-33.

45. Xu Y, Derbyshire V, Ng K, Sun XC, Grindley ND, Joyce CM. Biochemical and

mutational studies of the 5'-3' exonuclease of DNA polymerase I of *Escherichia coli*. J

Mol Biol. 1997;268: 284-302.

46. Hayner JN. and Bloom LB. The β sliding clamp closes around DNA prior to release by

the *Escherichia coli* clamp loader γ complex. J Biol Chem. 2013;288: 1162-70.

47. Turner RM Jr, Grindley ND, Joyce CM. Interaction of DNA polymerase I (Klenow

fragment) with the single-stranded template beyond the site of synthesis. Biochem.

2003;42: 2373-2385.

48. Hall KB, McLaughlin LW. Thermodynamic and structural properties of pentamer

DNA.DNA, RNA.RNA, and DNA.RNA duplexes of identical sequence. Biochem.

1991;30: 10606-10613.

49. Ratmeyer L, Vinayak R, Zhong YY, Zon G, Wilson WD. Sequence specific

thermodynamic and structural properties for DNA.RNA duplexes. Biochem. 1994;33:

5298-5304.

50. Onrust R, Finklestein J, Naktinis V, O'Donnell M. Assembly of a chromosomal

replication machine: two DNA polymerases, a clamp loader, and sliding clamps in one

holoenzyme particle. I. Organization of the clamp loader. J Biol Chem. 1995;270: 13366–

13377.

51. Maki H, Maki S, Kornberg A. DNA Polymerase III Holoenzyme of *Escherichia coli* IV.

The holoenzyme is an asymmetric dimer with twin active sites. J Biol Chem. 1988;263:

6570-6578.

52. Duzen JM, Walker GC, Sutton MD. Identification of specific amino acid residues in the

E. coli β processivity clamp involved in interactions with DNA polymerase III, UmuD

and UmuD'. DNA repair 2004;3: 301-312.

53. Derbyshire V, Freemont PS, Sanderson MR, Beese L, Friedman JM, Joyce CM, et al.

Genetic and crystallographic studies of the 3',5'-exonucleolytic site of DNA polymerase I.

Science 1998;240: 199-201.

54. Zhao G, Tang S, Li J, Hu T, Guan Y. Effects of cations on small fragment of DNA

Polymerase I using a novel FRET assay. J Biol Chem. 2014;46: 659-667.

Supporting information

S1 Fig. Gel shift assay shows correct primer annealing during preparation of template.

The native gel shows the separation of 19 nucleotides (lane 1), 28 nucleotides and 67

nucleotides (lane 3) long oligonucleotides. When they mixed in an annealing reaction, the

individual oligonucleotides disappeared from their original position and gave rise to a higher

band (Lane 4), suggesting that annealing was 100% efficient. However, non-complementary

oligonucleotides (19 and 28 bases) did not found to be annealed (lane 5). On the other hand,

the 67 bases long oligonucleotide, which has complementary regions for both 19 and 28

bases long oligonucleotides, has also completely annealed to these 19 and 28 bases

oligonucleotides (lanes 6 and 7, respectively).

S2 Fig. β-clamp does not affect the rate of nick translation through the RNA substrates.

The fraction of the initial counts that appeared at 67 nucleotide position were plotted. The

rate of the nick translation through RNA substrate was calculated from the fitted curve shown

above (also see S1 text). The negligible difference between the rate of nick translation in the

presence and absence of β-clamp, suggests that β-clamp has no influence on nick translation

through RNA substrates.

S3 Fig. Interaction of the loaded-β clamp on the primed template. (A) The lanes 2-5 are

showing purified individual proteins that are used in the pull down assay. The streptavidin-

bound primed biotinylated template could pull-down β-clamp (marked by white arrow),

while clamp loader proteins washed away (compare unwashed and washed lanes 6 and 7, respectively). (**B**) The his-tag β clamp pull down primed template, as indicated by the pulled down radiolabeled 19nt long oligonucleotide, which was the integral part of the template.

S4 Fig. Clamp loader proteins in the reaction mixture do not interfere with nick translation. The urea denaturing gel indicating that washing the streptavidin-bound reaction products, which washes out the clamp loader protein, but retains β clamp (S3 Fig), generates similar pauses (+wash lanes) as observed for standard reaction (-wash lanes) performed in the presence of β clamp loader proteins.

S5 Fig. Comparison between Pol I and 3'-exo⁻ Pol I mediated nick translation. (A) The autoradiogram represents nick translation product formation at three different time points by Pol I and 3'-exo⁻ Pol I**. (B)** The rate of nick translation was calculated as described in S1 text, from three independent autoradiograms and plotted. The initial rate of nick translation was found to be faster in case of 3'-exo⁻ Pol I.

S6 Fig. Influence of β-clamp on nick translation coupled ligation. (A). A new template was assembled using 67 bases, 5'-phosphorylated 38 bases and 5'-radiolabelled (asterisk) 19 bases oligonucleotides, so that a 10 nucleotide long 3' overhang was generated as shown. **(B)** The autoradiogram represents Pol I mediated nick translation-coupled ligation in the presence or absence of β-clamp. Visibly, 77 nucleotide ligation product that appears at lower time points, degraded quickly due to 3' exonuclease activity of intact Pol I. **(C, D)** The representative autoradiograms show that 3'-exo⁻ Pol I and exo⁻ Pol I produced stable nick translation-coupled ligation products in the presence or absence of β-clamp in three different

time points. (E) RLFs calculated at different time points in the presence or absence of β-

clamp from panels S6C and S6D are shown.

S7 Fig. The template-loaded β-clamp enhances the efficiency of ligation. The nick

translation coupled ligation assay was performed on streptavidin beads to remove clamp

loader proteins after loading of the β-clamp. Addition of 3' exo- Pol I and dNTPs exhibited

pausing at the 39[th] and downstream locations at two different time points. Addition of the Pol

I and ligase efficiently produces 67nt products, suggesting that in the absence of clamp loader

complex the nick translation coupled ligation reactions are similar as demonstrated in Fig.

1D.

S8 Fig. β-clamp increases flap endonuclease activity by Pol I. (A). Single nucleotide

cleavage was shown in Fig 4C. The single nucleotide fraction cleavage values at different

time points were plotted to calculate the rate. The presence of β-clamp was found to modestly

increase the single nucleotide cleavage. **(B)** Similarly, flap cleavage was shown in Fig 4D and

the flap fraction cleavage values at different time points were plotted to calculate the flap

fraction cleavage rates. Interestingly, β-clamp greatly enhanced the flap cleavage rate.

S9 Fig. Model for PolK/ β-clamp/Nicked DNA ternary complex. (A) Cartoon

representation of the docked model of Klenow (cyan) /β-clamp (gray and blue) protein

complex. The possible protein-protein interaction interfaces are encircled in red and blue. **(B)**

Molecular dynamics simulations for the docked Klenow/β-clamp protein-protein complex.

(C) Surface representation of the modelled Klenow (grey) /β-clamp (yellow and blue)/ DNA

(orange and green) ternary complex.

S10 Fig. Final model for PolI/clamp/Nicked DNA complex. The structural model of *E. coli*

small domain of Pol I (light pink) was built using Prime module in the Schrodinger software

suite using 1BGX as the template. The models were manually placed near the Klenow/nicked

DNA (A) and Klenow/ β-clamp/nicked DNA (B) models using PyMOL. The predicted

flexible linker region connecting small domain and Klenow fragment has been shown by

undulating lines in different grey shades. The conformational space available in β-clamp-

bound and β-clamp free Klenow/nicked DNA structures are shown by multiple copies of

Psmall domains. From this model, we hypothesize that the presence of β- clamp could

sterically restrict (shown by red T blocker) the conformational freedom of small domain

hence increasing the chance of accessing 5' exonuclease site.

S1 Text. Supporting methods and results

Figure 1

A

28 nt downstream DNA

Klenow										
(exo-)	−	+	+	+	+	−	+	+	+	+
Ligase	−	−	−	+	+	−	−	−	+	+
β-clamp	−	−	−	−	−	−	+	+	+	+
Time (Sec)	0	30	60	30	60	0	30	60	30	60

67nt

39nt

20nt

19nt

Lanes:1 2 3 4 5 6 7 8 9 10

Relative ligation efficiency
Without clamp = 0.05 ± 0.02
With clamp   = 0.82 ± 0.03

B

28 nt downstream DNA

PolI										
(exo⁻)	−	+	+	+	+	−	+	+	+	+
Ligase	−	−	−	+	+	−	−	−	+	+
β-clamp	−	−	−	−	−	−	+	+	+	+
Time (Sec)	0	30	60	30	60	0	30	60	30	60

67nt

39nt

20nt

19nt

Lanes:1 2 3 4 5 6 7 8 9 10

Relative ligation efficiency
Without clamp = 0.15 ± 0.05
With clamp   = 0.71 ± 0.09

Figure 2

Figure 3

**Relative rate of single nt. cleavage
(Fraction counts/second)**
Without clamp = 0.029 ± 0.0009
With clamp = 0.040 ± 0.0062

**Relative rate of 10 nt. flap cleavage
(Fraction counts/second)**
Without clamp = 0.0039 ± 0.0001
With clamp = 0.0096 ± 0.001

Figure 4

A

Active strand displacement by Pol I
Slower Flap endonuclease
Faster nick translation

Slower ligation

B

β-clamp

Passive strand displacement
Faster flap endonuclease
Slower nick translation

Quicker and efficient ligation

Lig

Figure 5