Biological Sciences: Biochemistry

Physical Sciences: Earth, Atmospheric, and Planetary Sciences

Multiple prebiotic metals mediate translation

Marcus S. Bray[a*], Timothy K. Lenz[b*], Jay William Haynes[b], Jessica C. Bowman[b], Anton S. Petrov[b], Amit R. Reddi[b], Nicholas V. Hud[b], Loren Dean Williams[b,1] and Jennifer B. Glass[c,1]

[a]School of Biological Sciences, Georgia Institute of Technology, 950 Atlantic Dr, Atlanta, GA, USA, 30332

[b]School of Chemistry and Biochemistry, Georgia Institute of Technology, 901 Atlantic Dr, Atlanta GA, USA, 30332

[c]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, 311 Ferst Dr, Atlanta, GA, US, 30332

[*]These two people have contributed equally to this work.

[1]To whom correspondence should be addressed:
Jennifer B. Glass: School of Earth & Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332; jennifer.glass@eas.gatech.edu; Tel. 404- 894-3942; Fax. 404-894-5638
Loren Dean Williams: School of Chemistry & Biochemistry, Georgia Institute of Technology, Atlanta, GA 30332; loren.williams@chemistry.gatech.edu; Tel. 404-385-6258; Fax. 404-894-2295

Keywords: Translation, SHAPE, iron, manganese, magnesium, ribosome, ribosome function, ribosome structure, protein synthesis

Running title: Multiple prebiotic metals mediate translation

ABSTRACT

Today, Mg^{2+} is an essential cofactor with diverse structural and functional roles in life's oldest
macromolecular machine, the translation system. We tested whether ancient Earth conditions (low O_2,
high Fe^{2+}, high Mn^{2+}) can revert the ribosome to a functional ancestral state. First, SHAPE (Selective 2'-
Hydroxyl Acylation analyzed by Primer Extension) was used to compare the effect of Mg^{2+}, Fe^{2+}, and
Mn^{2+} on the tertiary structure of rRNA. Then, we used *in vitro* translation reactions to test whether Fe^{2+} or
Mn^{2+} could mediate protein production, and quantified ribosomal metal content. We found that: (i) Mg^{2+},
Fe^{2+}, and Mn^{2+} had strikingly similar effects on rRNA folding; (ii) Fe^{2+} and Mn^{2+} can replace Mg^{2+} as the
dominant divalent cation during translation of mRNA to functional protein; (iii) Fe and Mn associate
extensively with the ribosome. Given that the translation system originated and matured when Fe^{2+} and
Mn^{2+} were abundant, these findings suggest that Fe^{2+} and Mn^{2+} played a role in early ribosomal evolution.

SIGNIFICANCE

Ribosomes are found in every living organism where they are responsible for the translation of messenger
RNA into protein. The ribosome's centrality to cell function is underscored by its evolutionary
conservation; the core structure has changed little since its inception ~4 billion years ago when
ecosystems were anoxic and metal-rich. The ribosome is a model system for the study of bioinorganic
chemistry, owing to the many highly coordinated divalent metal cations that are essential to its function.
We studied the structure, function, and cation content of the ribosome under early Earth conditions (low
O_2, high Fe^{2+}, high Mn^{2+}). Our results expand the roles of Fe^{2+} and Mn^{2+} in ancient and extant
biochemistry as cofactors for ribosomal structure and function.

\body
Life arose around 4 billion years ago on an anoxic Earth with abundant soluble Fe^{2+} and Mn^{2+} (1-5).
Biochemistry had access to vast quantities of these metals for over a billion years before biological O_2
production was sufficient to oxidize and precipitate them. The pervasive use of these "prebiotic" metals in
extant biochemistry, despite current barriers to their biological acquisition, likely stems from their
importance in the evolution of the early biochemical systems.
38
The translation system, which synthesizes all coded protein (6, 7), originated and matured during the
Archean Eon (4-2.5 Ga) in low-O_2, high-Fe^{2+}, and high-Mn^{2+} conditions (8). The common core of the
ribosome, and many other aspects of the translation system, have remained essentially frozen since the
last universal common ancestor (9). In extant biochemistry, Mg^{2+} ions are essential for both structure and
function of the ribosome (10) and other enzymes involved in translation (11). In ribosomes, Mg^{2+} ions
engaged in a variety of structural roles (**Table 1**), including in Mg^{2+}-rRNA clamps (12, 13) (**Fig. 1a**), in
dinuclear microclusters that frame the peptidyl transferase center (PTC) (13) (**Fig. 1b**), and at the small
subunit-large subunit (SSU-LSU) interface (14) (**Fig. 1c**). Functional Mg^{2+} ions stabilize a critical bend in
mRNA between the P-site and A-site codons (15) (**Fig. 1d**), and mediate rRNA-tRNA and rRNA-mRNA
interactions (16) (**Fig. 1e, f**). Mg^{2+} ions also interact with some rProteins (17). Additionally, accessory
enzymes needed for translation – aminoacyl tRNA synthetases, methionyl-tRNA transformylase, creatine
kinase, myokinase, and nucleoside-diphosphate kinase – require Mg^{2+} ions as cofactors (**Table 1**).
51
Multiple types of cationic species can interact productively with RNAs in a variety of systems (18-20).
Recent results support a model in which Fe^{2+} and Mn^{2+}, along with Mg^{2+}, were critical cofactors in ancient
nucleic acid function (21). As predicted by this model, functional Mg^{2+}-to-Fe^{2+} substitutions under anoxic
conditions were experimentally verified to support RNA folding and catalysis by ribozymes (22, 23), a
DNA polymerase, a DNA ligase, and an RNA polymerase (24). Functional Mg^{2+}-to-Mn^{2+} substitution has
long been known for DNA polymerases (24-26). For at least some nucleic acid processing enzymes,
optimal activity is observed at lower concentrations of Fe^{2+} than Mg^{2+} (22, 24). Based on these previous
results, we hypothesized that Fe^{2+} and Mn^{2+} could partially or fully replace Mg^{2+} during translation. In
this study, we relocated the translation system to the low-O_2, Fe^{2+}-rich, or Mn^{2+}-rich environment of its
ancient roots, and compared its structure, function, and cation content under modern vs. ancient
conditions.
63
64   **Results**
65
Fe^{2+} and Mn^{2+} fold LSU rRNA to a near-native state. To test whether Fe^{2+} or Mn^{2+} can substitute for
Mg^{2+} in folding rRNA to a native-like state, we compared folding of LSU rRNA of the bacterial ribosome
in the presence of Mg^{2+}, Fe^{2+}, or Mn^{2+} by SHAPE (Selective 2'-Hydroxyl Acylation analyzed by Primer
Extension). SHAPE provides quantitative, nucleotide-resolution information about RNA flexibility, base
pairing, and 3D structure, and has previously been used to monitor the influence of cations, small
molecules, or proteins on RNA structure (27-32). We previously used SHAPE to show that the LSU
rRNA adopts a near-native state in the presence of Mg^{2+}, with the core inter-domain architecture of the
assembled ribosome and residues positioned for interactions with rProteins (33). Here, SHAPE
experiments were performed in an anoxic chamber to maintain the oxidation state of the metals and to
prevent Fenton cleavage. The minimum concentration required to fully fold rRNA (10 mM Mg^{2+}, 2.5 mM
Fe^{2+}, or 2.5 mM Mn^{2+}) was used for all SHAPE experiments (**Datasets S1 and S2**).
Addition of Mg^{2+}, Fe^{2+}, or Mn^{2+} induced widespread structural changes in the LSU rRNA in the presence
of Na^+, as reflected in SHAPE profiles (see **Materials and Methods**) and displayed as 'heat maps' on the
LSU rRNA secondary structure (**Fig. 2; *SI Appendix*, Fig. S1**). Among the nucleotides forming the PTC,
similar SHAPE profiles were obtained in the presence of Mg^{2+}, Fe^{2+}, or Mn^{2+} (***SI Appendix*, Fig. S1**). The
ΔFe^{2+} and ΔMg^{2+} heat maps obtained for the entire 23S rRNA are nearly identical in most regions (**Fig.**

2d, **e**). As expected for conversion of secondary structure to fully folded tertiary structure, helices tended
to be invariant, whereas loops and bulges were impacted by addition of Mg^{2+}, Fe^{2+}, or Mn^{2+}. For the 23S
rRNA, 86% of nucleotides (43/50) that exhibited a significant response (>0.3 SHAPE units) to Mg^{2+} also
exhibited a similar trend with Fe^{2+}. The greatest discrepancy between Fe^{2+} and Mg^{2+} was observed in the
L11 binding domain (**Fig. 2d, e**).

Fe^{2+} and Mn^{2+} mediate translation. Translation reactions were performed in an anoxic chamber in the
presence of various cations and cation concentrations. Production of the protein dihydrofolate reductase
(DHFR) from its mRNA was used to monitor translational activity. Protein synthesis was assayed by
measuring the rate of NADPH oxidation by DHFR. These reactions were conducted in a small
background of 2.5 mM Mg^{2+} (*SI Appendix*, **Fig. S2a**). This background is below the requirement to
support translation, consistent with previous findings that a minimum of ~5 mM Mg^{2+} is needed for
assembly of mRNA onto the SSU (34, 35). As a control, we recapitulated the previously established Mg^{2+}
dependence of the translation system, and then repeated the assay with Fe^{2+}.

Activity of the translation system with variation in [Fe^{2+}] closely tracks activity with variation in [Mg^{2+}]
(**Fig. 3**). Below 7.5 mM, total divalent cation concentration, minimal translation occurred with either Fe^{2+}
or Mg^{2+}, as expected (36). Activity peaked at 9.5 mM for both cations and decreased modestly beyond the
optimum. At a given divalent cation concentration, Fe^{2+} supported around 50-80% of activity with Mg^{2+}
(**Fig. 4**). This result was observed with translation reactions run for 15, 30, 45, 60, 90 and 120 min at the
optimal divalent cation concentrations. Mn^{2+} also supported similar translation activity to Fe^{2+} at optimal
divalent concentrations (*SI Appendix*, **Fig. S3**). Along with Mg^{2+}, Fe^{2+}, and Mn^{2+} we investigated
whether other divalent cations could support translation. No translation activity was detected with Co^{2+},
Cu^{2+}, or Zn^{2+} (*SI Appendix*, **Fig. S3**).

To test whether alternative divalent cations could completely replace Mg^{2+} in translation, we decreased
the background Mg^{2+} from 2.5 to 1 mM by thoroughly washing the ribosomes prior to translation
reactions with 7-11 mM Fe^{2+} or Mn^{2+} (*SI Appendix*, **Fig. S2b**). With 1 mM background Mg^{2+}, Fe^{2+}
supported 12-23% of the activity with Mg^{2+} over the concentrations tested, while Mn^{2+} supported 43-50%
activity relative to Mg^{2+} (**Fig. 5a**). Washing the factor mix allowed us to decrease the background Mg^{2+} in
translation reactions to ~4-6 µM (*SI Appendix*, **Fig. S2c**). At this level, minimal protein production was
observed with Fe^{2+}, while Mn^{2+} supported 29-38% of the activity measured with Mg^{2+} (**Fig. 5b**).

Fe and Mn associate extensively with the ribosome. To experimentally confirm that Fe and Mn associate
with the assembled ribosome, we analyzed the total Fe or Mn content of ribosomes after incubation in
anoxic reaction buffer containing 7 mM Fe^{2+} or 7 mM Mn^{2+}. Under the conditions of our translation
reactions, 584 ± 9 Fe atoms or 507 ± 28 Mn atoms associate with each ribosome.

Finally, we computationally investigated whether Mg^{2+}, Fe^{2+}, and Mn^{2+} might be interchangeable during
translation using quantum mechanical characterization of M^{2+}-rRNA clamps (**Fig. 1a;** *SI Appendix*, **Fig.**
S4), which are abundant in the ribosome (12, 13). The geometries of Mg^{2+}-rRNA, Fe^{2+}-rRNA and Mn^{2+}-
rRNA clamps, are nearly identical (*SI Appendix*, **Table S1-S3**). However, due to the accessibility of their
d-orbitals, more charge is transferred to Fe^{2+} or Mn^{2+} (*SI Appendix*, **Table S2**) than to Mg^{2+} (*SI*
Appendix, **Table S3**). The effect of the modestly greater radius of Mn^{2+} (*SI Appendix*, **Table S1**) is offset
by d-orbital charge transfer (*SI Appendix*, **Table S2**), leading to elevated stability of Fe^{2+}-rRNA clamp
over the Mn^{2+}-rRNA clamp (*SI Appendix*, **Table S3).**

Discussion

In this study, we successfully replaced ribosomal Mg^{2+} with Fe^{2+} or Mn^{2+} under conditions mimicking the
anoxic Archean Earth. Previously, the only divalent cation known to mediate rRNA folding and function
was Mg^{2+}. We found that isolated rRNA folds to essentially the same global state (37, 38) with Mg^{2+},

Fe^{2+}, or Mn^{2+} under anoxia. We then revealed that Fe^{2+} or Mn^{2+} can serve as the dominant divalent cation
during translation. Mg^{2+} at 2.5 mM was insufficient to mediate protein synthesis; 5 mM additional Mg^{2+},
Fe^{2+}, or Mn^{2+} restored translational activity. These findings suggest that functional substitutions of Mn^{2+}
or Fe^{2+} for Mg^{2+} can occur in large ribozymes, similar to previous reports for protein enzymes and small
ribozymes (24-26, 39, 40). Near-complete removal of Mg^{2+} prevented Fe^{2+}-mediated translation and
partially inhibited Mn^{2+}-mediated translation, suggesting that Mg^{2+} is optimal for some specific roles in
the translation system. Regardless, the general effectiveness of Mn^{2+} or Fe^{2+} for Mg^{2+} substitutions in the
translation system is astounding considering the enormous number of divalent cations associated with a
given ribosome, and the broad scope of their structural and functional roles (10, 11) (**Table 1, Fig. 1**).

The observation that >500 Fe or Mn ions can associate with a bacterial ribosome is consistent with the
number of Mg^{2+} ions observed by X-ray diffraction (100-1000 Mg^{2+} per ribosome (41)), and supports a
model in which Fe^{2+} or Mn^{2+} has replaced Mg^{2+} as the dominant divalent cation in our experiments. The
high capacity of ribosomes for Fe^{2+} and Mn^{2+} reflects all rRNA-associated divalent cations, including
condensed, glassy and chelated divalent cations (42), and in addition, we presume that Fe^{2+} or Mn^{2+} can
associate with a variety of rProteins, including those previously shown to bind Zn^{2+} (e.g. S2, S15, S16,
S17, L2, L13, L31, L36 in *E. coli*) (43).

The differences in protein production observed among the three divalent cations likely arise from a
variety of evolutionary and physiological factors. For instance, *E. coli* ribosomes may be evolutionarily
adapted to Mg^{2+} instead of Fe^{2+} or Mn^{2+}. The difference in translational activity between Mn^{2+} and Fe^{2+},
particularly when background Mg^{2+} was removed, suggests that Mn^{2+} is more viable than Fe^{2+} upon full
substitution. Mn^{2+}/Mg^{2+} interchangeability may depend on relative stabilities of Mn^{2+} and Mg^{2+} in M^{2+}-
rRNA clamps (*SI Appendix*, **Fig. S3**). Besides the ribosome, our translation reactions utilize many
accessory proteins such as elongation factors and aminoacyl-tRNA synthetases that also have divalent
cation requirements. Decreased activity of any of one these systems with Mn^{2+} and Fe^{2+} would cause a
pinch point in an otherwise fully functional translation system. Indeed, the relative activity of myokinase
and arginine t-RNA synthetase are both lower with Mn^{2+} or Fe^{2+} than with Mg^{2+} (44, 45).

While intracellular Mg^{2+} is around 10^{-3} M (46), specific physiological or environmental conditions can
significantly elevate intracellular Fe^{2+} and Mn^{2+}. Under oxidative stress, some microbes accumulate
excess Mn^{2+}. For example, radiation-tolerant *Deinococcus radiodurans* contains ~10 times higher Mn^{2+}
than *E. coli* (~10^{-5} M Mn^{2+} (47, 48)). In the absence of O_2, *E. coli* contains ~10 times higher labile Fe^{2+}
(~10^{-4} M) than in the presence of O_2 (~10^{-5} M (49)). Thus, it is possible that the absence of Fe^{2+} and Mn^{2+}
in experimentally determined ribosomal structures is reflective of culturing, purification, or crystallization
conditions (high O_2, high Mg^{2+}, low Fe^{2+}, low Mn^{2+}), and that other cations may also be present under
diverse physiological conditions.

We have shown that the translation system functions with mixtures of divalent cations, which are variable
during long-term evolutionary history and during short term changes in bioavailability and oxidative
stress. When combined with previous results that DNA replication and transcription can be facilitated by
Fe^{2+} and Mn^{2+} (18-20, 22-26, 39, 40), our findings that both Fe^{2+} and Mn^{2+} can mediate rRNA folding and
translation of active protein has revealed that these "prebiotic" divalent metals can facilitate the entire
central dogma of molecular biology (DNA→RNA→protein). These findings raise important questions
about evolutionary and physiological roles for Fe^{2+} and Mn^{2+} in ancient and extant biological systems.
Were Mg^{2+}, Fe^{2+}, and Mn^{2+} collaborators as cofactors on the ancient Earth, when Fe^{2+} and Mn^{2+} were
more abundant (1-5), and Mg^{2+} was less abundant (2), than today? What was the role of Fe^{2+} and Mn^{2+} in
the origin and early evolution of the translational system? Finally, what are the implications for ribosome-
bound Fe^{2+} in oxidative damage and disease (50, 51)?

Materials and Methods
179

rRNA folding via SHAPE. SHAPE (28, 32, 33) was conducted on the ~2900 nt *Thermus thermophilus*
23 rRNA (LSU) in 250 mM monovalent cation (Na$^+$ or K$^+$) to favor formation of secondary structure, and
in 250 mM Na$^+$ or K$^+$ plus various divalent cation (10 mM MgCl$_2$, 2.5 mM FeCl$_2$, or 2.5 mM MnCl$_2$) to
favor tertiary interactions. These divalent cation concentrations are sufficient to fold rRNA. To keep
rRNA samples from O$_2$, solutions of rRNA alone or 200 mM NaOAc or KOAc plus 50 mM Na-HEPES
(pH 8) or K-HEPES (pH 8) and divalent cations were lyophilized and transferred into an anoxic chamber
with a 98% Ar and 2% H$_2$ atmosphere. The rRNA solutions were rehydrated with nuclease-free, degassed
water, and added to the dried salts to achieve the appropriate concentrations. After rRNA modification
reactions, divalent cations were removed by chelating beads. Samples were removed from the anoxic
chamber before reverse transcription and analysis by capillary electrophoresis as in ref. (33). Essentially
identical SHAPE profiles were observed with Na$^+$ or K$^+$ alone (***SI Appendix,* Fig. S1**), as previously
described (32, 52); and for monovalent cations in combination Mg^{2+}, Fe^{2+}, or Mn^{2+} (**Fig. 2; *SI Appendix,***
Fig. S1). Nucleotides were classified as exhibiting a significant change in SHAPE reactivity if the
difference between the initial reactivity (in Na$^+$) and final reactivity (in Na$^+$/Mg^{2+}, Na$^+$/Fe^{2+}, or Na/Mn^{2+})
was >0.3 SHAPE units. To compare the Mg^{2+}-, Fe^{2+}-, and Mn^{2+}- responsiveness of specific nucleotides,
we binned nucleotides into three categories (increased, decreased, or little/no change) based on their
general SHAPE reactivity response to each divalent cation (SHAPE data are found in **Datasets S1 and**
S2).

***In vitro* translation.** Each 30 µL reaction contained 2 µM (4.5 µL of 13.3 µM stock) *E. coli* ribosomes in
10 mM Mg^{2+} (New England Biolabs, Ipswich MA, USA; catalog # P0763S), 3 µL factor mix (with RNA
polymerase, and transcription/translation factors in 10 mM Mg^{2+}) from the PURExpress® Δ Ribosome
Kit (New England Biolabs E3313S), 0.1 mM amino acid mix (Promega, Madison WI, USA; catalog #
L4461), and 0.2 mM tRNAs from *E. coli* MRE 600 (Sigma-Aldrich, St. Louis MO, USA; product #
TRNAMRE-RO). Thus, a total of 2.5 mM "background" Mg^{2+} was present in each reaction (***SI Appendix,***
Fig. S2a). To remove the background Mg^{2+}, we exchanged the buffer of the ribosome and factor mix
using centrifugal filter units. Thirty microliters of either ribosome solution or factor mix was added to an
Amicon Ultra 0.5 mL centrifugal filter (Millipore-Sigma), followed by 450 µL of divalent-free buffer (20
mM HEPES pH 7.6, 30 mM KCl, and 7 mM β-mercaptoethanol). Samples where spun at 14,000 x g at
4°C until the minimum sample volume (~15 µL) was reached. The samples were resuspended in 450 µL
of divalent-free buffer and centrifugation was repeated. The samples were then transferred to new tubes
and 15 µL of divalent-free buffer was added to bring the volume to 30 µL. This process decreased Mg^{2+}
concentrations in the ribosome and factor mix from 10 mM to 10-30 µM Mg^{2+}, resulting in 4-6 µM Mg^{2+}
in each reaction (***SI Appendix,* Figs. S2b, S2c**).

Translation experimental conditions. All reactions (30 µL total volume) were assembled and incubated
in an anoxic chamber. Divalent cation salts (MgCl$_2$, FeCl$_2$, MnCl$_2$, Zn(OAc)$_2$, CoCl$_2$, CuSO$_4$) were added
to 7 mM final concentration, with the exception of MgCl$_2$ and FeCl$_2$, which were tested over a range of
concentrations (1, 3, 4, 5, 6, 7, 8, 9, and 11 mM; ***SI Appendix,* Fig. S2**). Solutions were clear, with no
indication of metal precipitate, suggesting that reduced, divalent metals cations were the primary
chemical species. All experiments were assembled in the following order: dihydrofolate reductase
(DHFR) mRNA (~5 µg per 30 µL reaction), factor mix, ribosomes, amino acids, tRNA, nuclease-free
H$_2$O, reaction buffer (see ***SI Appendix*** for details on mRNA template and reaction buffer recipe).
Changing the order of reactant addition did not affect translational activity. Reactions were run in
triplicate on a 37°C heat block for up to 120 minutes. Reactions were quenched on ice and stored on ice
until they were assayed for protein synthesis.

Protein activity assay. Protein synthesis was measured using a DHFR assay kit (Sigma-Aldrich product
CS0340), which measures the oxidation of NADPH (60 mM) to NADP$^+$ by dihydrofolic acid (51 µM).
Assays were performed by adding 5 µL of protein synthesis reaction to 995 µL of 1x assay buffer. The

NADPH absorbance peak at 340 nm (Abs_{340}) was measured in 15 s intervals over 2.5 min. The slope of
the linear regression of Abs_{340} vs. time was used to determine protein activity (Abs_{340} min^{-1}). Different
counter ions (Cl^-, CH_3COO^-, SO_4^{2-}) had no effect on protein synthesis from mRNA. To our knowledge, no
dependence on, nor inhibitory effect of Mg^{2+} or Fe^{2+} exists for DHFR. We confirmed this by varying the
metal concentrations in our assay reaction, which had no effect on DHFR activity.
235
Ribosome metal content. The Fe and Mn content of *E. coli* ribosomes was measured by total reflection
X-ray fluorescence (TRXF) spectroscopy after the ribosomes were incubated in 7 mM $FeCl_2$ or 7 mM
$MnCl_2$. See *SI Appendix* for additional details.

Quantum mechanical calculations. The atomic coordinates of a Mg^{2+}-rRNA clamp were initially
extracted from the X-ray structure of the *Haloarcula marismortui* LSU (PDB 1JJ2) (53). The free 5′ and
3′ termini of the phosphate groups were capped with methyl groups in lieu of the remainder of the RNA
polymer, and hydrogen atoms were added, where appropriate (*SI Appendix,* **Fig. S4**). Additional details
on calculations adapted from previous publications (12, 22) are described in *SI Appendix*.

ACKNOWLEDGMENTS. This research was supported by the National Aeronautics and Space
Administration grants NNX14AJ87G, NNX16AJ28G, and NNX16AJ29G. We acknowledge helpful
discussions with Corinna Tuckey of New England Biolabs, and with Eric B. O'Neill and Claudia
Montllor Albalate of Georgia Institute of Technology. We thank Michael Goodisman for access to a
capillary electrophoresis instrument.
249
Competing interests. The authors declare that they have no competing interests with the contents of this
article.

Author contributions: MSB, TKL, JCB, ARR, NVH, LDW, and JBG conceived and designed the
experiments. MSB, TKL, and JWH collected data and performed analysis. ASP contributed to analysis.
MSB, TKL, LDW, and JBG wrote the paper with editorial input from all authors.

Table 1. Structural and functional roles for select divalent cations in the translation system. All
biomolecules in the table have been shown to require Mg^{2+} and may also be active with Fe^{2+} or Mn^{2+}.
"n.a." indicates that data are not available.

Translation system component(s)	Location of divalent ion	Role of divalent cation	Optimal $[Mg^{2+}]$ (mM)
Ribosome			
LSU/SSU	M^{2+}-rRNA clamps (12)	Mediates and maintains folding/structure of rRNAs	~10 (34)
LSU	Dinuclear microclusters (13)	Frames peptidyl transferase center (PTC)	~10 (34)
LSU/SSU	LSU/SSU interface (27)	Mediates docking of mRNA to SSU and association of SSU with LSU	~10 (34)
SSU/mRNA	Critical bend in mRNA between the P-site and A-site codons (16, 54)	Maintains correct reading frame mRNA	~10 (34)
A-site tRNA/ P-site tRNA	tRNA-tRNA interface (27)	Stabilize tRNAs in the PTC	~10 (34)
LSU/tRNA	rRNA-tRNA interface (27)	Stabilize rRNA-tRNA in the PTC	~10 (34)
Auxiliary			
EF-Tu	GTP binding site (55)	Stabilizes the transition state	5-15 (56)
EF-G	GTP binding site (57)	Stabilizes the transition state	n.a.
Aminoacyl-tRNA synthetases	ATP binding site (58)	Stabilizes the transition state	>1 (59)
Methionyl-tRNA transformylase	ATP binding site (60)	Stabilizes the transition state	7 (60)
Creatine kinase	NTP binding site (61)	Stabilizes the transition state	~5 (61)
Myokinase	Acceptor NDP binding site (62)	Stabilizes the transition state	~3 (45)
Nucleoside-diphosphate kinase	NTP binding site (63)	Stabilizes the transition state	>1 (63)
Pyrophosphatase	Active site (64)	Stabilizes the transition state	>7 (65)

259
260

261   **References**

262   1.   Derry LA (2015) Causes and consequences of mid-Proterozoic anoxia. *Geophys. Res. Lett.*
263   42(20):8538-8546.
264   2.   Jones C, Nomosatryo S, Crowe SA, Bjerrum CJ, & Canfield DE (2015) Iron oxides, divalent
265   cations, silica, and the early earth phosphorus crisis. *Geology* 43(2):135-138.
266   3.   Poulton SW & Canfield DE (2011) Ferruginous conditions: a dominant feature of the ocean
267   through Earth's history. *Elements* 7(2):107-112.
268   4.   Holland HD (1984) *The Chemical Evolution of the Atmosphere and Oceans* (Princeton University
269   Press, Princeton, N.J.).
270   5.   Johnson JE, Webb SM, Ma C, & Fischer WW (2016) Manganese mineralogy and diagenesis in
271   the sedimentary rock record. *Geochem. Cosmochim. Acta* 173:210-231.
272   6.   Ramakrishnan V (2002) Ribosome structure and the mechanism of translation. *Cell* 108(4):557-
273   572.
274   7.   Yonath A (2002) The search and its outcome: high-resolution structures of ribosomal particles
275   from mesophilic, thermophilic, and halophilic bacteria at various functional states. *Annu. Rev.*
276   *Biophys. Biomol. Struct.* 31(1):257-273.
277   8.   Fox GE (2010) Origin and evolution of the ribosome. *Cold Spring Harb. Perspect.* 2(9):a003483.
278   9.   Bernier CR, Petrov AS, Kovacs NA, Penev P, & Williams LD (2018) Translation: the universal
279   structural core of life. *Mol. Biol. Evol.* 35(8):2065–2076.
280   10.   Klein DJ, Moore PB, & Steitz TA (2004) The contribution of metal ions to the structural stability
281   of the large ribosomal subunit. *RNA* 10(9):1366-1379.
282   11.   Cusack S (1997) Aminoacyl-tRNA synthetases. *Curr. Opin. Struct. Biol.* 7(6):881-889.
283   12.   Petrov AS, Bowman JC, Harvey SC, & Williams LD (2011) Bidentate RNA–magnesium clamps:
284   On the origin of the special role of magnesium in RNA folding. *RNA* 17(2):291-297.
285   13.   Hsiao C & Williams LD (2009) A recurrent magnesium-binding motif provides a framework for
286   the ribosomal peptidyl transferase center. *Nucleic Acid Res.* 37(10):3134-3142.
287   14.   Schuwirth BS, *et al.* (2005) Structures of the bacterial ribosome at 3.5 Å resolution. *Science*
288   310(5749):827-834.
289   15.   Demeshkina N, Jenner L, Westhof E, Yusupov M, & Yusupova G (2012) A new understanding
290   of the decoding principle on the ribosome. *Nature* 484(7393):256-259.
291   16.   Selmer M, *et al.* (2006) Structure of the 70S ribosome complexed with mRNA and tRNA.
292   *Science* 313(5795):1935-1942.
293   17.   Petrov AS, *et al.* (2012) RNA–magnesium–protein interactions in large ribosomal subunit. *J.*
294   *Phys. Chem. B* 116(28):8113-8120.
295   18.   Pyle A (2002) Metal ions in the structure and function of RNA. *J. Biol. Inorg. Chem.* 7(7):679-
296   690.
297   19.   Pyle AM (1993) Ribozymes: a distinct class of metalloenzymes. *Science* 261(5122):709-714.
298   20.   Ward WL, Plakos K, & DeRose VJ (2014) Nucleic acid catalysis: metals, nucleobases, and other
299   cofactors. *Chem. Rev.* 114(8):4318-4342.
300   21.   Okafor CD, Bowman JC, Hud NV, Glass JB, & Williams LD (2018) Folding and catalysis near
301   life's origin: support for Fe2+ as a dominant divalent cation. *Prebiotic Chemistry and Chemical*
302   *Evolution of Nucleic Acids*, (Springer), pp 227-243.
303   22.   Athavale SS, *et al.* (2012) RNA folding and catalysis mediated by iron (II). *PLoS One*
304   7(5):e38024.
305   23.   Popović M, Fliss PS, & Ditzler MA (2015) *In vitro* evolution of distinct self-cleaving ribozymes
306   in diverse environments. *Nucleic Acids Res.* 43(14):7070-7082.
307   24.   Okafor CD, *et al.* (2017) Iron mediates catalysis of nucleic acid processing enzymes: support for
308   Fe (II) as a cofactor before the great oxidation event. *Nucleic Acids Res.* 45(7):3634-3642.

309   25.   Tabor S & Richardson CC (1989) Effect of manganese ions on the incorporation of
310   dideoxynucleotides by bacteriophage T7 DNA polymerase and *Escherichia coli* DNA polymerase
311   I. *Proc. Natl. Acad. Sci.* 86(11):4076-4080.
312   26.   Litman RM (1971) The differential effect of magnesium and manganese ions on the synthesis of
313   poly (dGd· C) and *Micrococcus luteus* DNA by *Micrococcus luteus* DNA polymerase. *J. Mol.*
314   *Biol.* 61(1):1-23.
315   27.   Dann III CE*, et al.* (2007) Structure and mechanism of a metal-sensing regulatory RNA. *Cell*
316   130(5):878-892.
317   28.   Mortimer SA & Weeks KM (2007) A fast-acting reagent for accurate analysis of RNA secondary
318   and tertiary structure by SHAPE chemistry. *J. Am. Chem. Soc.* 129(14):4144-4145.
319   29.   Athavale SS*, et al.* (2012) Domain III of the *T. thermophilus* 23S rRNA folds independently to a
320   near-native state. *RNA* 18(4):752-758.
321   30.   Hsiao C*, et al.* (2013) Molecular paleontology: a biochemical model of the ancestral ribosome.
322   *Nucleic Acids Res.* 41(5):3373-3385.
323   31.   Lanier KA, Athavale SS, Petrov AS, Wartell R, & Williams LD (2016) Imprint of ancient
324   evolution on rRNA folding. *Biochemistry* 55(33):4603-4613.
325   32.   Wilkinson K, Merino E, & Weeks K (2006) Selective 2'-hydroxyl acylation analyzed by primer
326   extension (SHAPE): quantitative RNA structure analysis at single nucleotide resolution. *Nat.*
327   *Protoc.* 1(3):1610-1616.
328   33.   Lenz TK, Norris AM, Hud NV, & Williams LD (2017) Protein-free ribosomal RNA folds to a
329   near-native state in the presence of Mg^{2+}. *RSC Adv.* 7(86):54674-54681.
330   34.   Kigawa T*, et al.* (1999) Cell-free production and stable-isotope labeling of milligram quantities of
331   proteins. *FEBS Lett.* 442(1):15-19.
332   35.   Goldberg A (1966) Magnesium binding by *Escherichia coli* ribosomes. *J. Mol. Biol.* 15(2):663-
333   673.
334   36.   Pronczuk A, Baliga B, & Munro H (1968) Effect of nucleoside triphosphate and magnesium ion
335   concentration on the stability and function of rat liver polysomes *in vitro*. *Biochem. J.*
336   110(4):783-788.
337   37.   Brion P & Westhof E (1997) Hierarchy and dynamics of RNA folding. *Annu. Rev. Biophys.*
338   *Biomol. Struct.* 26:113-137.
339   38.   Tinoco I, Jr. & Bustamante C (1999) How RNA folds. *J. Mol. Biol.* 293(2):271-281.
340   39.   Bock CW, Katz AK, Markham GD, & Glusker JP (1999) Manganese as a replacement for
341   magnesium and zinc: functional comparison of the divalent ions. *J. Amer. Chem. Soc.*
342   121(32):7360-7372.
343   40.   Imlay JA (2014) The mismetallation of enzymes during oxidative stress. *J. Biol. Chem.*
344   289(41):28121-28128.
345   41.   Zheng H, Shabalin IG, Handing KB, Bujnicki JM, & Minor W (2015) Magnesium-binding
346   architectures in RNA crystal structures: validation, binding preferences, classification and motif
347   detection. *Nucleic Acids Res.* 43(7):3789-3801.
348   42.   Bowman JC, Lenz TK, Hud NV, & Williams LD (2012) Cations in charge: magnesium ions in
349   RNA folding and catalysis. *Curr. Opin. Struct. Biol.* 22(3):262-272.
350   43.   Hensley MP, Tierney DL, & Crowder MW (2011) Zn (II) binding to Escherichia coli 70S
351   ribosomes. *Biochemistry* 50(46):9937-9939.
352   44.   Craine J & Peterkofsky A (1976) Studies on arginyl-tRNA synthetase from *Escherichia coli* B.
353   Dual role of metals in enzyme catalysis. *J. Biol. Chem.* 251(1):241-246.
354   45.   Ito Y, Tomasselli AG, & Noda LH (1980) ATP: AMP phosphotransferase from baker's yeast:
355   purification and properties. *Eur. J. Biochem.* 105(1):85-92.
356   46.   Da Silva JF & Williams RJP (2001) *The Biological Chemistry of the Elements: the Inorganic*
357   *Chemistry of Life* (Oxford University Press) Second Ed.
358   47.   Daly MJ*, et al.* (2004) Accumulation of Mn(II) in *Deinococcus radiodurans* facilitates gamma-
359   radiation resistance. *Science* 306(5698):1025-1028.

360   48.   Outten CE & O'Halloran TV (2001) Femtomolar sensitivity of metalloregulatory proteins
361   controlling zinc homeostasis. *Science* 292(5526):2488-2492.
362   49.   Beauchene NA, *et al.* (2017) O2 availability impacts iron homeostasis in *Escherichia coli*. *Proc.*
363   *Natl. Acad. Sci.* 14 (46):12261-12266.
364   50.   Honda K, *et al.* (2005) Ribosomal RNA in Alzheimer disease is oxidized by bound redox-active
365   iron. *J. Biol. Chem.* 280(22):20978-20986.
366   51.   Zinskie JA, *et al.* (2018) Iron-dependent cleavage of ribosomal RNA during oxidative stress in
367   the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 293(37):14237-14248.
368   52.   Holmstrom E, Fiore J, & Nesbitt D (2012) Thermodynamic origins of monovalent facilitated
369   RNA folding. *Biochemistry* 51(18):3732-3743.
370   53.   Ban N, Nissen P, Hansen J, Moore PB, & Steitz TA (2000) The complete atomic structure of the
371   large ribosomal subunit at 2.4 Å resolution. *Science* 289(5481):905-920.
372   54.   Keedy HE, Thomas EN, & Zaher HS (2018) Decoding on the ribosome depends on the structure
373   of the mRNA phosphodiester backbone. *Proc Natl Acad Sci*:DOI: 10.1073/pnas.1721431115.
374   55.   Kjeldgaard M, Nissen P, Thirup S, & Nyborg J (1993) The crystal structure of elongation factor
375   EF-Tu from *Thermus aquaticus* in the GTP conformation. *Structure* 1(1):35-50.
376   56.   Campuzano S & Modolell J (1981) Effects of antibiotics, N‑Acetylaminoacyl‑tRNA and other
377   agents on the Elongation‑Factor‑Tu dependent and ribosome‑dependent GTP hydrolysis
378   promoted by 2′(3′)‑O‑l‑Phenylalanyladenosine. *Eur. J. Biochem.* 117(1):27-31.
379   57.   Chen Y, Feng S, Kumar V, Ero R, & Gao Y-G (2013) Structure of EF-G–ribosome complex in a
380   pretranslocation state. *Nature Struct. Mol. Biol.* 20(9):1077-1084.
381   58.   Lovgren TNE, Heinonen J, & Loftfield RB (1978) Mechanism of aminoacylation of transfer
382   ribonucleic-acid. Role of magnesium and spermine in synthesis of isoleucyl-transfer RNA. *J.*
383   *Biol. Chem.* 253(19):6702-6710.
384   59.   Thiebe R (1975) Aminoacylation of tRNA. Magnesium requirement and spermidine effect. *FEBS*
385   *Lett.* 51(1-2):259-261.
386   60.   Blanquet S, Dessen P, & Kahn D (1984) Properties and specificity of methionyl-tRNAfMet
387   formyltransferase from *Escherichia coli*. *Methods Enzymol.* 106:141-152.
388   61.   Morin L (1977) Creatine kinase: stability, inactivation, reactivation. *Clin. Chem.* 23(4):646-652.
389   62.   Vasavada K, Kaplan J, & Rao BN (1984) Analysis of phosphorus-31 NMR spectra of enzyme-
390   bound reactants and products of adenylate kinase using density matrix theory of chemical
391   exchange. *Biochemistry* 23(5):961-968.
392   63.   Munoz-Dorado J, Inouye S, & Inouye M (1990) Nucleoside diphosphate kinase from *Myxococcus*
393   *xanthus*. II. Biochemical characterization. *J. Biol. Chem.* 265(5):2707-2712.
394   64.   Kankare J, *et al.* (1994) The structure of *E. coli* soluble inorganic pyrophosphatase at 2.7 Å
395   resolution. *Protein Eng. Des. Sel.* 7(7):823-830.
396   65.   Bloch-Frankenthal L (1954) The role of magnesium in the hydrolysis of sodium pyrophosphate
397   by inorganic pyrophosphatase. *Biochem. J.* 57(1):87.
398   66.   Bernier CR, *et al.* (2014) RiboVision suite for visualization and analysis of ribosomes. *Faraday*
399   *Disc* 169:195-207.

400   **Figure Legends**
401
Fig. 1. Divalent cations serve many structural and functional roles in the ribosome. Mg^{2+} ions: **a)** form
bidentate clamps with adjacent phosphate groups of rRNA, **b)** form dinuclear microclusters that frame the
rRNA of the PTC, **c)** stabilize the LSU-SSU interface, **d)** stabilize a functional kink in mRNA, **e)** stabilize
association of tRNA (teal) with 23S rRNA (beige carbon atoms), and **f)** stabilize association of mRNA
(green) with 16S rRNA (beige carbon atoms). Thick dashed lines are first shell RNA interactions of Mg^{2+}.
Dotted lines indicate second shell interactions. Images are of the *Thermus thermophilus* ribosome (PDB
ID: 1VY4). This figure was generated with the program RiboVision (66).

Fig. 2. SHAPE reactivities mapped onto the *Thermus thermophilus* LSU rRNA secondary structure in **a)**
Na^+, **b)** Na^+/Fe^{2+}, or **c)** Na^+/Mg^{2+}. Key functional elements are labeled in panel a, and the scale in panel
(a) applies to panels b and c. **d)** Fe^{2+}-induced changes (ΔFe^{2+}) in SHAPE reactivity calculated by
subtracting Na^+ data from Na^+/Fe^{2+} data for each nucleotide, and **e)** Mg^{2+}-induced changes (ΔMg^{2+}) in
SHAPE reactivity calculated by subtracting Na^+ data from Na^+/Mg^{2+} data for each nucleotide. The scale
shown for panel d also applies to panel e. Positive values indicate increased SHAPE reactivity in presence
of the divalent cation, while negative values denote decreased reactivity. Regions where data are not
available (5' and 3' ends) are grey. These figures were generated with the program RiboVision (66). The
L11 binding region, where the greatest discrepancy between Fe^{2+} and Mg^{2+} is observed, is indicated with
an arrow.

Fig. 3. Mg^{2+} and Fe^{2+} stimulate translational activity over a range of concentrations. The activity of the
translation product (dihydrofolate reductase, which catalyzes the oxidation of NADPH, with a maximum
absorbance at 340 nm) was used as a proxy for protein production. Translation reactions were run for 120
minutes. All translation reactions contained 2.5 mM background Mg^{2+}, to which varying amounts of
additional Mg^{2+} or Fe^{2+} were added. The error bars for triplicate experiments (n=3) are plotted as the
standard error of the mean.

Fig. 4. Fe^{2+} consistently supports 50-80% of the translational activity as Mg^{2+} when the translation
experiments are run for 15-120 minutes. The activity of the translation product (dihydrofolate reductase,
which catalyzes the oxidation of NADPH, with a maximum absorbance at 340 nm) was used as a proxy
for protein production. All translation reactions contained 2.5 mM background Mg^{2+}, to which 7 mM
additional Mg^{2+} or Fe^{2+} were added, totaling to 9.5 mM divalent cation. The error bars for triplicate
experiments (n=3) are plotted as the standard error of the mean.

Fig. 5. Mn^{2+} can support translation after removal of background Mg^{2+}. **a)** Reactions prepared with
washed *E. coli* ribosomes, reducing the background Mg^{2+} to 1 mM, to which 7, 9, or 11 mM additional
Mg^{2+}, Fe^{2+}, or Mn^{2+} were added, totaling 8, 10, or 12 mM divalent cation (M^{2+}). **b)** Reactions prepared
using washed *E. coli* ribosomes and washed factor mix, which reduced the background Mg^{2+} to the low
μM level, to which 8, 10, or 12 mM additional, Mg^{2+}, Fe^{2+}, or Mn^{2+} were added. The activity of the
translation product (dihydrofolate reductase, which catalyzes the oxidation of NADPH, with a maximum
absorbance at 340 nm) was used as a proxy for protein production. The error bars for triplicate
experiments   (n=3)   are   plotted   as   the   standard   error   of   the   mean.

a) Mg²⁺-rRNA clamp

b) dinuclear microcluster

c) LSU-SSU interface

d) mRNA bend

e) rRNA-tRNA interface

f) rRNA-mRNA interface