**A systematic review of Hepatitis B virus (HBV) drug and vaccine escape mutations in Africa: a call for urgent action**

Jolynne Mokaya[1], Anna L McNaughton[1], Martin J Hadley[2], Apsotolos Beloukas[3], Anna-Maria Geretti[3], Dominique Goedhals[4], Philippa C Matthews[1,5*]

[1] Nuffield Department of Medicine, Medawar Building,

South Parks Road, Oxford OX1 3SY, UK

[2] Oxford University Academic IT Department,

13 Banbury Road, Oxford OX2 6NN, UK

[3] Institute of Infection and Global Health, University of Liverpool, Liverpool L69 7BE,

UK

[4] Division of Virology, University of the Free State/National Health Laboratory Service,

PO Box 339(G23), Bloemfontein 9300, Republic of South Africa

[5] Department of Microbiology and Infectious Diseases, Oxford University Hospitals

NHS Foundation Trust, Headley Way, Headington, Oxford OX3 9DU, UK

**\*Corresponding author**: philippa.matthews@ndm.ox.ac.uk

**Short title:** Drug and vaccine resistance in HBV in Africa

**Keywords:** Hepatitis B virus, treatment, antiviral, nucleotide analogues, mutations, drug resistance, vaccine escape, Africa

**ABSTRACT:**

International sustainable development goals for the elimination of viral hepatitis as a public health problem by 2030 highlight the pressing need to optimize strategies for prevention, diagnosis and treatment. Selected or transmitted resistance associated mutations (RAMs) and vaccine escape mutations (VEMs) in hepatitis B virus (HBV) may reduce the success of existing treatment and prevention strategies. These issues are particularly pressing for many settings in Africa where there is high HBV prevalence and co-endemic HIV infection, but lack of robust epidemiological data and limited education, diagnostics and clinical care. The prevalence, distribution and impact of RAMs and VEMs in these populations are neglected in the current literature. We therefore set out to assimilate data for sub-Saharan Africa through a systematic literature review and analysis of published sequence data, and present these in an on-line database (https://livedataoxford.shinyapps.io/1510659619-3Xkoe2NKkKJ7Drg/). The majority of the data were from HIV/HBV coinfected cohorts. The commonest RAM was rtM204I/V, either alone or in combination with compensatory mutations, and reported in both treatment-naïve and treatment-experienced adults. We also identified the suite of mutations rtM204V/I + rtL180M + rtV173L that is reported in association with vaccine escape, in over 1/3 of cohorts. Although tenofovir has a high genetic barrier to drug resistance, it is of concern that emerging data suggest polymorphisms that may be associated with resistance, although the precise impact of these is unknown. Overall, there is an urgent need for improved diagnostic screening, enhanced laboratory assessment of HBV before and during therapy, and sustained roll out of tenofovir in preference to lamividine. Further data are urgently needed in order to inform population and individual approaches to HBV diagnosis, monitoring and therapy in these highly vulnerable settings.

**Author's summary**

The Global Hepatitis Health Sector Strategy is aiming for the elimination of viral hepatitis as a public health threat by 2030. However, drug and vaccine resistance may reduce the success of existing treatment and prevention strategies. In the current literature, the prevalence, distribution and impact of hepatitis B drug and vaccine resistance in many settings in Africa are neglected, despite the high prevalence of hepatitis B and co-endemic HIV infection. This systematic review describes the frequency, prevalence and co-occurrence of mutations associated with hepatitis B drug and vaccine resistance in Africa. The findings suggests a high prevalence of these mutations in some populations in sub-Saharan Africa. Scarce resources have contributed to the lack of diagnostic screening, inconsistent supply of hepatitis B drugs, and poor access to clinical monitoring, all of which contribute to drug and vaccine resistance. Sustainable long-term investment is required to expand consistent drug and vaccine supply, to provide screening infection and for drug resistance, and to provide appropriate targeted clinical monitoring for treated patients.

**ABBREVIATIONS**

- 3TC - lamivudine

- ADV – Adefovir

- ART – antiretroviral therapy

- AZT – zidovudine

- CHB – chronic hepatitis B infection

- DNA – deoxyribonucelic acid

- ETV – entecavir (ETV)

- HBsAg – Hepatitis B surface antigen

- HBV – Hepatitis b virus

- LdT – telbivudine

- NA – Nucleos(t)ide analogue

- ORF – overlapping reading frame

- Pol – polymerase

- RAM – Resistance Associated Mutation

- RNA – ribonucleic acid

- RT – reverse transcriptase

- TAF – tenofovir alafenamide

- TDF – tenofovir disoproxil fumarate

- VEM – Vaccine escape mutation

- WHO – World Health Organisation

**INTRODUCTION**

In 2015, the World Health Organisation (WHO) estimated that 3.5% of the world's population (257 million people) were living with Hepatitis B virus (HBV) infection, resulting in 887,000 deaths each year, mostly from complications including cirrhosis and hepatocellular carcinoma (HCC) (1). United Nations Sustainable Development Goals set out the challenge of 'elimination of viral hepatitis as a public health threat by the year 2030' (2). One of the existing strategies in the elimination toolbox is use of antiviral drugs in the form of nucleos(t)ide analogues (NAs). Suppression of viraemia not only reduces inflammatory liver disease in the individual receiving treatment but also reduces the risk of transmission. However, the emergence of HBV resistance-associated mutations (RAMs) is a potentially significant concern for the success of this strategy.

Africa is the continent with the second largest number of individuals with chronic HBV infection (CHB), having 6.1% of the adult population infected (1). However, there is very little commitment and resources invested into the burden of this disease, and many barriers are contributing to the epidemic (3,4). Globally, less than 10% of the population with CHB infection are diagnosed, with an even smaller proportion on treatment (1,4). This proportion is likely to be even lower in Africa. The situation in Africa is further complicated by the substantial public health challenge of coendemic human immunodeficiency virus (HIV) and HBV; high coinfection rates worsen the prognosis in dually infected individuals (5). There is also a lack of robust epidemiological data on HBV from Africa (3,4).

Widespread use of antiretroviral therapy (ART) for HIV, incorporating NAs that also

have activity against HBV, may have an impact on HBV both through improved rates

of viraemic suppression, and also potentially by driving the selection of RAMs. The

WHO recommends HBsAg screening for all HIV-1 infected individuals prior to ART

initiation and for all pregnant women during antenatal visits, to improve the clinical

outcomes of people living with CHB and to reduce the incidence of new cases (6).

However, screening of HBsAg is not routinely performed in many settings in Africa

due to cost and lack of availability. HBV infected patients either remain untreated (in

the setting of monoinfection), or are exposed to antiviral drugs without proper

monitoring, putting them at risk of developing RAMs (in the setting of HIV

coinfection) (4,7–10).

HBV is a DNA virus that replicates via an RNA intermediate, with reverse

transcriptase (RT) catalysing the transcription of RNA into DNA (7). NAs that inhibit

RT are therefore used to prevent HBV replication, including lamivudine (3TC),

entecavir (ETV) and tenofovir disoproxil fumarate (TDF), with historical use of other

agents including telbivudine (LdT) and adefovir (ADV)(6,11). Choice of agent is

determined by a number of factors including potency, availability, cost, side-effects,

drug interactions, safety in pregnancy, and genetic barrier to resistance (4).

Emergence of mutations happens because the RT enzyme lacks proofreading

function required to repair errors during transcription (7); when these mutations

confer a selective advantage by allowing the virus to escape the effect of drug

therapy, they will become amplified in the viral population. Some RAMs confer

resistance to one agent only, while others are associated with resistance to several

agents (Fig 1). In Africa, the choice of agent is usually limited to 3TC and/or TDF.

**Fig 1: HBV drug resistance associated mutations (RAMs), vaccine escape mutations (VEMs) and mutations associated with Hepatitis B immunoglobulin (HBIg) resistance.** HBV genes are shown in the coloured ovals. TDF = tenofovir, ETV = entecavir, 3TC = lamivudine. This figure incorporates data from eight studies; three were identified by the systematic review presented in this manuscript (45,49,59) and five from the wider literature (7,13,14,20,27).

3TC was originally seen as a major breakthrough in treating HBV (12). However, it is now known to have a low genetic barrier to resistance and its long-term effectiveness is limited as a result of resistance mutations in the 'YMDD' motif (tyrosine, methionine, aspartate, aspartate; amino acids 203-206) in domain C of the viral polymerase (Pol) and associated upstream compensatory mutations in Pol domains A, B and in the B-C interdomain (7,13,14). Among chronic HBV monoinfected patients, incidence of HBV resistance to 3TC is as high as 20% per year, whereas in HIV/HBV coinfected patients, this can reach 90% over 5 years of treatment(5). 3TC has also been associated with the induction of cross-resistance to emtricitabine (FTC), LdT, and ETV, thus reducing the options for subsequent treatment (10).

TDF is widely used in treatment of both HIV and HBV and is generally well tolerated. TDF has a high genetic barrier to resistance and maintains effective suppression of HBV in both monoinfected and HIV/HBV coinfected individuals (5,7,10,15,16). Although it has a recognised association with nephrotoxicity in HIV treatment, current literature suggests it may be better tolerated in HBV infection (11). African populations have a higher background of renal disease (17) and could be potentially

more vulnerable to nephrotoxitiy from TDF(18). A tenofovir prodrug, tenofovir alafenamide (TAF), delivers equally potent viraemic suppression at lower plasma levels, and is therefore thought to be associated with reduced nephrotoxicity(19). However, TAF is not widely available. TDF resistance is not well characterised, but there are emerging data from *in vitro* studies associating Pol mutations rtA194T and rtN236T with decreased susceptibility (11,15). Virological breakthrough on TDF therapy has been reported in two patients harbouring rtS78T/sC69 mutations (20), and in another patient with multisite polymerase mutations; rtL80M, rtL180M, rtM204V/I, rtA200V, rtF221Y, rtS223A, rtT184A/L, rtR153Q, and rtV191I (21). The significance of these mutations needs to be further explored in clinical studies.

Current guidelines recommend the use of first-line ART regimens containing TDF in patients coinfected with HIV and HBV (22). In accordance with this, there is evidence of a recent increase in the prescription of TDF in sub-Saharan Africa (sSA) (23). However, it is likely that many ART regimens in sSA continue to include 3TC as the only agent active against HBV (7,8). In South Africa, second line ART substitutes TDF for Zidovudine (AZT) leaving only 3TC coverage for HBV. Among HBV/HIV coinfected children in South Africa treated with regimens including 3TC and/or TDF, HBV viraemia has been demonstrated highlighting potential underlying HBV drug resistance (24).

ETV is another active agent, and is safe and well tolerated. However it is not active against HIV and therefore has to be added to ART regimens rather than being part of the primary backbone, is not recommended in pregnancy, and is not routinely available in most African settings(25). Resistance arises more commonly in the

context of prior 3TC exposure (11,26), which may limit its future potential in Africa,

particularly in HIV endemic populations.

As a component of the Expanded Programme on Immunization (EPI), HBV

preventive vaccines have been rolled out in Africa since 1995 (4). Mutations in HBV

Pol can also lead to amino acid changes in the Surface (S) protein due to

overlapping reading frames (ORFs) in the genome (14). Changes in the S protein

can result in vaccine escape mutants (VEMs) which can cause infection among

vaccinated individuals(14,27), and also diagnostic escape mutations which result in

false negative HBsAg testing (14). S protein mutation sG145R has been identified as

the major VEM and recently other mutations in S protein have been associated with

immune escape (14) (Fig 1).  There are no data for VEMs in Africa, but in other

settings of high endemicity VEMs can be common, as evidenced by a reported

prevalence of 28% in vaccinated HBV-infected children in Taiwan (28).

To date, no systematic review has assessed the geography and prevalence of HBV

RAMs and VEMs in Africa. An understanding of the extent to which these mutations

circulate in Africa is essential to improving HBV therapy in patients with and without

HIV coinfection. We therefore set out to describe the frequency, co-occurrence and

distribution of RAMs and VEMs in Africa, and to suggest whether changes are

needed in recommendations for laboratory diagnostics and/or approaches to drug

therapy or vaccine deployment. This will underpin further research to identify and

track relevant mutations in these populations.

**METHODS**

**Search strategy**

Between October 2017 and January 2018, we searched the published literature, in MEDLINE (PubMed; https://www.ncbi.nlm.nih.gov/pubmed), SCOPUS (https://www.elsevier.com/solutions/scopus) and EMBASE (https://www.elsevier.com/en-gb/solutions/embase-biomedical-research). Our search strategy is detailed in S1 Table and S1 Fig. The earliest paper we identified on HBV drug resistance in Africa was published in 2007. We reviewed the titles and abstracts matching the search terms and only included those relating to drug or vaccine resistance in HBV infection, including only those that presented original data and had undergone peer review. All retrieved articles were in English, therefore no exclusion in relation to language was required.

For each publication we recorded reference, publication year, study design, sample size, study population, proportion of participants who tested HBsAg+ or HBV DNA+, country, year(s) of specimen collection, genotype identified, antiviral treatment, sequencing method, gene sequenced, number of sequenced samples, participant recruitment site and sequence accession number. Data were curated using MS Excel software (Microsoft, Redmond, WA).

**RAMs reported in published sequences not represented in primary studies**

We expanded our search for evidence of RAMs by identifying publicly available HBV sequences from Africa, that had not been included in the results of our primary literature search. We used both the Hepatitis B Virus database

(https://hbvdb.ibcp.fr/HBVdb/)(29) and Hepatitis Virus Diversity Research Alignments

database (http://hvdr.bioinf.wits.ac.za/alignments/)(30).

**Analysis**

In order to determine the prevalence of RAMs and VEMs, we first reported these

using the denominator (total number of HBV positive patients) and numerator (total

number of HBV positive patients with the specified mutation) as reported in

published studies. We also pooled data by country in order to provide regional

estimates. Downloaded sequences were managed using Sequence editor, database

and analysis platform, SSE version 1.3, for analysis (31).

**Data visualisation**

We developed an R package, gene.alignment.tables, for the visualisation of the

sequence data in this study; this is available on Github (32) and can be used for

visualising generic gene sequence datasets. The package was developed by

University of Oxford's Interactive Data Network and a specific instance of the

visualisation is hosted as a Shiny app which can be viewed here:

https://livedataoxford.shinyapps.io/1510659619-3Xkoe2NKkKJ7Drg/ (33).

**RESULTS**

The initial search yielded 56 articles in MEDLINE, 150 in SCOPUS and 150 in

EMBASE. Of these; 32, 136 and 119 were excluded from search results of

MEDLINE, SCOPUS and EMBASE respectively, as they did not did not meet the

inclusion criteria. After de-duplication, 37 articles were included: 27 articles identified

from MEDLINE, SCOPUS and EMBASE were identical; five unique articles were

included from EMBASE, four from SCOPUS and one from MEDLINE. A total of 37

articles were downloaded in full (S1 Fig; S2 Table).

**Study characteristics**

Epidemiological data for HBV represented by the 37 studies we identified are

summarised in Table 1. Studies included were from Southern Africa (Botswana,

Mozambique, South Africa, Zambia and Zimbabwe), East Africa (Ethiopia, Kenya,

Malawi, Sudan and Uganda), West Africa (Cote d'Ivoire, Gambia, Ghana, Guinea-

Bissau and Nigeria) and Central Africa (Cameroon, Gabon). There was considerable

heterogeneity in recruitment protocols and exposure to anti-viral treatment. Twenty-

six studies recruited from hospitals, three studies recruited from the community

(8,34,35) and eight studies did not specify where recruitment was undertaken

(10,36–42). All studies were observational.

 **Table 1: Prevalence of HBsAg, HBeAg and HDV coinfection**

| | Study (Reference) | Country | Characteristics of study population | HIV co-infection status of cohort[a] | HBV prevalence in this cohort (reported as HBsAg prevalence unless otherwise specified) | HBeAg prevalence among HBV-positive individuals |
|---|---|---|---|---|---|---|
| **East Africa** | Deressa et al 2017 (42) | Ethiopia | Patients attending outpatient ART clinic at tertiary referral university hospital | + | 17/308 (6%) | Not reported |
| | Hundie et al 2016 (33) | Ethiopia | Stored plasma samples from HBV infected blood donors obtained from blood bank centres | ± | 391/391 (100%) | Not reported |
| | Day et al 2013 (7) | Kenya | Longitudinal cohort study of female sex workers in an urban setting | + | 11/159 (7%) | 6/11 (55%) |
| | Kim et al 2011 (8) | Kenya | Individuals from an urban centre enrolled in randomised controlled trial of adherence to ART | + | 27/389 (7%) | 24/27 (89%) |
| | Mabeya et al 2017 (44) | Kenya | Individuals seeking treatment at the comprehensive HIV Clinic at tertiary referral university hospital | + | 29/400 (7%) | Not reported |
| | Auodjane et al 2014 (45) | Malawi | Individuals starting ART treatment at a tertiary referral university hospital | + | 133/1117 (12%) | 67/133 (50%) |
| | Galluzzo et al 2012 (37) | Malawi | Pregnant women enrolled in a PMTCT study on safety and pharmacokinetics of antiretroviral drugs | + | 21/21 (100%) | 7/21 (33%) |
| | Mahgoub et al 2011 (34) | Sudan | Plasma samples from blood donors from capital city in Sudan | ± | 16/404 (4%) | Not reported |
| | Yousif et al 2014 (39) | Sudan | Individuals seeking treatment at a AIDS care unit and HIV treatment centre | + | 96/358 (27%)[b] | 32/ 50 (64%) |
| | Calisti et al 2015 (46) | Uganda | All HIV patients attending a regional referral hospital | + | 109/2820 (4%) | Not reported |

| West Africa | Boyd et al 2015 (47) | Cote d'Ivoire | Individuals enrolled in randomised multi centre trials of benefits and risks of early ART initiation | + | 259/ 2465 (11%) | 39/168 (23%) |
|---|---|---|---|---|---|---|
| | Archampon g et al 2017 (48) | Ghana | Serum samples from HBV-HIV co-infected patients collected at tertiary referral university hospital | + | 235/235 (100%) | Not reported |
| | Chadwick et al 2012 (49) | Ghana | Stored sera from all adult patients attending the HIV clinic at a tertiary referral university hospital | + | 143/371 (39%) | Not reported |
| | Geretti et al 2010 (50) | Ghana | Paired serum and plasma samples collected from HIV-infected patients attending a tertiary referral university hospital | + | 140/838 (17%) | 37/140 (26%) |
| | Ndow et al 2017 (51) | Gambia | Individuals attending HIV clinic | + | 23/187 (12%) | Not reported |
| | Stewart et al 2011 (36) | Gambia | Individuals receiving HAART; recruitment site not specified | + | 70/ 570 (12%) | 6/21 (29%) |
| | Langhoff Honge et al 2014 (52) | Guinea Bissau | Patients attending outpatient ART clinic at tertiary referral university hospital | + | 94/576 (16%) | 16/94 (17%), HDV prevalence : 18/72 (25%) |
| | Faleye et al 2015 (64) | Nigeria | Pregnant women attending antenatal clinics from two tertiary university hospitals | ± | 15/272 (6%) | Not reported |
| Central Africa | Gachara et al 2017 (53) | Cameroon | Patients attending outpatient ART health centre | + | 20/337 (6%)[c] | Not reported |
| | Kouanfack et al 2012 (54) | Cameroon | Patients attending outpatient ART clinic at tertiary hospitals | + | 54/552 (10%) | Not reported |
| | Magoro et al 2016 (55) | Cameroon | Patients attending outpatient ART health centre | + | 116/445 (26%) | 16/102 (16%) |
| | Bivigou-Mboumba et al 2016 (56) | Gabon | Patients attending outpatient ART clinic | + | 71/762 (9%) | Not reported |
| | Bivigou-Mboumba et al 2018 (43) | Gabon | Patients attending HIV care centers | + | 43/487 (9%) | Not reported |

| | | | | | | |
|---|---|---|---|---|---|---|
| **Souther n Africa** | Anderson et al 2015 (40) | Botswan a | Stored plasma samples of HIV/HBV co-infected individuals collected from studies conducted in a Research Institution | + | 81/81 (100%) | Not reported |
| | Matthews et al 2015 (62) | Botswan a | Women attending antenatal and paediatric clinics | ± | 17/443 (4%) | 16/60 (27%); HDV: negative |
| | Chambal et al 2017 (57) | Mozamb ique | Patients attending outpatient ART health centre | + | 47/518 (9%) | Not reported |
| | Wandeler et al 2016 (58) | Mozamb ique | Individuals starting ART treatment at urban clinic in Mozambique and rural clinic in Zambia | + | 78/1032 (8%) | 24/168 (14%) |
| | Andersson et al 2013 (63) | South Africa | Stored serum of women infected with HIV enrolled in an Antenatal Sentinel HIV and Syphilis Prevalence Survey | ± | 97/3089 (3%) | 17/94 (18%); HDV: negative |
| | Amponsah-Dacosta et al 2015 (35) | South Africa | Stored serum of individuals exposed to HBV participating in a health facility-based hepatitis B serosurvey conducted at a provincial level. | ± | 33/201 (16%) | Not reported |
| | Amponsah-Dacosta et al 2016 (41) | South Africa | Individuals due to HAART initiation enrolled in longitudinal study | + | 5/5 (100%) | 5/5 (100%) |
| | Hamers et al 2013 (9) | South Africa | Individuals enrolled in a multicentre prospective study of ART resistance monitoring | + | 37/175 (21%) | Not reported |
| | Gededzha et al 2016 (65) | South Africa | Stored sera from HBV infected individuals attending a tertiary referral university hospital | ± | 8/9 (89%) | Not reported |
| | Makondo et al 2012 (38) | South Africa | Stored sera from HIV infected individuals prior to ART initiation, recruitment site not specified | + | 71/298 (24%)[b] | Not reported |

| | | | | | | |
|---|---|---|---|---|---|---|
| | Matthews et al 2015 (62) | South Africa | Women attending antenatal and paediatric clinics in South Africa and Botswana | ± | 49/507 (10%) | Not reported; HDV: negative |
| | Powell et al 2015 (61) | South Africa | Stored serum samples of individuals infected with HIV receiving care at a tertiary university hospital | + | 37/394 (9%) | Not reported |
| | Selabe et al 2007 (66) | South Africa | Individuals infected with HBV admitted at tertiary University hospital | ± | 35/35 (100%) | Not reported |
| | Selabe et al 2009 (67) | South Africa | Individuals infected with HBV admitted at tertiary University hospital | - | 17/17 (100) | 9/17 (53%) |
| | Wandeler et al 2016 (58) | Zambia | Individuals starting ART treatment at urban clinic in Mozambique and rural clinic in Zambia | + | 90/797 (11%) | 24/168 (14%) |
| | Hamers et al 2013 (9) | Zambia | Individuals enrolled in a multicentre prospective study of ART resistance monitoring | + | 55/523 (11%) | Not reported |
| | Baudi et al 2017 (59) | Zimbabwe | Stored plasma samples of individuals attending HIV support clinic | + | 19/176 (11%) | Not reported |

HBsAg, HBeAg and HDV coinfection prevalence across 37 studies identified by a systematic literature search of HBV resistance associated mutations (RAMs) and vaccine escape mutations (VEMs) from African cohorts published between 2007 and 2017 (inclusive).

[a] HIV status is designated '+' whole cohort HIV-positive; '±' some of cohort HIV-positive; '-'none of cohort HIV-positive

[b] HBV prevalence in these cohorts was reported using HBV DNA detection rather than HBsAg

[c] Occult HBV prevalence reported in these cohorts

Study populations were categorised as follows:

- HBV/HIV coinfected patients: (n=28 studies), (8–10,37–40,42–62);

- HBV infected with and without HIV coinfection: (n=8 studies), (34–36,63–67);

- Chronic HBV monoinfection: (n=1 study), (68).

Antiviral treatment exposure varied as follows:

• Treatment-naïve: (n=8 studies),  (39–41,58–60,63,67);

• 3TC-based regimen only: (n=11 studies),  (8,9,37,38,42,43,46,50,55,57,67);

• Regimens including 3TC or TDF: (n=8 studies), (9,42, 43,44,46,47,51,63);

• Mixed regimen where some received 3TC, others TDF, while others left

untreated; (n=4 studies),(44,49,56,66);

• Treatment regimen not specified: (n=6 studies), (33–35,53,61,64).

HBV amino acid polymorphisms were studied from within the following proteins;

• Pol only (n=13 studies), (8,9,37,45,46,49,50,52,53,55,59,63,67); only one of

these used a deep sequencing method (46);

• S only (n=3 studies), (36,64,65);

• Pol and S (n=12 studies),(10,34,38,41,44,47,48,51,54,56,58,62);

• Polymerase, S and PC/BCP (n=4 studies), (40,43,57,68);

• S and PC/BCP region (n=3 studies), (35,39,60);

• Whole genome (n=2 studies), (42,66).

All studies, except for two (50,67), specified the HBV genotype (S2 Table & S3A

Fig).

**Prevalence of HBsAg, HBeAg and HDV co-infection**

The prevalence rates of HBsAg in these study cohorts ranged from 3%-26%;

however, the populations included were highly selective and therefore not

necessarily representative of the general population, particularly as a result of a

strong bias towards HIV-infection (Table 1). Only three studies included in this

review reported on HDV prevalence: two studies did not detect any HDV antibodies

(63,64), whereas the other study reported a HDV prevalence of 25% in Guinea-

Bissau (53).

**RAMs identified in African cohorts**

The co-occurrence and distribution of HBV RAMs and VEMs are summarised

according to the region where they were identified (Fig 2). This illustrates the patchy

and limited data that are available, with South Africa, Ghana and Cameroon best

represented, but with large areas (especially in northern and central Africa) not

represented at all in the literature.

**Figure 2: Annotated map to summarise HBV drug Resistance Associated**

**Mutations (RAMs) and Vaccine Escape Mutations (VEMs).** Mutations identified

from 33 studies of African cohorts published between 2007 and 2017 (inclusive).

Four studies identified by our systematic literature review were not represented here

as they did not report any RAMs.

Although 35 studies specified the HBV genotype, it was only possible to group RAMs

according to genotype in fourteen studies (8,9,37,39,40,43–45,47,53,56,57,59,68)

(S3B Fig).The remaining 21 studies generally reported the genotypes identified,

however they were not specific on which genotype inidividuals with RAMs were

infected with.

We have developed an interactive tool to display the genomic positions of RAMs

identified through our literature review alongside relevant metadata. This can be

accessed        on-line        here:        https://livedataoxford.shinyapps.io/1510659619-

3Xkoe2NKkKJ7Drg/ (33).

Overall, the most prevalent RAM was rtM204V/I in both treatment experienced and

treatment naïve patients and occurring either alone or in combination with other

polymorphisms rtL80I/V, rtV173L, rtL180M, rtA181S, rtT184S, rtA200V and/or

rtS202S (Fig 3, S3 Table & S4 Table). This mutation was present in 29 studies at a

highly variable prevalence of between 0.4% (49) and 76% (68). Across all cohorts,

the mutation was present in 208/2569 (8%) of all individuals represented. The

mutation, by itself, was most prevalent in South Africa; on pooling data for three

studies from this setting, it was present in both treatment experienced and treatment

naïve patients (n=13/17, 76% (68) and n=16/72, 22% (62,67) respectively). In

addition to South Africa, rtM204I/V was also frequent in Malawi among treatment

experienced patients (n=24/154, 16% (38,46)) (Fig 3), and in genotype non-A

infection: in this setting, the mutation was detected in genotype C infection (n=2/17,

12% (68)) (S3B Fig).

**Figure 3: Prevalence of HBV resistance associated mutations (RAMs) in Pol/RT**

**proteins among HBV infected patients in Africa.**

These data are derived from 27 studies of HBV drug resistance in Africa published

between 2007 and 2017 (inclusive). The countries represented are listed in

alphabetical order. A detailed summary of RAMs identified from each study is

presented in fig 2, and in Suppl Tables 3 and 4. Prevalence of RAMs for a specific

country was determined by grouping all studies from that country that reported a

specific mutation; the denominator was the total number of individuals infected with

HBV from these studies and the numerator was the total number of individuals with

that specific mutation from these studies.

Panel A: treatment naïve;

Panel B: treatment experienced.

rtM204I/V mutation by itself confers resistance to 3TC, and may also be associated

with reduced efficacy to TDF when in combination with A194T, and with vaccine

escape in combination with L180M and V173L. Although TDF has a high genetic

barrier to resistance, and is associated with reliable suppression of HBV viraemia

(7,10,15,16), mutations rtN236T and rtA194T, which have been linked with

resistance to both TDF and ADV (7), have been identified in Southern Africa in both

treatment naïve (59) and treatment experienced (10) patients.

WHO guidelines recommend a first-line regimen including TDF in HIV/HBV

coinfected patients(6), and the South African Department of Health HIV/AIDS

treatment guideline included TDF as first-line regimen from 2010 (69), however we

found a minority of studies (9/37, 24%) reporting TDF-containing regimens for

HIV/HBV coinfected individuals. As anticipated, most of the studies that did use TDF

were carried out after 2010, whereas those that used 3TC were generally earlier (S2

Table).

From this dataset, it is difficult to ascertain whether RAMs are genuinely more

prevalent in genotype A infection, or this simply reflects enrichment of genotype A in

sub-Saharan African populations (S3 Fig). Interpreting RAMs according to sub-

genotypes was difficult since most studies did not specify sub-genotype and others

did not indicate which RAMs were identified in which genotype. Of concern is the

detection of RAMs even in treatment naïve individuals (Fig 3 & S3 Table),

suggesting that RAMs are being transmitted. A study in South Africa that recruited

3TC-naïve HBV infected adults with or without HIV, reported rtM204I in 13/35 (37%)

individuals (67).

**HBV RAMs in published sequences from Africa**

We searched the Hepatitis B Virus database and GenBank to identify HBV

sequences derived from Africa, from studies not already included in our review. We

identified additional 69 isolates: 23 had undergone full length genome sequencing

whereas 46 isolates represented either the polymerase (n=3) or S region (n= 43) of

the HBV genome (Table 2). We excluded fourteen studies already identified by our

literature review that had submitted their sequences to GenBank (S2 Table) to avoid

duplication of results. RAMs in the additional 69 isolates were as follows:

• rtM204V in genotype A (2/69, 2.9% of sequences), this occurred in

combination with rtL180M;

• rtM204V + rtL180M in genotype E (1/69, 1.5%);

• rt180M + rtA181V in genotype E (1/69 (1.5%);

• rtQ215S identified in genotype D (4/69, 5.8%).

All these mutations are associated with 3TC resistance, rtA181V has also been

associated with reduced susceptibility to TDF (7,13).

In the S gene, the most prevalent mutations were:

• sD144A/E/G occurring in genotype A (6/69, 8.7%), D (10/69, 14.5%) and E

(7/69, 10.1%) associated with VEM;

• sI110L occurring in genotype A (3/69, 4.3%), D (4/69, 5.8%) and E (11/69,

15.9%) associated with immunoglobulin resistance.

**Table 2: HBV drug resistant mutations (RAMs) identified from HBV genome**

**sequences from Africa downloaded from the Hepatitis B Virus database**

**(https://hbvdb.ibcp.fr/HBVdb/)** (31) **and GenBank database**

**(http://hvdr.bioinf.wits.ac.za/alignments/)** (32)

| HBV Protein | HBV Genotype | Position and nature of the mutation | Number (%) of HBV sequences with mutation | Accession number | Region sequenced | Country of origin |
|---|---|---|---|---|---|---|
| Polymerase (Pol) | A | rtL180M | 2/69 (2.9%) | KM519454 | Full length | South Africa |
| | | | | FM199980 | Full length | Rwanda |
| | | rtM204V | 2/69 (2.9%) | KM519454 | Full length | South Africa |
| | | | | FM199980 | Full length | Rwanda |
| | | rtI233V | 1/69 (1.4%) | HM535205 | Full length | Zimbabwe |
| | D | rtV214A | 1/69 (1.4%) | FJ904395 | Polymerase | Tunisia |
| | | rtQ215S | 4/69 (5.8%) | FJ904414 | Full length | Tunisia |
| | | | | FJ904431 | Full length | Tunisia |
| | | | | FJ904436 | Full length | Tunisia |
| | | | | FJ904438 | Full length | Tunisia |
| | E | rtV173L | 1/69 (1.4%) | KF849723 | Full length | Angola |
| | | rtL180M | 2/69 (2.9%) | KF849720 | Full length | Angola |
| | | | | KF849723 | Full length | Angola |
| | | rtA194T | 1/69 (1.4%) | GQ161771 | Full length | Guinea |
| | | rtM204V | 1/69 (1.4%) | KF849723 | Full length | Angola |
| | | rtA181V | 1/69 (1.4%) | KF849720 | Full length | Angola |
| | | rtN238D | 2/69 (2.9%) | HM363566 | Polymerase | Nigeria |
| | | | | HM363587 | Polymerase | Nigeria |
| Surface (S) | A | sI110L | 3/69 (4.3%) | KY493896 | S | Cameroon |
| | | | | KP168431 | Full length | Kenya |
| | | | | AY233286 | Full length | South Africa |
| | | sP120S | 1/69 (1.4%) | KX648547 | S | Zimbabwe |
| | | sG129R | 1/69 (1.4%) | FN547352 | S | Cameroon |
| | | sT126A | 1/69 (1.4%) | JN182330 | S | South Africa |
| | | sD144A/E/G | 6/69 (8.7%) | FN547249 | S | Cameroon |
| | | | | KX493873 | S | Cameroon |
| | | | | FM199980 | Full length | Rwanda |
| | | | | FM200180 | S | Rwanda |
| | | | | FM200189 | S | Rwanda |
| | | | | KF467020 | S | South Africa |
| | | sG145R | 1/69 (1.4%) | FM200185 | S | Rwanda |
| | | sC149R | 1/69 (1.4%) | KF476024 | S | South Africa |

| | | | | | | |
|---|---|---|---|---|---|---|
| | D | sI110L | 4/69 (5.8%) | AB561830 | S | Egypt |
| | | | | KX357627 | Full length | Ethiopia |
| | | | | FJ904429 | S | Tunisia |
| | | | | KJ416196 | S | Tunisia |
| | | sP120S | 1/69 (1.4%) | KX357636 | Full length | Ethiopia |
| | | sM133T | 1/69 (1.4%) | KM108592 | S | Sudan |
| | | sD144E | 10/69 (14.5%) | FJ904427 | Full length | Tunisia |
| | | | | FN547165 | S | Cameroon |
| | | | | FN547179 | S | Cameroon |
| | | | | FN547239 | S | Cameroon |
| | | | | FN547255 | S | Cameroon |
| | | | | FN547258 | S | Cameroon |
| | | | | FN547262 | S | Cameroon |
| | | | | FN547281 | S | Cameroon |
| | | | | FN547318 | S | Cameroon |
| | | | | FN547319 | S | Cameroon |
| | E | sI110L | 11/69 (15.9%) | KY494047 | S | Cameroon |
| | | | | AM494711 | S | Central Africa Republic |
| | | | | AM494720 | S | Central Africa Republic |
| | | | | AM494725 | S | Central Africa Republic |
| | | | | AM494727 | S | Central Africa Republic |
| | | | | JQ972822 | S | Central Africa Republic |
| | | | | AB205190 | Full length | Ghana |
| | | | | GQ161756 | S | Guinea |
| | | | | GQ161768 | Full length | Guinea |
| | | | | GQ161795 | | Guinea |
| | | | | DQ060822 | Full length | South Western Africa and Madagascar |
| | | sT126N | 1/69 (1.4%) | HM363608 | S | Nigeria |
| | | sP120S | 1/69 (1.4%) | HM363599 | Full length | Nigeria |
| | | sM133T | 3/69 (4.3%) | HM363603 | S | Nigeria |
| | | | | KF170751 | S | Sudan |
| | | | | KF170752 | S | Sudan |
| | | sD144E | 7/69 (10.1%) | FN547300 | S | Cameroon |
| | | | | KY494047 | S | Cameroon |

| | | Accession | Type | Country |
|---|---|---|---|---|
| | | AM494719 | S | Central African Republic |
| | | AM494726 | S | Central African Republic |
| | | FN594756 | Full length | Niger |
| | | HM363565 | Full length | Nigeria |
| | | HM363590 | S | Nigeria |
| G145A/R | 3/69 (4.3%) | KY493921 | S | Cameroon |
| | | AB205327 | S | Ghana |
| | | AM494741 | S | Central African Republic |

**VEMs**

VEMs were identified in Central, East, West and Southern Africa (Fig. 2). However, it was not possible to ascertain whether individuals harbouring these mutations had been vaccinated against HBV infection. The most common VEM was the triple mutation rtV173L + rtL180M + rtM204I/V, found in the Pol gene. This suite of mutations was identified in 14 studies (37,43–52,54–56), at a pooled prevalence of 4% (57/1462). Another significant VEM, sG145K/R (14), was identified in six studies (35,40,44,49,56,65) and sM133L/T, associated with VEM, immunoglobulin and diagnostic escape mutation(41,49), was identified in seven studies (34,40,41,44,49,60,65) (Fig. 2).

**DISCUSSION**

**Summary**

To our knowledge, this is the first systematic review that assesses RAMs and VEMs for HBV in Africa. The high rates of HBV infection among HIV infected individuals in some locations including Cameroon (56) and South Africa (10) could be an indication that HBV infection has been previously under-reported, possibly due to lack of

routine screening, poor awareness, stigma, high costs and limited clinical and

laboratory infrastructure (4,8–10,38,50). The literature suggests a widespread

exposure of the HIV-infected population to 3TC-based treatment. Although this may

be changing over time, the introduction of TDF-based regimens for HIV treatment

has been slow and inconsistent, and TDF monotherapy is not consistently available

for HBV infection in the absence of HIV.

In keeping with other settings, the most common RAM identified here was rtM204I/V,

either alone or in combination with compensatory mutations rtL180M ± rtV173L. Of

concern, rtM204V/I was seen in 76%, (68) of treatment experienced patients and

22% of treatment naïve patients (62,67) in South Africa. A review of worldwide

incidence of RAMs among treatment naïve patients also described rtM204V/I as the

most frequent, but with a much lower prevalence of 5% (70). Likewise, a European

study demonstrated that the most frequent primary mutation was rtM204V/I, found in

49% of treatment experienced patients (71), while in China rtM204I, rtN236T and

rtL180M+rtM204V+rtV173L/rtS202G were also the most prevalent RAMs (72).

The triple mutation rtM204V + rtL180M + rtV173L has been identified in East, West

and Central Africa (37,46–48,50,51,55). This combination of polymorphisms is

associated with both vaccine escape and resistance to 3TC and other □-nucleoside

analogues (37,46,47,51,55,56). Interestingly, this triple mutation has not been

reported in the Southern African region to date. It could be a matter of time before it

is reported; larger studies and data on the vaccination status of research participants

are required.

**Clinical and public health significance of RAMs**

The spread of these mutations may lead to a rise in drug resistance in treatment naïve chronic HBV infection, representing a substantial challenge for Africa and highlighting an imperative to ensure routine use of TDF in preference to 3TC-based therapy. Although these data provide a preliminary picture of the prevalence of RAMs in some settings, there are no recommendations to stipulate any specific prevalence threshold above which HBV drug resistance mutations represent a significant barrier to successful treatment at a population level, and/or RAM prevalence thresholds that should trigger a switch to alternative first-line therapy. For HIV, surveillance for transmission of RAMs is based on screening recently infected, treatment naive individuals, and classifies drug resistance using thresholds of <5%, 5-15%, and >15% to stratify the risk to public health (73). Similar thresholds and recommendations for HBV could help to underpin the assimilation of epidemiological data and unify treatment approaches.

**TDF resistance**

The identification of mutations associated with reduced TDF susceptibility are of concern, as they suggest the potential for increasing prevalence of polymorphisms that confer partial or complete viral escape from a drug that to date has not been widely associated with resistance. There is now potential for increasing selection of TDF resistance as this drug becomes more widely used. However, as a new first line single tablet option incorporating 3TC, TDF and Dolutegravir ('LTD') emerges for HIV treatment, surveillance is needed to determine the clinical outcomes for HBV.

If clinically significant, TDF resistance mutations may still represent a particular

problem for many African settings, as resource constraints make it unrealistic to

provide baseline screening for RAMs, or to monitor patients on treatment with serial

viral load measurements. Despite these potential concerns, it has been shown that

TDF is effective even in the presence of RAMs and that there is comparable efficacy

among 3TC-experienced and NA-naïve patients (74).

**Limitations of current data**

Screening for HBV infection is not routinely performed in many African settings and

therefore the true prevalence and characteristics of HBV infection are not known

(4,7–10). We identified very few published studies; only a minority of patients had

HBV sequencing undertaken, and there were no data from certain regions of Africa.

This highlights the substantial problem of HBV neglect in Africa , and a specific blind-

spot relating to sequence data (4). Identifying the true prevalence of resistance

mutations, and characterising the populations in which these are selected and

enriched, is currently not possible due to sparse data and lack of clear descriptions

of the denominator population. Most such studies do not perform a truly systematic

assessment, but focus on high risk groups – particularly including those with

HIV/HBV coinfection: of the 37 studies included here, only one exclusively reported

on participants who were HBV mono-infected (68). Although we have made every

effort to assimilate the relevant data to build up a regional picture for Africa, the

heterogeneity between studies makes it difficult to draw robust conclusions from

pooled data. These findings are a reflection of the little attention paid towards the

burden of this disease in Africa and the neglect in robust epidemiological data.

Only nine studies undertook a longitudinal approach to detection of drug resistance

(8–10,37,38,42,46,48,50). The results of the other 28 studies that undertook a cross-

sectional approach could be skewed by the timing of recruitment of study

participants, with a risk of under-representation of drug resistance if screening is

undertaken only at baseline, and potentially an over-representation if screening is

undertaken in patients with HIV coinfection, who are more at risk of advanced

disease and prolonged drug exposure. As most of these studies recruited individuals

from hospital settings, this raises the latter possibility.

Mutations across the whole genome might be relevant in determining resistance

(75). However, most of the included studies analysed only defined genes from within

the HBV genome; only two sequenced the whole genome, and these determined

consensus sequence. This potentially results in an under-representation of RAMs

and VEMs that may be present as low numbers of quasispecies, but could become

significant if selected out by exposure to drug or vaccine.

In studies that reported RAMs among treatment naïve individuals, the literature

suggests that identification of RAMs and VEM was performed prior to ART initiation.

However, we cannot exclude the possibility that that some of these participants had

prior ART exposure. Due to the nature of the cohorts that have been studied, most of

the RAMs identified were from HIV/HBV coinfected individuals. It is possible that HIV

increases the risk of HBV RAMs both in terms of drug exposure, and also as a

function of increased HBV viral loads. A study carried out in Italy revealed that

patients with HIV coinfection were more likely to harbour the rtM204V mutation and

to show multiple mutations compared to HBV monoinfected patients (76). It would be

worth exploring this hypothesis in Africa, as there are currently no such data.

**Challenges and opportunities for Africa**

A major challenge for Africa is to deploy more widespread screening for HBV, to

identify and treat infections when required, but also in order to avoid exposure to

inappropriate HBV monotherapy in HIV-coinfected individuals starting ART (16).

Despite WHO recommendations to monitor for the development of drug resistance

once on therapy (6), implementation remains challenging as viral load monitoring

and sequencing are both rarely available (7). Development of a cheap rapid test for

the detection of the most frequently observed RAMs and VEMs should be

considered as potentially cost-effective alternative for Africa. New metagenomic

sequencing platforms, such as Illumina and Nanopore, provide the opportunity for

whole and deep genome sequencing, which reveal the full landscape of HBV

mutations in individual patients, quantify the prevalence of drug resistance mutations

among HBV quasi-species, and determine the relationship between these

polymorphisms and treatment outcomes (76). Nanopore technology also has the

potential to develop into an efficient point of care test that could detect viral infection

and coinfection, as well as determining the presence of VEMs and RAMs (77), but is

currently limited by cost and concerns about high error rates.

There have been few studies looking at the correlation between genotype, clinical

outcomes of disease, response to antiviral therapy and RAMs/VEMs, but none from

Africa. Studies outside Africa have shown that genotype A is more prone to

immune/vaccine escape mutants, pre-S mutants associated with immune

suppression, drug associated mutations and HCC in HIV/HBV coinfected participants

(39,76,78). Studies investigating the role of genotypes in predicting response to

antiviral therapy and their association with various types of mutations are urgently

needed in Africa, particularly in light of the high frequency of genotype A infection

and high population exposure to antiviral agents that have been rolled out over the

past two decades as a component of first-line ART.

Existing infrastructure for diagnosis, clinical monitoring and drug therapy for HIV

represents an opportunity for linkage with HBV care; particularly in settings of limited

resource, joining up services for screening and management of blood-borne virus

infection could be a cost-effective pathway to service improvements.

**Conclusions**

This review highlights the very limited data that are available from Africa. Scarce

resources resulting in lack of diagnostic screening, inconsistent supply of HBV drugs

and vaccines, and poor access to clinical monitoring contribute to drug and vaccine

resistance, potentially amplifying the risk of ongoing transmission and adding to the

long-term burden of HBV morbidity and mortality in Africa. We call for urgent action

to gather and analyse better data, particularly representing the HBV monoinfected

population, and for improved access to TDF.

HBV RAMs and VEMs have been identified in several African countries among

HIV/HBV coinfected and HBV monoinfected patients, before and during treatment

with NAs but the data are currently insufficient to allow us to form a clear picture of

the prevalence, distribution or clinical significance of these mutations.  Overall, the

data we describe suggest a significantly higher prevalence of drug resistance in

some African populations than has been described elsewhere, and that is not

confined only to drug-exposed populations, highlighting an urgent need for better

population screening, assessment of HBV infection before and during therapy, and

increasing roll out of TDF in preference to 3TC.  At present, TDF accessibility is

largely confined to HIV/HBV coinfected individuals; we now need to advocate to

make monotherapy available for HBV infected individuals. However, there are

uncertainties as to whether its long-term use might result in nephrotoxicity, and

potentially in an increase in selection of TDF RAMs. We should ideally aim for the

goals of a combined HBV test that includes diagnosis of infection, genotype and

presence of RAMs/VEMs; new sequencing platforms such as Nanopore make this

technically possible, although cost remains a significant barrier at present.

Sustainable long-term investment is required to expand consistent drug and vaccine

supply, to provide screening infection and for drug resistance, and to provide

appropriate targeted clinical monitoring for treated patients.

**FUNDING**

JM is funded by a Leverhulme Mandela Rhodes Doctoral Scholarship. PCM is

funded by a Wellcome Trust Intermediate Fellowship (grant number 110110). The

funders had no role in study design, data collection and analysis, decision to publish,

or preparation of the manuscript.

**References**

1. WHO Global Hepatitis Programme. Global hepatitis report. 2017.

   http://www.who.int/hepatitis/publications/global-hepatitis-report2017/en/

2. WHO Combating Hepatitis B and C to reach elimination by 2030.  Advocacy brief.

   2016.

   http://apps.who.int/iris/bitstream/10665/206453/1/WHO_HIV_2016.04_eng.pdf

3. Lemoine M, Thursz MR. Battlefield against hepatitis B infection and HCC in Africa.

   J Hepatol. 2017.66: 645-654.

4. O'Hara GA, McNaughton AL, Maponga T, Jooste P, Ocama P, Chilengi R, et al.

   Hepatitis B virus infection as a neglected tropical disease. PLoS Negl Trop Dis.

   2017.11:10.

5. Ocama P, Seremba E, Apica B, Opio K. Hepatitis B and HIV co-infection is still

   treated using lamivudine-only antiretroviral therapy combination in Uganda. Afr

   Health Sci. 2015.15(2):328–33.

6. WHO Hepatitis B treatment guidelines. 2015

   http://www.who.int/mediacentre/news/releases/2015/hepatitis-b-guideline/en/

7. Beloukas A, Geretti AM. Hepatitis B Virus Drug Resistance. In: Antimicrobial Drug

   Resistance; 2017.p.1227–42.

8. Day SL, Odem-davis K, Mandaliya KN, Jerome KR, Cook L, Masese LN, et al.

   Prevalence, Clinical and Virologic Outcomes of Hepatitis B Virus Co-Infection in

   HIV-1 Positive Kenyan Women on Antiretroviral Therapy. PloS One. 2013.8(3):1–

   5.

9. Kim HN, Scott J, Cent A, Cook L, Morrow RA, Richardson B, et al. HBV

   lamivudine resistance among hepatitis B and HIV coinfected patients starting

   lamivudine, stavudine and nevirapine in Kenya. J Viral Hepat. 2011.18(10):447-52.

10. Hamers RL, Zaaijer HL, Wallis CL, Siwale M, Ive P, Botes ME, et al. HIV-HBV

coinfection in Southern Africa and the effect of lamivudine- versus tenofovir-

containing cART on HBV outcomes. J Acquir Immune Defic Syndr. 2013

64(2):174–82.

11. Fung J, Lai C-L, Seto W-K, Yuen M-F. Nucleoside/nucleotide analogues in the

treatment of chronic hepatitis B. J Antimicrob Chemother. 2011.66(12):2715–25.

12. Fischer KP, Gutfreund KS, Tyrrell DL. Lamivudine resistance in hepatitis B:

mechanisms and clinical implications. Drug Resist Updat. 2001.4(2):118–28.

13. Warner N, Locarnini S. Mechanisms of hepatitis B virus resistance development.

Intervirology. 2014.57(3–4):218–24.

14. Caligiuri P, Cerruti R, Icardi G, Bruzzone B. Overview of hepatitis B virus

mutations and their implications in the management of infection. World J

Gastroenterol. 2016.22(1):145–54.

15. Singhatiraj E, Suri J, Goulston C. HIV Co-Infections with Hepatitis B and C. J AIDS

Clin Res. 2013.1(S3):1–12.

16. Sheldon J, Camino N, Rodés B, Bartholomeusz A, Kuiper M, Tacke F, et al.

Selection of hepatitis B virus polymerase mutations in HIV-coinfected patients

treated with tenofovir. Antivir Ther. 2005.10(6):727–34.

17. Zoulim F. Hepatitis B virus resistance to antiviral drugs: where are we going? Liver

Int. 2011.31(s1):111–6.

18. Patzer RE, McClellan WM. Influence of race, ethnicity and socioeconomic status

on kidney disease. Nat Rev Nephrol. 2012.8(9):533–41.

19. Agbaji OO, Agaba PA, Idoko JA, Taiwo B, Murphy R, Kanki P, et al. Temporal

changes in renal glomerular function associated with the use of Tenofovir

Disoproxil Fumarate in HIV-infected Nigerians. West Afr J Med. 2011.30(3):164–8.

20. Buti M, Riveiro-Barciela M, Esteban R. Tenofovir Alafenamide Fumarate: A New

Tenofovir Prodrug for the Treatment of Chronic Hepatitis B Infection. J Infect Dis.

2017.216(s8):S792–6.

21. Shirvani-Dastgerdi E, Winer BY, Celià-Terrassa T, Kang Y, Tabernero D, Yagmur

E, et al. Selection of the highly replicative and partially multidrug resistant rtS78T

HBV polymerase mutation during TDF-ETV combination therapy. J Hepatol. 2017.

67(2):246–54.

22. Lee HW, Chang HY, Yang SY, Kim HJ. Viral evolutionary changes during tenofovir

treatment in a chronic hepatitis B patient with sequential nucleos(t)ide therapy. J

Clin Virol. 2014.60(3):313–6.

23. WHO Guidelines for the prevention, care and treatment of persons with chronic

hepatitis B infection. 2017.

http://apps.who.int/iris/bitstream/10665/154590/1/9789241549059_eng.pdf

24. Coffie PA, Egger M, Vinikoor MJ, Zannou M, Diero L, Patassi A, et al. Trends in

hepatitis B virus testing practices and management in HIV clinics across sub-

Saharan Africa. BMC Infect Dis. 2017.17(S1):706.

25. Jooste P, van Zyl A, Adland E, Daniels S, Hattingh L, Brits A, et al. Screening,

characterisation and prevention of Hepatitis B virus (HBV) co-infection in HIV-

positive children in South Africa. J Clin Virol. 2016.85:71–4.

26. McMahon MA, Jilek BL, Brennan TP, Shen L, Zhou Y, Wind-Rotolo M, et al. The

HBV Drug Entecavir — Effects on HIV-1 Replication and Resistance. N Engl J

Med. 2007.356(25):2614–21.

27. Locarnini SA. The Hepatitis B Virus and Antiviral Drug Resistance: Causes,

Patterns and Mechanisms. In: Antimicrobial Drug Resistance; 2017.p. 565–77.

28. Cooreman MP, Leroux-Roels G, Paulij WP. Vaccine - and hepatitis B immune

globulin-induced escape mutations of hepatitis B virus surface antigen. J Biomed

Sci. 2001.8(3):237–47.

29. Hsu H-Y, Chang M-H, Liaw S-H, Ni Y-H, Chen H-L. Changes of hepatitis B

surface antigen variants in carrier children before and after universal vaccination in

Taiwan. Hepatology.1999.30(5):1312–7.

30. Hayer J, Jadeau F, Deléage G, Kay A, Zoulim F, Combet C. HBVdb: a knowledge

database for Hepatitis B Virus.Nucleic Acids Res. 2013.41: D566-D570.

31. Bell TG, Yousif M, Kramvis A. Bioinformatic curation and alignment of genotyped

hepatitis B virus (HBV) sequence data from the GenBank public database.

Springerplus. 2016.5(1):1896.

32. Simmonds P. SSE: a nucleotide and amino acid sequence analysis platform. BMC

Res Notes. 2012.5(1):50.

33. Mokaya J, Hadley M, Matthews P. gene.alignment.tables. Figshare. 2017.

doi.org/10.6084/m9.figshare.5729229

34. Mokaya J, Hadley M, Matthews P. On-line tool to visualise sites of drug and

vaccine escape mutations within the HBV genome. 2018.

https://livedataoxford.shinyapps.io/1510659619-3Xkoe2NKkKJ7Drg/

35. Hundie GB, Raj VS, Michael DG, Pas SD, Osterhaus ADME, Koopmans MP, et al.

Molecular epidemiology and genetic diversity of hepatitis B virus in Ethiopia. J

Med Virol. 2016.88(6):1035–43.

36. Mahgoub S, Candotti D, El Ekiaby M, Allain J-P. Hepatitis B Virus (HBV) Infection

and Recombination between HBV Genotypes D and E in Asymptomatic Blood

Donors from Khartoum, Sudan. J Clin Microbiol. 2011.49(1):298–306.

37. Amponsah-Dacosta E, Lebelo RL, Rakgole JN, Selabe SG, Gededzha MP,

Mayaphi SH, et al. Hepatitis B virus infection in post-vaccination South Africa:

Occult HBV infection and circulating surface gene variants. J Clin Virol. 2015.

63:12–7.

38. Stewart B, Jobarteh ML, Sarge-njie R, Alabi A, Silva T De, Peterson K, et al.

Emergence of HBV resistance to lamivudine (3TC) in HIV / HBV co-infected

patients in The Gambia, West Africa. BMC Res Notes. 2011.4(1):561.

39. Galluzzo C, Liotta G, Andreotti M, Luhanga R, Jere H, Mancinelli S, et al.

Emergence of lamivudine resistance hepatitis B virus mutations in pregnant

women infected with HBV and HIV receiving antiretroviral prophylaxis for the

prevention of mother-to-infant transmission in Malawi. J Med Virol.

2012.84(10):1553–7.

40. Makondo E, Bell TG, Kramvis A. Genotyping and molecular characterization of

hepatitis B virus from human immunodeficiency virus-infected individuals in

southern Africa. PLoS One. 2012.7(9): e46345.

41. Yousif M, Mudawi H, Hussein W, Mukhtar M, Nemeri O, Glebe D, et al.

Genotyping and virological characteristics of hepatitis B virus in HIV-infected

individuals in Sudan. Int J Infect Dis. 2014.29:125–32.

42. Anderson M, Gaseitsiwe S, Moyo S, Wessels MJC, Mohammed T, Sebunya TK,

et al. Molecular characterisation of hepatitis B virus in HIV-1 subtype C infected

patients in Botswana. BMC Infect Dis. 2015.15(1):335.

43. Amponsah-Dacosta E, Rakgole JN, Gededzha MP, Lukhwareni A, Blackard JT,

Selabe SG, et al. Evidence of susceptibility to lamivudine-based HAART and

genetic stability of hepatitis B virus (HBV) in HIV co-infected patients: A South

African longitudinal HBV whole genome study. Infect Genet Evol. 2016.43:232–8.

44. Deressa T, Damtie D, Fonseca K, Gao S, Abate E, Alemu S, et al. The burden of

hepatitis B virus (HBV) infection, genotypes and drug resistance mutations in

human immunodeficiency virus-positive patients in Northwest Ethiopia. PLoS One.

2017.12(12):e0190149.

45. Bivigou-Mboumba B, Amougou-Atsama M, Zoa-Assoumou S, M'boyis Kamdem H,

Nzengui-Nzengui GF, Ndojyi-Mbiguino A, et al. Hepatitis B infection among HIV

infected individuals in Gabon: Occult hepatitis B enhances HBV DNA prevalence.

PLoS One. 2018.13(1): e0190592.

46. Mabeya SN, Ngugi C, Lihana RW, Khamadi SA, Nyamache AK. Predominance of

Hepatitis B Virus Genotype A Among Treated HIV Infected Patients Experiencing

High Hepatitis B Virus Drug Resistance in Nairobi, Kenya. AIDS Res Hum

Retroviruses. 2017.33(9):966–9.

47. Aoudjane S, Chaponda M, Gonzalez Del Castillo AA, O'Connor J, Noguera M,

Beloukas A, et al. Hepatitis B virus sub-genotype A1 infection is characterized by

high replication levels and rapid emergence of drug resistance in HIV-positive

adults receiving first-line antiretroviral therapy in Malawi. Clin Infect Dis. 2014.

59(11):1618–26.

48. Calisti G, Muhindo R, Boum Y 2nd, Wilson LA, Foster GM, Geretti AM, et al.

Epidemiology of HBV infection in a cohort of Ugandan HIV-infected patients and

rate and pattern of lamivudine-resistant HBV infection in patients receiving

antiretroviral therapy. Trans R Soc Trop Med Hyg. 2015.109(11):723–9.

49. Boyd A, Moh R, Gabillard D, le Carrou J, Danel C, Anglaret X, et al. Low risk of

lamivudine-resistant HBV and hepatic flares in treated HIV-HBV-coinfected

patients from Cote d'Ivoire. Antivir Ther. 2015.20(6):643–54.

50. Archampong TNA, Boyce CL, Lartey M, Kwamena W, Obo-akwa A, Kenu E, et al.

HBV genotypes and drug resistance mutations in antiretroviral treatment-naïve

and treatment-experienced HBV-HIV co-infected patients. Antivir Ther.

2017.22(1):13-20.

51. Chadwick D, Ankcorn M, Sarfo F, Phillips R, Fox Z, Garcia A, et al. Outcomes of

starting first-line antiretroviral therapy in hepatitis B virus / HIV-coinfected patients

in Ghana. J Antimicrob Chemother. 2012.67: 2939–2942

52. Geretti AM, Patel M, Sarfo FS, Chadwick D, Verheyen J, Fraune M, et al.

Detection of highly prevalent hepatitis B virus coinfection among HIV-seropositive

persons in Ghana. J Clin Microbiol. 2010.48(9):3223–30.

53. Ndow G, Gore ML, Shimakawa Y, Suso P, Jatta A, Tamba S, et al.  Hepatitis B

testing and treatment in HIV patients in The Gambia—Compliance with

international guidelines and clinical outcomes. PLoS One. 2017.12(6): e0179025.

54. Hønge BL, Jespersen S, Medina C, Té D da S, da Silva ZJ, Lewin S, et al.

Hepatitis B and Delta Virus Are Prevalent but Often Subclinical Co-Infections

among HIV Infected Patients in Guinea-Bissau, West Africa: A Cross-Sectional

Study. PLoS One. 2014.9(6): e99971.

55. Gachara G, Magoro T, Mavhandu L, Lum E, Kimbi HK, Ndip RN, et al.

Characterization of occult hepatitis B virus infection among HIV positive patients in

Cameroon. AIDS Res Ther. 2017.14(1):11.

56. Kouanfack C, Aghokeng AF, Mondain A, Bourgeois A, Kenfack A, Ducos J, et al.

Lamivudine-resistant HBV infection in HIV-positive patients receiving antiretroviral

therapy in a public routine clinic in Cameroon. Antivir Ther. 2012.326:321–6.

57. Magoro T, Gachara G, Mavhandu L, Lum E, Kimbi HK, Ndip RN, et al. Serologic

and genotypic characterization of hepatitis B virus in HIV-1 infected patients from

South West and Littoral Regions of Cameroon. Virol J. 2016.13(1):178.

58. Bivigou-Mboumba B, Francois-Souquiere S, Deleplancque L, Sica J, Mouinga-

Ondeme A, Amougou-Atsama M, et al. Broad Range of Hepatitis B Virus (HBV)

Patterns, Dual Circulation of Quasi-Subgenotype A3 and HBV/E and

Heterogeneous HBV Mutations in HIV-Positive Patients in Gabon. PLoS One.

2016.11(1):e0143869.

59. 795    Chambal LM, Samo Gudo E, Carimo A, Corte Real R, Mabunda N, Maueia C, et

796    al. HBV infection in untreated HIV-infected. PLoS One. 2017.12(7): e0181836.

60. 797    Wandeler G, Musukuma K, Zurcher S, Vinikoor MJ, Llenas-Garcia J, Aly MM, et

798    al. Hepatitis B Infection, Viral Load and Resistance in HIV-Infected Patients in

799    Mozambique and Zambia. PLoS One. 2016.11(3): e0152043.

61. 800    Baudi I, Iijima S, Chin'ombe N, Mtapuri-Zinyowera S, Murakami S, Isogawa M, et

801    al. Molecular epidemiology of co-infection with hepatitis B virus and human

802    immunodeficiency virus (HIV) among adult patients in Harare, Zimbabwe. J Med

803    Virol. 2017.9(2):257–66.

62. 804    Anderson M, Gaseitsiwe S, Moyo S, Wessels MJC, Mohammed T, Sebunya TK,

805    et al. Molecular characterisation of hepatitis B virus in HIV-1 subtype C infected

806    patients in Botswana. BMC Infect Dis. 2015.15:335.

63. 807    Powell EA, Gededzha MP, Rentz M, Rakgole NJ, Selabe SG, Seleise TA, et al.

808    Mutations associated with occult hepatitis B in HIV-positive South Africans. J Med

809    Virol. 2015.87(3):388–400.

64. 810    Matthews PC, Beloukas A, Malik A, Carlson JM, Jooste P, Ogwu A, et al.

811    Prevalence and characteristics of hepatitis B virus (HBV) coinfection among HIV-

812    Positive women in South Africa and Botswana. PLoS One. 2015.10(7):1–11.

65. 813    Andersson MI, Maponga TG, Ijaz S, Barnes J, Theron GB, Meredith SA, et al. The

epidemiology of hepatitis B virus infection in HIV-infected and HIV-uninfected

pregnant women in the Western Cape, South Africa. Vaccine. 2013.31(47):5579–

84.

66. Faleye TOC, Adewumi MO, Ifeorah IM, Omoruyi EC, Bakarey SA, Akere A, et al.

Detection of hepatitis B virus isolates with mutations associated with immune

escape mutants among pregnant women in Ibadan, southwestern Nigeria.

Springerplus. 2015.4:43.

67. Gededzha MP, Muzeze M, Burnett RJ, Amponsah-Dacosta E, Mphahlele MJ,

Selabe SG. Complete genome analysis of hepatitis B virus in human

immunodeficiency virus infected and uninfected South Africans. J Med Virol. 2016.

88(9):1560–6.

68. Selabe SG, Lukhwareni A, Song E, Leeuw YGM, Burnett RJ, Mphahlele MJ.

Mutations associated with lamivudine-resistance in therapy-naive hepatitis B virus

(HBV) infected patients with and without HIV co-infection: implications for

antiretroviral therapy in HBV and HIV co-infected South African patients. J Med

Virol. 2007.79(11):1650–4.

69. Selabe SG, Song E, Burnett RJ, Mphahlele MJ. Frequent detection of hepatitis B

virus variants associated with lamivudine resistance in treated South African

patients infected chronically with different HBV genotypes. J Med Virol.

2009.81(6):996–1001.

70. WHO Clinical guidelines for the management of HIV & AIDS in adults and

adolescents. 2010.

http://www.who.int/hiv/pub/guidelines/south_africa_art.pdf

71. Zhang Q, Liao Y, Cai B, Li Y, Li L, Zhang J, et al. Incidence of natural resistance

mutations in naïve chronic hepatitis B patients: A systematic review and meta-

analysis. J Gastroenterol Hepatol. 2015.30(2):252–61.

72. Hermans LE, Svicher V, Pas SD, Salpini R, Alvarez M, Ben Ari Z, et al. Combined

analysis of the prevalence of drug-resistant Hepatitis B virus in antiviral therapy-

experienced patients in Europe (CAPRE). J Infect Dis. 2016.213(1):39–48.

73. Meng T, Shi X, Gong X, Deng H, Huang Y, Shan X, et al. Analysis of the

prevalence of drug-resistant hepatitis B virus in patients with antiviral therapy

failure in a Chinese tertiary referral liver centre (2010–2014). J Antimicr Res. 2017.

8, p.74–81

74. WHO Surveillance of transmitted HIV drug resistance. 2013.

http://www.who.int/hiv/topics/drugresistance/surveillance/en/

75. Baran B, Soyer OM, Ormeci AC, Gokturk S, Evirgen S, Bozbey HU, et al.  Efficacy

of tenofovir in patients with Lamivudine failure is not different from that in

nucleoside/nucleotide analogue-naive patients with chronic hepatitis B. Antimicrob

Agents Chemother. 2013.57(4):1790–6.

76. Betz-Stablein BD, Töpfer A, Littlejohn M, Yuen L, Colledge D, Sozzi V, et al.

Single-Molecule Sequencing Reveals Complex Genome Variation of Hepatitis B

Virus during 15 Years of Chronic Infection following Liver Transplantation. J Virol.

2016.90(16):7171–83.

77. Iacomi F, Vincenti D, Vairo F, Solmone M, Mariano A, Piselli P, et al.  Effect of HIV

co-infection on mutation patterns of HBV in patients with lamivudine-resistant

chronic hepatitis B. J Med Virol. 2009.81(7):1151–6.

78. Oxford Nanopore Technologies: https://nanoporetech.com/

79. Kramvis A, Kew MC. Epidemiology of hepatitis B virus in Africa, its genotypes and

clinical associations of genotypes. J Hep Res. 2007.37 (s1): 9–19.

80. Mokaya J, Hadley M, Matthews P. A systematic review of Hepatitis B virus (HBV)

drug and vaccine escape mutations in Africa - Supplementary data. Figshare.

2018. doi.org/ 10.6084/m9.figshare.5774091

**Suppl Figures**

**S1 Fig: Flow diagram illustrating identification and inclusion of studies for a systematic review of drug and vaccine resistance mutations in Africa, based on PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) criteria.**

**S2 Fig: HBV drug Resistance Associated Mutations (RAMs) grouped according to genotype.** Data summarised from fourteen studies published between 2009-2017 (inclusive). 21 studies were not represented here as they did not specifically indicate which genotype individuals with RAMs belonged to.

**S3 Fig: Distribution of HBV genotypes and prevalence of HBV resistance associated mutations (RAMs) in Pol/RT proteins in geno-A and geno-non-A samples.**

Panel A: Distribution of HBV genotypes derived from 35 studies reporting resistance associated mutations (RAMs) in Africa published between 2009 to 2017 (inclusive); Panel B: Prevalence of HBV resistance associated mutations (RAMs) in Pol/RT proteins in geno-A and geno-non-A samples. These data are derived from 14 studies of HBV drug resistance in Africa published between 2007 and 2017 (inclusive). 21 studies were not represented here as they did not specifically indicate which genotype individuals with RAMs belonged to. We had more geno-A samples represented than other samples, we therefore combined samples from other genotypes that had RAMs (B, C, D, E, D/E) to form geno-non-A samples. We then compared prevalence of Pol/RT mutation between geno-A samples to geno-non-A samples.

Prevalence of RT/Pol mutations for a specific genotype(geno-A/geno-non-A) was

determined by grouping all studies with geno-A/geno-non-A infection that reported a

specific mutation; the denominator was the total number of individuals infected with

geno-A/geno-non-A  from these studies and the numerator was the total number of

individuals infected with geno-A/geno-non-A with that specific mutation.

**Suppl Tables**

**S1 Table: Details of search strategy used to identify studies on HBV resistance**

**associated mutations (RAMs) and vaccine escape mutations (VEMs)**

**conducted in Africa.** A: PubMed database; B: SCOPUS and EMBASE database.

**S2 Table: Full details of 37 studies identified by a systematic literature search**

**of HBV resistance associated mutations (RAMs) and vaccine escape mutations**

**(VEMs) from African cohorts published between 2007 and 2017 (inclusive).**

**S3 Table: HBV Pol/RT mutations among treatment-naïve HBV infected patients**

**in Africa from 12 studies published between 2007 and 2017 (inclusive).**

**S4 Table: HBV Pol/RT mutations among treatment-experienced HBV infected**

**patients in Africa, from 25 studies published between 2009 and 2017 (inclusive).**

# Fig 1: Resistant mutations associated with drug, vaccine and immunoglobulin resistance

•Beloukas A, Geretti AM (2017) Hepatitis B Virus Drug Resistance. In: Antimicrobial Drug Resistance. p. 1227–42.

•Shirvani-Dastgerdi E, Winer BY, Celià-Terrassa T, Kang Y, Tabernero D, Yagmur E, et al. (2017) Selection of the highly replicative and partially multidrug resistant rtS78T HBV polymerase mutation during TDF-ETV combination therapy. J Hepatol. 67(2):246–54.

•Mabeya SN, Ngugi C, Lihana RW, Khamadi SA, Nyamache AK (2017) Predominance of Hepatitis B Virus Genotype A Among Treated HIV Infected Patients Experiencing High Hepatitis B Virus Drug Resistance in Nairobi, Kenya. AIDS Res Hum Retroviruses ;33(9):966–9.

•Wandeler G, Musukuma K, Zurcher S, Vinikoor MJ, Llenas-Garcia J, Aly MM, et al. (2016) Hepatitis B Infection, Viral Load and Resistance in HIV-Infected Patients in Mozambique and Zambia. PLoS One 11(3):e0152043.

•Cooreman MP, Leroux-Roels G, Paulij WP (2001) Vaccine- and hepatitis B immune globulin-induced escape mutations of hepatitis B virus surface antigen. J Biomed Sci ;8(3):237–47.

• Archampong TNA, Boyce CL, Lartey M, Kwamena W, Obo-akwa A, Kenu E, et al. (2017) HBV genotypes and drug resistance mutations in antiretroviral treatment-naïve and treatment-experienced HBV-HIV co-infected patients. Antivir Ther 22(1):13-20.

•Warner N, Locarnini S (2014) Mechanisms of hepatitis B virus resistance development. Intervirology 57(3–4):218–24.

•Caligiuri P, Cerruti R, Icardi G, Bruzzone B (2016) Overview of hepatitis B virus mutations and their implications in the management of infection. World J Gastroenterol 22(1):145–54.

# Fig 2: HBV drug Resistance Associated Mutations (RAMs) and Vaccine Escape Mutations (VEMs) mutations identified in 33 studies from Africa

**Guinea Bissau**
*RAM*
rtV173L#
rtV214A#
rtN238D#
rtL180M+rtM204V
*(31) Langhoff Honge et al 2014*

**Gambia**
*RAM*
rtM204I/V (33)
rtM204V+rtL180M (32, 33)
rtI169L+rtL180M+rtM204V (32)
rtM204V+rtV173L+rtL180M (32, 33)
*VEM*
*rt*M204V+rtV173L+rtL180M (32, 33)
*(32) Stewart et al 2011*
*(33) Ndow et al 2017*

**Sudan**
*RAM*
rtA181# (1)
rtQ215S# (1)
rtI233# (1)
*VEM*
sG145R (1,2)
*HBIg mutations*
sM133T (1)
sG145R (1,2)
*(1) Yousif et al 2014*
*(2) Mahgoub et al 2011*

**Ethiopia**
*RAM*
rtI169T (4); rtT184A (4);
rtA194T (4); rtL180M (3)
rtL180M+rtM204V (3,4)
rtV173L+rtL180M+rtM204V (3)
*VEM*
rtV173L+rtL180M+rtM204V (3)
sP120S (4); sT126I (4); sQ129R/H (4);
sM133T/I (4); sN/T131S/I (4)
sY134F (4); sT140I (4); sT143M (4)
sD144E (4)
*(3) Deressa et al 2017*
*(4) Hundie et al 2016*

**Uganda**
*RAM*
rtL180M/+rtM204V
rtM204I+rtV173L
rtM204V+rtL80I@
rtV173L+rtL180M+rtM204V@
rtM204I+ rtL80I+rtL80V
*VEM*
*rt*V173L+rtL180M+rtM204V
*(5) Calisti et al 2015*

**Kenya**
*RAM*
rtM204I (6,7);rtA181T# (7)
rtI233V# (7); rtV207I# (7)
rtA200V+rtM204I (7)
rtM204V+rtL180M (8)
rtM204V+rtV173L+rtL180M (8)
*VEM*
*rt*M204V+rtV173L+rtL180M (8)
*(6)Day et al 2013*
*(7)Kim et al 2011*
*(8)Mabeya et al 2017*

**Ghana**
*RAM*
rtL180M (28)
rtL180M+rtM204I/V (28)
rtV173L+rtA194T (20)
rtM204I+rtL80I (28)
rtV173L+rtL180M+rtM204V/I (28, 30)
rtL80I+rtL180M+rtM204V (28)
rtM204I/V+/-rtL80I+/-rtV173L+/-rtL180M (29)
*VEM*
rtV173L+rtL180M+rtM204V/I (28, 29, 30)
sG145R (28); sT126I (28); sM133L (28)
sD144E/H (28); sP120S (28)
*HBIg mutations*
sM133L (28); sD144E/H (28)
sG145R (28)
*(28) Archampong et al 2017*
*(29) Chadwick et al 2012*
*(30) Geretti et al 2010*

**Cote d'I voire**
*RAM*
rtV173L+rtL180M+rtM204V
*VEM*
rtV173L+rtL180M+rtM204V
sQ129P/R
*(27) Boyd et al 2015*

**Nigeria**
*VEM*
sM133T
sG145K
*HBIg mutations*
sM133T
sG145K
*(26) Faleye et al*

**Cameroon**
*RAM*
rtL180M+rtM204V (23, 24)
rtV173L+rtL180M+rtM204V@ (23, 24, 25)
rtL180M+rtM204V+rtT184S @ (23,25)
rtL180M+rtS202G+rtM204V (24)
*VEM*
*rt*V173L+rtL180M+rtM204V@ (23, 24, 25)
sQ129H (23); sG145A (25)
*HBIg mutations*
T131N (25); S143T (25); sG145A (25)
*(23) Gachara et al 2017*
*(24) Kouanfack et al 2012*
*(25) Magoro et al 2013*

**Gabon**
*RAM*
rtM204I (22); I169L (22)
rtL180M+rtM204V/I@ (21, 22)
rtV173L+rtL180M+rtM204V (22)
*VEM*
sT126I(21); sP120T (21); sG130N (22);s143T (22);
sG145K (22); sK141R (22); sM133I (22); sD144N (22)
*(21) Bivigou-Mboumba et al 2016*
*(22) Bivigou-Mboumba et al 2018*

**Zambia**
*RAM*
rtV173L#
rtL180M+rtM204V#
rtV173L+rtL180M#
*(11) Wandeler et al 2016*

**Malawi**
*RAM*
rtM204I/M/ V@ (9, 10)
rtL180M+rtM204V/I (9, 10)
rtM204V+rtL80I (10)
rtM204V/I+rtA181T (10)
rtM204I/V+rtV173L (10)
rtV173L+rtL180M+rtM204V (10)
rtM204V+rtL180M+rtA181S (10)
rtM204V+rtL80I+rtL180M (9)
*VEM*
rtV173L+rtL180M+rtM204V (10)
*(9) Galluzzo et al 2014*
*(10) Aoudjane et al 2014*

**Mozambique**
*RAM*
rtI169T/L# ;
rtN236T#; rtA194T#;
rtM250V/I#
rtL180M+rtS202G+rtM204V/I#
*(11) Wandeler et al 2016*

**South Africa**
*RAM*
rtM204I/V@*(13, 14, 16)
rtV173L#* (15, 17)
rtV214# (15)
rtL180M+rtM204V#(15)
*VEM*
sR48T (15)
sQ10R (15)
*(13) Selabe et al 2007;* *(14) Selabe et al 2009; (15) Makondo et al 2012*
*(16) Powell et al 2015*
*(17) Andersson et al 2013*

**South Africa & Zambia**
*RAM*
rtM204I; rtT184S#; rtN236T; rtI169L#
*(18) Hamers et al 2013*

**South Africa & Botswana**
*RAM*
rtM204I#
*(19) Matthews et al 2015*

**Zimbabwe**
*VEM & HBIg mutations*
sM133L; sD144G
*(12) Baudi et al 2017*

**Botswana**
*VEM & HBIg mutations*
sT126A; sQ129R; sM133L sF134V
*(20) Anderson et al 2015*

**Key**

Number inside the circle represents number of studies included from that country

- 🟠 studies with drug resistant mutations (DRM) and VEM
- 🟢 studies with drug resistant mutations only
- 🔵 studies VEM/ immunoglobulin mutations (HBIg)

#: Mutation identified among treatment naïve individuals
@: Mutation identified among treatment naïve and treatment experienced individuals
*: Mutation identified among HIV/HBV co-infected individuals and HBV mono-infected individuals

**Fig 3:Prevalence of HBV resistance associated mutations (RAMs) in Pol/RT proteins among HBV infected patients in Africa (A) treatment naïve; (B) treatment experienced**