1 **Genetic meta-analysis identifies 10 novel loci and functional pathways for Alzheimer's**
2 **disease risk**
3

Iris E Jansen[1,2], Jeanne E Savage[1], Kyoko Watanabe[1], Julien Bryois[3], Dylan M Williams[3], Stacy
Steinberg[4], Julia Sealock[5], Ida K Karlsson[3], Sara Hägg[3], Lavinia Athanasiu[6,7], Nicola Voyle[8],
Petroula Proitsi[9], Aree Witoelar[6,10], Sven Stringer[1], Dag Aarsland[11,12], Ina S Almdahl[13-15], Fred
Andersen[16], Sverre Bergh[17,18], Francesco Bettella[6,10], Sigurbjorn Bjornsson[19], Anne Brækhus[17,20,],
Geir Bråthen[21,22], Christiaan de Leeuw[1], Rahul S Desikan[23], Srdjan Djurovic[6,24], Logan
Dumitrescu[25], Tormod Fladby[13,14], Timothy Homan[25], Palmi V Jonsson[19,26], Arvid Rongve[27,28],
Ingvild Saltvedt[21,29], Sigrid B. Sando[21,22,], Geir Selbæk[17,30], Nathan Skenne[31], Jon Snaedal[19],
Eystein Stordal[21,32], Ingun D. Ulstein[10,15], Yunpeng Wang[6,10], Linda R White[21,22], Jens Hjerling-
Leffler[31], Patrick F Sullivan[3,33,34], Wiesje M van der Flier[2], Richard Dobson[11,35], Lea K. Davis[36,37],
Hreinn Stefansson[4], Kari Stefansson[4], Nancy L Pedersen[3], Stephan Ripke[38-40*], Ole A
Andreassen[6,10*], Danielle Posthuma[1,41,*#]
15
1. Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research,
Amsterdam Neuroscience, VU University, Amsterdam, The Netherlands.
2. Alzheimer Center and Department of Neurology, Amsterdam Neuroscience, VU University
Medical Center, Amsterdam, The Netherlands.
3. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,
Sweden.
4. deCODE Genetics/Amgen, Reykjavik, Iceland.
5. Interdisciplinary Graduate Program, Vanderbilt University, Nashville, USA.
6. NORMENT, K.G. Jebsen Centre for Psychosis Research, Institute of Clinical Medicine,
University of Oslo, Oslo, Norway.
7. Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway.
8. SGDP Centre, IoPPN, Denmark Hill, London, King's College London, London, UK.
9. Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and
Neuroscience, King's College London, London, UK.
10. Institute of Clinical Medicine, University of Oslo, Oslo, Norway
11. Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.
12. Center for Age-Related Diseases, Stavanger University Hospital, Stavanger, Norway.
13. Department of Neurology, Akershus University Hospital, Lørenskog, Norway.
14. AHUS Campus, University of Oslo, Oslo, Norway.
15. Department of Psychiatry of Old Age, Oslo University Hospital, Oslo, Norway.
16. Department of Community Medicine, University of Tromsø, Tromsø, Norway.
17. Norwegian National Advisory Unit on Ageing and Health, Vestfold Hospital Trust, Tønsberg,
Norway.
18. Centre for Old Age Psychiatry Research, Innlandet Hospital Trust, Ottestad, Norway.
19. Department of Geriatric Medicine, Landspitali University Hospital, Reykjavik, Iceland.
20. Geriatric Department, University Hospital Oslo and University of Oslo, Oslo, Norway.
21. Department of Neuroscience, Norwegian University of Science and Technology, Trondheim,
Norway.

22. Department of Neurology, St Olav's Hospital, Trondheim University Hospital, Trondheim,
Norway.
23. Neuroradiology Section, Department of Radiology and Biomedical Imaging, University of
California, San Francisco, USA.
24. Department of Medical Genetics, Oslo University Hospital, Oslo, Norway.
25. Vanderbilt Memory & Alzheimer's Center, Department of Neurology, Vanderbilt University
Medical Center, Nashville, USA.
26. Faculty of Medicine, University of Iceland, Reykjavik, Iceland.
27. Department of Research and Innovation, Helse Fonna, Oslo, Norway.
28. Department of Clinical Medicine, University of Bergen, Bergen, Norway.
29. Department of Geriatrics, St. Olav's Hospital, Trondheim University Hospital, Trondheim,
Norway.
30. Institute of Health and Society, University of Oslo, Oslo, Norway.
31. Laboratory of Molecular Neurobiology, Department of Medical Biochemistry and
Biophysics, Karolinska Institutet, Stockholm, Sweden.
32. Department of Psychiatry, Namsos Hospital, Namsos, Norway.
33. Department of Genetics, University of North Carolina, Chapel Hill, USA.
34. Department of Psychiatry, University of North Carolina, Chapel Hill, USA.
35. Farr Institute of Health Informatics Research, University College London, London, UK.
36. Department of Medicine, Division of Genetic Medicine, Vanderbilt University Medical
Center, Nashville, US.
37. Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Nashville, US.
38. Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, USA.
39. Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge,
USA.
40. Dept. of Psychiatry and Psychotherapy, Charité - Universitätsmedizin, Berlin, Germany.
41. Department of Clinical Genetics, VU University Medical Center, Amsterdam, The
Netherlands.
72
73 * These authors contributed equally to this work
74
#Correspondence to: Danielle Posthuma: Department of Complex Trait Genetics, VU
University, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands. Phone: +31 20 598
2823, Fax: +31 20 5986926, d.posthuma@vu.nl

Word count: Introductory paragraph: 206; main text: 3,825; Online methods: 5,412
Display items: 5 (Figures 4)
Includes **Supplementary Figures 1-7, Supplementary Tables 1-20.**
81
82

Abstract

Alzheimer's disease (AD) is the most frequent neurodegenerative disease with more than 35

million people affected worldwide, and no curative treatment currently available. AD is highly

heritable and recent genome-wide meta-analyses have conclusively identified over 20 genomic

loci associated with the late onset type of AD, yet only explaining a small proportion of the

genetic variance indicating that undiscovered loci exist. Here, we present the largest genome-

wide association study of AD and AD-by-proxy (71,880 AD cases, 383,378 controls) AD-by-proxy

status is based on parental AD diagnosis, and showed strong genetic correlation with AD (0.81),

Genetic meta-analysis identified 29 risk loci (confirming 19, 10 novel), implicating 215 potential

causative genes. Independent replication further supports the genetic involvement of the novel

loci in AD. Associated genes are strongly expressed in immune related tissues and cell types

(spleen, liver and microglia). Furthermore, gene-set analyses confirm the genetic contribution

of biological mechanisms involved in lipid-related processes and degradation of amyloid

precursor proteins. We show strong genetic correlations with multiple health-related

outcomes, and Mendelian randomisation results suggest a protective effect of cognitive ability

on AD risk. These results are a step forward in identifying the genetic factors that contribute to

AD risk, and add novel insights into the neurobiology of AD.

Main text

Alzheimer's disease (AD) is the most frequent type of dementia with roughly 35 million affected

to date.[1] Results from twin studies indicate that AD is highly heritable, with estimates ranging

between 60-80%.[2] Genetically, AD can be roughly divided into 2 subgroups: 1) familial early-

onset cases that are relatively often explained by rare variants with a strong effect,[3] and 2) late-

onset cases that are influenced by multiple common variants with low effect sizes.[4] Segregation

analyses have linked several genes to the first subgroup, including *APP*[5], *PSEN1*[6] and *PSEN2*[7].

The identification of these genes has resulted in valuable insights into a molecular mechanism

with an important role in AD pathogenesis, the amyloidogenic pathway,[8] providing a prominent

example of how gene discovery can add to biological understanding of disease aetiology.

Besides the identification of a few rare genetic factors (e.g. *TREM2*[9] and *ABCA7*[10]),

genome-wide association studies (GWAS) have mostly discovered common risk variants for the

more complex late-onset type of AD. *APOE* is the strongest genetic risk locus for late-onset AD,

where heterozygous and homozygous Apoε4 carriers are predisposed for a 3-fold and 15-fold

increase in risk, respectively.[11] A total of 19 additional GWAS loci have been described using a

discovery sample of 17,008 AD cases and 37,154 controls, followed by replication of the

implicated loci with 8,572 AD patients and 11,312 controls.[4] The current more than 20 loci do

not fully explain the heritability of AD and increasing the sample size is likely to lead to more

genome-wide significant risk loci, which will aid in understanding biological mechanisms

involved in the risk for AD.

In the current study, we included 455,258 individuals of European ancestry meta-

analysed in 3 stages (**Figure 1**). These consisted of 24,087 clinically diagnosed late-onset AD

cases, paired with 55,058 controls (phase 1). In phase 2, we analysed an AD-by-proxy

phenotype, based on individuals in the UK Biobank (UKB) for whom parental AD status was

available (N proxy cases=74,793; N proxy controls=328,320; **Online Methods**). In the UKB

sample, parental diagnosis for AD was available for N=376,113 individuals, of whom 393

individuals had a known diagnosis of AD themselves (identified from medical register data). The

high heritability of AD implies that case status for offspring can to some extent be inferred from

parental case status and that offspring of AD parents are likely enriched for a higher genetic AD

risk load. We thus defined individuals with one or two parents with AD as proxy cases

(N=47,793), while putting more weight on the proxy cases with 2 parents. Similarly, the proxy

controls include subjects with 2 parents without AD (N=328,320), where older cognitively

normal parents were up-weighted as proxy controls to account for the higher likelihood that

younger parents may still develop AD. As the proxy phenotype is not a pure measure of an

individual's AD status and may include individuals that never develop AD, genetic effect sizes

will be somewhat underestimated. However, the proxy case-control sample is very large (N

proxy cases=47,793; N proxy controls=328,320), and therefore adds a substantial amount of

power to detect genetic effects for AD. We first analysed the clinically defined case-control

samples separately from the by-proxy case control sample to allow investigation of overlap in

genetic signals for these two measurements of AD risk. Finally in phase 3, we meta-analysed all

individuals of phase 1 and phase 2 together.

Genome-wide meta-analysis for AD status

Phase 1 involved a genome-wide meta-analysis for AD case-control status using cohorts

collected as part of 3 independent main consortia (PGC-ALZ, IGAP and ADSP), totalling 79,145

individuals of European ancestry and 9,862,738 genetic variants passing quality control **(Figure**

1, Supplementary Table 1). The ADSP cohort obtained whole exome sequencing data from

4,343 cases and 3,163 controls, while the remaining datasets consisted of genotype SNP array

data. AD patients were diagnosed according to generally acknowledged diagnostic criteria, such

as the NINCDS-ADRDA (See **Methods**). All cohorts for which we had access to the raw genotypic

data were subjected to a standardized quality control pipeline, and GWA analyses were run per

cohort and then included in a meta-analysis, alongside one dataset (IGAP) for which only

summary statistics were available (see **Methods**). The full sample liability SNP-heritability

(h^2_{SNP}), estimated with the more conservative LDSC method, was 0.055 (SE=0.0099), implying

that 5.5% of AD heritability can be explained by the tested SNPs. This is in line with previous

estimates for IGAP (6.8%) also estimated by LDSC regression method, which is based on

summary statistics.[12,13] We do note that previously reported estimates using a method based

on raw genotypes (Genome-wide Complex Trait Analysis, GCTA), estimated that up to 53% of

total phenotypic variance in AD could be explained by common SNPs, of which up to 6% could

be explained by *APOE* alone, up to 13% by the then known variants, and up to 25% by

undiscovered loci.[14,15] The conservative LDSC estimate of h^2_{SNP} is presumably a consequence of

the underlying LDSC algorithm which is based on common HapMap SNPs and excludes all

variants with extreme associations.

The λ_{GC}=1.10 indicated the presence of inflated genetic signal compared to the null

hypothesis of no association. The linkage disequilibrium (LD) score intercept[13] was 1.044

(SE=0.0084) indicating that most inflation could be explained by polygenic signal

(**Supplementary Figure 1**). In the meta-analysis of AD case-control status, 1,067 variants

indexed by 51 lead SNPs in approximate linkage equilibrium (r^2<0.1) reached genome-wide

significance (GWS; P<5×10^{-8}) (**Supplementary Figure 1; Supplementary Table 2)**. These were

located in 18 distinct genomic loci **(Table 1)**. 15 of these loci confirmed previous findings

(Lambert et al[4]) in a partially overlapping sample of the current study. The 3 remaining loci

(lead SNPs* rs7657553, rs11257242 and rs2632516) have been linked more recently to AD in a

genetic study[16] of AD-related triglyceride levels while conditioning on lipid levels and in a

transethnic genome-wide association study of AD.[17]

We next (phase 2) performed a GWAS for AD-by-proxy using 376,113 individuals of

European ancestry from the UKB version 2 release using parental AD status weighted by age

and corrected for population frequency to construct an AD-by-proxy status (**Figure 1; see**

Methods). The LD score intercept was 1.022 (SE=0.0099) indicating that most of the inflation in

genetic signal (λ_{GC}=1.071) could be explained by polygenic signal (**Supplementary Figure 1b**).

For AD-by-proxy, 719 GWS variants were indexed by 61 lead SNPs in approximate linkage

equilibrium (r^2<0.1) reached genome-wide significance (P<5×10^{-8}), located in 13 loci

(**Supplementary Figure 1a**). Of these, 8 loci overlapped with the significantly associated loci

identified for clinical AD case control status (**Table 1**).

We observed a strong genetic correlation of 0.81 (SE= 0.185, using LDScore regression)

indicating substantial overlap between genetic effects on clinical AD and AD-by-proxy status,

beyond shared GWS SNPs. Sign concordance tests indicated that 50.4% of all LD-independent

(r^2<0.1) genome-wide SNPs (significant and non-significant) had consistent direction of effects

between the two phenotypes (N=344,581 overlapping SNPs), slightly greater than the chance

expectation of 50% (exact binomial test P=2.45×10^{-7}). Of the 51 lead SNPs identified by the

case-control meta-analysis, all were available in UKB and 96.1% were sign-concordant

(P=2.98x10^{-12}), while of the 61 GWS lead SNPs identified in UKB, 48 were available in the case-

*We choose to not report the gene that is in closest proximity to the lead SNP as the ID for the locus, as this incorrectly implies that the gene is the causal gene for AD pathogenesis. We therefore believe it is preferred to use the rs-number as an ID for the locus, and aim to highlight the most likely causal genes with more sophisticated functional interpretation analyses in later sections of this study.

control meta-analysis and 99.7% of these were sign-concordant (P=5.98×10^{-14}). Such

substantial overlap suggests that the AD-by-proxy phenotype captures a large part of the

associated genetic effects on AD.

Given the high genetic overlap, in phase 3, we conducted a meta-analysis on the clinical

AD case-control GWAS and the AD-by-proxy GWAS (**Figure 1**), comprising a total sample size of

455,258 (71,880 (proxy) cases and 383,378 (proxy) controls). The LD score intercept was 1.0018

(SE=0.0109) indicating again that most of the inflation in genetic signal (λ_{GC}=1.0833) could be

explained by polygenic signal (**Supplementary Figure 1b**). There were 2,357 GWS variants,

which were represented by 94 lead SNPs, located in 29 loci (**Table 1**, **Figure 2**). These included

15 of the 18 loci detected in our case-control analyses, all of the 13 detected in the AD-by-proxy

analyses, as well as 9 loci that were sub-threshold in both individual analyses, but reached

significance in the meta-analysis. All 2,160 GWS SNPs that were available in both the case-

control and AD-by-proxy sub-samples were sign concordant (exact binomial test P<1×10^{-300}),

including all of the 82 available independent lead SNPs (P=1.68x10^{-23}). There was evidence of

substantial association signal in both AD and AD-by-proxy for 22 (out of 27 overlapping) loci for

which SNP(s) in each locus had a robust P-value (P < .05/94 independent signals). Of the 29

associated loci, 16 loci were previously identified by the GWAS of Lambert et al.,[4] and 13 were

newly implicated AD loci in our meta-analysis. Three of these (with lead SNPs rs184384746,

rs187370608 and rs114360492) were only available in the UKB cohort (**Table 1**). Verifying our

results against other[9,18] and more recent[16,19] genetic studies on AD, 3 loci (rs11257238,

rs28394864 and rs187370608) were previously discovered, leaving 10 novel loci (rs4575098,

rs184384746, rs6448453, rs114360492, rs442495, rs117618017, rs59735493, rs113260531,

rs76726049 and rs76320948). Considering all loci of Lambert et al, we were unable to replicate

4 loci (*MEF2C, NME8, CELF1* and *FERMT2**) at a genome-wide significance level (observed *P*-

values were 1.6 x 10^{-5} to 0.0011), which was mostly caused by a lower association signal in the

UKB dataset (**Supplementary Table 3**). By contrast, Lambert et al. were unable to replicate the

DSG2 and *CD33* locus with their second stage of their study. In our study, *DSG2* locus also

lacked significance (meta-analysis *P* = 0.030; UKB analysis *P* = 0.766; **Table 1**), implying

invalidation of this locus, while we did observe a significant association (meta-analysis *P* = 6.34

x 10^{-9}; UKB analysis *P* = 4.97 x 10^{-5}) for *CD33* (rs3865444 in **Table 1***),* implying a genuine genetic

association to AD risk.

Next, we aimed to find further support for the novel findings of the phase 3 meta-

analysis, by using an independent Icelandic cohort (deCODE[20,21]), including 6,593 AD cases and

174,289 controls (**Figure 1; see Methods**; **Supplementary Table 4**). We were unable to test

support of two loci as the lead SNPs (and SNPs in high LD) were missing in deCODE, which is

most likely due to imputation differences (HRC reference for UKB dataset vs. 1000G reference

for deCODE dataset). For 7 of the 8 novel loci tested for replication, we observed the same

direction of effect in the deCODE cohort. Furthermore, 4 loci (rs6448453, rs442495,

rs117618017, rs76320948) showed nominal significant association results (P < 0.05) for the

same SNP or a SNP in high LD (r^2 > 0.9) within the same locus (two-tailed binomial test

P=3.7x10^{-4}). The locus on chromosome 1 (rs45759098) was very close to significance (P =

0.053). Apart from the novel loci, we also observed sign concordance for 95.6% of the 90 lead

SNPs in all loci from the meta-analysis (*P*=1.60x10^{-20}) that were available in deCODE (out of 94).

* For straightforward comparison to this GWAS, we do here report the genes in closest
proximity to the lead SNP. However, we would like to point out that GWAS findings implicate a
genomic locus, and that the closest gene is not necessarily the causal gene.

As an additional method of testing for replication using genome-wide polygenic score

prediction,[22] the current results explain 7.1% of the variance in clinical AD at a low best fitting

P-threshold of 1.69×10^{-5} ($P=1.80\times10^{-10}$) in an independent sample of 761 individuals. When

excluding the *APOE*-locus (chr19: 45020859-45844508), the results explain 3.9% of the variance

with a best fitting P-threshold of 3.5×10^{-5} ($P=1.90\times10^{-6}$).

Functional interpretation of genetic variants contributing to AD and AD-by-proxy

Next, we conducted a number of in silico follow-up analyses to interpret our findings in a

biological context. Functional annotation of all GWS SNPs ($n=2,178$) in the associated loci

showed that SNPs were mostly located in intronic/intergenic areas, yet in regions that were

enriched for chromatin states 4 and 5, implying effects on active transcription **(Figure 3A, 3B**

and 3C; Supplementary Table 5). 24 GWS SNPs were exonic non-synonymous (ExNS) (**Figure**

3A; Supplementary Table 6) with likely deleterious implications on gene function. Converging

evidence of strong association ($Z> |7|$) and a high observed probability of a deleterious variant

effect (CADD[23] score≥30) was found for rs75932628 (*TREM2*), rs142412517 (*TOMM40*) and

rs7412 (*APOE*). The first two missense mutations are rare (MAF=0.002 and 0.001, respectively)

and the alternative alleles were associated with higher risk for AD*.* The latter *APOE* missense

mutation is the well-established protective allele Apoε2*.* The effect sizes for ExNS ranged from

moderate to high. **Supplementary Tables 5 and 6** present a detailed annotation catalogue of

variants in the associated genomic loci. Partitioned analysis,[24] excluding SNPs with extremely

large effect sizes (i.e. *APOE* variants) showed enrichment for h^2_{SNP} for variants located in

H3K27ac marks (Enrichment=3.18, $P=9.63\times10^{-5}$), which are associated with activation of

transcription, and in Super Enhancers (Enrichment=3.62, P=2.28×10^{-4}), which are genomic

regions where multiple epigenetic marks of active transcription are clustered (**Figure 3D;**

Supplementary Table 7). Heritability was also enriched in variants on chromosome 17

(Enrichment=3.61, P=1.63x10^{-4}) and we observed a trend of enrichment for variants with high

minor allele frequencies (Enrichment=3.31, P=2.85x10^{-3}), (**Supplementary Figure 3;**

Supplementary Tables 8 and 9). Although a large proportion (23.9%) of the heritability can be

explained by SNPs on chromosome 19, this enrichment is not significant, due to the large

standard errors around this estimate (**Supplementary Table 8**). Overall these results suggest

that, despite some nonsynonymous variants likely contributing to AD risk, most of the GWS

SNPs are located in non-coding regions, and are enriched for regions that have an activating

effect on transcription of coding regions.

Implicated genes

To link the associated variants to genes, we applied three gene-mapping strategies

implemented in FUMA[25] (**Online Methods**). We used all SNPs with a P-value < 5x10-8 and r^2 of

0.6 with one of the independently associated SNPs for gene-mapping. *Positional* gene-mapping

aligned SNPs to 100 genes by their location within or immediately up/downstream (+/-10kb) of

known gene boundaries, *eQTL (expression quantitative trait loci)* gene-mapping matched cis-

eQTL SNPs to 170 genes whose expression levels they influence in one or more tissues, and

chromatin interaction mapping linked SNPs to 21 genes based on three-dimensional DNA-DNA

interactions between each SNP's genomic region and nearby or distant genes, which we limited

to include only interactions between annotated enhancer and promotor regions (**Figure 3B and**

3C; Supplementary Figure 4; Supplementary Tables 10 and 11). This resulted in 192 uniquely

mapped genes, 80 of which were implicated by at least two mapping strategies and 17 by all 3

(**Figure 4E**). Eight genes (*HLA-DRB5, HLA-DRB1, HLA-DQA, HLA-DQB1, KAT8, PRSS36, ZNF232*

and *CEACAM19*) are particularly notable as they are implicated via eQTL association in the

hippocampus, a brain region highly affected early in AD pathogenesis (**Supplementary Table**

10). Of special interest is the locus on chromosome 8 (rs4236673). In the GWAS by Lambert et

al.[4], this locus was defined as 2 distinct loci (*CLU* and *PTK2B*), while our meta-analysis specified

this locus as a single locus based on LD-patterns. This is also supported by a chromatin

interaction between the two regions (**Figure 3E**), which is observed in two immune-related

tissues – the spleen and liver (**Supplementary Table 11**). Chromosome 16 contains a locus

implicated by long-range eQTL association (**Figure 3F**) clearly illustrating more distant genes can

be affected by a genetic factor (**Figure 3F**) and emphasising the relevance of considering

putative causal genes or regulatory elements not solely on the physical location but also on

epigenetic influences. **Supplementary Figure 4** displays chromatin interactions for all

chromosomes containing significant GWAS loci.

Although these gene-mapping strategies imply multiple putative causal genes per GWAS

locus, several of these genes in the novel loci (and significantly replicated by the deCODE

cohort) are of particular interest, as the genes have functional or previous genetic association

to AD. For locus 1 in **Supplementary Table 10**, *ADAMTS4* encodes a protein of the ADAMTS

family which has a function in neuroplasticity and has been extensively studied for their role in

AD pathogenesis.[26] For locus 19, the obvious most likely causal gene is *ADAM10,* as this gene

has been conclusively associated to AD through the effect of rare coding variants. However this

is the first time that this gene is implicated as a common risk factor for AD. The lead SNP for

locus 20 is a nonsynonymous variant in exon 1 of *APH1B,* which encodes for a protein part of

the *γ*-secretase complex cleaving *APP.*[27] Although previously reported functional information

on genes can be of great value, it is preferable to consider all implicated genes as putative

causal factors to guide potential functional follow-up experiments.

We next performed genome-wide gene-based association analysis (GWGAS) using

MAGMA.[28] This method annotates SNPs to known protein-coding genes to estimate aggregate

associations based on all SNPs in a gene. It differs from the gene-mapping strategies in FUMA as

it provides a statistical gene-based test, whereas FUMA maps individually significant SNPs to

genes. With GWGAS, we identified 97 significantly associated genes (**Supplementary Figure 5;**

Supplementary Table 12), of which 74 were also mapped by FUMA **(Figure 4E)**. In total, 16

genes were implicated by all four strategies (**Supplementary Table 13),** of which 7 genes (*HLA-*

DRA, HLA-DRB1, PTK2B, CLU, MS4A3, SCIMP and *RABEP1*) are not located in the *APOE*-locus,

and therefore of high interest for further investigation.

Gene-sets implicated in AD and AD-by-proxy

Using the gene-based P-values, we performed gene-set analysis for 6,994 biological-pathway-

based gene-sets, 53 tissue expression-based gene-sets and 39 brain single cell expression based

gene-sets (24 derived from mouse data and 15 derived from human data). We found 4 Gene

Ontology[19] gene-sets that were significantly associated with AD risk: *Protein lipid complex*

($P=3.93\times10^{-10}$), *Regulation of amyloid precursor protein catabolic process* ($P=8.16\times10^{-09}$), *High*

density lipoprotein particle ($P=7.81\times10^{-8}$), and *Protein lipid complex assembly* ($P=7.96\times10^{-7}$)

(Figure 4A; Supplementary Tables 14 and 15). Conditional analysis on the *APOE* locus showed

associations with AD for these four gene-sets independent of the effect of *APOE*, as they

remained significantly associated (P<0.0125), yet less strong, suggesting that *APOE* is

contributing a substantial part to the association signal, but does not completely drive the

signal. There was overlap between genes included in the 4 gene-sets, and conditioning on each

significant gene-set association showed that 3 gene-sets were associated with AD

independently of each other (**Supplementary Table 14 and 15**). All 25 genes of the *High density*

lipoprotein particle pathway are also part of the *Protein lipid complex* (conditional analysis

P=0.18)*,* and these pathways are therefore not interpretable as independent associations.

Linking gene-based *P*-values to tissue- and cell-type-specific gene-sets**,** no association

survived the stringent Bonferroni correction, which corrected for all tested gene-sets (i.e. 6,994

GO categories, 54 tissues and 39 cell types). However, we did observe associations when

correcting only for the number of tests within all tissue types or cell-types. This was the case for

gene expression across immune-related tissues (**Figure 4C; Supplementary Table 16**),

particularly whole blood (P=5.61×10^{-6}), spleen (P=1.50x10^{-5}) and lung (P=4.67x10^{-4}). In brain

single-cell expression gene-set analyses, we found associations for microglia, both in the

mouse-based expression dataset (P=1.96x10$^{-3)}$ (**Figure 4B; Supplementary Table 17)** and the

human-based expression dataset (P=2.56x10^{-3}) (**Supplementary Figure 6; Supplementary Table**

18).

Cross-trait genetic influences

For a more comprehensive understanding of the genetic background of AD, we next tested

whether AD is likely to share genetic factors with other phenotypes. This might reveal some

functional insights about the genetic aetiology of AD. We conducted bivariate LDscore[13]

regression to test for genetic correlations between AD and 41 other traits for which large GWAS

summary statistics were available. We observed significant negative genetic correlations with

adult cognitive ability (r_g=-0.22, P=7.28x10-5), age of first birth (r_g=−0.33, P=1.22×10^{-4}),

educational attainment (r_g=−0.25, P=5.01×10^{-4}), and confirmed a very strong positive

correlation with previous GWAS of Alzheimer's disease (r_g=0.90, P=3.29x10^{-16}) (**Figure 4D;**

Supplementary Table 19).

We then used Generalised Summary-statistic-based Mendelian Randomisation[29] (GSMR;

see **Methods**) to test for potential credible causal associations of genetically correlated

outcomes which may directly influence the risk for AD. Due to the nature of AD being a late-

onset disorder and summary statistics for most other traits being obtained from younger

samples, we do not report tests for the opposite direction of potential causality (i.e. we did not

test for a causal effect of a late-onset disease on an early onset disease). In this set of analyses,

SNPs from the summary statistic of genetically correlated phenotypes were used as

instrumental variables to estimate the putative causal effect of these "exposure" phenotypes

on AD risk by comparing the ratio of SNPs' associations with each exposure to their associations

with AD outcome (see **Methods**). Association statistics were standardized, such that the

reported effects reflect the expected difference in odds ratio (OR) for AD as a function of every

SD increase in the exposure phenotype. We observed a protective effect of cognitive ability

(OR=0.89, 95% confidence interval[CI]: 0.85-0.92, P=5.07x10^{-9}), educational attainment

(OR=0.88, 95%CI: 0.81-0.94, *P*=3.94×10^{-4}), and height (OR=0.96, 95%CI: 0.94-0.97, *P*=1.84x10^{-8})

on risk for AD (**Supplementary Table 20; Supplementary Figure 7)**. No substantial evidence of

pleiotropy was observed between AD and these phenotypes, with <1% of overlapping SNPs

being filtered as outliers (**Supplementary Figure 7**).

Discussion

In conclusion, by using a non-conventional approach of including a by-proxy phenotype for AD

to increase sample size, we have identified novel loci and gained novel biological knowledge on

AD aetiology. Both the high genetic correlation between the standard case-control status and

the UKB by proxy phenotype (*r$_g$*=0.81) and the high rate of novel loci replication in the

independent deCODE cohort, suggest that this strategy is robust. Through extensive in silico

functional follow-up analysis, and in line with previous research,[19,30] we emphasise the crucial

causal role of the immune system - rather than immune response as a consequence of disease

pathology - by establishing variant enrichments for immune-related body tissues (whole blood,

spleen, liver) and for the main immune cells of the brain (microglia). Furthermore, we observe

informative eQTL associations and chromatin interactions within immune-related tissues for

identified genomic risk loci. Together with the AD-associated genetic effects on lipid

metabolism in our study, these biological implications strengthen the hypothesis that AD

pathogenesis involves an interplay between inflammation and lipids, as lipid changes might

harm immune responses of microglia and astrocytes, and vascular health of the brain.[31]

In accordance with previous clinical research, our study suggests an important role for

protective effects of several human traits on AD. As an example, cognitive reserve has been

proposed as a protective mechanism in which the brain aims to control brain damage with prior

existing cognitive processing strategies.[32] Our findings imply that some component of the

genetic factors for AD might affect cognitive reserve, rather than being involved in AD-

pathology-related damaging processes, influencing AD pathogenesis in an indirect way through

cognitive reserve. Similarly, in a largescale community-based study it was observed that AD

incidence rates declined over decades, which was specific for individuals with at minimum a

high school diploma.[33] Combined with our Mendelian randomization results for educational

attainment, this suggests that the protective effect of educational attainment on AD is

influenced by genetics.

The results of this study could furthermore serve as a valuable resource (e.g.

Supplementary Tables 10 and 13) for selection of promising genes for functional follow-up

experiments. We anticipate that functional interpretation strategies and follow-up experiments

will result in a comprehensive understanding of sporadic late-onset AD aetiology, which will

serve as a solid foundation for future AD drug development and stratification approaches.

URLs:

http://ukbiobank.ac.uk

https://www.ncbi.nlm.nih.gov/gap

http://fuma.ctglab.nl

http://ctg.cncr.nl/software/magma

http://genome.sph.umich.edu/wiki/METAL_Program

https://github.com/bulik/ldsc

http://ldsc.broadinstitute.org/

https://data.broadinstitute.org/alkesgroup/LDSCORE/

http://www.genecards.org

http://www.med.unc.edu/pgc/results-and-downloads

http://software.broadinstitute.org/gsea/msigdb/collections.jsp

https://www.ebi.ac.uk/gwas/

https://github.com/ivankosmos/RegionAnnotator

http://cnsgenomics.com/software/gsmr/

Acknowledgments: This work was funded by The Netherlands Organization for Scientific

Research (NWO VICI 453-14-005) and the Sophia Foundation for Scientific Research (grant nr:

S14-27). The analyses were carried out on the Genetic Cluster Computer, which is financed by

the Netherlands Scientific Organization (NWO: 480-05-003), by the VU University, Amsterdam,

The Netherlands, and by the Dutch Brain Foundation, and is hosted by the Dutch National

Computing and Networking Services SurfSARA. This research has been conducted using the UK

Biobank resource under application number 16406 and the public ADSP dataset, obtained

through the Database of Genotypes and Phenotypes (dbGaP) under accession number

phs000572. We thank the International Genomics of Alzheimer's Project (IGAP) for providing

summary results data for these analyses. The investigators within IGAP contributed to the

design and implementation of IGAP and/or provided data but did not participate in analysis or

writing of this report. IGAP was made possible by the generous participation of the control

subjects, the patients, and their families. The i–Select chips was funded by the French National

Foundation on Alzheimer's disease and related disorders. EADI was supported by the LABEX

(laboratory of excellence program investment for the future) DISTALZ grant, Inserm, Institut

Pasteur de Lille, Université de Lille 2 and the Lille University Hospital. GERAD was supported by

the Medical Research Council (Grant n° 503480), Alzheimer's Research UK (Grant n° 503176),

the Wellcome Trust (Grant n° 082604/2/07/Z) and German Federal Ministry of Education and

Research (BMBF): Competence Network Dementia (CND) grant n° 01GI0102, 01GI0711,

01GI0420. CHARGE was partly supported by the NIH/NIA grant R01 AG033193 and the NIA

AG081220 and AGES contract N01–AG–12100, the NHLBI grant R01 HL105756, the Icelandic

Heart Association, and the Erasmus Medical Center and Erasmus University. ADGC was

supported by the NIH/NIA grants: U01 AG032984, U24 AG021886, U01 AG016976, and the

Alzheimer's Association grant ADGC–10–196728. This paper represents independent research

funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at

South London and Maudsley NHS Foundation Trust and King's College London. The views

expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the

Department of Health.

Genotyping for the Swedish Twin Studies of Aging was supported by NIH/NIA grant R01

AG037985. Genotyping in TwinGene was supported by NIH/NIDDK U01 DK066134. WvdF is

recipient of Joint Programming for Neurodegenerative Diseases (JPND) grants PERADES (ANR-

13-JPRF-0001) and EADB (733051061). JB was supported by a grant from the Swiss National

Science Foundation. JHL was supported by the Swedish Research Council (Vetenskapsrådet,

award 2014-3863), the Wellcome Trust (108726/Z/15/Z), and the Swedish Brain Foundation

(Hjärnfonden). NS was supported by the Wellcome Trust (108726/Z/15/Z). RD was supported

by National Institute for Health Research University College London Hospital's Biomedical

Research Centre, Arthritis Research UK, the British Heart Foundation, Cancer Research UK, the

Chief Scientist Office, the Economic and Social Research Council, the Engineering and Physical

Sciences Research Council, the National Institute for Social Care and Health Research, and the

Wellcome Trust (grant number MR/K006584/1).

We thank the numerous participants, researchers, and staff from many studies who collected

and contributed to the data. Additional acknowledgments can be found in the Supplementary

Information file. Summary statistics will be made available for download upon publication from

http://ctglab.vu.nl.

Author Contributions: I.E.J. and J.E.S. performed the analyses. D.P. and O.E.A. conceived the

idea of the study. D.P. and S.R. supervised analyses. S.St. performed QC on the UK Biobank data

and wrote the analysis pipeline. K.W. constructed and applied the FUMA pipeline for

performing follow-up analyses. J.B. conducted the single cell enrichment analyses. J.H.L and

N.S. contributed data. D.P. and I.E.J. wrote the paper. All authors critically reviewed the paper.

Author Information: PF Sullivan reports the following potentially competing financial interests:

Lundbeck (advisory committee), Pfizer (Scientific Advisory Board member), and Roche (grant

recipient, speaker reimbursement). JHL: Cartana (Scientific Advisor) and Roche (grant recipient).

All other authors declare no financial interests or potential conflicts of interest.

Correspondence and requests for materials should be addressed to d.posthuma@vu.nl.

475
476

477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524

Table 1. Summary statistics for the meta-analysis of case-control status, by proxy phenotype and both.
Note: Independent lead SNPs are defined by r2 < .1; distinct genomic loci are >250kb apart. Allele1 is the effect allele for the meta association statistic. Meta-analysis effect direction (column AL) is in the following order: ADSP, IGAP, UKB, PGC-ALZ, note that the first cohort is often missing as this concerns exome sequencing data. Corrected P value for significance = 5E-08 (marked as boldfaced and underlined values). Note that the lead SNP can differ between the distinct analyses, while it tags the same locus

		Case-control status						By proxy AD						Overall						
Locus	Chr	SNP	bp	Allele 1	Allele 2	Z	p	SNP	bp	Allele 1	Allele 2	Z	p	SNP	bp	Allele 1	Allele 2	Z	p	direction
1	1	rs4575098	161155392	A	G	3.78	0.000157	rs4575098	161155392	A	G	5.39	6.88E-08	rs4575098	161155392	A	G	6.36	**2.05E-10**	?+++
2	1	rs6656401	207692049	A	G	8.54	**1.39E-17**	rs679515	207750568	T	C	6.13	**8.85E-10**	rs2093760	207786828	A	G	8.82	**1.10E-18**	++++
3	2	rs4663105	127891427	C	A	11.21	**3.58E-29**	rs4663105	127891427	C	A	10.54	**5.46E-26**	rs4663105	127891427	C	A	13.94	**3.38E-44**	?+++
4	2	rs10933431	233981912	G	C	-4.79	1.67E-06	rs10933431	233981912	G	C	-4.71	2.51E-06	rs10933431	233981912	G	C	-6.13	**8.92E-10**	?---
5	3	NA						rs184384746	57226150	T	C	5.70	**1.24E-08**	rs184384746	57226150	T	C	5.69	**1.24E-08**	??+?
6	4	rs6448453	11026028	A	G	2.26	0.023641	rs6448451	11024682	G	C	5.70	**1.19E-08**	rs6448453	11026028	A	G	6.00	**1.93E-09**	?++
7	4	rs7657553	11723235	A	G	5.60	**2.16E-08**	rs7657553	11723235	A	G	-0.27	0.79	rs7657553	11723235	A	G	1.95	0.051	?-+
8	6	rs9269853	32550322	A	C	5.56	**2.66E-08**	rs6931277	32583357	T	A	-5.22	1.78E-07	rs6931277	32583357	T	A	-6.49	**8.41E-11**	?---
9	6	NA						rs187370608	40942196	A	G	8.26	**1.45E-16**	rs187370608	40942196	A	G	8.26	**1.45E-16**	??+?
10	6	rs9381563	47432637	C	T	5.84	**5.35E-09**	rs9381563	47432637	C	T	4.46	8.10E-06	rs9381563	47432637	C	T	6.33	**2.52E-10**	?+++
11	7	rs1859788	99971834	A	G	-5.82	**6.05E-09**	rs7384878	99932049	C	T	-6.34	**2.38E-10**	rs1859788	99971834	A	G	-7.93	**2.22E-15**	----
12	7	rs11763230	143108841	T	C	-6.67	**2.58E-11**	rs7810606	143108158	T	C	-4.89	1.01E-06	rs7810606	143108158	T	C	-6.62	**3.59E-11**	?---
13	7	NA						rs114360492	145950029	T	C	5.99	**2.1E-09**	rs114360492	145950029	T	C	5.99	**2.10E-09**	??+?
14	8	rs4236673	27464929	A	G	-9.14	**6.36E-20**	rs1532278	27466315	T	C	-5.78	**7.45E-09**	rs4236673	27464929	A	G	-8.98	**2.61E-19**	----
15	10	rs11257242	11721119	C	G	-5.58	**2.38E-08**	rs11257238	11717397	C	T	4.02	5.84E-05	rs11257238	11711397	C	T	5.69	**1.26E-08**	?+++
16	11	rs7935829	59942815	G	A	-7.16	**8.21E-13**	rs1582763	60021948	A	G	-5.86	**4.72E-09**	rs2081545	59958380	A	C	-7.97	**1.55E-15**	----
17	11	rs10792832	85867875	A	G	-8.56	**1.12E-17**	rs3844143	85850243	C	T	-6.56	**5.31E-11**	rs867611	85776544	G	A	-8.75	**2.19E-18**	?+++
18	11	rs11218343	121435587	C	T	-6.55	**5.57E-11**	rs11218343	121435587	C	T	-4.68	2.81E-06	rs11218343	121435587	C	T	-6.79	**1.09E-11**	?---
19	14	rs12590654	92938855	A	G	-5.61	**1.98E-08**	rs12590654	92938855	A	G	-4.63	3.70E-06	rs12590654	92938855	A	G	-6.39	**1.65E-10**	?---
20	15	rs442495	59022615	C	T	-3.61	0.000309	rs442495	59022615	C	T	-5.15	2.65E-07	rs442495	59022615	C	T	-6.07	**1.31E-09**	?---
21	15	rs117618017	63569902	T	C	2.29	0.022178	rs117618017	63569902	T	C	5.15	2.64E-07	rs117618017	63569902	T	C	5.52	**3.35E-08**	+++
22	16	rs59735493	31133100	A	G	-3.34	0.000825	rs59735493	31133100	A	G	-4.63	3.72E-06	rs59735493	31133100	A	G	-5.49	**3.98E-08**	?---
23	17	rs113260531	5138980	A	G	4.66	3.21E-06	rs9916042	4984447	A	G	5.46	**4.73E-08**	rs113260531	5138980	A	G	6.12	**9.16E-10**	?+++
24	17	rs28394864	47450775	A	G	3.97	7.29E-05	rs28394864	47450775	A	G	4.50	6.80E-06	rs28394864	47450775	A	G	5.62	**1.87E-08**	?+++
25	17	rs2632516	56409089	C	G	-6.05	**1.42E-09**	rs2632516	56409089	C	G	-2.79	0.005288	rs2632516	56409089	C	G	-4.90	9.66E-07	?---
26	18	rs8093731	29088958	T	C	-5.46	**4.63E-08**	rs8093731	29088958	T	C	-0.30	0.7656	rs8093731	29088958	T	C	-2.17	0.030	?-?
27	18	rs76726049	56189459	C	T	2.06	0.039261	rs76726049	56189459	C	T	5.22	1.83E-07	rs76726049	56189459	C	T	5.52	**3.30E-08**	?+++
28	19	rs4147929	1063443	A	G	5.76	**8.64E-09**	rs3752241	1053524	G	C	-5.55	**2.87E-08**	rs111278892	1039323	G	C	6.50	**7.93E-11**	?+++
29	19	rs41289512	45351516	G	C	29.74	**2.70E-194**	rs75627662	45413576	T	C	36.79	**9.5E-296**	rs41289512	45351516	G	C	35.50	**5.79E-276**	?+++
30	19	rs76320948	46241841	T	C	4.32	1.54E-05	rs76320948	46241841	T	C	4.29	1.80E-05	rs76320948	46241841	T	C	5.46	**4.64E-08**	?++?
31	19	rs3865444	51727962	A	C	-5.48	**4.25E-08**	rs3865444	51727962	A	C	-4.06	4.97E-05	rs3865444	51727962	A	C	-5.81	**6.34E-09**	?---
32	20	rs6014724	54998544	G	A	-5.35	8.72E-08	rs6014724	54998544	T	C	-4.52	6.32E-06	rs6014724	54998544	G	A	-6.18	**6.56E-10**	?---

525 **Figure 1. Overview of analyses steps.** The main genetic analysis encompasses the procedures to detect
526 GWAS risk loci for AD. The functional analysis part includes the *in silico* functional follow-up procedures
527 with the aim to put the genetic findings in biological context. The Mendelian randomisation analysis has
528 been performed on the results of phase 1 to account for sample overlap between our study and other
529 traits for which they have used the UKB dataset.
530
531

541 **Figure 2. GWAS results for AD risk (N=455,258).** Manhattan plot displays all associations per variant
542 ordered according to their genomic position on the x-axis and showing the strength of the association
543 with the –log10 transformed P-values on the y-axis. The y-axis is limited to enable visualization of non-
544 APOE loci.
545

Figure 3. Functional annotation of association results. a) Heritability enrichment of 28 functional
variant annotations calculated with stratified LD score regression. UTR=untranslated region;
CTCF=CCCTC-binding factor; DHS=DNaseI Hypersensitive Site; TFBS=transcription factor binding site;
DGF=DNAaseI digital genomic footprint; **b)** Functional effects of variants in genomic risk loci of the
meta-analysis – the second bar shows distribution for exonic variants only; **c)** Distribution of
RegulomeDB score for variants in genomic risk loci, with a low score indicating a higher probability of
having a regulatory function. **d)** Distribution of minimum chromatin state across 127 tissue and cell
types for variants in genomic risk loci, with lower states indicating higher accessibility and states 1-7
referring to open chromatin states. **e)** Zoomed-in circos plot of chromosome 8. **f) Z**oomed-in circos plot
of chromosome 16. Circos plots show implicated genes by significant loci, where blue areas indicate
genomic risk loci, green indicates eQTL associations and orange indicates chromatin interactions. Genes
mapped by both eQTL and chromatin interactions are red. The outer layer shows a Manhattan plot
containing the negative log10-transformed P-value of each SNP in the GWAS meta-analysis of AD. Full
circos plots of all autosomal chromosomes are provided in Supplementary Figures 4.

Figure 4. Functional implications based on gene-set analysis, genetic correlations and functional
annotations. The gene-set results are displayed per category of biological mechanisms (A), brain cell-
types (B) and tissue types (C). The red horizontal lines indicates the significance threshold corrected for
all gene-set tests of all categories, while the blue horizontal lines display the significance threshold
corrected only for the number of tests within the three categories (i.e. gene-ontology, tissue expression,
single cell expression). (D) Genetic correlations between AD and other heritable traits. (E) Venn diagram
showing the number of genes mapped by four distinct strategies.
573

Online methods

<u>1.1 Study Cohorts</u>

1.1.1 PGC-ALZ cohorts

Three non-public datasets (the Norwegian DemGene network, The Swedish Twin Studies of

Aging and TwinGene) were meta-analyzed as part of the Alzheimer workgroup initiative of the

Psychiatric Genomic Consortium (PGC-ALZ).

We collected genotype data from the Norwegian DemGene Network consisting of 2,224

cases and 1,855 healthy controls. The DemGene Study is a Norwegian network of clinical sites

collecting cases from Memory Clinics based on standardised examination of cognitive,

functional and behavioural measures and data on progression of most patients. We diagnosed

2,224 cases of AD from 7 studies: the Norwegian Register of persons with Cognitive Symptoms

(NorCog), the Progression of Alzheimer's Disease and Resource use (PADR), the Dementia Study

of Western Norway (DemVest), the AHUS study, the Dementia Study in Rural Northern Norway

(NordNorge), HUNT Dementia Study and Nursing Home study, and the TrønderBrain study.

These cases were diagnosed according to the recommendations from the National Institute on

Aging–Alzheimer's Association (NIA/AA) (AHUS), the NINCDS-ADRDA criteria (DemVest and

TrønderBrain) or the ICD-10 research criteria (NorCog, PADR, NordNorge and HUNT). The

controls from Norway were obtained through the AHUS, NordNorge, HUNT and TrønderBrain

studies. Controls were screened with standardized interview and cognitive tests. Genotypes of

the 4079 individuals from the DemGene Study were obtained with Human Omni Express-24

v.1.1 (Illumina Inc., San Diego, CA, USA) at deCODE Genetics (Reykjavik, Iceland). To increase

the statistical power of our association analysis, the controls were combined with additional

5786 population controls from Norwegian blood donor samples (Oslo University Hospital,

Ullevål Hospital, Oslo) and controls from Thematically Organized Psychosis Research (TOP)

Study27 (between 25-65 years). Control subjects of TOP Study were of Caucasian origin without

history of moderate/severe head injury, neurological disorder, mental retardation and were

excluded if they or any of their close relatives had a lifetime history of a severe psychiatric

disorder, a history of medical problems thought to interfere with brain function or significant

illicit drug use.

The Swedish Twin Studies of Aging (STSA) (n cases = 398, n controls = 1079) includes

three sub-studies of aging within the Swedish Twin Registry[34]: The Swedish Adoption/Twin

Study of Aging (SATSA)[35], Aging in Women and MEN (GENDER)[36], and The Study of Dementia in

Swedish Twins (HARMONY)[37]. Informed consent was obtained from all participants and the

studies were approved by the Regional Ethics Board in Stockholm and the Institutional Review

Board at the University of Southern California. DNA was extracted from blood samples and

genotyped using Illumina Infinium PsychArray. Alzheimer's disease patients were diagnosed as

part of the studies according to the NINCDS/ADRDA criteria[38]. In addition, information on

disease after last study participation was retrieved from three population-based health care

registers: The National Patient Register, the Causes of Death Register, and the Prescribed Drug

Register.

TwinGene[34] is a population-based study of older twins drawn from the Swedish Twin

Registry. Written informed consent was obtained from all participants and the study was

approved by the Regional Ethics Board in Stockholm. DNA was extracted from blood samples

and genotyped using Illumina Human OmniExpress for 1791 individuals. Information about

Alzheimer's disease (n cases = 343, n controls = 9070) was extracted from the National Patient

Register, the Causes of Death Register, and the Prescribed Drug Register, all of which are

population-based health care registers with nationwide coverage.

1.1.2 IGAP

Publically available (http://web.pasteur-lille.fr/en/recherche/u744/igap/igap_download.php)

genome-wide association analysis results of the International Genomics of Alzheimer's Project

(IGAP)[4] were included as one of the four cohorts that were meta-analysed in our effort. IGAP is

a large two-stage study based upon genome-wide association studies (GWAS) on individuals of

European ancestry. We focused on results of stage 1, for which IGAP used genotyped and

imputed data of 7,055,881 single nucleotide polymorphisms (SNPs) to meta-analyse four

previously-published GWAS datasets consisting of 17,008 Alzheimer's disease cases and 37,154

controls (The European Alzheimer's disease Initiative – EADI the Alzheimer Disease Genetics

Consortium – ADGC The Cohorts for Heart and Aging Research in Genomic Epidemiology

consortium – CHARGE The Genetic and Environmental Risk in AD consortium – GERAD). As the

purpose of stage 2 (2, 11,632 SNPs were genotyped and tested for association in an

independent set of 8,572 Alzheimer's disease cases and 11,312 controls) was replication of the

significantly associated loci of stage 1, we limited the inclusion of the summary statistics for our

own analyses to stage 1. Written informed consent was obtained from study participants or, for

those with substantial cognitive impairment, from a caregiver, legal guardian or other proxy,

and the study protocols for all populations were reviewed and approved by the appropriate

institutional review boards.

1.1.3 ADSP

The Alzheimer's Disease Sequencing Project (ADSP) collaboration has the aim to identify novel

genetic factors that contribute to AD risk by studying genetic sequencing data. ADSP has made

their sequencing data available through the Genotypes and Phenotyps database (dbGaP) under

the study accession: phs000572.v7.p (https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi?study_id=phs000572.v1 .p1). We have obtained access to 10,907 individuals

(5,771 cases, 5,136 controls) with whole-exome sequencing data to include as the second

cohort within our meta-analysis. A substantial proportion of the ADSP individuals were

previously also included in IGAP. We applied two strategies to prevent inflated meta-analysis

results due to sample overlap: (1) exclusion of ADSP individuals that were duplicates based on

genotype data comparison of individual level genetic data between IGAP and ADSP, (2) perform

meta-analysis while correcting for cross-study LD score regression intercept (see section 1.4.).

To accomplish the first approach we obtained access for all IGAP datasets for which individual

level genotype data was available through dbGaP (phs000160.v1.p1 - https://

www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=   phs000160.v1.p1;

phs000219.v1.p1   -   https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi?study_id=phs000219.v1.p1;   phs000372.v1.p1   -

https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000372.v1   .p1;

phs000168.v2.p2 - https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=

phs000168.v2.p2; phs000234.v1.p1 - https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi? study_id=phs000234.v1.p1) or NIAGADS (NG00026 -

https://www.niagads.org/datasets/ng00026; NG00028 -

https://www.niagads.org/datasets/ng00028; NG00029 - https://www.niagads.org/

datasets/ng00029; NG00031 - https://www.niagads.org/datasets/ng00030 ; NG00031 -

https://www.niagads.org/datasets/ng00031; NG00034 -

https://www.niagads.org/datasets/ng00034). By calculating identity-by-descent using PLINK[39],

we identified duplicates, which were excluded from the ADSP WES dataset for subsequent

analyses.

1.1.1 UK Biobank study

The current study used data from the UK Biobank[40] (UKB; www.ukbiobank.ac.uk), a large

population-based cohort that includes over 500,000 participants and aims to improve insight

into a wide variety of health-related determinants and outcomes across the UK. Between 2006

and 2010, approximately 9.2 million invitations to participate in the study were sent to

individuals aged 40-69 years who were registered with the National Health Service (NHS) and

were living within 25 miles from one of the 22 study research centers. In total, 503,325

participants were recruited in the study, from which we used a subsample of individuals of

European ancestry with available phenotypic and genotypic data (*M* age = 56.5, 54.0% female),

described in more detail below. Besides phenotypic information obtained from the NHS

registries and associated medical records, participants completed an in-person visit at one of

the study research centers where extensive self-report data were collected by questionnaire in

addition to anthropometric assessments, DNA collection from blood samples, and magnetic

resonance imaging of body and brain. All participants provided written informed consent; the

UKB received ethical approval from the National Research Ethics Service Committee North

West-Haydock (reference 11/NW/0382), and all study procedures were in accordance with the

World Medical Association for medical research. Access to the UK Biobank data was obtained

under application number 16406.

<u>1.2 UKB by proxy phenotype</u>

A proxy phenotype for Alzheimer's disease case-control status in UKB was assessed as part of

the self-report questionnaire administered during the in-person assessment. Participants were

asked to report whether their biological mother or father ever suffered from Alzheimer's

disease/dementia, and to report each parent's current age (or age at death, if applicable). Of

376,113 individuals in our analytic subsample who completed these questions, a diagnosis was

reported for 32,327 mothers (8.6%) and 17,014 fathers (4.5%), resulting in 47,793 participants

(12.7%) with one or both parents affected. We created a proxy phenotype from these questions

to index genetic risk for Alzheimer's based on parents' diagnoses. The phenotype was

constructed as a linear count of the number of affected biological parents (0, 1, or 2). The

contribution for each unaffected parent to this count was weighted by the parent's age/age at

death to account for the fact that they may not yet have passed through the period of risk for

this late-onset disease. Specifically, each affected parent contributed one full unit of "risk" to

the count, while each unaffected parent contributed a proportion of one unit of "risk" inversely

related to their age. This was calculated as the ratio of parent's age to age 100 (approximately

the 95th percentile for life expectancy in developed countries, such that weight=(100-age)/100.

The weight for unaffected parents was capped at 0.32, corresponding to a risk equivalent to

that of the maximum population prevalence of AD.[41] The phenotype thus ranged from

approximately 0-2, with values near zero when both parents were unaffected (lower for older

parents and possible values below zero if both parents were over age 100) and values of two

when both parents were affected. Participants who were uncertain or chose not to answer

questions about either parent's disease status or age were excluded from analyses, resulting in

a final N=364,859.

Additional information on Alzheimer's disease risk was obtained from national medical

records linked to participant data. This information pertained to the participants themselves

(not their parents), and was extracted from hospital records obtained between 1996 and the

present or from national death registries in the case of participants who passed away after

initial enrolment in the study, as described in more detail in the UKB resources

(http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=146641;

http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=115559). Briefly, primary and secondary

diagnoses from inpatient hospital stays and primary and secondary causes of death from death

records were recorded using ICD-10 codes. Participants with a diagnosis of "Alzheimer's

disease" (diseases of the nervous system chapter; code G30) or "Dementia in Alzheimer's

disease" (mental and behavioral disorders chapter; code F00) from any record of a hospital stay

or as a cause of death were treated as Alzheimer's cases as given the maximum possible "risk"

score of 2, regardless of affectation status of their parents. The reported rate of Alzheimer's in

parents of cases (27.4%) was more than double that of non-cases (12.7%; $\chi^2(1)$=71.7, P=2.45E-

17). There were 393 individuals in the analytic subsample classified as affected by these

records; due to the small number of cases and the limited representativeness of these types of

health records, we use this information to supplement the proxy parent phenotype rather than

as a primary outcome. This information reduces the possibility of misclassification in the proxy

phenotype method, and also allows us to evaluate the performance of the proxy phenotype

method.

<u>1.3 Genome-wide association analysis</u>

Except for IGAP (obtained summary statistics), we performed genome-wide association

analyses for the ADSP, PGC-ALZ and UKB cohorts. For the UKB dataset, quality control and

imputation procedures were slightly different, and therefore described separately in the

sections below.

1.3.1a Quality control and imputation procedures for ADSP and PGC-ALZ datasets

Prior to individual quality control steps, all datasets were filters on a max missingness of 5%.

Individuals were excluded when identified as a low quality sample (individual call rate < 0.98),

heterozygosity outlier (F +/-.20), gender mismatch (females: F >0.2, males: F < 0.2) when

comparing phenotypic and genotypic data, population outlier (defined by principal component

boundaries of 1000 Genomes European samples) or being related (PI_HAT > 0.2). Inclusion

criteria for variants encompassed a call rate > 0.98, a case-control missingness difference <

0.02, a Hardy-Weinberg equilibrium *p*-value < 10e-6 for controls (<10e-10 for cases) and a valid

association *p*-value (excluding the variants with low allele frequencies).

Pre-imputation, the ADSP and PGC-ALZ datasets were checked for palindromic variants

with allele frequency close to 0.5, incorrect reference allele definitions, false strand designation

and extreme deviations from expected allele frequencies. Subsequently the ADSP and PGC-ALZ

datasets were imputed with the 1000 Genomes Phase 3[42] reference panel. The reported SNPs

all have a considerable imputation quality (INFO score>0.591) and variants with a low allele

frequency (MAF<0.01) were excluded, resulting in a total of 7508 individuals (4343 cases and

3165 controls) and 260,934 variants for the ADSP cohort and 17477 individuals (2,736 cases and

14,471 controls) and 9,629,492 variants for the PGC-ALZ cohort.

1.3.1b Quality control and imputation for UKB dataset

We used second-release genotype data that were made available by UKB in July 2017.

Genotype data collection and processing are described by the UKB in a previous overview

paper[43]. DNA was extracted from blood samples and genotyping was completed for 488,366

individuals on one of two Affymetrix genotyping arrays with custom content, the UK BiLEVE

Axiom array (n=49,949) or UK Biobank Axiom array (n=438,417), covering 812,428 genetic

markers common to both arrays. Of these, 488,377 individuals and 805,426 markers passed

genotype quality control checks conducted by UKB (see

http://www.biorxiv.org/content/early/2017/07/20/166298 for details). Samples were excluded

for low DNA concentration, call rate < 95%, excess heterozygosity, sex chromosome

abnormality, or sample duplication. Variants were excluded if they exhibited poor clustering of

allele calls, batch, plate, array, or sex effects, departures from HWE, or discordance between

technical replicate samples.

After quality control, samples were imputed to approximately 92 million SNPs using

both the reference panel of the Haplotype Reference Consortium (HRC)[44] as well as a combined

reference panel of the 1000 Genomes Project[42] and UK10K. As recommended by UKB, we

removed variants that were not imputed on the HRC reference panel due to technical errors in

the imputation process of the combined panel. We converted imputed variants to hard calls

(certainty > 0.9), filtered by imputation quality (INFO score >0.9), and excluded multi-allelic

SNPs, indels, SNPs without unique rsID, and SNPs with minor allele frequency (MAF) <0.0001,

resulting in 10,847,151 SNPs available for analysis.

For the present study, we selected unrelated individuals of European ancestry. To

empirically determine ancestry, we projected genetic principal components from known

ancestral populations in the 1000 Genomes Project onto the UKB genotypes and assigned

individuals to the continental ancestral superpopulation with the closest Mahalanobis

distance.[45] Within-ancestry principal components were created using FlashPCA2[46] to correct for

any residual population stratification within the European ancestry subset. Unrelated

individuals (less than 3rd degree relatives, as indicated by genomic relatedness coefficients

calculated by UKB) were selected by sequentially removing participants with the greatest

number of relatives until no related pairs remained. After applying these filtering criteria and

removing any participants with missing phenotypic or covariate data and participants who

withdrew consent, 364,859 individuals remained for analysis in the UKB sample.

1.3.2 Single-marker association analysis

Genome-wide association analysis (GWAS) for the ADSP, PGC-ALZ and UKB datasets was

performed in PLINK[39], using logistic regression for dichotomous phenotypes (cases versus

controls for ADSP and PGC-ALZ cohorts), and linear regression for phenotypes analysed as

continuous outcomes (by proxy parental AD phenotype for UKB cohort). For the ADSP and PGC-

ALZ cohorts, association tests were adjusted for gender, batch (if applicable), and the first 4

principal components. Twenty principal components were calculated, and depending on the

dataset being tested, additional principal components (on top of the standard inclusion of 4

PCAs) were added if significantly associated to the phenotype. Furthermore, for the PGC-ALZ

cohorts age was included as a covariate. For 4,537 controls of the DemGene cohort, no detailed

age information was available, besides the age range the subjects were in (20-45 years). We

therefore set the age of these individuals conservatively to 20 years. For the ADSP dataset, age

was not included as a covariate due to the enrichment for older controls (mean age cases =

73.1 years (SE=7.8); mean age controls = 86.1 years (SE=4.5)) in their collection procedures.

Correcting for age in ADSP would remove a substantial part of genuine association signals (e.g.

well-established *APOE* locus rs11556505 is strongly associated to AD ($P=1.08 \times 10^{-99}$), which is

lost when correcting for age ($P=0.0054$). For the UKB dataset, 12 components were included as

covariates, as well as age, sex, genotyping array, and assessment centre. We used the genome-

wide threshold for significance of P<5×10-8).

1.3.3 *Multivariate genome-wide meta-analysis*

Two meta-analyses were performed, including: 1) cohorts with case-control phenotypes (IGAP,

ADSP and PGC-ALZ datasets), 2) all cohorts, also including the by proxy phenotype of UKB.

The per SNP test statistics is defined by

$$Z_k = \frac{\sum_i w_i Z_i}{\sqrt{\sum_i w_i^2 + \sum_i \sum_j w_i w_j |CTI_{ij}| \, (i \neq j)}}$$

835   where w_i and Z_i are squared root of the sample size and the test statistics of SNP k of cohort i,

836   respectively. CTI is cross trait LD score intercept estimated by LDSC using genome-wide

837   summary statistics as

$$CTI_{ij} = \frac{N_{sij}\rho_{ij}}{\sqrt{N_i N_j}}$$

where N_{sij} and r_{ij} are overlapping samples and phenotypic correlation between cohort i and j,

respectively.[13] Test statistics per SNP per GWAS was converted from P-value by taking sign of

either beta or odds ratio. When direction is aligned the conversion is two-sided. To avoid

infinite value, we replaced P-value 1 with 0.999999 and P-value < 1e-323 to 1e-323 (the

minimum >0 value in Python).

The effective sample size (N_{eff}) is computed for each SNP k from the matrix M,

containing the sample size N_i of each cohort i on the diagonal and the estimated number of

shared data points $N_{sij} \times \rho_{ij} = CTI_{ij} \times \sqrt{N_i N_j}$ for each pair of cohorts i and j as the off-diagonal

values. N_{eff} is computed recursively as follows. Starting with the first cohort in M, N_{eff} is first

increased by $M_{1,1}$, corresponding to the sample size of that cohort. The proportion of samples

shared between cohort 1 and each other cohort j is them computed as $p_{1,j} = M_{1,j}/M_{j,j}$, and M is

then adjusted to remove this overlap, multiplying all values in each column j by $1-p_{1,j}$. This

amounts to reducing the sample size of each other cohort j by the number of samples it shares

with cohort 1, and reducing the shared samples between cohort j and subsequent cohorts by

the same proportion. After this, the first row and column of M are discarded, and the same

process is applied to the new M matrix. This is repeated until M is empty. The script for the

multivariate GWAS is available from https://github.com/Kyoko-wtnb/mvGWAMA.

1.5 Replication of meta-analysis result in Icelandic sample

The study group included 6,593 Alzheimer's disease cases (4,923 of whom were chip-typed) and

174,289 controls (88,581 of whom were chip-typed). In 16% of patients, the diagnosis of

Alzheimer's disease was established according to the criteria for definite, probable, or possible

Alzheimer's disease of the National Institute of Neurological and Communicative Disorders and

Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA). In 77%

of patients, the diagnosis was established according to the criteria for code 331.0 in ICD-9, or

for F00 and G30 in ICD-10. Seven percent of the patients were identified in the Directorate of

Health medication database as having been prescribed Donepezil (Aricept), a palliative

treatment for Alzheimer's disease. The controls were drawn from various research projects at

deCODE Genetics, excluding those in whom Alzheimer's disease had been diagnosed.

The study was approved by the National Bioethics Committee and the Icelandic Data Protection

Authority. Written informed consent was obtained from all participants or their guardians

before blood samples were drawn. All sample identifiers were encrypted in accordance with

the regulations of the Icelandic Data Protection Authority.

Chip-typing and long-range phasing of 155,250 individuals was carried out as described

previously.[20] Imputation of the variants found in 28,075 whole-genome sequenced individuals

into the chip-typed individuals and 285,664 close relatives was performed as detailed earlier.[20]

Association analysis was carried out using logistic regression with Alzheimer's disease status as

the response and genotype counts and a set of nuisance variables including sex, county of birth,

and current age as predictors.[21] Correction for inflation of test statistics due to relatedness and

population stratification was performed using the intercept estimate from LD score regression

(1.29) as described in.[13]

1.6 Genomic risk loci definition

We used FUMA[25], an online platform for functional mapping and annotation of genetic variants,

to define genomic risk loci and obtain functional information of relevant SNPs in these loci. We

first identified independent significant SNPs that have a genome-wide significant P-value

(<5×10-8) and are independent from each other at r^2<0.6. These SNPs were further represented

by lead SNPs, which are a subset of the independent significant SNPs that are in approximate

linkage equilibrium with each other at r^2>0.6. We then defined associated genomic risk loci by

merging any physically overlapping lead SNPs (LD blocks <250kb apart). Borders of the genomic

risk loci were defined by identifying all SNPs in LD (r^2>0.6) with one of the independent

significant SNPs in the locus, and the region containing all these candidate SNPs was considered

to be a single independent genomic risk locus. LD information was calculated using the UK

Biobank genotype data as a reference.

1.7 Cohort Heritability and Genetic Correlation

LD score regression[13] was used to estimate genomic inflation and heritability of the AD in each

of the 7 cohorts (PGC-ALZ, ADSP, IGAP, UKB, DemGene, STSA, TwinGene) using their post-

quality control summary statistics, and to estimate the cross-cohort genetic correlations.[47] Pre-

calculated LD scores from the 1000 Genomes European reference population were obtained

from   https://data.broadinstitute.org/alkesgroup/LDSCORE/.   Genetic   correlations   were

calculated on HapMap3 SNPs only. LD score regression was also used on the case-control and

by-proxy phenotype result to estimate heritability and genetic correlations for the two

phenotype definitions.

1.8 Polygenic risk scoring

We calculated polygenic scores (PGS) based on the SNP effect sizes of meta-analyses. PGS were

calculated using an independent genotype dataset of 761 individuals (379 cases and 382

controls) from the ADDNeuroMed study.[48] The same QC and imputation approach was applied

as for the other datasets with genotype-level data (see Method section 1.3.1a). PRSice PGS

were calculated on hard-called imputed genotypes using P-value thresholds from 0.0 to 0.5 in

steps ranging from 5×10^{-8} to 0.001. The explained variance (ΔR^2) was derived from a linear

model in which the AD phenotype was regressed on each PGS while controlling for the same

covariates as in each cohort-specific GWAS, compared to a linear model with GWAS covariates

only.

1.9 Stratified Heritability

To test whether specific categories of SNP annotations were enriched for heritability, we

partitioned SNP heritability for binary annotations using stratified LD score regression

(https://github.com/bulik/ldsc)[13]. Heritability enrichment was calculated as the proportion of

heritability explained by a SNP category divided by the proportion of SNPs that are in that

category. Partitioned heritability was computed by 28 functional annotation categories, by

minor allele frequency (MAF) in six percentile bins and by 22 chromosomes. Annotations for

binary categories of functional genomic characteristics (e.g. coding or regulatory regions) were

obtained from the LD score website (https://github.com/bulik/ldsc). The Bonferroni-corrected

significance threshold for 56 annotations was set at: $P<0.05/56=8.93\times10^{-4}$.

<u>1.10 Functional Annotation of SNPs</u>

Functional annotation of SNPs implicated in the meta-analysis was performed using FUMA[25]

(http://fuma.ctglab.nl/). We selected all *candidate SNPs* in associated genomic loci having an

$r^2 \geqq 0.6$ with one of the independent significant SNPs (see above), a *P*-value ($P<1\times10^{-8}$) and a

MAF>0.0001 for annotations. Functional consequences for these SNPs were obtained by

matching SNPs' chromosome, base-pair position, and reference and alternate alleles to

databases containing known functional annotations, including ANNOVAR[49] categories,

Combined Annotation Dependent Depletion (CADD) scores[23], RegulomeDB[50] (RDB) scores, and

chromatin states[51,52]. ANNOVAR annotates functional consequence of SNPs on genes (e.g.

intron, exon, intergenic). CADD scores predict how deleterious the effect of a SNP with higher

scores referring to higher deleteriousness. A CADD score above 12.37 is the threshold to be

potentially pathogenic[53]. The RegulomeDB score is a categorical score based on information

from expression quantitative trait loci (eQTLs) and chromatin marks, ranging from 1a to 7 with

lower scores indicating an increased likelihood of having a regulatory function. Scores are as

follows: 1a=eQTL + Transciption Factor (TF) binding + matched TF motif + matched DNase

Footprint + DNase peak; 1b=eQTL + TF binding + any motif + DNase Footprint + DNase peak;

1c=eQTL + TF binding + matched TF motif + DNase peak; 1d=eQTL + TF binding + any motif +

DNase peak; 1e=eQTL + TF binding + matched TF motif; 1f=eQTL + TF binding / DNase peak;

2a=TF binding + matched TF motif + matched DNase Footprint + DNase peak; 2b=TF binding +

any motif + DNase Footprint + DNase peak; 2c=TF binding + matched TF motif + DNase peak;

3a=TF binding + any motif + DNase peak; 3b=TF binding + matched TF motif; 4=TF binding +

DNase peak; 5=TF binding or DNase peak; 6=other;7=None. The chromatin state represents the

accessibility of genomic regions (every 200bp) with 15 categorical states predicted by a hidden

Markov model based on 5 chromatin marks for 127 epigenomes in the Roadmap Epigenomics

Project[39]. A lower state indicates higher accessibility, with states 1-7 referring to open

chromatin states. We annotated the minimum chromatin state across tissues to SNPs. The 15-

core chromatin states as suggested by Roadmap are as follows: 1=Active Transcription Start Site

(TSS); 2=Flanking Active TSS; 3=Transcription at gene 5' and 3'; 4=Strong transcription; 5= Weak

Transcription;   6=Genic   enhancers;   7=Enhancers;   8=Zinc   finger   genes   &   repeats;

9=Heterochromatic; 10=Bivalent/Poised TSS; 11=Flanking Bivalent/Poised TSS/Enh; 12=Bivalent

Enhancer; 13=Repressed PolyComb; 14=Weak Repressed PolyComb; 15=Quiescent/Low.

1.11 Gene-mapping

Genome-wide significant loci obtained by GWAS were mapped to genes in FUMA[25] using three

strategies:

1.   Positional mapping maps SNPs to genes based on physical distance (within a 10kb

window) from known protein coding genes in the human reference assembly

(GRCh37/hg19).

2.   eQTL mapping maps SNPs to genes with which they show a significant eQTL association

(i.e. allelic variation at the SNP is associated with the expression level of that gene).

eQTL mapping uses information from 45 tissue types in 3 data repositories (GTEx[54],

Blood eQTL browser[55], BIOS QTL browser[56]), and is based on cis-eQTLs which can map

SNPs to genes up to 1Mb apart. We used a false discovery rate (FDR) of 0.05 to define

significant eQTL associations.

3.   Chromatin interaction mapping was performed to map SNPs to genes when there is a

three-dimensional DNA-DNA interaction between the SNP region and another gene

region. Chromatin interaction mapping can involve long-range interactions as it does not

have a distance boundary. FUMA currently contains Hi-C data of 14 tissue types from

the study of Schmitt et al[57]. Since chromatin interactions are often defined in a certain

resolution, such as 40kb, an interacting region can span multiple genes. If a SNPs is

located in a region that interacts with a region containing multiple genes, it will be

mapped to each of those genes. To further prioritize candidate genes, we selected only

genes mapped by chromatin interaction in which one region involved in the interaction

overlaps with a predicted enhancer region in any of the 111 tissue/cell types from the

Roadmap Epigenomics Project[52] and the other region is located in a gene promoter

region (250bp up and 500bp downstream of the transcription start site and also

predicted by Roadmap to be a promoter region). This method reduces the number of

genes mapped but increases the likelihood that those identified will have a plausible

biological function. We used a FDR of 1×10^{-5} to define significant interactions, based on

previous recommendations[44] modified to account for the differences in cell lines used

here.

1.12 Gene-based analysis

To account for the distinct types of genetic data in this study, genotype array (PGC-ALZ, IGAP,

UKB) and whole-exome sequencing data (ADSP), we first performed two gene-based genome-

wide association analysis (GWGAS) using MAGMA[28], followed by a meta-analysis. SNP-based P-

values from the meta-analysis of the 3 genotype-array-based datasets were used as input for

the first GWGAS, while the unimputed individual-level sequence data of ADSP was used as

input for the second GWGAS. 18,233 protein-coding genes (each containing at least one SNP in

the GWAS) from the NCBI 37.3 gene definitions were used as basis for GWGAS in MAGMA.

Bonferroni correction was applied to correct for multiple testing ($P<2.74\times10^{-6}$).

1.13 Gene-set analysis

Results from the GWGAS analyses were used to test for association in three types of 7,087

predefined gene-sets:

1. 6,994 curated gene-sets representing known biological and metabolic pathways derived

from Gene Ontology (5917 gene-sets), Biocarta (217 gene-sets), KEGG (186 gene-sets),

Reactome (674 gene-sets) catalogued by and obtained from the MsigDB version 6.1[58]

(http://software.broadinstitute.org/gsea/msigdb/collections.jsp)

2. Gene expression values from 54 (53 + 1 calculated 1^{st} PC of three tissue subtypes)

tissues obtained from GTEx[54], log2 transformed with pseudocount 1 after winsorization

at 50 and averaged per tissue.

3. Cell-type specific expression in 173 types of brain cells (24 broad categories of cell types,

'level 1' and 129 specific categories of cell types 'level 2'), which were calculated

following the method described in [30]. Briefly, brain cell-type expression data was drawn

from single-cell RNA sequencing data from mouse brains. For each gene, the value for

each cell-type was calculated by dividing the mean Unique Molecular Identifier (UMI)

counts for the given cell type by the summed mean UMI counts across all cell types.

Single-cell gene-sets were derived by grouping genes into 40 equal bins based on

specificity of expression.

4. Nucleus specific gene expression of 15 distinct human brain cell-types of study

described in[59]. The value for each cell-type was calculated with the same method as

explained in point 3 above.

These gene-sets were tested using MAGMA. We computed competitive P-values, which

represent the test of association for a specific gene-set compared with genes not in the gene-

set to correct for baseline level of genetic association in the data. The Bonferroni-corrected

significance threshold was 0.05/7,087 gene-sets=$7.06×10^{-6}$. The suggestive significance

threshold was defined by the number of tests within the category. Conditional analyses were

performed as a follow-up using MAGMA to test whether each significant association observed

was independent of all others and of *APOE* (a gene-set including all genes within genomic

region chr19:45,020,859-45,844,508). Furthermore, the association between each of the

significant gene-set was tested conditional on each of the other significantly associated gene-

sets. Gene-sets that retained their association after correcting for other sets were considered to

represent independent signals. We note that this is not a test of association per se, but rather a

strategy to identify, among gene-sets with known significant associations and overlap in genes,

which set (s) are responsible for driving the observed association.

1033

1.14 Cross-Trait Genetic Correlation

Genetic correlations (r_g) between AD and 41 phenotypes were computed using LD score

regression[13], as described above, based on GWAS summary statistics obtained from publicly

available databases (http://www.med.unc.edu/pgc/results-and-downloads; http://

ldsc.broadinstitute.org/; **Supplementary Table 19**). The Bonferroni-corrected significance

threshold was 0.05/41 traits=1.22×10^{-3}.

1040

1.15 Mendelian Randomisation

To infer credible causal associations between AD and traits that are genetically correlated with

AD, we performed Generalised Summary-data based Mendelian Randomisation[29] (GSMR;

http://cnsgenomics.com/software/gsmr/). This method utilizes summary-level data to test for

putative causal associations between a risk factor (exposure) and an outcome by using

independent genome-wide significant SNPs as instrumental variables as an index of the

exposure. HEIDI-outlier detection was used to filter genetic instruments that showed clear

pleiotropic effects on the exposure phenotype and the outcome phenotype. We used a

threshold p-value of 0.01 for the outlier detection analysis in HEIDI which removes 1% of SNPs

by chance if there is no pleiotropic effect. To test for a potential causal effect of various

outcomes on risk for AD, we selected phenotypes in non-overlapping samples that showed

(suggestive) significant ($P<0.05$) genetic correlations (r_g) with AD. With this method it is typical

to test for bi-directional causation by repeating the analyses while switching the role of the

exposure and the outcome; however, because AD is a late-onset disease, it makes little sense to

estimate its causal effect on outcomes that develop earlier in life, particularly when the

summary statistics for these outcomes were derived mostly from younger samples than those

of AD cases. Therefore, we conducted these analyses only in one direction. For genetically

correlated phenotypes, we selected independent (r^2=<0.1), GWS lead SNPs as instrumental

variables in the analyses. The method estimates a putative causal effect of the exposure on the

outcome (b_{xy}) as a function of the relationship between the SNPs' effects on the exposure (b_{zx})

and the SNPs' effects on the outcome (b_{zy}), given the assumption that the effect of non-

pleiotropic SNPs on an exposure (x) should be related to their effect on the outcome (y) in an

independent sample only via mediation through the phenotypic causal pathway (b_{xy}). The

estimated causal effect coefficients (b_{xy}) are approximately equal to the natural log odds ratio

(OR)[29] for a case-control trait. An OR of 2 can be interpreted as a doubled risk compared to the

population prevalence of a binary trait for every SD increase in the exposure trait. For

quantitative traits the b_{zx} and b_{zy} can be interpreted as a one standard deviation increase

explained in the outcome trait for every SD increase in the exposure trait. This method can help

differentiate the causal direction of association between two traits, but cannot make any

statement about the intermediate mechanisms involved in any potential causal process.

1071

1072   *Data availability*

1073   Summary statistics will be made available for download upon publication (https://ctg.cncr.nl).

1074
1075   **References**
1076

1.   Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W, Ferri CP. The global prevalence of
dementia: a systematic review and metaanalysis. *Alzheimer's & dementia : the journal of the*
Alzheimer's Association 2013; **9**(1): 63-75.e2.
2.   Gatz M, Reynolds CA, Fratiglioni L, et al. Role of genes and environments for explaining
Alzheimer disease. *Archives of general psychiatry* 2006; **63**(2): 168-74.
3.   Cacace R, Sleegers K, Van Broeckhoven C. Molecular genetics of early-onset Alzheimer's
disease revisited. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 2016;
12(6): 733-48.
4.   Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals
identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics* 2013; **45**(12): 1452-
8.
5.   Goate A, Chartier-Harlin MC, Mullan M, et al. Segregation of a missense mutation in the
amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 1991; **349**(6311): 704-
6.
6.   Sherrington R, Rogaev EI, Liang Y, et al. Cloning of a gene bearing missense mutations in
early-onset familial Alzheimer's disease. *Nature* 1995; **375**(6534): 754-60.
7.   Sherrington R, Froelich S, Sorbi S, et al. Alzheimer's disease associated with mutations in
presenilin 2 is rare and variably penetrant. *Human molecular genetics* 1996; **5**(7): 985-8.
8.   Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for Alzheimer's
disease: an appraisal for the development of therapeutics. *Nature reviews Drug discovery* 2011;
10(9): 698-712.
9.   Jonsson T, Stefansson H, Steinberg S, et al. Variant of TREM2 associated with the risk of
Alzheimer's disease. *The New England journal of medicine* 2013; **368**(2): 107-16.
10.   Steinberg S, Stefansson H, Jonsson T, et al. Loss-of-function variants in ABCA7 confer risk
of Alzheimer's disease. *Nature genetics* 2015; **47**(5): 445-7.
11.   Liu CC, Liu CC, Kanekiyo T, Xu H, Bu G. Apolipoprotein E and Alzheimer disease: risk,
mechanisms and therapy. *Nature reviews Neurology* 2013; **9**(2): 106-18.
12.   Zheng J, Erzurumluoglu AM, Elsworth BL, et al. LD Hub: a centralized database and web
interface to perform LD score regression that maximizes the potential of summary level GWAS

data for SNP heritability and genetic correlation analysis. *Bioinformatics (Oxford, England)* 2017; **33**(2): 272-9.

13.   Bulik-Sullivan BK, Loh PR, Finucane HK, et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature genetics* 2015; **47**(3): 291-5.

14.   Ridge PG, Mukherjee S, Crane PK, Kauwe JS. Alzheimer's disease: analyzing the missing heritability. *PloS one* 2013; **8**(11): e79771.

15.   Ridge PG, Hoyt KB, Boehme K, et al. Assessment of the genetic variance of late-onset Alzheimer's disease. *Neurobiology of aging* 2016; **41**: 200.e13-.e20.

16.   Desikan RS, Schork AJ, Wang Y, et al. Polygenic Overlap Between C-Reactive Protein, Plasma Lipids, and Alzheimer Disease. *Circulation* 2015; **131**(23): 2061-9.

17.   Jun GR, Chung J, Mez J, et al. Transethnic genome-wide scan identifies novel Alzheimer's disease loci. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 2017; **13**(7): 727-38.

18.   Guerreiro R, Wojtas A, Bras J, et al. TREM2 variants in Alzheimer's disease. *The New England journal of medicine* 2013; **368**(2): 117-27.

19.   Sims R, van der Lee SJ, Naj AC, et al. Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. *Nature genetics* 2017; **49**(9): 1373-84.

20.   Gudbjartsson DF, Helgason H, Gudjonsson SA, et al. Large-scale whole-genome sequencing of the Icelandic population. *Nature genetics* 2015; **47**(5): 435-44.

21.   Steinthorsdottir V, Thorleifsson G, Sulem P, et al. Identification of low-frequency and rare sequence variants associated with elevated or reduced risk of type 2 diabetes. *Nature genetics* 2014; **46**(3): 294-8.

22.   Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics (Oxford, England)* 2015; **31**(9): 1466-8.

23.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nature genetics* 2014; **46**(3): 310-5.

24.   Finucane HK, Bulik-Sullivan B, Gusev A, et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nature genetics* 2015; **47**(11): 1228-35.

25.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nature communications* 2017; **8**(1): 1826.

26.   Gurses MS, Ural MN, Gulec MA, Akyol O, Akyol S. Pathophysiological Function of ADAMTS Enzymes on Molecular Mechanism of Alzheimer's Disease. *Aging and disease* 2016; **7**(4): 479-90.

27.   Dries DR, Yu G. Assembly, maturation, and trafficking of the gamma-secretase complex in Alzheimer's disease. *Current Alzheimer research* 2008; **5**(2): 132-46.

28.   de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS computational biology* 2015; **11**(4): e1004219.

29.   Zhu Z, Zheng Z, Zhang F, et al. Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nature communications* 2018; **9**(1): 224.

30. Skene NG, Grant SG. Identification of Vulnerable Cell Types in Major Brain Disorders
Using Single Cell Transcriptomes and Expression Weighted Cell Type Enrichment. *Frontiers in*
neuroscience 2016; **10**: 16.
31. Loewendorf A, Fonteh A, Mg H, Me C. Inflammation in Alzheimer's Disease: Cross-talk
between Lipids and Innate Immune Cells of the Brain; 2015.
32. Stern Y. Cognitive reserve in ageing and Alzheimer's disease. *The Lancet Neurology*
2012; **11**(11): 1006-12.
33. Satizabal C, Beiser AS, Seshadri S. Incidence of Dementia over Three Decades in the
Framingham Heart Study. *The New England journal of medicine* 2016; **375**(1): 93-4.
34. Magnusson PK, Almqvist C, Rahman I, et al. The Swedish Twin Registry: establishment of
a biobank and other recent developments. *Twin research and human genetics : the official*
journal of the International Society for Twin Studies 2013; **16**(1): 317-29.
35. Finkel D, Pedersen NL. Processing Speed and Longitudinal Trajectories of Change for
Cognitive Abilities: The Swedish Adoption/Twin Study of Aging. *Aging, Neuropsychology, and*
Cognition 2004; **11**(2-3): 325-45.
36. Gold CH, Malmberg B, McClearn GE, Pedersen NL, Berg S. Gender and health: a study of
older unlike-sex twins. *The journals of gerontology Series B, Psychological sciences and social*
sciences 2002; **57**(3): S168-76.
37. Gatz M, Fratiglioni L, Johansson B, et al. Complete ascertainment of dementia in the
Swedish Twin Registry: the HARMONY study. *Neurobiology of aging* 2005; **26**(4): 439-47.
38. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis
of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of
Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;
34(7): 939-44.
39. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK:
rising to the challenge of larger and richer datasets. *GigaScience* 2015; **4**: 7.
40. Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying
the causes of a wide range of complex diseases of middle and old age. *PLoS medicine* 2015;
12(3): e1001779.
41. Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010-
2050) estimated using the 2010 census. *Neurology* 2013; **80**(19): 1778-83.
42. The Genomes Project C. A global reference for human genetic variation. *Nature* 2015;
526: 68.
43. Davies G, Marioni RE, Liewald DC, et al. Genome-wide association study of cognitive
functions and educational attainment in UK Biobank (N=112 151). *Molecular psychiatry* 2016;
21(6): 758-67.
44. McCarthy S, Das S, Kretzschmar W, et al. A reference panel of 64,976 haplotypes for
genotype imputation. *Nature genetics* 2016; **48**(10): 1279-83.
45. Peterson RE, Edwards AC, Bacanu SA, Dick DM, Kendler KS, Webb BT. The utility of
empirically assigning ancestry groups in cross-population genetic studies of addiction. *The*
American journal on addictions 2017; **26**(5): 494-501.
46. Abraham G, Qiu Y, Inouye M. FlashPCA2: principal component analysis of Biobank-scale
genotype datasets. *Bioinformatics (Oxford, England)* 2017; **33**(17): 2776-8.

47.   Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across
human diseases and traits. *Nature genetics* 2015; **47**(11): 1236-41.
48.   Lovestone S, Francis P, Kloszewska I, et al. AddNeuroMed--the European collaboration
for the discovery of novel biomarkers for Alzheimer's disease. *Annals of the New York Academy*
of Sciences 2009; **1180**: 36-46.
49.   Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from
high-throughput sequencing data. *Nucleic acids research* 2010; **38**(16): e164.
50.   Boyle AP, Hong EL, Hariharan M, et al. Annotation of functional variation in personal
genomes using RegulomeDB. *Genome research* 2012; **22**(9): 1790-7.
51.   Ernst J, Kellis M. ChromHMM: automating chromatin-state discovery and
characterization. *Nature methods* 2012; **9**(3): 215-6.
52.   Kundaje A, Meuleman W, Ernst J, et al. Integrative analysis of 111 reference human
epigenomes. *Nature* 2015; **518**(7539): 317-30.
53.   Amendola LM, Dorschner MO, Robertson PD, et al. Actionable exomic incidental findings
in 6503 participants: challenges of variant classification. *Genome research* 2015; **25**(3): 305-15.
54.   Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue
gene regulation in humans. *Science (New York, NY)* 2015; **348**(6235): 648-60.
55.   Westra HJ, Peters MJ, Esko T, et al. Systematic identification of trans eQTLs as putative
drivers of known disease associations. *Nature genetics* 2013; **45**(10): 1238-43.
56.   Zhernakova DV, Deelen P, Vermaat M, et al. Identification of context-dependent
expression quantitative trait loci in whole blood. *Nature genetics* 2017; **49**(1): 139-45.
57.   Schmitt AD, Hu M, Jung I, et al. A Compendium of Chromatin Contact Maps Reveals
Spatially Active Regions in the Human Genome. *Cell reports* 2016; **17**(8): 2042-59.
58.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of*
the National Academy of Sciences of the United States of America 2005; **102**(43): 15545-50.
59.   Habib N, Avraham-Davidi I, Basu A, et al. Massively parallel single-nucleus RNA-seq with
DroNc-seq. *Nature methods* 2017; **14**(10): 955-8.
1220

Supplementary Figure 1. Manhattan and QQ plots of single variant association results per main cohort. For each cohort, Manhattan and QQ plots are shown. A) The Manhattan plot displays all associations per variant ordered according to their genomic position on the x-axis and showing the strength of the association with the −log10 transformed *P*-values on the y-axis. The y-axis is limited to enable visualization of non-*APOE* loci. B) The QQ plot displays the expected −log10 transformed *p*-values on the x-axis and the observed −log10 transformed *p*-values on the y-axis.

B

Supplementary Figure 2. Regional plot for the 29 significant loci of the meta-analysis. Every point represents a SNP, which are colour-coded based on the highest r2 to one of the most significant SNPs, if greater or equal to r2 of 0.6. Other SNPs are coloured in grey.

Supplementary Figure 3. Partitioned heritability results for the meta-analysis. Variants were binned by chromosome or minor allele frequency and tested for a significant over- or underrepresentation as to what is expected by chance. A) Enrichment results for heritability calculations where variants have been partitioned per chromosome. B) Enrichment results for heritability calculations where variants have been partitioned into multiple categories based on minor allele frequency.

A

B

Supplementary Figure 4. Full circos plots of chromatin interactions and eQTLs for all chromosomes with significantly associated loci. The distinct layers and colors correspond to various features. The outer layer contains zoomed in Manhattan plots containing only SNPs with P < 0.05. SNPs in genomic risk loci are color-coded as a function of their maximum r2 to the one of the independent significant SNPs in the locus, as follows: red (r2 > 0.8), orange (r2 > 0.6), green (r2 > 0.4) and blue (r2 > 0.2). SNPs that are not in LD with any of the independent significant SNPs (with r2 ≤ 0.2) are grey. The second layer displays the position of the genomic risk loci in blue. The third layer contains the mapped genes that are implicated by chromatin interactions and/or eQTL analysis (orange = chromatin interaction; green = eQTL; red = chromatin interaction and eQTL).

Supplementary Figure 5. Gene-based association results with MAGMA. A) The Manhattan plot displays all associations per gene ordered according to their genomic position (start of gene) on the x-axis and showing the strength of the association with the –log10 transformed *P*-values on the y-axis. B) The QQ plot displays the expected –log10 transformed *p*-values on the x-axis and the observed –log10 transformed *p*-values on the y-axis.

Supplementary Figure 6. Single-cell expression gene-set results of human brain tissue. The black vertical line indicates the significance threshold correcting for number of tests within category. MG = microglia; ExCA1 = Hippocampal CA 1 pyramidal neurons; END = Endothelial cells; ExPFC2 = Prefrontal glutamergic neurons 2; ExPFC1 = Prefrontal glutamergic neurons 1; ODC1 = Oligodendrocytes; ASC2 = Astrocytes 2; OPC = Oligodendrocyte precursor cells 1; GABA1 = GABAergic interneurons 1; NSC = Neuronal stem cells; GABA2 = GABAergic interneurons 2; OPC2 = Oligodendrocyte precursor cells 2; ExCA3 = Hippocampal CA 3 pyramidal neurons; ASC1 = Astrocytes 1; ExDG = Dentate gyrus granule neurons.

Supplementary Figure 7. Mendelian Randomization tests for the effect of correlated phenotypes on risk for Alzheimer's disease. For independent significant SNPs from each correlated phenotype, effect sizes of the SNPs for Alzheimer's disease (b_{zy}) are shown on the x-axis and effect sizes for correlated phenotypes are on the y-axis (b_{zx}). The dotted line represents a line with slope of (b_{xy}) and an intercept of 0.Red dots represent outliers that were excluded for the Mendelian Randomization analysis.