An important class of intron retention events in human erythroblasts is regulated by cryptic exons proposed to function as splicing decoys

Marilyn Parra[1], Ben W. Booth[1], Richard Weiszmann[1], Brian Yee[3], Gene W. Yeo[3], Susan E. Celniker[1], James B. Brown[2], John G. Conboy[1]

[1]Biological Systems and Engineering Division and [2]Environmental Genomics and Systems Biology, Lawrence Berkeley National Laboratory, Berkeley, CA; [3]Cellular and Molecular Medicine, University of California at San Diego, La Jolla, CA

Abstract

During terminal erythropoiesis, the splicing machinery in differentiating erythroblasts executes a robust intron retention (IR) program that impacts expression of hundreds of genes. We studied IR mechanisms in the SF3B1 splicing factor gene, which expresses ~50% of its transcripts in late erythroblasts as a nuclear isoform that retains intron 4. RNA-seq analysis of nonsense-mediated decay (NMD)-inhibited cells revealed splice junctions connecting constitutive exons 4 and 5 to cryptic cassette exons encoded in highly conserved intron sequences. In minigene splicing reporter assays, these cassette exons functioned to promote IR. Genome-wide analysis of splice junction reads demonstrated that cryptic noncoding cassettes are much more common in large (>1kb) retained introns than they are in small retained introns or in non-retained introns. Functional assays showed that heterologous cassettes could promote retention of intron 4 in the SF3B1 splicing reporter. Although most of these cryptic exons were spliced inefficiently, they exhibited substantial binding of U2AF1 and U2AF2 adjacent to their splice acceptor sites. We propose that these exons function as decoys that engage the intron-terminal splice sites, blocking cross-intron interactions required for excision. Developmental regulation of decoy function underlies a major component of the erythroblast IR program.

Introduction.

Intron retention (IR) is a common variant of alternative splicing in which selected intron(s) are specifically retained in an otherwise spliced and polyadenylated transcript. IR can be developmentally or physiologically regulated as an important component of gene regulation in many cell types (Jacob and Smith 2017). In some contexts, pre-existing IR transcripts are relatively stable and can be stored as a source of new mRNA, to be spliced in response to appropriate signals (Boothby et al. 2013; Mauger et al. 2016). Partial diversion of transcriptional output into IR isoforms also functions as a post-transcriptional pathway to modulate expression levels (Wong et al. 2013; Braunschweig et al. 2014; Shalgi et al. 2014; Boutz et al. 2015; Ni et al. 2016). Some IR transcripts can be recruited to ribosomes to function in translation (Li et al. 2016), while other might serve as miRNA sponges in the nucleus (Schmitz et al. 2017).

The diversity of IR programs and functions suggests that a number of regulatory pathways exist, allowing cells to integrate multiple inputs in order to independently regulate different subsets of IR transcripts. Consistent with this notion, coherent subsets of genes can be regulated by IR, especially those encoding RNA binding proteins and spliceosomal factors (Shalgi et al. 2013; Wong et al. 2013; Braunschweig et al. 2014; Boutz et al. 2015; Pimentel et al. 2016). How these sub-programs are regulated is not well understood. Recent studies have implicated transcription rate/pausing (Braunschweig et al. 2014), specific splicing factors (SRSF4), and DNA and protein methylation factors (by MECP2, in granulocytes, and by PRMT5, in glioblastoma) as important effectors of IR (Boutz et al. 2015; Braun et al. 2017; Wong et al. 2017). In the brain, signaling-dependent splicing was observed to require NMDA-type glutamate receptor or calmodulin-dependent protein kinase pathways for removal of retained introns upon neuronal activation (Mauger et al. 2016). There are also a few recent examples showing that individual IR events can be regulated by feedback mechanisms to control physiological pathways (Park et al. 2017; Pendleton et al. 2017; Pirnie et al. 2017).

Terminal erythropoiesis is an excellent model system for studies of IR. Primary erythroblasts differentiating in culture carry out a robust IR program (Edwards et al. 2016; Pimentel et al. 2016) impacting the expression of many important erythroid proteins. In mature erythroblasts, IR transcripts comprise 25-50% of steady state RNA for genes encoding essential splicing factors (SF3B1 and others), mitochondrial iron importers required for heme biosynthesis (mitoferrins, encoded by SLC25A37 and SLC25A28), and major cytoskeletal proteins (alpha spectrin, encoded by SPTA1). Moreover, the cellular complement of IR events is continuously remodeled in a differentiation stage-dependent manner through the combined effects of differentiation-independent and –dependent IR networks (Pimentel et al. 2016).

In the current study, we focused on regulatory mechanisms for a subset of genes represented by SF3B1. SF3B1 is an essential splicing factor that functions in normal 3' splice site regulation. Mutations in the SF3B1 gene are found in many MDS (myelodysplastic syndrome) patients, where they induce RNA processing errors due to altered 3' splice site selection (Obeng et al. 2016) and changes in exon skipping (Jin et al. 2017). A combination of comparative genomics, RNA-seq and RT-PCR analysis, and minigene splicing reporters was used to discover highly conserved decoy exons that strongly influenced the level of intron 4 retention. Bioinformatics analysis of RNA-seq data showed that decoy exons are a common feature of large (>1kb) introns. We propose that

decoy exons interfere with function of intron-terminal splice sites required for intron excision, and that this mechanism regulates regulate a critical subset of IR events in differentiating erythroblasts.

Results.

<u>SF3B1 retained intron 4 (i4) harbors cryptic exon(s) that are highly conserved.</u>
Comparative genomics showed that SF3B1 i4 is extremely conserved among vertebrate
genomes (Figure 1A). Three 125-200nt regions are 93-98% identical from chicken to
human, and core areas of two are 79-94% identical from zebrafish to man (Figure S1).
Sequence inspection revealed several pairs of consensus 3' and 5' splice sites in these ultra

conserved regions, as well as
in three additional conserved
regions, predicting six short
exons of 29-56nt (Figure 1A,
E4a-E4f). The extraordinary
conservation of these exons
to fish (E4d and E4e),
reptiles (E4b, E4c, and E4f),
or mammals (E4a) is shown
in Figure S1, and suggests
that these cryptic exons
might have important
function(s) in SF3B1
regulation. Consistent with
the idea that these exons
might be recognized by the
splicing machinery, the 3'
splice site factors U2AF1 and
U2AF2 cross-linked to five of
the six exons in K562 eCLIP
experiments (Figure 1a,
lower panel) and to all six
exons in HEPG2 cells (not
shown). Since all except E4c
would introduce premature
termination codons (PTCs),
we hypothesized a non-
coding function for these
exons.

Figure 1. Structure and expression of SF3B1 intron 4. A. Top
portion, genome browser tracks showing exons, conservation, and
RNA-seq coverage of the SF3B1 gene. Bottom portion, close-up of i4
region showing predicted cryptic exons E4a-E4f and binding data for
U2AF subunits. B. RNA-seq coverage in control cells vs NMD-
inhibited cells. C. RT-PCR analysis of cryptic exon expression. D. Plots
showing RNA-seq junctions in NMD-inhibited D16 erythroblasts.

In order to study expression
of cryptic exons in SF3B1
intron 4, we performed
transcriptome analysis of
early erythroblast
progenitors (culture day 9,
D9) and mature
erythroblasts (culture day 16, D16) that had been treated with inhibitors of nonsense-
mediated decay (NMD). This strategy, by increasing the relative abundance of transcripts
containing premature termination codons (PTC), validated expression of two cryptic exons
(E4b and E4e) in the RNA-seq profiles and by RT-PCR (Figures 1B-C). Furthermore,
analysis of individual RNA-seq reads revealed new splice junctions that validate expression

of all six predicted cassettes, and indicate extensive connections between and among these cryptic exons and the flanking constitutive exons 4 and 5 (Figure 1D). Two important features were noted in this analysis. First, each cryptic exon was represented by at least one unique RNA-seq read that connected it to both upstream and downstream exons, confirming its ability to be recognized and spliced as a discrete exon. Second, splice junctions that link constitutive exons 4 and 5 with intron-internal sites were surprisingly abundant. In the D16 sample, E4-E4b and E4-E4e junctions together represented ~33% of splice junction reads (261/778) that connect E4 to downstream sequences, while ~30% of E5 upstream splice junction reads (223/739) involved E4b or E4e. Splice junctions that connect E4b to E4e were also common. At lower frequency, splice junctions connected the other cryptic exons to each other and to the flanking constitutive exons (Figure 1D).

Together these experiments confirmed that the intron-terminal splice sites of i4 interact with internal splice sites associated with cryptic exons. We hypothesized that the cryptic exons might function as decoys whose splice sites could compete with the cross-intron interactions required for intron excision, and thereby might promote intron retention. We reasoned that many of these interactions might be "leaky", leaving behind splice junctions that serve as indirect evidence for these interactions. This model suggested several testable predictions: that deleting decoy exons should reduce intron retention; that known 3' splice site factors should cross-link to the cryptic junctions; and that analogous decoy exons / splice junctions should be common in retained introns.

Figure 2. Intron retention assays in minigene splicing reporters. A. Structure of minigene splicing reporters spanning the E3-E6 region of SF3B1. Shown are the wild type reporter (WT) and a series of variants with deletions of candidate decoy exons. B. Splicing assay performed with the wild type reporter in K562 and HEK293 cells. C. Splicing assays show that single-decoy deletions have mild effects on IR. Numbers below the lanes indicate apparent IR percent as determined by densitometry, corrected for differences in band size. D. Splicing assays with multi-decoy deletions show that even a single decoy (E4e) can promote IR.

Decoy exons promote SF3B4 i4 retention. We investigated the sequence requirements for intron retention using a series of minigene splicing reporters. The wild type (WT) construct contains a 4.7kb fragment of the SF3B1 gene that spans exons 3-6 and includes full length intron sequences in this region (Figure 2A). This reporter was spliced to produce two major products when expressed in K562 cells: a fully spliced RNA containing exons 3-6,

and a larger transcript that has excised introns 3 and 5 but specifically retained i4 (Figure 2B, left lane). In contrast, little intron retention was observed in HEK (human embryonic kidney) cells (Figure 2B, right lane), demonstrating cell type-specific regulation of i4 retention.

Deletion of individual decoy exons resulted in slightly reduced IR compared with the WT construct, with exon 4e having the greatest effect (Figure 2C). Deletion of all six decoys resulted in a smaller intron with little or no IR (Figure 2D, compare first two lanes), while adding back a single copy of E4e restored substantial retention (3rd lane). As a control, we showed that loss of IR in the decoy-deficient reporter was not due to the smaller size of the intron, since E4e-containing constructs with similar sized introns retained significant IR (last two lanes).

Figure 3. **Testing IR activity of a decoy exon at heterologous sites**. A. Structure of minigene splicing reporters testing function of E4e in a different region of i4 (mut21) or in i5 (mut24.2, mut31). B. RT-PCR analysis of intron retention showing that decoy exons can promote retention at heterologous sites in the same intron or in another intron. Numbers below the lanes indicate apparent IR percent as determined by densitometry, corrected for differences in band size.

Decoy exon E4e can promote IR at heterologous sites. We next investigated whether E4e could promote IR in different intronic contexts. First, the ability of E4e to function at a heterologous position in i4 was tested. Indeed, the reduction in IR observed when E4e was deleted from the wild type construct (Figure 3B, compare WT with Δ4e) was rescued when E4e was substituted at the E4d site (mut21).

We then asked whether IR-promoting elements in i4 could induce IR in a non-retained intron. Intron 5 (i5) in the SF3B1 gene ordinarily exhibits negligible IR in either the endogenous gene (not shown), or in splicing reporters (Figure 3B, right gel, lanes WT). In construct mut31, most of i5 (~1.1kb) was replaced with a similar length of i4 including E4a through E4e, but ~160nt of the natural i5 sequence was maintained at either end. This arrangement induced substantial retention of i5. When E4e alone was inserted at a random site within i5 (mut24.2), modest IR activity was observed, but it was accompanied by substantial inclusion of E4e. Thus, decoy exons can promote IR in a heterologous intron, independent of specific sequences at the intron-terminal splice sites or flanking constitutive exons. The neighboring sequence environment of a decoy is critical, however, since a "permissive" sequence context in intron 5 apparently enhanced exon inclusion rather than the IR phenotype observed in intron 4. In fact, for this experiment it was necessary to mutate a cryptic 5' splice site in intron 5 (marked by the "X"), because otherwise the major product was strong inclusion of an aberrant E4e-related exon.

Candidate decoy exons occur in many retained introns. Studies of IR-promoting elements in introns of SF3B1, OGT (Park et al. 2017), and ARGLU1 (Pirnie et al. 2017) suggested that

decoy-mediated IR might be a common regulator of IR. In the OGT gene, an intron splicing silencer (ISS) inhibits intron 4 removal, thus favoring intron retention and contributing to O-GlcNAc homeostasis (Park et al. 2017). OGT i4 was strongly retained in erythroblasts, and splice junction data revealed a novel ~146nt cassette exon that overlapped the ISS (Figures S2 and S3). In ARGLU1, a 58nt exon previously shown to promote IR in HeLa cells (Pirnie et al. 2017) was also expressed in erythroblasts (Figures S2 and S3). Further inspection of splice junction reads revealed candidate decoy cassettes in retained introns predicted earlier in DDX39B, SPTA1, KEL, SNRNP70, and FUS transcripts ((Edwards et al. 2016; Pimentel et al. 2016); Figure S3). In each case, the putative decoy is flanked on both sides by retained intron(s), analogous to the configuration we reported earlier in which PTC-containing alternative exons are often situated between two consecutive retained introns (Pimentel et al. 2016). These observations suggested that decoy cassettes might be a common phenomenon among retained introns.

To address the genome-wide potential for decoy exon-mediated IR, we correlated IR and splice junction data for all introns in erythroid-expressed genes. Among 20,534 retained introns identified in the early erythroblast D9 sample, 770 predicted cassettes were identified (Supplemental Table 1). Late erythroblast D16 cells, that have down-regulated expression of many genes, exhibited 411 cassettes in 9677 retained introns. Interestingly, the incidence of candidate decoys was strongly dependent on intron length; both D9 and D16 transcriptomes exhibited a much higher frequency of cassettes in longer retained introns (≥ 1kb; 12-17%) than in shorter retained introns (~2%). Consistent with this observation, the median length of retained introns with cassettes was much greater (1176-1280nt) than that of retained introns lacking cassettes (258-311nt). These results provided the first clue that distinct IR mechanisms might preferentially regulate introns of different lengths.

Table 1.

sample	R-introns>1kb		% Cassette	R-introns<1kb		% Cassette
	total	+cass.		total	+cass.	
D9	4038	482	11.9	16,496	288	1.7
D16	1436	238	16.6	8241	173	2.1
sample	N-introns>1kb		% Cassette	N-introns<1kb		% Cassette
	total	+cass.		total	+cass.	
D9	82,853	3600	4.3	36,664	179	0.5
D16	78,454	3443	4.4	41,360	244	0.6

However, the analysis above does not take into account that some of cassettes are likely to represent "conventional" NMD-inducing exons that play no role in IR. We reasoned that the frequency of IR-promoting cassettes could be estimated by comparing cassette frequency in retained vs non-retained introns, since the latter group by definition lacks IR-promoting cassettes. Table I shows that the frequency of cassettes was 3-4-fold higher in the large retained introns (R-introns) than in the comparable non-retained introns (N-introns), for both D9 and D16 transcriptomes. Based on the larger overall numbers of N-introns, and the

greater frequency of cassettes in R-introns, we conclude (1) that most noncoding cassette exons in erythroblasts are located in non-retained introns and are not relevant to IR, and (2) that a majority of cassettes located in R-introns likely do act to promote IR.

These observations support the concept of two mechanistically distinct classes of retained introns: longer introns in which decoy exons often promote IR, and shorter introns that are retained via decoy-independent mechanism(s). Overall, the data indicates that several hundred introns may be regulated by the decoy mechanism in human erythroblasts. The true number could be higher, if some functional decoys were not detected because they are poor spliced and did not generate splice junction reads.

Heterologous decoy exons can promote intron 4 retention. We proposed that heterologous IR-associated cassette exons could substitute for SF3B1 E4e to promote retention of i4. To explore this hypothesis, we assayed their function in minigene splicing reporters (Figure 4A). Each candidate decoy was inserted into reporter ΔE4e, which has low intrinsic IR, and changes in retention were assessed in transfected K562 cells (Figure 4B). As a positive control we tested the 58nt ARGLU1 exon, and in parallel tested the 146nt OGT exon. In the minigene assay, both exhibited stronger IR activity than SF3B1 E4e, as indicated by increased intensity of the IR bands and decreased intensity of the spliced products (Figure 4B, lanes 3-4). Next we assayed four candidate decoys that have not been studied previously. DDX39B intron 6 was predicted to encode a decoy of 122-369nt, depending on use of alternative splice sites; this decoy exhibited strong IR activity (lane 5). A 60nt cassette in SNRNP70 intron 7 showed variable but relatively low activity (lane 6). In contrast, neither an 85nt predicted cassette in FUS intron 7, nor a 33nt cassette in KEL intron 6, had detectable IR activity (lanes 7-8).

Figure 4. **A. Splicing reporters used to assay heterologous decoys**. Construct "decoy" had E4e deleted and replaced with candidate decoys from other genes. B. RT-PCR analysis of intron retention activity associated with heterologous decoys in K562 cells. Lane 1, wild type; lane 2, ΔE4e; lanes 3-8, replacement of E4e with candidate decoys from ARGLU1 (lane 3); OGT (lane 4); DDX39B (lane 5); SNRNP70 (lane 6); FUS (lane 7); and KEL (lane 8). IR, intron retention product; spl, spliced product. Numbers below the lanes indicate apparent IR percent as determined by densitometry, corrected for differences in band size.

Careful analysis of the splice junction data revealed that most of the validated IR-promoting decoys have multiple alternative 5' and/or 3' splice sites (Figure S2). For ARGLU1, splice junction reads in NMD-inhibited erythroblasts confirmed four of the 5' splice sites identified earlier (Pirnie et al. 2017), and found two additional 3' splice sites. Similarly, we detected splice junctions corresponding to multiple sites in OGT (two 5' and two 3' splice sites), DDX39B (three 5' and two 3' splice sites), and SNRNP70 (three 5' splice sites) Figure S3. In contrast, candidate decoy exons with minimal IR activity in this assay (FUS, KEL33) lacked apparent competing splice sites.

Interestingly, although no alternative 3' splice sites were evident at SF3B1 E4e, mutation

analysis suggested that nearby AG dinucleotides can influence IR activity. Two AG dinucleotides were located 13-16nt upstream of the functional 3' splice site, in a region normally devoid of competing AGs. Mutation of these sites to TTTT (mutTT) or ACAC (mutAC) converted E4e splicing into a strongly included exon, and reduced but did not eliminate IR (Fig. 5, lanes 3-6; compare to wild type in lanes 1-2). Mutation of the functional 3' splice site AG (mut3') yielded a relative reduction in IR but without E4e inclusion (lanes 7-8). Mutation of all three AGs (mut3ag) yielded inclusion of a smaller exon E4e' by use of

Figure 5. **Splice site mutations at E4e affect the balance of IR and decoy exon inclusion**. Top region shows the mutations tested near the 3' and 5' splice sites. Gels below demonstrate the effects on splicing in K562 cells. WT, wild type; mutTT and mutAC, mutations of the upstream AG dinucleotides; mut3', mutation of the 3' splice site; mut3ag, mutation of all three AG dinucleotides; Δexon, deletion of E4e; mut5's, strong 5' splice site; mut5'w, mutated 5' splice site.

an internal AG 22nt downstream of the normal site (lanes 9-10). Together these results suggest that a key component of non-productive splice site interactions might be the presence of competing splice sites.

5' splice site strength also affected IR. For example, a C->A substitution at position +4 at the E4e 5' splice site modestly increased its splice site strength score, and noticeably increased E4e inclusion (lane 13) relative to that observed for WT (lane 11) and ΔE4e (lane 12). In contrast, a GT to CT mutation predicted to destroy the 5' splice site resulted in substantially reduced IR (lane 14). We conclude that strong splice sites favor decoy exon inclusion, thus reducing both productive splicing of E4 to E5 and retention of the intron. Conversely, splice sites that are too weak decrease decoy function, leading to increased productive splicing at the expense of competing pathways.

Binding of U2AF1 and U2AF2 to decoy exons. If decoy exons compete with intron terminal splice sites to enhance IR, the decoy model predicts that decoy splice junctions must be recognized by the splicing machinery as an essential early step. We examined this question by taking advantage of ENCODE eCLIP data that defined U2AF1 and U2AF2 binding sites in the K562 transcriptome. Figure S3 showed that U2AF1 and 2 binding sites align with the OGT, ARGLU1, and DDX39B decoy exons that have documented IR activity, but less or no binding was associated with decoys that did not promote IR. When analyzed for the full set of decoy exons, good peaks were observed for both U2AF1 and U2AF2 upstream of the decoy exons (Figure 6). These patterns were very similar to, although slightly weaker than, profiles for known expressed cassettes in K562 cells (native cassette exons), despite the fact that most candidate decoys exhibit very low splicing efficiencies in control erythroid cells.

Figure 6. U2AF binding profiles in retained introns with candidate decoys. ENCODE binding profiles for U2AF1 and U2AF2 in K562 erythroleukemia cells. Profiles show enriched binding for U2AF factors near candidate decoys was similar to that observed for a control set of cassette exons in the same cells.

Discussion

Experimental and computational data support the concept that decoy-mediated IR is an novel mechanism for regulating an important component of the erythroid IR program. Experimental analysis of SF3B1 minigene splicing reporters demonstrated that highly conserved decoy exons are required for optimum retention, that decoy exons can induce IR in an ordinarily non-retained intron, and that heterologous decoys from other retained introns can promote SF3B1 i4 retention. In parallel, computational analysis of RNA-seq data from NMD-inhibited cells suggested a much broader program of decoy exon-regulated IR is executed during terminal erythropoiesis. According to this model, intron-terminal splice sites in many >1kb introns interact with competing intron-internal splice sites at decoy exons, blocking the cross-intron interactions needed to facilitate intron excision (Figure 7). In human erythroblasts, our data indicate that decoy-regulated RIs comprise several hundred erythroblast retention events that are enriched in RNA binding proteins, and are mechanistically distinct from the bulk population of smaller RIs that generally lack decoys. Notably, the decoy exon mechanism greatly extends the model reported recently

for ARGLU1, where it was proposed that unproductive splicing complexes assembled at the alternative exon disfavor intron splicing so as to promote its retention (Pirnie et al. 2017). The model is also consistent with previous reports that RIs are enriched adjacent to alternative exons (Braunschweig et al. 2014).

Figure 7. Decoy splicing model. In the gene region shown above, retention of the red intron can be regulated by use of competing splice sites associated with the red decoy exon. If the decoy is not engaged, intron-terminal splice sites interact to promote excision of the intron. If the decoy is recognized, possibly aided by intron splicing enhancers (ISE), then competing splice site interactions prevent excision to yield intron retention.

The role of decoy-mediated IR in terminal erythropoiesis remains to be investigated. Given our earlier finding that several major spliceosomal factors possess highly retained introns (Pimentel et al. 2016), and new data showing that some of these possess IR-promoting decoys, we suggest this mechanism could modulate changes in splicing capacity of late stage erythroblasts as they reduce gene expression in preparation for enucleation. Another possibility is that regulated IR could contribute to balanced expression of competing or cooperating genes. For example, modulation of IR plays an important role in regulating expression for competing OGT and OGA enzymes to ensure O-GlcNAc homeostasis (Park et al. 2017), and the documented decoy exon in OGT likely contributes to that control. Speculatively, IR could help ensure comparable expression of alpha and beta spectrin subunits, two structural polypeptides that form long heterodimers and provide mechanical support to the red cell membrane skeleton; a decoy exon in SPTA1 intron 20 could play a role in balancing expression.

The molecular mechanism by which decoy sites can non-productively engage intron-terminal splice sites requires further study. Decoy splice site function must be carefully tuned, since strong splice sites favor decoy exon inclusion over intron retention, while splice site-inactivating mutations can abrogate IR-promoting activity. One feature that may be critical is the presence of multiple competing splice sites at the decoy exon. Multiple 5' and/or 3' splice sites are a feature of the decoy exon in ARGLU1 (Pirnie et al. 2017) and other transcripts. Moreover, even when explicit splice junction variation is not evident, some decoys possess potentially competing splice site motifs in the proximal introns, exemplified by the extra AG dinucleotides upstream of SF3B1 E4e (Figure 5). On the other hand, cassette exons encoded in retained introns from FUS and KEL did not appear to have

alternative splice sites, and did not exhibit IR activity in the splicing reporter assay.

Competition between splice sites is a fundamental governing principle of alternative splicing. The concept that competing sites can be spliced inefficiently, functioning mainly as decoys to block functional use of other sites, is also well grounded in previous work. An early example was the Drosophila P-element gene, where U1 snRNP binding to an exonic pseudo-5' splice site was shown to compete with the normal downstream 5' splice site to promote retention of adjacent intron in somatic cells (Siebel et al. 1992). Decoy 3' acceptors can engage the 5' splice site of an upstream exon to promote its skipping in caspase-2 (Cote et al. 2001) and other transcripts (Havlioglu et al. 2007). The concept that a noncoding exon might engage splice sites of both flanking exons to induce IR was suggested recently (Pirnie et al. 2017) and the decoy exon data presented here greatly expands the potential role of decoy splice sites in regulation of splicing outcomes. Interestingly, the appreciation of decoy exons might help to explain the observation that U2AF binds to numerous intronic locations not corresponding to known 3' splice sites (Shao et al. 2014). Some of these sites might compete with nearby exons to influence exon skipping (Shao et al. 2014), and our data suggest that other sites might promote intron retention. Moreover, based on our finding that retained introns sometimes possess "extra" U2AF binding not aligned with observed splice junctions, we speculate that a subset of intronic U2AF sites function as decoys without ever being spliced themselves.

In conclusion, the decoy model represents a new and distinctive component of the erythroid alternative splicing program. We speculate that developmentally regulated IR events in other cell types may also be regulated by decoy exons, and that physiological control of such programs could be mediated by context-dependent combinations of splicing enhancer and silencer proteins that impact recognition of the decoys. We anticipate that analogous subsets of IR events might be regulated by decoy mechanisms in other developmental or physiological contexts. Possible candidates could include differentiating granulocytes (Wong et al. 2013; Wong et al. 2017), activated T cells (Ni et al. 2016), stimulated neurons (Mauger et al. 2016), cells subjected to proteotoxic stress (Shalgi et al. 2014), proliferating vs differentiated muscle cells (Llorian et al. 2016), differentiating germ cells (Naro et al. 2017), etc. Identifying the RBPs that regulate these decoy-dependent programs will be an important goal of future studies in this area.

Methods.

Erythroblast culture.

Cells were cultured as described previously (Hu et al. 2013). For NMD experiments, cultures were divided and one half of each culture was incubated with 100ug/ml emetine for 8 hours and 100ug/ml cycloheximide for 4 hours. Experiments were done in biological duplicates for a total of eight samples (two differentiation stages, two conditions plus or minus NMD inhibitors, and two biological replicates.

RNA and RNA library Preparation

RNA was isolated from erythroblasts ($4x10^6$ cells from day 9 and $16x10^6$ cells on day 16) according to the manufacturer's instructions (Qiagen). Sequencing libraries were prepared

from 500 ng of total RNA using the NEBNext Poly(A) mRNA Magnetic Isolation Module (catalog number E7490, protocol revision 5.0), NEBNext Ultra Directional RNA Library Preparation Kit for Illumina (catalog number E7420, protocol revision 6.0), and NEBNext Multiplex Oligos for Illumina (catalog number E7600, protocol revision 2.0) with the following modifications: E7420, Section1.2 ("mRNA Isolation, Fragmentation and Priming Total RNA") Step 37, we decreased the incubation time from 15 min to 5 min; E7420, Section 1.3 ("First Strand cDNA Synthesis"), Step 2,we increased the incubation time from 15 min to 50 min; for the size selection we used 40ul AMPure XP beads for the 1st bead selection and 20ul AMPure XP beads for the 2nd bead selection (targeting an insert size of 300-450bp and a final library size of 400-550bp); and in E7420, Section 1.9A ("PCR library Enrichment"), Step 2 we used 14 cycles for PCR cycling and dual index primers (i507 and i705-i712). Individual libraries were normalized to 10nM and eight samples were pooled per lane. Sequencing was performed at UC Berkeley's QB3 Vincent J. Coates Genomics Sequencing Laboratory on an Illumina HiSeq4000 instrument, generating 150bp paired end reads.

RNA-seq analysis.

For each sample we produced 13-60M total reads and 4-30M mapped reads. Replicates were merged and aligned to the GRCh38.p8 version of the human genome using TopHat version: 2.1.1 (Trapnell et al. 2009; Kim et al. 2013). BAM files generated were used for the alternative splicing analysis and for the cassette reannotation. All the splicing-junction data formatted as bigBed and bigWig files obtained in this study were uploaded onto the UCSC genome browser; it can be accessed by copying the following link into a web browser: http://genome.ucsc.edu/cgi-bin/hgTracks?db=hg38&hubUrl=https://sina.lbl.gov/seqdata/conboy-ucsc/hub.txt. The FASTQ files were run through the STAR aligner (Dobin et al. 2013) to produce expression scores as input for DEXSeq (Anders et al. 1984; Reyes et al. 2013)

Custom Cassette Reannotation Scripts

A custom reannotation tool was written using the Rust language to reannotate an input annotation to include additionally discovered cassette exons. The source code can be found at GitHub: https://github.com/bdgp/cassette_reannotation. The reannotation tool was given a curated subset of the NCBI RefSeq annotation version GCF_000001405.34 based on GRCh38.p8. This curated subset only included features beginning with NM, NR, and YP. TopHat version: 2.1.1 (Trapnell et al. 2009; Kim et al. 2013) generated alignment files of the sample reads were also passed in to the re-annotation tool. The tool finds constitutive splice pairs in the input annotation, then searches for overlapping reads in the input read alignment files that provide evidence for unannotated cassette exons. The tool requires at least two paired-end fragment splices for both the start and the stop of the cassette which also must splice at least one of the annotated flanking exons. The tool also requires contiguous read coverage throughout the discovered cassette. The tool then produces a re-annotation that includes all of the features of the input annotation, along with newly created transcript features. The newly created transcript features are based on transcript features existing in the input annotation, but with the discovered cassettes added.

Alternative Splicing Analysis

The re-annotated annotation including discovered cassettes, along with the alignment files, were passed to SplAdder (Kahles et al. 2016) http://github.com/ratschlab/spladder to discover retained introns and alternative exons in the annotation. Some minor bug fixes, disabling of code assertions and parameter tuning were required for SplAdder to perform the analysis. We discovered that the least-stringent default parameter set was still to stringent to detect many retained introns in our sample data, so the individual parameters were made even less stringent. Details of the parameters used in our run of SplAdder can be found Supplementary Materials.

The retained intron adjusted PSI values returned by SplAdder unfortunately did not account for exons overlapping the retained intron, so a custom tool was written to correct the adjusted PSI values. SplAdder computes the adjusted PSI values using the average per-base coverage over the intron (intron_cov) and the number of reads with splices confirming the intron (intron_conf). The tool written to correct the adjusted PSI values recomputes the average per-base coverage by subtracting out the coverage of exonic regions. The tool uses this corrected intron_cov value along with the intron_conf directly from SplAdder to compute the corrected adjusted PSI value for the retained intron.

Splicing reporter assays.

A 4.7 kb region of the human SF3B1 gene extending from the 3' end of intron 2 to the 5' end of intron 6, was amplified using the following primers: F: 5'-tggaattctgcagatAAGGAGGGCTTAGACATCACAC-3'; R: 5'-gccagtgtgatggatCTATGGCAACCCAAGCAGA-3'. The fragment was cloned into pcDNA3.0 using In-Fusion methods (Gibson 2011) with 15nt in lower case sequence representing overlap with the ends of EcoRV-linearized vector. The splicing reporter was transfected into K562 using Fugene HD according to the manufacturer's instructions (Promega). RNA was harvested after 48 hrs and purified according to the manufacturer's instructions (Qiagen), but with the addition of a DNase step to eliminate potential contamination by genomic DNA. RNA was reverse transcribed with Superscript III (Invitrogen) into cDNA using the BGH reverse primer in the vector (5'-tagaaggcacagtcgagg-3'). Spliced products were amplified using a forward primer in exon 3 (5'-catcatctacgagtttgcttgg-3') and a reverse primer in the vector (5'-atttaggtgacactatagaatagggc-3'). PCR reaction conditions were adjusted to allow for amplification of IR products ≥ 3 kb in length (denaturation at 95° C for 20sec , annealing at 56° C for 10sec, extension at 70° C for 2min 30sec; 35 cycles) using KOD polymerase in the presence of betaine to enhance amplification. PCR products were analyzed on either 2% agarose gels or 4.5% acrylamide gels. All PCR products discussed in the manuscript were confirmed by DNA sequencing. This strategy amplified minigene-derived transcripts but not endogenous SF3B1 mRNA, as confirmed using RNA from untransfected or empty vector-transfected cells.

CLIP-seq analysis.

Splicing maps were generated using U2AF1 (ENCSR862QCH) and U2AF2 (ENCSR893RAV) eCLIP normalized densities overlapped with selected retained introns containing candidate decoy exons at two time points (D9, D16). These density values were normalized first using an RPM transformation to account for variation in sequencing depth, then normalized

again using equivalent densities from a size-matched input sample. To perform this second normalization, both IP and equivalent input signals were transformed into their probability densities to preserve overall shape of binding and to reduce signal dominance from few events. Input densities for each event were then subtracted from the corresponding IP to remove background signal. Each line represents the mean of these normalized densities devoid of any value exceeding the 95% median at each position to reduce confounding outlier effects.

In addition to candidate decoy exons, normalized densities were also overlapped with a set of cassette exons, derived from a subset of Gencode (v19) constitutive exons. Within these annotations, we define a 'cassette' as any exon found between 10% and 90% spliced in at least 50% of all shRNA knockdown control data (encodeproject.org, all non-specific target controls, aligned to hg19 with TopHat), filtering any region that is not supported by at least 30 reads. From this set, regions with the highest inclusion average between two replicates were chosen if any regions overlapped to remove any possibility of double counting eCLIP signal.

ACKNOWLEDGEMENTS

This work was funded by NIH grant 5R01DK108020 (JGC).

REFERENCES

Anders RF, Coppel RL, Brown GV, Saint RB, Cowman AF, Lingelbach KR, Mitchell GF, Kemp DJ. 1984. Plasmodium falciparum complementary DNA clones expressed in Escherichia coli encode many distinct antigens. *Mol Biol Med* **2**: 177-191.

Boothby TC, Zipper RS, van der Weele CM, Wolniak SM. 2013. Removal of retained introns regulates translation in the rapidly developing gametophyte of Marsilea vestita. *Dev Cell* **24**: 517-529.

Boutz PL, Bhutkar A, Sharp PA. 2015. Detained introns are a novel, widespread class of post-transcriptionally spliced introns. *Genes Dev* **29**: 63-80.

Braun CJ, Stanciu M, Boutz PL, Patterson JC, Calligaris D, Higuchi F, Neupane R, Fenoglio S, Cahill DP, Wakimoto H et al. 2017. Coordinated Splicing of Regulatory Detained Introns within Oncogenic Transcripts Creates an Exploitable Vulnerability in Malignant Glioma. *Cancer Cell* **32**: 411-426 e411.

Braunschweig U, Barbosa-Morais NL, Pan Q, Nachman EN, Alipanahi B, Gonatopoulos-Pournatzis T, Frey B, Irimia M, Blencowe BJ. 2014. Widespread intron retention in mammals functionally tunes transcriptomes. *Genome Res* **24**: 1774-1786.

Cote J, Dupuis S, Jiang Z, Wu JY. 2001. Caspase-2 pre-mRNA alternative splicing: Identification of an intronic element containing a decoy 3' acceptor site. *Proc Natl Acad Sci USA* **98**: 938-943.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**.

Edwards CR, Ritchie W, Wong JJ, Schmitz U, Middleton R, An X, Mohandas N, Rasko JE, Blobel GA. 2016. A dynamic intron retention program in the mammalian megakaryocyte and erythrocyte lineages. *Blood* **127**: e24-e34.

Gibson DG. 2011. Enzymatic assembly of overlapping DNA fragments. *Methods in Enzymology* **498**: 349-361.

Havlioglu N, Wang J, Fushimi K, Vibranovski MD, Kan Z, Gish W, Fedorov A, Long M, Wu JY. 2007. An intronic signal for alternative splicing in the human genome. *PLoS ONE* **2**: e1246.

Hu J, Liu J, Xue F, Halverson G, Reid M, Guo A, Chen L, Raza A, Galili N, Jaffray J et al. 2013. Isolation and functional characterization of human erythroblasts at distinct stages: implications for understanding of normal and disordered erythropoiesis in vivo. *Blood* **121**: 3246-3253.

Jacob AG, Smith CWJ. 2017. Intron retention as a component of regulated gene expression programs. *Hum Genet* **136**: 1043-1057.

Jin S, Su H, Tran N-T, Song J, Lu SS, Li Y, Huang S, Abdel-Wahab O, Liu Y, Zhao X. 2017. Splicing factor SF3B1K700E mutant dysregulates erythroid differentiation via aberrant alternative splicing of transcription factor TAL1. *PLOS ONE* **12**: e0175523.

Kahles A, Ong CS, Zhong Y, Rätsch G. 2016. SplAdder: identification, quantification and testing of alternative splicing events from RNA-Seq data. *Bioinformatics* **32**: 1840-1847.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology* **14**: R36.

Li Y, Bor YC, Fitzgerald MP, Lee KS, Rekosh D, Hammarskjold ML. 2016. An NXF1 mRNA with a retained intron is expressed in hippocampal and neocortical neurons and is translated into a protein that functions as an Nxf1 cofactor. *Mol Biol Cell* **27**: 3903-3912.

Llorian M, Gooding C, Bellora N, Hallegger M, Buckroyd A, Wang X, Rajgor D, Kayikci M, Feltham J, Ule J et al. 2016. The alternative splicing program of differentiated smooth muscle cells involves concerted non-productive splicing of post-transcriptional regulators. *Nucleic Acids Res* **44**: 8933-8950.

Mauger O, Lemoine F, Scheiffele P. 2016. Targeted Intron Retention and Excision for Rapid Gene Regulation in Response to Neuronal Activity. *Neuron* **92**: 1266-1278.

Naro C, Jolly A, Di Persio S, Bielli P, Setterblad N, Alberdi AJ, Vicini E, Geremia R, De la Grange P, Sette C. 2017. An Orchestrated Intron Retention Program in Meiosis Controls Timely Usage of Transcripts during Germ Cell Differentiation. *Developmental cell* **41**: 82-93 e84.

Ni T, Yang W, Han M, Zhang Y, Shen T, Nie H, Zhou Z, Dai Y, Yang Y, Liu P et al. 2016. Global intron retention mediated gene regulation during CD4+ T cell activation. *Nucleic Acids Res* **44**: 6817-6829.

Obeng EA, Chappell RJ, Seiler M, Chen MC, Campagna DR, Schmidt PJ, Schneider RK, Lord AM, Wang L, Gambe RG et al. 2016. Physiologic Expression of Sf3b1(K700E) Causes Impaired Erythropoiesis, Aberrant Splicing, and Sensitivity to Therapeutic Spliceosome Modulation. *Cancer Cell* **30**: 404-417.

Park SK, Zhou X, Pendleton KE, Hunter OV, Kohler JJ, O'Donnell KA, Conrad NK. 2017. A Conserved Splicing Silencer Dynamically Regulates O-GlcNAc Transferase Intron Retention and O-GlcNAc Homeostasis. *Cell Rep* **20**: 1088-1099.

Pendleton KE, Chen B, Liu K, Hunter OV, Xie Y, Tu BP, Conrad NK. 2017. The U6 snRNA m6A Methyltransferase METTL16 Regulates SAM Synthetase Intron Retention. *Cell* **169**: 824-835.e814.

Pimentel H, Parra M, Gee SL, Mohandas N, Pachter L, Conboy JG. 2016. A dynamic intron retention program enriched in RNA processing genes regulates gene expression during terminal erythropoiesis. *Nucleic Acids Res* **44**: 838-851.

Pirnie SP, Osman A, Zhu Y, Carmichael GG. 2017. An Ultraconserved Element (UCE) controls homeostatic splicing of ARGLU1 mRNA. *Nucleic Acids Res* **45**: 3473-3486.

Reyes A, Anders S, Weatheritt RJ, Gibson TJ, Steinmetz LM, Huber W. 2013. Drift and conservation of differential exon usage across tissues in primate species. *Proceedings of the National Academy of Sciences* **110**: 15377.

Schmitz U, Pinello N, Jia F, Alasmari S, Ritchie W, Keightley MC, Shini S, Lieschke GJ, Wong JJ, Rasko JEJ. 2017. Intron retention enhances gene regulatory complexity in vertebrates. *Genome Biol* **18**: 216.

Shalgi R, Hurt JA, Krykbaeva I, Taipale M, Lindquist S, Burge CB. 2013. Widespread regulation of translation by elongation pausing in heat shock. *Mol Cell* **49**: 439-452.

Shalgi R, Hurt JA, Lindquist S, Burge CB. 2014. Widespread Inhibition of Posttranscriptional Splicing Shapes the Cellular Transcriptome following Heat Shock. *Cell reports* **7**: 1362-1370.

Shao C, Yang B, Wu T, Huang J, Tang P, Zhou Y, Zhou J, Qiu J, Jiang L, Li H et al. 2014. Mechanisms for U2AF to define 3' splice sites and regulate alternative splicing in the human genome. *Nat Struct Mol Biol* **21**: 997-1005.

Siebel CW, Fresco LD, Rio DC. 1992. The mechanism of somatic inhibition of Drosophila P-element pre-mRNA splicing: multiprotein complexes at an exon pseudo-5' splice site control U1 snRNP binding. *Genes Dev* **6**: 1386-1401.

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* **25**: 1105-1111.

Wong JJ, Gao D, Nguyen TV, Kwok CT, van Geldermalsen M, Middleton R, Pinello N, Thoeng A, Nagarajah R, Holst J et al. 2017. Intron retention is regulated by altered MeCP2-mediated splicing factor recruitment. *Nature communications* **8**: 15134.

Wong JJ, Ritchie W, Ebner OA, Selbach M, Wong JW, Huang Y, Gao D, Pinello N, Gonzalez M, Baidya K et al. 2013. Orchestrated intron retention regulates normal granulocyte differentiation. *Cell* **154**: 583-595.

Table 1. Splicing reporters

Splicing reporter	Genome coordinates (GRCh38/hg38)
Wild type	chr2:197,416,658-197,421,257
ΔE4ab	WT Δ chr2:197,420,152-197,420,358
ΔE4c	WT Δ chr2:197,419,814-197,419,929
ΔE4d	WT Δ chr2:197,419,575-197,419,677
ΔE4e	WT Δ chr2:197,418,852-197,419,014
ΔE4f	WT Δ chr2:197,418,719-197,418,815
OGT	chrX:71,546,208-71,546,632
ARGLU1	chr13:106,557,456-106,557,720
DDX39B	chr6:31,534,442-31,534,912
SNRNP70	chr19:49,101,950-49,102,379
KEL	chr7:142,956,987-142,957,311
FUS	chr16:31,187,786-31,188,063
mut24.2	E4e chr2:197,418,851-197,419,013 inserted in i5 after position chr2:197,417,934
mut31	1.5kb of i4 sequences (chr2:197,418,852-197,420,358) inserted in place of 1.25 kb of i5 sequence (chr2:197,418,326-197,417,086)
Decoy exon regions from OGT, ARGLU1, DDX39B, SNRNP70, KEL, and FUS were inserted in place of the E4e region chr2:197,418,795-197,419,034	

Figure S1. Conservation of SF3B1 exon 4a in most mammals.

Figure S1, cont. Conservation of SF3B1 exon 4b to reptiles.

Figure S1, cont. Conservation of SF3B1 exon 4c to reptiles.

Figure S1, cont. Conservation of SF3B1 exon 4d to fish genomes.

Figure S1, cont. Conservation of SF3B1 exon 4e to fish genomes.

Figure S1, cont. Conservation of SF3B1 exon 4f to fish genomes.

Figure S2

Figure S2. Splice junctions defining candidate decoy exons in selected erythroblast retained introns.

Figure S3

Figure S3. Genome browser tracks showing gene structure and expression feature of representative retained introns. Red boxes highlight predicted cassette exons that reside in retained introns. Considerable overlap with U2AF binding peaks was observed for some decoys (ARGLU1, OGT, DDX39B), but overlap was reduced (SNRNP70, KEL) or absent (FUS) in others.