How to make use of ordination methods to identify local adaptation: a comparison of genome scans based on PCA and RDA.

Thibaut CAPBLANCQ[1*], Keurcien LUU[2], Michael G.B. BLUM[2] and Eric BAZIN[1,#]

[1] *Univ. Grenoble Alpes, CNRS, LECA UMR 5553, Grenoble, France*
[2] *Univ. Grenoble Alpes, CNRS, TIMC-IMAG UMR 5525, Grenoble, France*
[#]*Deceased*

* Corresponding author: Thibaut Capblancq, 2233 Rue de la Piscine, 38041 Grenoble Cedex, France. Tel: +33 (0)4 76 51 36 71, e-mail: thibaut.capblancq@univ-grenoble-alpes.fr

Date: January 30[th], 2018

Running title: RDA-based procedure for genome scans

ABSTRACT

Ordination is a common tool in Ecology that aims at representing complex biological information on a reduced space. For instance, it is frequently used to study geographic distribution pattern of species diversity and to study the link between ecological variable such as temperature, drought, etc, and the species turnover. Recently, ordination methods such as PCA have been used as a framework to detect adaptive variation based on genomic data. Taking advantage of environmental data in addition to genotype data, redundancy analysis (RDA) is another ordination approach useful to detect adaptive variation. These methodologies are specially used in Landscape Genomic where one wants to study the link between environmental variable and the distribution pattern of genome wide diversity, especially its adaptive component. This paper aims at proposing a test based on RDA approaches to search for genes under selection and to compare it to the PCA method implemented in the *pcadapt* package. Individual based simulations identify evolutionary scenarios where RDA genome scan has a greater statistical power than PCA genome scan. By constraining the analysis with environmental variables, RDA performs better than PCA on identifying adaptive variation when selection gradients are weakly correlated with population structure. Additionally, we show that RDA is efficient to identify the main selective gradients among a set of environmental variables. Finally, to give a concrete illustration of RDA in population genomics, we

apply this method to the detection of outliers and selective gradients on a SNP data

set of *Populus trichocarpa* in North America (Geraldes et al, 2013). RDA-based

approach identifies the main selective gradient that corresponds to a temperature

gradient contrasting southern and coastal populations to northern and continental

populations in North Western American coast.

KEYWORDS: Genome scans - Multivariate analysis - Redundancy Analysis –

pcadapt - Selection.

INTRODUCTION

Natural selection is the result of a complex set of environmental pressures.

Environment acts on several characters simultaneously and these adaptive

characters can be determined by a large number of alleles of small effects (Pritchard

and Di Rienzo, 2010). Patterns of adaptive variation within species are usually

studied through physiological, morphometric and fitness comparisons in common

gardens experiments. However, this is a difficult and constraining task and is even

unrealistic on species with long generation time. The recent development of Next

Generation Sequencing (NGS) technics has opened the possibility to access to a large

amount of genetic variation across the genome. Therefore, relationships between

genetic polymorphism, phenotypic variation and environmental variables can now

be studied and quantified by in situ approaches (Steane *et al.*, 2014, Vangestel *et al.*,

2018).

The search of adaptive genetic variation can be performed by genome scan

methods (Foll and Gaggiotti 2008, Frichot *et al.*, 2013, Vatsiou *et al.*, 2015, Duforet-

Frebourg *et al.*, 2016). Among the different genome scan procedures, ecological

association methods aim at detecting alleles that are associated with environmental

variables (e.g. temperature, drought) with the idea that these alleles may confer a

selective advantage in some environment (Coop *et al.*, 2010, Frichot *et al.*, 2013).

Finally, methods based on genetic differentiation assume that adaptive alleles have

increased genetic differentiation. A common statistic to perform genome scans

based on genetic differentiation is F_{st} (Lewontin and Krakauer, 1973). Many

likelihood or Bayesian methods use F_{st}-related statistics to scan genomes for local

adaptation (Bazin *et al.*, 2010, De Villemereuil and Gaggiotti, 2015, Foll and

Gaggiotti, 2008, Whitlock and Lotterhos, 2015).

F_{st}-based approaches require to define discrete populations. However, many

species are not structured in different populations but show a pattern of isolation by

distance and display continuous population structure (Martins *et al.* 2016). In these

cases, defining populations can be challenging and can directly interfere with the

result of the analysis. Instead of defining populations, multivariate methods such as

PCA can be used to ascertain population structure. Following this analysis, outlier

loci are detected if they have extremely high correlation with one or more ordination

axes (Duforet-Frebourg *et al.*, 2016). Simulations have shown that they have similar

power than model-based methods under island models and perform better when the

simulated demographic model drift from the island model, for instance under

hierarchical population structure or isolation by distance patterns (Luu *et al.*, 2017).

Nevertheless, one conundrum of such approaches is the difficulty to interpret

ordination axes in term of ecological meanings. These are usually tight to

geographical axes (latitudinal or longitudinal) but they are not necessary linked to

an environmental variable such as Temperature, drought, diet habit, etc. Therefore,

when environmental information exists, it has to be used a posteriori as a mean of

interpretation but it is not involved in the inference process.

However, it is possible to detect the adaptive fraction among genetic variation,

which is explained by environmental variation. Some authors (Lasky *et al.*, 2012,

DeKort *et al.*, 2014; Steane *et al.*, 2014, Hecht *et al.*, 2015) have proposed to use

ordination technics such as RDA (Redundancy Analysis) or CAPs (Canonical Analysis

of Principal coordinates) to subsequently align adaptive variation with

environmental data in order to be able to identify the environmental gradients that

are the most correlated with adaptive variation. In *Arabidopsis thaliana,* loci

involved in adaptation to climate have been found using RDA (Lasky *et al.*, 2012).

Outliers were identified as SNPs with the greatest squared loadings along the first

RDA axes (i.e. those in the 0.5 % tail). RDA can also be used to derive an adaptive

index that predicts the performance of individuals in different environmental

conditions. Steane *et al.* (2014) applied this approach to *Eucalyptus trichocarpa*

populations and showed that this in-situ estimation of adaptive variation is

correlated to what can be measured in common garden experiments. Finally, in two

recent studies, Forester and colleagues (2015 and 2017) tested the power of RDA

based method for detecting signature of local adaptation in a heterogeneous

landscape (Forester *et al*, 2015) and in comparison with other constrained

ordination method (Forester *et al*, 2017). Using simulations, they found that a RDA

based method shows a great potential to identify adaptive loci and, has a superior

combination of low false positive and high true positive rates when compared to

several other approaches.

In this paper, we build on these previous applications of RDA to develop a

statistical test for searching loci under selection. We use simulations and real data

sets to shed light on the conditions within which RDA is the most efficient and to

document the possibilities given by RDA when studying landscape genomics. We

also compare results of genome scans based on RDA and based on PCA as it is

implemented in the *pcadapt* package (Luu *et al.,* 2017).

First, we show that genome scans based on RDA allows to recover more

outliers than a genome scan based on PCA because it accounts for environmental

variation. Second, we show that RDA can identify important environmental

gradients that explain adaptive variation in the data. It is therefore a proof of

concept of the idea of using constrained ordination method as an environmental

genomic tool to identify relevant selective gradients in the environmental data

(DeKort *et al.*, 2014, Steane *et al.*, 2014). Finally, to illustrate the pertinence of such

method in concrete example, we apply it to a *Populus trichocarpa* SNP dataset

obtained from Geraldes et al (2013). Our results mostly concord with the ones

obtained with the *Fst* based methods used in their study but we also show some new

genetic markers specifics to the RDA-based approach. Additionally, RDA identifies

the main selective gradients along the natural populations of *Populus trichocarpa* in

the north-western coast of North America.

MATERIALS AND METHODS

1.   Genome scan using a RAD based approach

Redundancy analysis (RDA) was first introduced by Rao (1964). It is the

extension of multiple regressions to the modelling of multivariate response data

(Legendre and Legendre 2012, section 11.1). The data are separated in two sets, a

response matrix Y of variable to be explained (e.g. species abundance in a set of

sites; m sites and n species) and an explanatory matrix X (e.g. a set of environmental

variable within each site; m sites and p environment). In the following analysis, loci

replace species abundances and individuals replace sites. RDA seeks to project on a

reduced space the genetic variation between individuals that is explained by

environmental data.

The analysis starts using a classical RDA procedure performed with the

function *rda* of the package R *"vegan"* (Oksanen *et al.*, 2015). A genetic dataset (e.g.

SNPs genotypes 0, 1, 2) is used as response matrix Y and a set of environmental

variables is used as explanatory matrix X. RDA amounts at constructing a matrix Y' of

predicted genotypes based on the environmental variables and at performing

principal component analysis on the matrix Y' (Legendre and Legendre 2012, section

11.1). After this constrained ordination step, we follow the methodology

implemented in *pcadapt* to find outliers loci (Luu *et al.,* 2017). First we recover from

the RDA analysis the z-scores obtained when regressing the SNPs by the RDA axes.

We obtained vectors of z-scores denoted $z_j = (z_{j1}, ..., z_{jK})$ where z_{jk} corresponds to the

z-score obtained when regressing the j-th SNP by the K-th ordination axis. Only the

z-scores of the most informative ordination axis are kept for the rest of the

procedure. This number (K) is determined by looking at the amount of information

retained on the different axes of the RDA. A robust Mahalanobis distance D is then

computed for each locus to identify outlier vectors of z-scores. Computation of the

robust Mahalanobis distance uses the *covRob* function of the "robust" R package

(Wang et al., 2014). Mahalanobis distances are distributed as Khi2 distribution with

K degrees of freedom after correcting with the genomic inflation factor (Luu et al.,

2017). We use the False-Discovery-Rate (FDR) approach to control for false

positives. Qvalue are computed with "qvalue" R package (Dabney and Storey, 2011)

and a locus is considered as an outlier if its qvalue is less than 10%. The complete

procedure and our script are available in supplementary material (Script S1).

To evaluate the capacity of the RDA based approach to detect loci underlying

environmental selective pressure we first test the method with a simulated dataset.

We look at its capacity to detect positive loci (loci simulated under selection) and to

return a controlled number of false positives (neutral loci detected as outliers by the

analysis). Then, to emphasize the utility of RDA, we compared it to the PCA-based

procedure implemented in *pcadapt* that does not use environmental variables (Luu

et al., 2017).

2.   Simulated data

To test for the efficiency of RDA in population genomics, we used simulations

performed with the simuPop python library (Peng *et al.*, 2005). A lattice of 8x8

populations is simulated (i.e. 64 populations in total). Each population is initialized

with 200 diploid individuals with random genotypes. Migration is set to 0.1 resulting

in an isolation-by-distance pattern across the species range. Loci are assumed to be

biallelic SNPs. Allele frequency of the whole population is initialized at 0.5. 1000 loci

are simulated in total and they are separated in 200 chunks of 5 SNPs in physical

linkage with recombination rate between adjacent loci fixed at 0.1. Three different

traits are coded by a group of 10 different loci. The first trait is coded by the loci 1,

11, 21, ..., 91. The trait value is simply the sum of the genotype values and therefore

can take value between 0 and 20. We add to each trait a random noise (non heritable

variation) drawn from a normal distribution $N(0,2)$. The second trait is coded by loci

101, 111, ..., 191 and the third is coded by loci 201, 211, ..., 291. Each trait is

therefore coded by 10 independent SNP loci resulting in a total of 30 causal SNPs

among 1000. Selection can have an effect on linked loci, for instance, loci 2, 3, 4 and 5

can be impacted by selection on locus 1. However, recombination is high enough

(0.1) to expect a limited linkage effect. We have defined 10 different environmental

variables. The first one determines the selective pressure on trait 1, the second one

on trait 2 and the third one on trait 3. The first environment variable is a quadratic

gradient coded by function $env1$ = $-(\cos(\theta)*(i-3.5))^2 -(\sin \theta)*(j-3.5))^2 + 18$, $\theta = \pi/2$, i

and j being the population indicator on the 8x8 lattice. The second one is a linear

plan gradient coded by function $env2$ = $h*\cos(\theta)*(i-1) + h*\sin(\theta)*(j-1) + k$ with $h=2$,

$\theta = \pi/4$ and $k=3$. The third environment variable simulates a coarse environment

with value $env3$ = 2 for all populations except population (i,j) = (2,2), (2,3), (3,2),

(3,3), (6,2), (6,3), (7,2), (7,3), (2,6), (2,7), (3,6), (3,7), (6,6), (6,7), (7,6), (7,7) for

which $env3$ = 18. Env4, env5 and env6 have exactly the same equation than env1,

env2 and env3 respectively except for a noise term explained below. The remaining

4 environment variables are similar to env2 but with different value of h and θ. Env7

has h=2, θ = 0 and k=3. Env8 has h=2, $\theta = \pi/4$ and k=0. Env9 has h=1, $\theta = \pi/4$ and

k=4. Env10 has h=0.5, $\theta = \pi/4$ and k=8. Graphical representation of the mean

environmental value is given in Fig. 1. Environmental variables 4, 5 and 6 have

respectively the same mean value spatial distribution than environmental variables

1, 2 and 3. Environmental equation gives a mean value of the environment variable.

To avoid colinearity between environmental variable, we added noise by drawing an

environment value within a normal distribution N(μ = env, σ = 1). Fitness for each

trait is set to be –exp(x-env)^2/(2*ω^2), x being the quantitative trait value, env the

environmental value and ω is defining selection strength and has been set to 20

which was found to be enough for loci to be often detected. To get the overall fitness

for a given individual, fitness associated to each trait are multiplied. Fitness values

are used to determine the number of offsprings during the simulations, which have

been run during 500 generations. At the end of simulation, we sample 10 individuals

per population resulting in 640 individuals with 1000 SNP-like loci.

3.   **Real dataset**

The *Populus trichocarpa* dataset is a sample of 424 individuals genotyped on

33,070 SNPs from 25 drainages (i.e., topographic units separated by watershed

barriers) (Geraldes *et al.*, 2014). Genotyping of each accession was performed with a

34K Populus SNP array targeting 34,131 SNPs mostly within (plus 2kb upstream and

downstream) 3543 genes (Geraldes *et al.*, 2013). A total of 21 climatic variables are

available on each sampling site (Table S2). From the 33,070 SNPs, we removed the

SNPs with missing values and a minor allele frequency below 5% resulting in 17,224

SNPs.

We explored the adaptive genetic variability of *Populus trichocarpa* dataset

using both PCA-based (*pcadapt*) and RDA-based method using the 21 environmental

variables as explanatory variables (Table S2). We finally compared the list of outliers

found with these two ordination approaches to the ones found using *Fst* based

methods (Geraldes *et al.*, 2014).

We then performed a second RDA with only the loci previously found as

outliers (q.values < 0.1). This set of outliers gives an "adaptively enriched genetic

space" (Steane et al., 2014) and a second RDA on these specific loci allows to identify

the environmental variables that are the most correlated with "purely" adaptive

variation. In the same idea, we used the scores of the different *Populus trichocarpa*

populations along the first two RDA axes to build composite indexes that predict the

performance of individuals in the environment. The population's scores along one

axe of the adaptively enriched genetic RDA can be interpreted as a cursor of

population genetic adaptation to the environmental variables associated with this

axis. A map representation of these indexes allows for a better understanding of the

adaptive landscape in the sampling area.

RESULTS

1.   Genome scan on simulated dataset using RDA or PCA based methods

The procedure was first performed on one simulation (sim1) with both PCA

and RDA based approaches. To perform RDA, we considered environmental

variables 1-10 as explanatory variables. For the two approaches, we retained the

first four ordination axes to compute Mahalanobis distances as indicated by the

scree plots (Fig. S1).

The software package *pcadapt* is successful at detecting QTL2 SNPs (Fig. 2A)

but fails entirely at detecting QTL1 and QTL3 SNPs. The RDA-based approach also

detects QTL2 SNPS but also detects most of the QTL1 SNPs (Power=75%) by

contrast to *pcadapt* (Fig. 2B). For the SNPs associated with QTL3, both methods have

negligible power to detect them (PCA Power=0%, RDA Power=25%).

When looking at the projection of each SNP in the RDA space we can detect

links between QTL SNPs and environmental variables (Fig. 3). RDA1 is strongly

correlated to env2 and env5 and the QTL2 SNP show extreme scores on this axis

comparing to the other SNPs, especially the neutral ones (Fig. 3A). RDA2 is also

associated to env7, env8 and env9, which had not been used as direct selective

drivers during the simulations. RDA3 is correlated to env1 and env4 corresponding

to QTL1 SNPs and there is also a weak association between RDA4 and the SNPs

associated with QTL3 (Fig. 3B & 3C).

When looking at the percentage of environmental variance explained by the

RDA and PCA axes we find that the RDA first four axes explain more variance of each

environmental variable than the first four axes of the PCA (Fig. 4). The difference is

especially true for env1 and env4, which are strongly associated with the first four

axes of the RDA whereas PCA axes show an almost null correlation. The variables

env7, env8, env9 and env10 are also correlated to the first fourth axes of the RDA

whereas PCs show far less correlation with these environments. Finally, both RDA

axes and PCs fail to correlate with the complete variability of env3 and env6 even if

RDA (R^2 = ~0.25) is slightly more efficient than PCA (R^2 = ~0). Furthermore, we can

notice in Fig. S2 that each RDA axis, from the first one to the fourth, is associated to

one or several environmental variables whereas only one PC (PC1) is associated to

env2 and env5.

Finally, over the 100 simulations, we have measured the average FDR and

power for both *pcadapt* and RDA (Fig. 5). The two methods identify correctly almost

all the QTL2 SNPs (Fig. 5A). RDA-based analysis is more powerful than *pcadapt* to

identify QTL1 SNPs with respectively 75% and less than 10 % of true identification.

And for QTL3, *pcadapt* is not able to find any of the SNPs under selection whereas

RDA reaches 24 % of true identification. Both method properly control for the false

discovery rate because the proportion of false discoveries obtained with *pcadapt*

and the RDA-based method are respectively of 5% and 2.3% when controlling for a

nominal error rate of 5%

2.   *Populus trichocarpa*

For the RDA and *pcadapt* analysis on *P. trichocarpa* dataset, we retained

respectively $K=3$ and $K=6$ axes as indicated by the screeplots (Fig. S2). Analysis with

a FDR of 0.1 leads to a list of 151 outliers for RDA and 117 for *pcadapt*. Interestingly,

both methods have 29 outliers in common so that 122 and 88 SNPs are outliers

specific to respectively RDA and *pcadapt*.

When we compared RDA genome scan results with the outliers found by

Geraldes *et al.* (2014) based on Fdist, bayescan and bayenv methods, we found a

substantial overlap between them (Fig. 6). Among the 151 outliers found with the

RDA-based method, 69 have already been picked up by *Fst*-based methods (see

Table S3). Interestingly, the loci showing the lowest p-values with the RDA analysis

are also pointed out as outliers by *Fst* and *pcadapt* methods (Fig. 6). However, RDA

based genome scan found 72 SNPs that no other methods including *pcadapt* had

detected as outlier.

When we performed a new RDA analysis on the set of 151 outliers loci as

detected by RDA based genome scan method (adaptive enriched genetic space), we

found that the two first axes keep most of the adaptive genetic variance among the

populations (Fig. 7A). These two axes correspond to the two principal selective

pressures in the sampling area. RDA1 is correlated with several environmental

variables used in the analysis (Fig. 7A) including DD_0, DD18, MAT, MCMT, EMT,

which correspond to temperature indexes. RDA1 is also correlated to moisture

variables (e. g. AHM or MAP), continentality (TD) and flowering period length (bFFP

and eFFP). This axis could summarize a gradient of global climatic variation

encountered in the sampling area with southern and coastal populations opposed to

northern and continental populations (Fig. 7B).   On the other side RDA2 also

correlates with moisture and temperature variables such as AHM, MAP, TD, MCMT,

eFFP and contrasts individuals from coastal locations to individuals from inland

areas (Fig. 7C).

DISCUSSION

Genetic differentiation approaches, as implemented in *pcadapt*, detect true

positives when environmental gradients related to adaption are correlated to

population structure. For instance, QTL2 SNPs are adaptive along an environmental

gradient that is correlated with population structure (see Fig. 1) and they are

detected by *pcadapt* (Fig. 5). By contrast, when the environmental gradient is not

correlated to population structure (QTL3 and QTL1 to a lesser extent) it fails to

detect adaptive SNPs. In this case, PCA ordination is not able to orientate genetic

variation into the direction of environment 1 or 3. Figure 6 shows that RDA has a

larger statistical power than *pcadapt* to detect QTL1 SNPs by taking advantage of

information from environmental local conditions. It can be attributed to a better

alignment between the genetic space and the environmental variable improving the

power to detect true positives. This better alignment is easily visible when looking at

the PCA and RDA correlation with environmental variables (Fig. 4 & Fig. S2). In these

figures we notice that PCA is only capable of identify the preponderant genetic

variance correlated with geographic distance among populations (PC1 and PC2)

whereas RDA is able to also capture the genetic variability associated to other

environmental variables such as env1 (RDA3) and to a lesser extent env3 (RDA4).

However, RDA fails to detect QTL3 outliers in 75 % of the simulations (Fig. 5)

possibly because of the additional difficulty of detecting adaptive SNPs on a coarse

environment compared to a smooth environmental gradient as environment 1 and 2.

RDA-based genome scan relies on linear association between environment and

allelic abundance (0, 1 or 2). Patchy selective pressure like env3 is difficult to

identify with correlative approach. Forester and colleagues (2015) already found

that heterogeneous environmental selection decrease the capacity of RDA to identify

the loci under selection but they also showed that RDA remains the best method

among the ones they tested. The strength of selection and the level of allelic

dispersion during the simulations strongly influence the statistical power of RDA;

weak selective strength and strong dispersion being the less favourable conditions

(Forester *et al.*, 2015). Nonetheless, our simulations plead in favour of using

constrained ordination method when relevant environmental variable are available

in order to orientate ordination axis in the direction of environmental gradients.

This result concords with a very recent study of Forester and colleagues (2017) who

conclude that constrained ordination shows a superior combination of low false

positive and high true positives rates across all level of selection.

During our RDA-based procedure we use Mahalanobis distances as a test

statistic to identify genetic variation significantly associated with environmental

gradients. This distance is computed based on the scores of more than one RDA axis

and can thus condense in one statistic the information of several RDA axes. Some

other studies have used different RDA-based statistics to identify the loci under

putative selection. They often used a statistic estimated independently for the

different RDA axes such as the loadings of the loci along the axis (e. g. extreme

squared loadings for Latsky *et al.*, 2012) and considering a threshold such as ± 3 SD

(Forester *et al*, 2017). Mahalanobis distances can capture complex selection patterns

where allelic variation depends on more than one environmental gradient. This

statistic is appropriate to identify selective pressures on natural populations, which

usually encompass multi-factorial gradients of selection (Salmela 2014).

One application of ordination methods is to detect selective gradients. The

RDA-based approach succeeded very well to decompose the QTL SNPs-

environments associations along the different axes, at least in our simulations. In our

example, we succeeded to dissociate the outliers linked to the different

environmental variables and thus to precisely identify which set of variables drives

each part of the adaptive genetic variability (Fig. 3). RDA1 is strongly associated with

envir2, RDA2 with envir7 (collinear with isolation by distance pattern in the grid),

RDA3 with envir1 and RDA4 with envir3. As expected, the correlated environments

are also strongly associated with this respective axes. This is reflecting the fact that it

is difficult to distinguish among several collinear variables which environmental

variable has a causal effect on the individual fitness. However, it is often sufficient to

identify combination of environment variables having a strong association with

adaptive variation without knowing precisely the underlying mechanical process.

These simulations serve as a proof of concept to show that ordination axes can

provide holistic measures of genomic adaptation (Steane *et al.,* 2014). Under this

idea, scores of individuals on one RDA axis reflect their degree of adaptation along

this axis. It allows to produce useful spatial representations of the adaptive

constraint in the landscape (see Fig. 7) and to visualize a gradient of genetic

adaptation.

From the analysis of *Populus trichocarpa*, RDA found numerous outlier loci

(72) that are not detected by any other genome scan method, including *pcadapt* (Fig.

7 and table S2). These results support the simulation analysis conclusions that RDA

based genome scan is able to pick up more genes in relation to environmental

gradients than *pcadapt*. It is however noticeable that a proportion of outliers are

shared between PCA and RDA based methods (29), highlighting the fact that both

approaches are partially similar. The lack of interpretability of PCA method is

compensated by the fact that it is a blind method in regard to environmental data so

it can pick up genes that we will miss using RDA because some crucial

environmental data are lacking.

After focusing the RDA analysis on the loci showing significant association

with environmental variables (151 outliers), we identified the predominant

environmental gradients structuring the genetic adaptation of *Populus trichocarpa.*

The main one is a global climatic variation (RDA1 in Fig. 7), gathering precipitation,

temperature and seasonality variations. It clearly contrasts the populations coming

from the extreme northern part of the sampling (Alaska) and the population coming

from the inland locations (higher altitudes) to the populations close to the coast in

British Columbia (Canada) and the southern populations, near Portland (USA). This

results concord with Geraldes et al. (2014) conclusions. In their study they point out

the need of adaptation to seasonality, photoperiod and frost events for *Populus*

trichocarpa along its continuous distribution from 44 to almost 60 degree of latitude.

RDA2 is associated to moisture regimes and points out a gradient of adaptation

between coastal locations, strongly watered, and inland populations, receiving less

precipitation (Fig. 7). These two environmental gradients are the strongest selective

pressures driving *Populus trichocarpa* climatic adaptation in the North west of

America, considering the set of environmental variables available and the scale of

the analysis. To go further, these composite indexes could also serve to predict (i)

provenances that would perform well in common gardens, (ii) patterns of

adaptation to local climate across the geographic range or (iii) future adaptive

landscape in a context of climate change.

ACKNOWLEDGEMENTS

We would like to dedicate this paper to the memory of our colleague and

beloved friend Eric Bazin (1977-2017) who passed away too soon.

TC acknowledges support from the ANR project APPATS (ANR-15-CE02-

0004) and MB acknowledges Grenoble Alpes Data Institute, which is supported by

the French National Research Agency under the "Investissements d'avenir" program

(ANR-15-IDEX-02).

REFERENCES

Bazin, Eric, Kevin J. Dawson, and Mark A. Beaumont. 2010. "Likelihood-Free Inference of
Population Structure and Local Adaptation in a Bayesian Hierarchical Model." *Genetics* 185
(2): 587–602.

Coop, Graham, David Witonsky, Anna Di Rienzo, and Jonathan K. Pritchard. 2010. "Using
Environmental Correlations to Identify Loci Underlying Local Adaptation." *Genetics* 185
(4): 1411–23.

Dabney, Alan, and John Storey. 2011. "Bioconductor's Qvalue Package." *Gene*, no. 2001: 1–6.

Kort, Hanne De, Katrien Vandepitte, Hans Henrik Bruun, Déborah Closset-Kopp, Olivier
Honnay, and Joachim Mergeay. 2014. "Landscape Genomics and a Common Garden Trial
Reveal Adaptive Differentiation to Temperature across Europe in the Tree Species Alnus
Glutinosa." *Molecular Ecology* 23 (19): 4709–21.

Villemereuil, Pierre de, and Oscar E. Gaggiotti. 2015. "A New FST-Based Method to Uncover
Local Adaptation Using Environmental Variables." *Methods in Ecology and Evolution* 6
(11): 1248–58. doi:10.1111/2041-210X.12418.

Duforet-Frebourg, Nicolas, Keurcien Luu, Guillaume Laval, Eric Bazin, and Michael G B Blum.
2016. "Detecting Genomic Signatures of Natural Selection with Principal Component
Analysis: Application to the 1000 Genomes Data." *Molecular Biology and Evolution* 33 (4):
1082–93.

Foll, Matthieu, and Oscar Gaggiotti. 2008. "A Genome-Scan Method to Identify Selected Loci
Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective."
Genetics 180 (2): 977–93.

Forester, Brenna R, Jesse R Lasky, Helene H Wagner, and Dean L Urban. 2017. "Comparing
Methods for Detecting Multilocus Adaptation with Multivariate Genotype," no. 970: 1–42.

Forester, Brenna R., Matthew R. Jones, Stéphane Joost, Erin L. Landguth, and Jesse R. Lasky.
2016. "Detecting Spatial Genetic Signatures of Local Adaptation in Heterogeneous
Landscapes." *Molecular Ecology* 25 (1): 104–20.

Frichot, Eric, Sean D. Schoville, Guillaume Bouchard, and Olivier François. 2013. "Testing for
Associations between Loci and Environmental Gradients Using Latent Factor Mixed
Models." *Molecular Biology and Evolution* 30 (7): 1687–99.

Geraldes, A., S. P. Difazio, G. T. Slavov, P. Ranjan, W. Muchero, J. Hannemann, L. E. Gunter, et
al. 2013. "A 34K SNP Genotyping Array for Populus Trichocarpa: Design, Application to
the Study of Natural Populations and Transferability to Other Populus Species." *Molecular
Ecology Resources* 13 (2): 306–23.

Geraldes, Armando, Nima Farzaneh, Christopher J. Grassa, Athena D. Mckown, Robert D. Guy,
Shawn D. Mansfield, Carl J. Douglas, and Quentin C B Cronk. 2014. "Landscape Genomics
of Populus Trichocarpa: The Role of Hybridization, Limited Gene Flow, and Natural
Selection in Shaping Patterns of Population Structure." *Evolution* 68 (11): 3260–80.

Hecht, Benjamin C., Andrew P. Matala, Jon E. Hess, and Shawn R. Narum. 2015.
"Environmental Adaptation in Chinook Salmon (Oncorhynchus Tshawytscha) throughout
Their North American Range." *Molecular Ecology* 24 (22): 5573–95.

Lasky, Jesse R., David L. Des Marais, John K. McKay, James H. Richards, Thomas E. Juenger,
and Timothy H. Keitt. 2012. "Characterizing Genomic Variation of Arabidopsis Thaliana:
The Roles of Geography and Climate." *Molecular Ecology* 21 (22): 5512–29.

Luu, Keurcien, Eric Bazin, and Michael G B Blum. 2017. "Pcadapt: An R Package to Perform
Genome Scans for Selection Based on Principal Component Analysis." *Molecular Ecology
Resources* 17 (1): 67–77.

Martins, Helena, Kevin Caye, Keurcien Luu, Michael G.B. Blum, and Olivier François. 2016.
"Identifying Outlier Loci in Admixed and in Continuous Populations Using Ancestral
Population Differentiation Statistics." *Molecular Ecology* 25 (20): 5029–42.

Oksanen, Jari, F Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan
Mcglinn, Peter R Minchin, et al. 2015. "Vegan: Community Ecology Package." *R Package
Version 2.3-2*, https://CRAN.R-project.org/package=vegan.

Peng, Bo, and Marek Kimmel. 2005. "simuPOP: A Forward-Time Population Genetics
Simulation Environment." *Bioinformatics* 21 (18): 3686–87.

Pritchard, Jonathan K., and Anna Di Rienzo. 2010. "Adaptation - Not by Sweeps Alone." *Nature
Reviews Genetics* 11 (10). Nature Publishing Group: 665–67.

Salmela, Matti J. 2014. "Rethinking Local Adaptation: Mind the Environment!" *Forest Ecology*
and Management 312. Elsevier B.V.:271–81.

Savolainen, Outi, Martin Lascoux, and Juha Merilä. 2013. "Ecological Genomics of Local
Adaptation." *Nature Reviews Genetics* 14 (11): 807–20.

Steane, Dorothy A., Brad M. Potts, Elizabeth McLean, Suzanne M. Prober, William D. Stock,
René E. Vaillancourt, and Margaret Byrne. 2014. "Genome-Wide Scans Detect Adaptation
to Aridity in a Widespread Forest Tree Species." *Molecular Ecology* 23 (10): 2500–2513.

Vangestel, Carl, Andrew J. Eckert, Jill L. Wegrzyn, J. Bradley St. Clair, and David B. Neale.
2018. "Linking Phenotype, Genotype and Environment to Unravel Genetic Components
Underlying Cold Hardiness in Coastal Douglas-Fir (Pseudotsuga Menziesii Var.
Menziesii)." *Tree Genetics & Genomes* 14 (1). Tree Genetics & Genomes:10.

Vatsiou, Alexandra I., Eric Bazin, and Oscar E. Gaggiotti. 2016. "Detection of Selective Sweeps
in Structured Populations: A Comparison of Recent Methods." *Molecular Ecology* 25 (1):
89–103.

Wang, Jiahui, Ruben Zamar, Alfio Marazzi, Victor Yohai, Matias Salibian-Barrera, Ricardo
Maronna, Eric Zivot, et al. 2014. "Robust: Robust Library." *R Package Version 0.4-16*.
http://cran.r-project.org/package=robust.

Whitlock, Michael C., and Katie E. Lotterhos. 2015. "Reliable Detection of Loci Responsible for
Local Adaptation: Inference of a Null Model through Trimming the Distribution of F_{ST}."
The American Naturalist 186 (S1): S24–36.

DATA ACCESSIBILITY STATEMENT

484 • Simulations datasets are available at Dryad Digital Repository at the following
485   address: https://doi.org/10.5061/dryad.1s7v5
486
487 • Scripts used for simulations and *Populus trichocarpa* analysis are available on
488   Github: https://github.com/Capblancq/RDA-genome-scan
489
490 • *Populus trichocarpa* SNP data are available on Dryad
491   (http://datadryad.org/resource/doi:10.5061/dryad.1051d)

AUTHOR CONTRIBUTIONS

494   E. Bazin designed the study. E. B. and T. Capblancq performed the analysis and

495   treatments. M. Blum, E. B. and T. C. wrote the manuscript, and all authors

496   contributed substantially to revisions.

TABLES AND FIGURES

Figure 1: Graphical representations of mean environmental value for environment 1

to 10. Equation determining the environment is given in the main manuscript. Mean

environmental value for environment 4, 5 and 6 are respectively equivalent to

environments 1, 2 and 3.

Figure 2: Manhattan plots of the *pcadapt* (A) and RDA (B) results.

Figure 3: Projection of SNPs and environmental variables into the RDA space.

Figure 4: Proportion of environmental variance explained by the first four

ordination axis using RDA or PCA. The values correspond to the r-squared of the

regression between the environmental variable and the first four axes of the

multivariate analysis.

Figure 5: Performance results of RDA and *pcadapt* methods. Each performance

value is averaged over 100 simulated dataset (error bars are displayed). Power (A)

is given separately for loci coding for quantitative traits 1, 2 and 3. False Discovery

Rate (FDR) is given for the two methods (B).

Figure 6: Scatter plot of the p-values returned by RDA and *pcadapt* procedure for

the *Populus trichocarpa* dataset. Black dots correspond to loci identified as outliers

by *Fst* methods in Geraldes et al. (2014).

Figure 7: Projection of SNPs and environmental variables in the RDA performed on
the adaptively enriched genetic space (151 loci) of *P. trichocarpa* dataset (A) and
spatial representation of the population average scores on RDA1 (B) and RDA2 (C).
519

SUPPORTING INFORMATION

Script S1: Detailed script of the RDA-based genome scan analysis.

Table S1: Environmental variables used in the *Populus trichocarpa* analysis.

Table S2: Comparison between RDA, *pcadapt*, *fdist* and *bayescan* outliers loci.

Figure S1: Scree plot for PCA and RDA analysis of simulation 1 dataset

Figure S2: Heatmap of the R^2 resulting from linear regression between the fourth
first axes of PCA and RDA analysis and the environmental variables.

Figure S3: Scree plot for PCA and RDA analysis of *Populus trichocarpa* dataset.

envir1

envir2

envir3

envir4

envir5

envir6

Z

1
0
−1
−2

envir7

envir8

envir9

envir10