Original Article

Sent to the Corner: xylem vessel anatomy not surface area determines megaphyll hydraulic

efficiency in *Cecropia obtusa* Trécul (Urticaceae)

Running title: Xylem vessel anatomy not surface area determines megaphyll hydraulic

efficiency

Sébastien Levionnois[1*], Sabrina Coste[2], Eric Nicolini[3], Clément Stahl[4], Hélène Morel[5],

Patrick Heuret[4*]

1. CNRS, UMR EcoFoG (AgroParisTech, CIRAD, CNRS, INRA, UA, UG), 97379 Kourou

Cedex France.

2. UG, UMR EcoFoG (AgroParisTech, CIRAD, CNRS, INRA, UA, UG), 97379 Kourou

Cedex France.

3. CIRAD, UMR AMAP (CIRAD, CNRS, INRA, IRD, Université Montpellier II), 34398

Montpellier, France.

4. INRA, UMR EcoFoG (AgroParisTech, CIRAD, CNRS, INRA, UA, UG), 97379 Kourou

Cedex France.

5. CIRAD, UMR EcoFoG (AgroParisTech, CIRAD, CNRS, INRA, UA, UG), 97379 Kourou,

France.

*Authors for correspondence:

Sébastien Levionnois, PhD student

Postal address: UMR EcoFog, Campus Agronomique, Avenue de France

BP709 – 97387 Kourou Cedex

French Guiana

Tel: +33 760 26 73 83

Email: sebastien_levionnois@live.fr

Patrick Heuret, PhD

Postal adress: UMR EcoFog, Campus Agronomique, Avenue de France

BP709 – 97387 Kourou Cedex

French Guiana

Tel: +33 694 38 12 19

Email: patrick.heuret@inra.fr

Footnotes

[1]Manuscript received _______; revision accepted _______

Author contributions

P.H. designed and lead the project; P.H., S.L., E.N., S.C., H.M. and C.S. collected field

samples and measured morphological traits; S.L. realised the anatomical cuts and images

analysis; S.L. and P.H performed data analysis; S.L. and P.H. wrote the paper; all authors

discussed   the   results   and   commented   on.

Abstract

Corner's rule predicts a positive correlation between leaf dimensions and the cross-sectional

area of the primary stem. Although this relationship is usually explained by hydraulic and

mechanical requirements, these hypotheses have never been tested empirically. However,

Corner's rule is tricky to investigate since rapid secondary growth of the stem prevents a

rigorous link being established between a given leaf and the supporting stem. We chose a

twig-like leaf model since petiole anatomy is only linked to the attached lamina. We tested the

hypothesis that anatomical adjustments to hydraulic requirements related to vessel size enable

reduced investment in tissue in the framework of Corner's rule. We conducted a functional,

mechanistic and intraspecific investigation of *Cecropia obtusa* Trécul, a Neotropical pioneer

tree, by integrating morphological, anatomical and theoretical hydraulic traits around the

lamina-petiole size relationship. The twig-like structure of the leaf and the strong lamina-

petiole correlation of this model tree species made it possible to use the leaf-level model for a

rigorous investigation of the functional implications of Corner's rule. We found a positive

correlation between petiole size, lamina size, the ratio of mean vessel area to vessel frequency

in the petiole xylem and theoretical specific conductivity in the petiole xylem. Hydraulic

function supports Corner's rule to a lesser extent than previously thought. Variations in vessel

dimensions mainly drive xylem hydraulic performances and avoid disproportionate petiole

cross-sections to answer to hydraulic requirements associated with lamina size.

Key words: allometry, *Cecropia obtusa*, Corner's rules, hydraulic, intraspecific, leaf-stem

relationship, petiole anatomy, lamina-petiole relationship

INTRODUCTION

Leaves have received a great deal of attention because of the major role they play in the plant carbon economy thanks to photosynthesis. At the interspecific level, the currently widely accepted paradigm relates to a global spectrum of leaf economy, running from quick to slow returns on investments (Wright et al. 2004). Leaf surface area encompasses six orders of magnitude (Niinemets et al. 2007, Milla and Reich 2007, Wright et al. 2017). One possible and pertinent way for improving our understanding of such variability is to integrate the stem and to look at the size and functional relationship between stem and leaf.

From an evolutionary perspective, in his seminal paper on "Durian Theory" E.J.H. Corner already made empirical observations on such relationship, as "The stouter the main stem, the bigger the leaves and the more complicated their form" (Corner 1949). He completed and extended this with a second point he called "diminution on ramification" as "The greater the ramification, the smaller become the branches and their appendages". These principles were referred to as Corner's rules by Hallé et al. (Hallé et al. 1978) and have been explored by several authors (White 1983a, 1983b, Brouat et al. 1998, Brouat and McKey 2001, Westoby and Wright 2003, Preston and Ackerly 2003, Sun et al. 2006, Normand et al. 2008, Liu et al. 2010). However, from a more developmental point of view, even before Corner, by drawing attention to the pith, E.W. Sinnott (Sinnott 1921) suggested that the size of organs (the stem, leaf, fruit) is correlated with the size of the meristem from which they develop. Back to the contemporary research, plenty of studies have investigated both stem and leaf in their particular size and functional relationships (White 1983a, 1983b, Brouat et al. 1998, Ackerly and Donoghue 1998, Brouat and McKey 2001, Westoby and Wright 2003, Preston and Ackerly 2003, Sun et al. 2006, Kleiman and Aarssen 2007, Normand et al. 2008, Yang et al. 2008, 2008, Tao et al. 2009, Olson et al. 2009, Milla 2009, Liu et al. 2010, Whitman and Aarssen 2010, Scott and Aarssen 2012, Dombroskie and Aarssen 2012, Osada et al. 2015, Huang et al. 2016, Fan et al. 2017), usually (but not always) referring to Corner's rule. Since a unifying framework is lacking, and based on the Corner's vision, we will refer in this study to the leaf-stem relationship (LSR), that we defined as (i) the spectrum of size and functional variation in the relationship, (ii) between a leaf and its associated internode, or between leaves and the stem of a first-year shoot, and (iii) across or within species.

Both at the intra- and interspecific level, the LSR is frequently and intuitively explained by functional requirements in terms of hydraulic and mechanical supplies (Hallé et al. 1978, White 1983a, 1983b, Brouat et al. 1998, Westoby and Wright 2003, Normand et al. 2008). Since the respective contributions of the hydraulic and mechanical functions have not

yet been tested, there is a need to (i) explore the diversity of anatomical structures that lead to

interspecific variation in the LSR and (ii) understand the anatomical adjustments that allow

for variations in the size of the stem and the leaf at the intraspecific level (Lehnebach et al.).

The aim of this study was thus to investigate the intraspecific mechanisms of the LSR from

the point of view of hydraulics and the aforementioned carbon allocation issues.

In angiosperms, xylem vessels drive water conduction through the soil-plant-

atmosphere continuum (Tyree and Zimmermann 2002) according to the cohesion-tension

theory (Dixon and Joly 1895, Steudle 2001). Vessel dimensions vary considerably at

interspecific level (Chave et al. 2009, Zanne et al. 2010, Olson and Rosell 2013), at

intraspecific level (Lens et al. 2011, Hajek et al. 2016) and at the intra-individual level

(Anfodillo et al. 2006, 2013, Bettiati et al. 2012, Schuldt et al. 2013), with critical

consequences for hydraulic performance at the organ, individual, population or species levels

(Hacke et al. 2016). According to the Hagen-Poiseuille law (Tyree and Zimmermann 2002),

vessel hydraulic conductivity is a fourth-power function of the vessel diameter. Thus, even

slight variations in vessel diameter have dramatic impacts on associated conductance. A

greater or lesser water demand with respect to the leaf lamina surface can theoretically be

satisfied either by increasing the surface area of the conductive tissues which supply it, or by

modifying the conductivity of the related tissues by means of their anatomy. Therefore, we

hypothesise that adjustments to stem vessel size first drive size-related hydraulic supplies,

thereby limiting carbon increments of xylem surface with an increase in leaf size.

Consequently, even if stem xylem surface area is positively correlated with leaf size, the

hydraulic function explains the LSR to a lesser extent than previously thought.

However, focusing on the leaf-stem system is hampered by several technical

limitations. One of the main difficulties in applying a rigorous approach to the LSR is

defining where to sample the last elongated shoot. On one hand, Corner (Corner 1949) refers

to primary diameter (Hallé et al., 1978), which means that no secondary growth has occurred.

Secondary growth occurs very soon, possibly at the same time as the elongation process,

meaning that LSR may vary over time. On the other hand, the internode of a first-year shoot

supplies not only the leaf at its apex, but all the leaves supported by internodes distal to it as

well as the bud content (Cochard et al. 2005). It would thus be more relevant to analyse the

relationship between the total surface area of leaves borne by the first-year shoot and the

cross-sectional area of the shoot (Brouat et al. 1998). A compromise for a rigorous functional

assessment would be to shift to a leaf-level model (i.e. considering the petiole to be

homologous to the stem and its associated lamina), since there are plant groups in which the

differences between leaf and shoot, or more precisely between the petiole and the stem remain
fuzzy (Givnish 1984, Poorter and Rozendaal 2008, Nicotra et al. 2011). Indeed, compound or
double-compound leaves, or large palmate or palmatilobate leaves look like more to a branch
in a physiognomic or functional sense. For these species, the whole bifurcation structure of
the tree for foraging space is realised by leaves and stems together. A major symptom of the
transfer of the bifurcation function from stems to leaves is less stem branching intensity in
comparison with simple leaved tree species (Aarssen 2012). Developmental genetics
approaches also strengthen our posture since numerous studies suggest that the petiole should
be considered separately from the leaf lamina, from morphogenesis and developmental
genetic points of view, and more as an stem-like axial organ (Foster 1936, Tsukaya et al.
1993, 1995, 2002, Tsukaya 1995, 2013, Busch et al. 2011).

143   Using a leaf model to investigate the LSR makes it possible to avoid the bias
frequently encountered at the phytomer level. Indeed (i) a petiole strictly supports mechanical
and hydraulic constraints associated with its single lamina since it represents the closest
structure to the transpiring compartments (Bucci et al. 2003) and (ii) even if a petiole
undergoes secondary growth, the growth still only concerns the functional requirements of the
associated lamina. Consequently, using a leaf model would make it possible to focus on
petiole or venation traits, for instance, and to rigorously investigate the functional adjustments
underlying the LSR. Finally, the lamina-petiole relationship has already been proven to be a
relevant tool for functional investigations, and, for instance, (Royer et al. 2007) showed that
this relationship is a significant predictor for inferring fossil functional traits.

153   In this context, the genus *Cecropia* (Urticaceae), an iconic pioneer tree in the
Neotropics, could provide several consistent tree models to improve our understanding of
structure-function relationships. These plants have a simple architecture (Heuret et al. 2002,
Zalamea et al. 2008) with few botanical entities (e.g. axes, leaves), but have large petioles,
reaching more than 1 m in length and up to 2 cm in diameter (Fig. 1b). Linked to its heliophile
habit, petioles have a somewhat branching function (Givnish 1984, Poorter and Rozendaal
2008) and can be considered as axes, since the hydraulic and mechanical features of petioles
are comparable to a branch with respect to the form of the leaf (palmatilobate to palmate
megaphylls with numerous lobes tens of centimetres in length) and its size (up to 0.4 m² for
Cecropia obtusa Trécul with 1-m long petiole).

163   Here, we provide a functional assessment of intraspecific LSR in *Cecropia obtusa* by
focusing on the lamina-petiole relationship in the megaphyll and its hydraulic implications.
First, we report an extensive investigation of the relationships between leaf-level traits to

validate the lamina-petiole relationship. We hypothesise that, at this level, correlation
properties are comparable to the LSR, thereby enabling further functional assessment.
Second, after providing the first description of the anatomy of the petiole, we focus on the
lamina-petiole relationship by integrating, according to leaf size (i) the surface areas and
proportions of the different component tissues of the petiole, (ii) structural anatomical traits
such as the number of xylem vessels, their hydraulic-weighted dimensions and their
frequency, and (iii) theoretical hydraulic traits (derived from the Hagen-Poiseuille law) such
as hydraulic conductivity and xylem- or leaf-specific hydraulic conductivity.

MATERIALS AND METHODS
Study site
This study was conducted in French Guiana along a forestry road in Counami (N5.41430°,
W53.17547°, geodesic system WGS84) where the entrance to the road is located 5 km to the
east of Iracoubo municipality. The warm wet tropical climate of French Guiana is highly
seasonal due to the north-south movement of the Inter-Tropical Convergence Zone. Annual
rainfall is 3041 mm year^{-1} and annual mean air temperature is 25.7 °C at Paracou
experimental station (Gourlet-Fleury et al. 2004). There is one long dry season lasting from
mid-August to mid-November, during which rainfall is < 100 mm month^{-1}.

Plant material, sampling and morphological measurements
In order to explore a wide range of leaf sizes, we studied 25 adult trees (Fig. 1a) and four
juvenile trees. Juvenile trees were less than 2.2 m in height and by counting the number of
nodes along the trunk (Heuret et al. 2002, Zalamea et al. 2012), their age was estimated at
between 8 and 13 months. The dimensions of adult trees ranged from 10.9 to 22.6 m in height,
6.21 to 30.49 cm in diameter at breast height, and the trees were from 7 to 21 years old.
Juvenile trees were studied in September 2014. Adult trees were felled regularly from
September 2014 to October 2016, between 0800 am and 1030 pm. After felling, all the axes
of the crown were tagged according to their position in the tree architecture and all the leaves
were tagged according to which axis they belonged, and their rank from the apex. All the
leaves in the crown were cut by sectioning the petiole as close as possible to its insertion point
on the stem and immediately placed in plastic bags and then in coolers. One to five leaves per
tree -hereafter called P-leaves- just fully expanded and positioned on the 3rd or 4th nodes
located under any apex, were put aside for the anatomical studies.

To enable a full understanding of the study, here we recall certain plant morphology

prerequisites. *Cecropia* leaves are palmatilobate (Fig. 1b and 2). Consequently, there is no

single midrib but as many midribs as there are lobes, i.e. a midrib departs from each lobe. We

use the term lamina to refer to the flat thin photosynthetic part of the leaf. We use the term

petiole to refer to the physical connection between the lamina and the stem. Finally, we use

the term leaf to refer to the global structure composed of a petiole and a lamina. In the

laboratory, all the leaves were processed at the same time of day to keep them as fresh as

possible. The petioles were cut as close to the lamina as possible. The length of the petiole

(L_{pet}, cm) and two median and orthogonal diameters (mm) of each petiole were measured

(Fig. 2a). From these two diameters, a cross-sectional area was derived in the shape of an

ellipse (A_{pet}, mm²; Fig. 2a). For each lamina, the length of the main lobe (i.e. the largest one

in the continuation of the petiole; LM_{lobe}, cm) was measured and the number of lobes (N_{lobe})

counted (Fig. 2b, c). Exactly 2.851-cm² of lamina in the form of three tablets were sampled

with a circular iron punch (diameter = 1.1 cm) taking care to avoid main nerves (Fig. 2d). In

the P-leaves, two 1-cm-long petiole segments in the median position were sampled and

weighed (Fig. 2e, f). The first segment was used to measure their specific density, expressed

as the ratio of dry mass to the fresh volume (PSG: Petiole Specific Gravity; (Williamson and

Wiemann 2010); Table 1). The fresh volume of the sample was calculated using an inverse

Archimedes principle and a precision balance (CP224S, Sartorius), as the difference between

the fresh mass and immersed mass of the sample. Dry mass was derived after drying at 103

°C for three days. The second segment was preserved in 70° alcohol and later used for

anatomical measurements (see the section headed 'Anatomical and hydraulic measurements').

Lamina area (A_{lamina}, cm²; Fig. 2g) was measured with a planimeter (LiCor 3000A,

LiCor Inc., Lincoln, NE, USA). The condition of the lamina (damaged or undamaged) was

described. Each leaf (leaf + petiole + 3 tablets) was dried at 70 °C for 72 hours. The dry mass

of the petiole (M_{pet}, g), the lamina (M_{lamina}, g) and the three tablets_taken together (g) was

measured with a precision scale. The lamina mass area (LMA, g cm^{-2}) was derived from the

ratio of the dry mass to the known surface area of the three tablets (Fig. 2a).

Data from 2112 mature fully-developed leaves (P and non-P-leaves) were used for

morphological validation of the LSR. When we analysed relationships, including lamina area,

we only considered leaves with undamaged lamina (n = 1072). We also investigated the

branching intensity-stem size trade-off (Osada et al. 2015), considering lobes (i.e. main

nerves) as homologues of the branching process, and by studying the ratio of the number of

lobes to the undamaged lamina area (number of lobes per lamina unit area, we refer to as

"lobing" intensity; n mm^{-2}) according to the leaf size.

Anatomical and hydraulic measurements

Because anatomical studies are time consuming, we conducted our study on a subset of 60 P-

leaves selected to represent (i) our 29 individual trees to minimise individual effects and (ii)

the range including the largest petiole diameters (from 2.59 to 15.41 mm). Anatomical

transversal cross-sections, 20 to 50 μm thick, were sampled from the petiole middle segment

with a manual microtome (Mikrot L, Schenkung Dapples, Switzerland). All cross-sections

were stained in a safranin/astra-blue solution to color unlignified cells blue and lignified cells

red.

Measuring anatomy and xylem conductivity in one point is, of course, restrictive to

understand the whole vascular architecture and the underlying hydraulic of the entire leaf

(Coomes et al. 2008). However, because the aim of this paper is to understand the anatomical

patterns underlying the SLR we choose to focus on the median part of the petiole using the

petiole-lamina model on megaphylls.

Each cross-section corresponding to a given petiole was digitised with an optical

microscope (Olympus BX60; Olympus Corporation; Tokyo; Japan) with x50 magnification

and a Canon camera EOS 500D (lens Olympus U-TVI-X ; F 0.0; ISO 100; speed 1/25). Three

or four close ups were taken of a petiole part at different depths of focus and stacked with the

Helicon Focus software (v.6.3.2.Pro, http://heliconsoft.com/). Output pictures were assembled

in a panorama using Kolor AutoPanoGiga software (v.3.0.0, http://kolor.com/autopano/) to

obtain a complete picture of the cross-section. The digitised cross-sections were processed

with CS5 Photoshop software (v.12.0, http://adobe.com/products/photoshop/html). We

distinguished eight tissue categories that comprise the petiole anatomy (see anatomical

description in the 'Results' section). We manually delineated the tissues on the photographs

and created layer masks. The masks of these layers were used to calculate the surface area of

each tissue and the whole petiole area with the ImageJ software (v.1.43u;

http://imagej.nih.gov/ij/). For the xylem, the cropped part of the image in which the vessels

were visible was also analysed with the ImageJ software to calculate theoretical xylem

hydraulic properties (Abramoff et al. 2004). For each vessel, we calculated its surface area

(μm²) and its elliptical diameters. These data were used to derive all the anatomical and

hydraulic traits listed in Table 1. To study variations in the dimensions of the vessel, we used

the mean hydraulic diameter (Dmh, Table 1), i.e. the diameter that all vessels, considered as

circles, in a given tissue would have to sustain exactly the same tissue hydraulic conductivity

(Tyree and Zimmermann 2002). The number of vessel was counted for primary and

secondary xylems and the vessel frequency calculated by dividing this number by the surface

area of the related xylem. The lumen fraction (Zanne et al. 2010) refers to the total fraction of

empty holes formed by lumens. The ratio of mean vessel area to frequency (Zanne et al. 2010)

as a vessel composition index, refers to the relative weight of vessel size versus vessel

frequency. Many vessels could be obstructed by tyloses, so we tested for any effect of leaf

size on the frequency of obstruction or the effect of obstruction on petiole conductivity.

Vessel obstruction was accounted for by removing obstructed vessels from the calculations.

Data analysis

We used a prediction model to derive estimated lamina area (A_{lamina}, cm²), since the

relationship between the length of the main lobe and the undamaged lamina area is very

informative ($R^2 = 0.942$, $P < 0.001$, Fig. 3). Indeed, laminae used in anatomical studies are

sometimes damaged (i.e. biased lamina measurement). We derived the estimated lamina area

as:

$$Estimated\ lamina\ area = 0.701 * Main\ lobe\ length^{2.180}$$

Only this measurement of lamina area was used to compare anatomical and morphological

traits.

All statistical analyses were performed with R software (http://CRAN-R-

project.org). After log transformation, the relationship between each paired trait was

determined with an SMA (Standardized major axis regression; (Warton et al. 2006)), which

allows measurement of the error on both the x-axis and y-axis (Harvey and Pagel 1991).

These correlation relationships are described as: $y = bx^a$, such as: $\log(y) = \log(b) + a * \log(x)$,

where *a* is the slope (or allometric exponent) and *b* the intercept (allometric coefficient). A

95% confidence interval was used to decide whether it is significantly correlated or not. A

slope test was performed to determine if the slope differed from 1 (H1: a $\neq$ 1 for an allometric

relationship) or not (H0: a = 1 for an isometric relationship). To avoid ambiguity when

comparing p-values and decision thresholds, "P_{cor}" refers to the correlation test, and "P_{slope}"

refers to the slope test (allo- or isometric relationship). SMA were carried out with the

(S)MATR package (Falster et al. 2006). The lamina area prediction from the main lobe length

was modeled from an NLS (non-linear least squares) with the STATS package.

A paramount *leitmotif* in our study are the scaling relationships between traits, i.e.,

either iso- or allometric. Indeed, the mathematical relationship has dramatic consequences for

data interpretation since it has major functional implications for organism development and

phylogenesis (Thompson 1917, Huxley 1932, Niklas 1994). That is why these relationships

are briefly examined here. A relationship is expressed as a power law equation, in which the

key feature is the scaling exponent characterising the form of the relationship. In biological

studies, the relationship generally refers to a size trait on the *x*-axis, whatever the level of

organization (cell, tissue, organ, individual, etc.). As the absolute value of the exponent is

equal to 1, the relationship is isometric: there is a strictly preserved proportional relationship

between the two studied traits independently of their size. In other words, size does not affect

the functional nature of the relationship. As the absolute value of the exponent is different to

1, the relationship is allometric: the proportional relationship is not preserved and the *y*-axis

variable fails to keep pace with the x-axis according to the variation in size. In other words,

size affects the functional nature of the relationship depending on whether the absolute value

of the exponent is greater or less than 1. In the case of a value greater than 1, the *y*-axis

variable increases relatively more rapidly than the *x*-axis variable. In the course of

development, the importance of the scaling exponent is such that it can be considered as a trait

and a performance condition in its own right since it exhibits its own genetic basis and genetic

variations and it is subject to strong heritability (Vasseur et al. 2012).

RESULTS

Morphological validation of the lamina-petiole size relationship

Overall, leaf traits were highly and positively correlated ($P < 0.001$; Table 2). There was a

significant and positive correlation of allometric form among most traits (A_{pet}, L_{pet}, A_{lamina},

N_{lobe}, M_{pet}, M_{lamina}, LMA; $P_{cor} < 0.001$; $P_{slope} < 0.001$; Table 2) except for (i) A_{lamina} which

was isometrically correlated with M_{lamina} ($P_{slope} = 0.210$) and (ii) LMA which was not

correlated with L_{pet} and A_{lamina} ($P_{cor} > 0.05$; Table 2). PSG was not correlated with A_{pet}, N_{lobe},

M_{pet} and M_{lamina} ($P_{cor} > 0.05$; Table 2) but was negatively and allometrically correlated with

the other measured traits (L_{pet}, A_{lamina}, N_{lobe}; $P_{cor} < 0.05$; $P_{slope} < 0.001$), except for LMA,

which was positively correlated.

The number of lobes per lamina unit area (N_{lobe}/A_{lamina}) was negatively correlated with

A_{pet} ($P_{cor} < 0.001$; Fig. 4). For a given lamina area, N_{lobes} increased with a decrease in A_{pet}.

Petiole anatomy

Petioles showed pronounced radial symmetry (Fig. 1c). The central parenchymatous pith

formed the main part of the cross-section (Fig. 1c, d, e). The numerous bundles (20 to 80)

were radially arranged in the pith periphery, mainly in only one cycle (Fig. 1c), but in a few

cases in two cycles (Fig. 1f, g). A cambium gave rise to increments of secondary xylem

internally and secondary phloem externally (Fig. 1d). The cambium showed gradual inter-

petiole diversity with respect to the arrangement of the bundles. From a strictly cyclic

structure to a wavy one, and at the extremity, we observed isolated bundles with complete

cambium discontinuities in a more cortical position (Fig. 1f, g). Primary and secondary xylem

and secondary phloem were easy to identify. Tylosis partially or totally obstructed the xylem

vessels (Fig. 1d). A sub-continuous sclerenchymatous shield was present at the interface

between the vascular bundles and the pith (Fig. 1d). The vascular bundles were separated by

interfascicular parenchyma. Occasional sclerenchyma were present between the secondary

phloem and cortical parenchyma. Depending on the extent of secondary growth, the primary

phloem was crushed between the secondary phloem and cortical parenchyma. In the most

external part, there was a ring of tangential collenchyma under the epidermis, fully contiguous

to the cortical parenchyma (Fig. 1d). Laticiferous canals were frequently visible, mainly in the

cortex (Fig. 1f) but also in the pith, but were also sometimes completely absent. We

distinguished eight component petiole tissues for further anatomical analysis (Fig. 1e): (i) the

pith, (ii) the sclerenchymatous shield associated with the vascular bundles, (iii) interfascicular

parenchyma, (iv) primary xylem, (v) secondary xylem, (vi) phloem (comprising primary and

secondary phloem), (vii) cortical parenchyma and (viii) cortical collenchyma.

Partitioning of petiole tissue

We now describe the petiole cross-sectional area (A_{pet}) both (i) as representative of the

variations in several main morphological leaf traits (A_{lamina}, M_{pet}, M_{lamina}, N_{lobe}) and (ii) as a

proxy of the total leaf size; since there are strong correlations between these traits (Fig. 3;

Table 2) and all the anatomical traits are described in relation to the petiole.

All the surface areas of the different tissues observed on the petiole cross-section were

significantly and positively correlated with A_{pet} ($P_{cor} < 0.001$; Fig. 5a; Supplementary Table

1). Most of the proportions of the different tissues in the petiole cross-section were

significantly and positively correlated with A_{pet} ($P_{cor} < 0.01$; Fig. 5b; Supplementary Table 1),

except the primary xylem area, which represented a fixed proportion of A_{pet} (4.89 ± 0.12 %;

$P_{cor} = 0.337$; Fig. 5b,c; Supplementary Table 1). Sclerenchyma, interfascicular parenchyma,

primary xylem and secondary xylem represented only small proportions of the petiole cross-

section (0.00 to 3.58%; 1.71 to 7.76%; 3.50 to 5.96%; and 2.03 to 9.21%, respectively; Fig.

5b, c; Supplementary Table 1). In contrast, pith, cortical parenchyma, phloem and

collenchyma represented significant proportions of the petiole cross-section (31.90 to 59.91%; 8.84 to 29.63%; 2.15 to 17.30%; 8.49 to 23.54%, respectively; Fig. 5b,c; Supplementary Table 1) with high variability depending on the section of the petiole concerned (Fig. 5c; Supplementary Table 1).

A mean of $22.08 \pm 1.17\%$ of tissues was calculated for conduction including $4.89 \pm 0.12\%$ of the primary xylem, $6.96 \pm 0.32\%$ of the secondary xylem and $10.23 \pm 0.99\%$ of all the phloem (Fig. 5c; Supplementary Table 1).

Anatomical and hydraulic considerations of the lamina-petiole size relationship

Hydraulic performances are regulated by three main parameters according to the size of the leaf: surface area of the xylem tissue, the number of vessels (N_{vessel}) and the mean hydraulic diameter (Dmh) (Fig. 6). These three traits were closely related ($P_{cor} < 0.001$), and significantly and positively correlated with A_{pet} ($P_{cor} < 0.001$; Fig. 6a, b; Supplementary Table 2). However, the frequency of the vessels (f_{vessel}, obtained by dividing the number of vessels by the xylem area; $n.mm^{-2}$) was negatively correlated with A_{pet} ($P_{cor} < 0.05$; Fig. 6c; Supplementary Table 2). F was independent of xylem area ($P_{cor} > 0.05$; Supplementary Table 2). S was closely correlated with A_{pet} ($P_{cor} < 0.001$; Fig. 6d; Supplementary Table 2). Specifically, even for a given fixed vessel frequency, Dmh remained positively correlated with A_{pet}.

All these coordinated anatomical adjustments affect theoretical hydraulic properties. There was a positive allometric correlation between the theoretical hydraulic conductivity (K_{th}) and A_{pet} with a slope greater than 1 ($P_{cor} < 0.001$; $P_{slope} < 0.001$; Fig. 7a; Supplementary Table 2). The same was true for A_{lamina} ($P_{cor} < 0.001$; Supplementary Table 2). Overall, for each petiole, tyloses did not significantly affect K_{th}, whether we took closed vessels into account in our calculation or not, ($P = 0.682$). There was no correlation between petiole size and the frequency of obstruction ($P = 0.636$). But there was a significant difference in vessel size between obstructed and unobstructed vessels ($P < 0.001$) but the opposite pattern was found between primary and secondary xylem. The obstructed vessels were larger in the secondary xylem but smaller in the primary xylem. There was a high positive correlation between K_s and A_{pet} ($P_{cor} < 0.001$; Fig. 7b; Supplementary Table 2): even with a given fixed xylem area, associated K_{th} increased with A_{pet}. K_{th} was also positively correlated with A_{lamina} with a slope greater than 1 ($P_{cor} < 0.001$; $P_{slope} < 0.001$; Fig 7c; Supplementary Table 2): the larger the leaf, the more supplies go to a given lamina surface area. Moreover, K_s was positively correlated with A_{lamina} ($P_{cor} < 0.001$; Supplementary Table 2).

DISCUSSION

Our study describes for the first time anatomical structures and mechanisms to better understand the nature of the first premise of Corner's rule and its underlying hydraulics and mechanical significance. Here, we discuss our three main results: (i) the shift to a leaf-level model for a rigorous functional assessment of the LSR at the intraspecific level, (ii) the pith, not the secondary xylem, as the main driver of variations in petiole size and (iii) and variations in xylem vessel anatomy, not its surface area, as the main determinant of hydraulic requirements.

Using the leaf model to extend the leaf-stem size relationship

Our results showed strong positive mainly allometric correlations between morphological traits, validating the lamina-petiole relationship as an extensive consideration of the LSR in *C. obtusa*. Most leaf traits were related, not only internode cross-sectional area and lamina area, which are the most frequently studied (White 1983a, 1983b, Brouat et al. 1998, Brouat and McKey 2001, Westoby and Wright 2003, Preston and Ackerly 2003, Sun et al. 2006, Normand et al. 2008, Liu et al. 2010). The allometric form of the relationship between A_{lamina} and A_{pet} should be interpreted with caution, in view of existing results on LSR both at intra- and interspecific levels (White 1983a, 1983b, Brouat et al. 1998, Brouat and McKey 2001, Westoby and Wright 2003, Preston and Ackerly 2003, Sun et al. 2006, Normand et al. 2008, Liu et al. 2010). We argue that confounding intra- and interspecific levels of variation is misleading since the former relates more to developmental processes and mechanisms of adjustment, while the latter relates more to a diversity of strategies, selection pressures and evolutionary pathways. In connection with this distinction, it is interesting to note that although Sinnott's work (Sinnott 1921) and the first premise of Corner's rule (Corner 1949) overlap in the global frame of the LSR, Sinnott's work is more in agreement with an intraspecific and developmental perspective while Corner's formulations concern an interspecific perspective. The latter assumption is also in agreement with a recent study by (Huang et al. 2016) in which scaling models produced contrasted results between intra- and interspecific data, due to dramatic interspecific variation in stem tissue density and leaf size. Reviewing data produced by White (White 1983a, 1983b), who was the first to test Corner's rules across species, (Brouat et al. 1998) demonstrated an isometric LSR at the intraspecific level. However, Brouat & McKey (Brouat and McKey 2001) found an allometric form related to the myrmecophytic habit, which is in agreement with our result. Note that even if we focus

on the leaf level (lamina-petiole relationship), *C. obtusa* is a hollow-stemmed myrmecophytic

tree. As discussed by Brouat & McKey (2001) in connection with the LSR, the allometric

form of our lamina-petiole relationship is linked to the pith, which accounts for the largest

proportion of the petiole cross-sectional area.

We complete this first point with the branching intensity-stem size trade-off. We

found a negative correlation between the number of lobes per lamina unit area and A_{pet}. For a

given leaf surface, the more numerous the lamina lobes, the thinner the petiole upstream from

the lamina. This result is also in agreement with Corner's rules as interpreted by (Hallé et al.

1978), as "the greater the ramification, the smaller become the branches and their

appendages" (Corner 1949). At the branched-complex level, this implies that for both fixed

foliar area and photosynthetic rates (Enquist et al. 1999), the degree of branching should be a

trade-off with the volume and/or density of the axes. In agreement with this hypothesis, we

found a weak but significant trend to decreasing PSG for larger leaves (notably with A_{lamina}, *P*

= 0.01; Table 2). This decreasing PSG was largely driven by the proportion of pith tissue (*P* <

0.001) and underlines the significance of geometry in the support function versus the decrease

in PSG in the search for lower volumetric construction costs. In the same line of thought,

(Olson et al. 2009) claimed that the negative trade-off between the volume of an axis and its

density is a corollary to Corner's rules and the LSR. This is in agreement with the recent work

of (Trueba et al. 2016) at an intraspecific level, since leaf area was shown to be a negative

trade-off with both leafing intensity and stem density according to canopy openness in a New-

Caledonian woody shrub species. We are first the show that this aspect of Corner's rules

applies to the leaf scale, considering lobes as counterparts of the branching process. Above

all, this functional dimension (branching and leafing intensities) is far from being marginally

studied (White 1983a, Ackerly and Donoghue 1998, Kleiman and Aarssen 2007, Yang et al.

2008, Tao et al. 2009, Olson et al. 2009, Milla 2009, Whitman and Aarssen 2010, Scott and

Aarssen 2012, Dombroskie and Aarssen 2012, Yan et al. 2013, Osada et al. 2015, Huang et al.

2016, Fan et al. 2017), but is far from being integrated in the leaf or global spectra of

economy (Westoby et al. 2002, Westoby and Wright 2003, Wright et al. 2004, Díaz et al.

2016), whereas it is directly linked to the optimization strategy of leaf area partitioning (Smith

et al. 2017) and hence light-interception performance (Duursma et al. 2012).

Finally, the anatomy of the petiole shows characteristics that are generally associated

with stem-level anatomy, like radial symmetry and secondary growth with wood. This is in

agreement with the axis-like function of the petiole in the architecture of *Cecropia* (Givnish

1984). In the context of our study, all these features strengthen the homology with a stem-like

branched system and the interest of shifting to a leaf model for the rigorous functional and
mechanistic assessment of the LSR, i.e. integrating anatomical and hydraulic aspects.
However, secondary growth potentially implies an alteration of observed relationships and
scaling with time, i.e. with the age of the leaf. This implies that secondary growth continues
throughout the lifespan of the leaf, which, however could result in ambiguity with respect to
secondary growth, which is generally associated with the renewal of conductive and/or
supporting potentials whereas the leaves of *C. obtusa* have a relatively short lifespan
(generally < 100 d). So does secondary growth continue throughout the lifespan or only
during the phase of primary elongation? This question deserves further investigation in the
light of mechanical requirements or for vulnerability to cavitation, for instance.

The three points (i) the positive correlation between A_{pet} and A_{lamina}, (ii) the trade-off
between leaf size and lobe intensity and (iii) the axis-like anatomy of *Cecropia* petioles, call
for a functional analogy between the LSR and the lamina-petiole relationship. This conclusion
led us to study the functional significance of anatomical adjustments in the frame of the LSR
using a leaf model.

Pith, not secondary xylem, is the main driver of variations in petiole size
Variations in leaf dimensions could be the result of different processes which take place
simultaneously at a macro-anatomical level or at the tissue and cell level. It seems logical that
variations in petiole dimensions are primarily driven by variations in tissue surface areas and
volumes. In our study, we provided a complete description of petiole anatomy for *C. obtusa*
by completing a previous succinct description (Bonsen and Welle 1983). The surface area of
all the tissues we studied was found to be positively correlated with A_{pet}. However, a
surprising result was that the proportions of the petiole tissue were not consistent with the leaf
dimensions. Only the proportion of primary xylem to its surface area remained constant. Even
if the proportion of variation in the secondary xylem can vary substantially linked to the
secondary growth processes, the striking feature is the marked tendency of the variation in the
pith to drive variation in the absolute size of the petiole.

Because of a high second moment of areas in flexion of more external supporting
tissues (i.e. secondary xylem, sclerenchyma and collenchyma) thanks to the pith, it seems that
Cecropia petioles have a specific and relatively cheap mechanical strategy for supporting
large laminae. The key point is that the proportion of pith is positively correlated with A_{pet},
implying that the second moment of areas of external tissues is positively correlated with
petiole size (under the hypothesis of conserved material properties). In other words, changing

the proportion of pith with an increase in leaf size allowed for a strong structural effect (from

the mechanical point of view) of support tissues (e.g. secondary xylem) as they are pushed

centrifugally and since a tube is stiffer than a solid cylinder (for a given materiel and a given

amount of this materiel) (Niklas 1992). Here, more research is needed for mechanical

quantification. If it holds true, this feature underlines the importance of distinguishing intra-

and interspecific levels in scaling models as briefly discussed above. Indeed, here we see how

a specific strategy relating to leaf size and form can determine allometric scaling.

Functional significance of anatomical adjustments by the petiole

513   The positive correlation we found between the leaf dimensions and K_{th} is in agreement

with the results of previous studies showing that larger appendages require larger hydraulic

supplies (Preston and Ackerly 2003, Normand et al. 2008, Fan et al. 2017), linked to

increasing evaporative demand. However, the allometric form of this relationship suggests

that a given lamina surface area for large leaves is supplied with disproportionate amounts of

water in comparison with the supply to small leaves. This property could leads to increased

evaporation and stomatal conductance but deserves more appropriate ecophysiological

measurements. This process is mainly driven by an increase in the vessel diameter (derived

from the mean hydraulic diameter, Dmh) rather than an increase in N_{vessel} or in the xylem

area. Dmh was positively correlated with A_{pet} whereas f_{vessel} was negatively correlated with

A_{pet}. Secondary xylem area and its proportion were positively correlated with A_{pet}, but it is

still difficult to disentangle if this process is driven by hydraulic or mechanical needs.

525   Overall, S was positively correlated with A_{pet}, reflecting the greater weight of

variations in Dmh compared to f_{vessel} and leading to a larger petiole K_s for large leaves. Thus,

xylem is hydraulically optimised through variation in Dmh, avoiding a costly and less

efficient strategy which would cause more xylem tissues to grow (to support more vessels),

and would result in a disproportionately large petiole relative to the lamina. This feature is in

agreement with the global interspecific pattern described by Zanne *et al.* (Zanne et al. 2010).

However, despite the increasing return on hydraulic performance enabled by the variation in

Dmh, there must be a trade-off between Dmh and f_{vessel}. This statement is in agreement with

the theoretical model of the packing rule (Sperry et al. 2008, Chen et al. 2012), which predicts

that f_{vessel} should isometrically scale with conduit size and thus allows for the area-preserving

branching of vessels across different branching levels.

536   Finally, such variability calls for an investigation of vulnerability to cavitation. Indeed,

537   Scoffoni *et al.* (2016) found that larger vessels in the leaf petiole and major veins are related

to vulnerability to cavitation. In our study, we showed that larger leaves have larger vessels suggesting higher vulnerability to cavitation, which would have substantial implications for carbon economy or for dealing with the alternation of dry and wet seasons in our study region or the potential intensification of dry seasons in Amazonia with climate change. Accordingly, we found that tyloses occur preferentially on large vessels in the secondary xylem, while tyloses are usually explained in the light of cavitation mitigation (Micco et al. 2016). Since knowledge is lacking on the intraspecific variability of vulnerability to cavitation and related factors (size, season, ontogeny), further investigations are clearly needed.

Conclusion

Our study demonstrates that the functional analogy between the LSR and the lamina-petiole relationship holds true based on (i) the positive correlation between A_{pet} and A_{lamina}, (ii) the trade-off between leaf size and lobe intensity and (iii) the described axis-like anatomy of *Cecropia* petioles. From this framework, we point that (i) the pith, not the secondary xylem, is the main driver of variation in petiole size, and (ii) variation in xylem vessel anatomy, not its surface area, is the main determinant of hydraulic requirements by allowing a size-dependent xylem hydraulic efficiency. Our results suggest that hydraulics are a relatively poor explanation as a cause of the Corner's rule and LSR. Thus, size-related mechanical requirements have to be investigated in the same way, to totally disentangle the frequent but speculative statement that Corner's rule emerged from hydraulic and mechanical requirements. Such size-dependent xylem efficiency also calls for studying the consequence on stomatal conductance and photosynthesis. Finally, future work will be needed to evaluate if such statement holds true at the interspecific level, by ideally integrating morphological, anatomical and physiological data related to stem, petiole and lamina.

ACKNOWLEDGEMENTS

The authors thank Julie Bossu, Coffi Belmys Cakpo, Henri Caron, Jocelyn Cazal, Saint-Omer Cazal, Aurélie Cuvelier, Bruno Clair, Aurélie Dourdain, Jean-Yves Goret, Marie Hartwig, Ariane Mirabel, Audin Patient, Pascal Petronelli, Laurent Risser, Dylan Taxile, Valérie Troispoux, Niklas Tysklind and Lore Verryckt for their assistance with field work and measurement of leaf traits. We thank Jacques Beauchêne, Christine Heinz and Nick Rowe for preliminary discussions around the project and first results. We thank Tancrède Alméras for critical and valuable comments on the manuscript.

FUNDING

S.L. was supported by a doctoral fellowship from CEBA (ref. ANR-10-LABX-0025). This

study benefited from an *Investissement d'Avenir* grant managed by the *Agence Nationale de*

la Recherche (CEBA, ref. ANR-10-LABX-0025).

LITERATURE CITED

Aarssen L (2012) Reducing size to increase number: a hypothesis for compound leaves. Ideas
Ecol Evol 5. https://queens.scholarsportal.info/ojs-
archive/index.php/IEE/article/view/4274 (9 February 2018, date last accessed).

Abramoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ. Biophotonics
Int 11:36–42.

Ackerly D, Donoghue MJ (1998) Leaf size, sapling allometry, and Corner's rules: phylogeny
and correlated evolution in maples (Acer). Am Nat 152:767–791.

Anfodillo T, Carraro V, Carrer M, Fior C, Rossi S (2006) Convergent tapering of xylem
conduits in different woody species. New Phytol 169:279–290.

Anfodillo T, Petit G, Crivellaro A (2013) Axial conduit widening in woody species: a still
neglected anatomical pattern. IAWA J 34:352–364.

Bettiati D, Petit G, Anfodillo T (2012) Testing the equi-resistance principle of the xylem
transport system in a small ash tree: empirical support from anatomical analyses. Tree
Physiol 32:171–177.

Bonsen K, Welle BJH ter (1983) Comparative wood and leaf anatomy of the Cecropiaceae
(Urticales). Bull Mus Natl Hist Nat Sect B Adansonia Bot Phytochim.
http://agris.fao.org/agris-search/search.do?recordID=US201302138000 (28 December
2016, date last accessed).

Brouat C, Gibernau M, Amsellem L, McKey D (1998) Corner's Rules Revisited: Ontogenetic
and Interspecific Patterns in Leaf-Stem Allometry. New Phytol 139:459–470.

Brouat C, McKey D (2001) Leaf-stem allometry, hollow stems, and the evolution of caulinary
domatia in myrmecophytes. New Phytol 151:391–406.

Bucci SJ, Scholz FG, Goldstein G, Meinzer FC, Sternberg LDSL (2003) Dynamic changes in
hydraulic conductivity in petioles of two savanna tree species: factors and mechanisms
contributing to the refilling of embolized vessels. Plant Cell Environ 26:1633–1645.

Busch BL, Schmitz G, Rossmann S, Piron F, Ding J, Bendahmane A, Theres K (2011) Shoot
Branching and Leaf Dissection in Tomato Are Regulated by Homologous Gene
Modules. Plant Cell 23:3595–3609.

Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a
worldwide wood economics spectrum. Ecol Lett 12:351–366.

Chen H, Niklas KJ, Sun S (2012) Testing the packing rule across the twig–petiole interface of
609  temperate woody species. Trees 26:1737–1745.

610 Cochard H, Coste S, Chanson B, Guehl J-M, Nicolini É (2005) Hydraulic architecture
611  correlates with bud organogenesis and primary shoot growth in beech (Fagus
612  sylvatica). Tree Physiol 25, 25. http://prodinra.inra.fr/?locale=fr#!ConsultNotice:8708
613  (7 March 2017, date last accessed).

614 Coomes DA, Heathcote S, Godfrey ER, Shepherd JJ, Sack L (2008) Scaling of xylem vessels
615  and veins within the leaves of oak species., Scaling of xylem vessels and veins within
616  the leaves of oak species. Biol Lett Biol Lett 4, 4:302, 302–306.

617 Corner EJH (1949) The Durian Theory or the Origin of the Modern Tree. Ann Bot 13:367–
618  414.

619 Díaz S, Kattge J, Cornelissen JHC, Wright IJ, Lavorel S, Dray S, Reu B, Kleyer M, Wirth C,
620  Prentice IC, Garnier E, Bönisch G, Westoby M, Poorter H, Reich PB, Moles AT,
621  Dickie J, Gillison AN, Zanne AE, Chave J, Wright SJ, Sheremet'ev SN, Jactel H,
622  Baraloto C, Cerabolini B, Pierce S, Shipley B, Kirkup D, Casanoves F, Joswig JS,
623  Günther A, Falczuk V, Rüger N, Mahecha MD, Gorné LD (2016) The global spectrum
624  of plant form and function. Nature 529:167–171.

625 Dixon H, Joly J (1895) On the ascent of sap. Philos Trans R Soc Lond

626 Dombroskie SL, Aarssen LW (2012) The leaf size/number trade-off within species and within
627  plants for woody angiosperms. Plant Ecol Evol 145:38–45.

628 Duursma RA, Falster DS, Valladares F, Sterck FJ, Pearcy RW, Lusk CH, Sendall KM,
629  Nordenstahl M, Houter NC, Atwell BJ, Kelly N, Kelly JWG, Liberloo M, Tissue DT,
630  Medlyn BE, Ellsworth DS (2012) Light interception efficiency explained by two
631  simple variables: a test using a diversity of small- to medium-sized woody plants. New
632  Phytol 193:397–408.

633 Enquist BJ, West GB, Charnov EL, Brown JH (1999) Allometric scaling of production and
634  life-history variation in vascular plants. Nature 401:907–911.

635 Falster DS, Warton DI, Wright II (2006) User's guide to SMATR: Standardised Major Axis
636  Tests & Routines Version 2.0, Copyright 2006. R Instr

637 Fan Z-X, Sterck F, Zhang S-B, Fu P-L, Hao G-Y (2017) Tradeoff between Stem Hydraulic
638  Efficiency and Mechanical Strength Affects Leaf–Stem Allometry in 28 Ficus Tree
639  Species. Front Plant Sci 8.
640  https://www.frontiersin.org/articles/10.3389/fpls.2017.01619/full (12 February 2018,
641  date last accessed).

642 Foster AS (1936) Leaf Differentiation in Angiosperms. Bot Rev 2:349–372.

643 Givnish TJ (1984) Leaf and Canopy Adaptations in Tropical Forests. In: Physiological
644  ecology of plants of the wet tropics. Dr. W. Junk Publisher, Hague, pp 51–84.
645  http://link.springer.com/chapter/10.1007/978-94-009-7299-5_6 (28 December 2016,
646  date last accessed).

Gourlet-Fleury S, Guehl JM, Laroussine O (2004) Ecology and management of a neotropical
rainforest : lessons drawn from Paracou, a long-term experimental research site in
French Guiana. Elsevier, Paris. http://publications.cirad.fr/une_notice.php?dk=522004

Hacke UG, Spicer R, Schreiber SG, Plavcová L (2016) An ecophysiological and
developmental perspective on variation in vessel diameter. Plant Cell Environ

Hajek P, Kurjak D, von Wühlisch G, Delzon S, Schuldt B (2016) Intraspecific Variation in
Wood Anatomical, Hydraulic, and Foliar Traits in Ten European Beech Provenances
Differing in Growth Yield. Front Plant Sci 7.
http://journal.frontiersin.org/article/10.3389/fpls.2016.00791/abstract (5 December
2016, date last accessed).

Hallé F, Oldeman RAA, Tomlinson PB (1978) Tropical Trees and Forests - An Architectural
Analysis. Springer-Verlag Berlin Heidelberg.
http://www.springer.com/us/book/9783642811920 (28 December 2016, date last
accessed).

Harvey PH, Pagel MD (1991) The comparative method in evolutionary biology. Oxford
University Press.

Heuret P, Barthélémy D, Guédon Y, Coulmier X, Tancre J (2002) Synchronization of growth,
branching and flowering processes in the South American tropical tree Cecropia
obtusa (Cecropiaceae). Am J Bot 89:1180–1187.

Huang Y, Lechowicz MJ, Zhou D, Price CA (2016) Evaluating general allometric models:
interspecific and intraspecific data tell different stories due to interspecific variation in
stem tissue density and leaf size. Oecologia 180:671–684.

Huxley JS (1932) Problems of Relative Growth. Methuen & Co. Ltd, London.

Kleiman D, Aarssen LW (2007) The leaf size/number trade-off in trees. J Ecol 95:376–382.

Lehnebach R, Beyer R, Letort V, Heuret P The pipe model theory half a century on: a review.
Ann Bot. https://academic.oup.com/aob/advance-
article/doi/10.1093/aob/mcx194/4820933 (14 February 2018, date last accessed).

Lens F, Sperry JS, Christman MA, Choat B, Rabaey D, Jansen S (2011) Testing hypotheses
that link wood anatomy to cavitation resistance and hydraulic conductivity in the
genus Acer. New Phytol 190:709–723.

Liu Z, Cai Y, Li K, Guo J (2010) The leaf size-twig size spectrum in evergreen broadleaved
forest of subtropical China. Afr J Biotechnol 9:3382–3387.

Micco VD, Balzano A, Wheeler EA, Baas P (2016) Tyloses and gums: A review of structure,
function and occurance of vessel occlusions. IAWA J 37:186–205.

Milla R (2009) The leafing intensity premium hypothesis tested across clades, growth forms
and altitudes. J Ecol 97:972–983.

Milla R, Reich PB (2007) The scaling of leaf area and mass: the cost of light interception
increases with leaf size. Proc R Soc Lond B Biol Sci 274:2109–2115.

Nicotra AB, Leigh A, Boyce CK, Jones CS, Niklas KJ, Royer DL, Tsukaya H (2011) The
evolution and functional significance of leaf shape in the angiosperms. Funct Plant
Biol 38:535–552.

Niinemets Ü, Portsmuth A, Tena D, Tobias M, Matesanz S, Valladares F (2007) Do we
Underestimate the Importance of Leaf Size in Plant Economics? Disproportional
Scaling of Support Costs Within the Spectrum of Leaf Physiognomy. Ann Bot
100:283.

Niklas KJ (1992) Plant Biomechanics: Engineering Approach to Plant Form and Function.
University of Chicago Press, Chicago.

Niklas KJ (1994) Plant Allometry: The Scaling of Form and Process. University of Chicago
Press.

Normand F, Bissery C, Damour G, Lauri P-É (2008) Hydraulic and mechanical stem
properties affect leaf–stem allometry in mango cultivars. New Phytol 178:590–602.

Olson ME, Aguirre-Hernández R, Rosell JA (2009) Universal foliage-stem scaling across
environments and species in dicot trees: plasticity, biomechanics and Corner's Rules.
Ecol Lett 12:210–219.

Olson ME, Rosell JA (2013) Vessel diameter–stem diameter scaling across woody
angiosperms and the ecological causes of xylem vessel diameter variation. New Phytol
197:1204–1213.

Osada N, Nabeshima E, Hiura T (2015) Geographic variation in shoot traits and branching
intensity in relation to leaf size in Fagus crenata: A common garden experiment. Am J
Bot. http://www.amjbot.org/content/early/2015/06/11/ajb.1400559 (28 December
2016, date last accessed).

Poorter, Rozendaal DMA (2008) Leaf size and leaf display of thirty-eight tropical tree
species. Oecologia 158:35–46.

Preston KA, Ackerly DD (2003) Hydraulic architecture and the evolution of shoot allometry
in contrasting climates. Am J Bot 90:1502–1512.

Royer DL, Coley PD, Cutter AD, Johnson KR, Labandeira CC, Moles AT, Palmer MB,
Valladares F, Sack L, Wilf P, Lusk CH, Jordan GJ, Niinemets U, Wright IJ, Westoby
M, Cariglino B (2007) Fossil leaf economics quantified: calibration, Eocene case
study, and implications. Paleobiology 33:574–589.

Scholz A, Klepsch M, Karimi Z, Jansen S (2013) How to quantify conduits in wood? Front
Plant Sci 4. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3600434/ (28 December
2016, date last accessed).

Schuldt B, Leuschner C, Brock N, Horna V (2013) Changes in wood density, wood anatomy
and hydraulic properties of the xylem along the root-to-shoot flow path in tropical
rainforest trees. Tree Physiol 33:161–174.

Scott SL, Aarssen LW (2012) Within-species leaf size–number trade-offs in herbaceous
angiosperms. Botany 90:223–235.

Sinnott EW (1921) The Relation between Body Size and Organ Size in Plants. Am Nat
55:385–403.

Smith DD, Sperry JS, Adler FR (2017) Convergence in leaf size versus twig leaf area scaling:
do plants optimize leaf area partitioning? Ann Bot 119:447–456.

Sperry JS, Meinzer FC, McCulloh KA (2008) Safety and efficiency conflicts in hydraulic
architecture: scaling from tissues to trees. Plant Cell Environ 31:632–645.

Steppe K, Lemeur R (2007) Effects of ring-porous and diffuse-porous stem wood anatomy on
the hydraulic parameters used in a water flow and storage model. Tree Physiol 27:43–
52.

Steudle E (2001) The cohesion-tension mechanism and the acqisition of water by plant roots.
Annu Rev Plant Physiol Plant Mol Biol 52:847–875.

Sun S, Jin D, Shi P (2006) The Leaf Size–Twig Size Spectrum of Temperate Woody Species
Along an Altitudinal Gradient: An Invariant Allometric Scaling Relationship. Ann Bot
97:97–107.

Tao L, JianMing D, GenXuan W, DongLiang C, ZeLong Y (2009) Isometric scaling
relationship between leaf number and size within current-year shoots of woody species
across contrasting habitats. Pol J Ecol 57:659–667.

Thompson DW (1917) On Growth and Form. Cambridge University Press, Cambridge.

Trueba S, Isnard S, Barthélémy D, Olson ME (2016) Trait coordination, mechanical behavior,
and growth form plasticity of Amborella trichopoda under variation in canopy
openness. AoB Plants:plw068.

Tsukaya H (1995) Developmental genetics of leaf morphogenesis in dicotyledonous plants. J
Plant Res 108:407–416.

Tsukaya H (2013) Leaf Development. Arab Book Am Soc Plant Biol 11.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3711357/ (12 February 2018, date
last accessed).

Tsukaya H, Inaba-Higano K, Komeda Y (1995) Phenotypic Characterization and Molecular
Mapping of an acaulis2 Mutant of Arabidopsis thaliana with Flower Stalks of Much
Reduced Length. Plant Cell Physiol 36:239–246.

Tsukaya H, Kozuka T, Kim G-T (2002) Genetic Control of Petiole Length in Arabidopsis
thaliana. Plant Cell Physiol 43:1221–1228.

Tsukaya H, Naito S, Redei GP, Komeda Y (1993) A new class of mutations in Arabidopsis
thaliana, acaulis1, affecting the development of both inflorescences and leaves.
Development 118:751–764.

Tyree, Zimmermann (2002) Xylem Structure and the Ascent of Sap. Springer.
http://www.springer.com/gp/book/9783540433545 (5 December 2016, date last
accessed).

Vasseur F, Violle C, Enquist BJ, Granier C, Vile D (2012) A common genetic basis to the
origin of the leaf economics spectrum and metabolic scaling allometry. Ecol Lett
15:1149–1157.

Warton DI, Wright IJ, Falster DS, Westoby M (2006) Bivariate line-fitting methods for
allometry. Biol Rev 81:259–291.

Westoby M, Daniel S. Falster DS, Angela T. Moles, Peter A. Vesk, Wright and IJ (2002)
Plant Ecological Strategies: Some Leading Dimensions of Variation Between Species.
Annu Rev Ecol Syst 33:125–159.

Westoby M, Wright IJ (2003) The Leaf Size-Twig Size Spectrum and Its Relationship to
Other Important Spectra of Variation among Species. Oecologia 135:621–628.

White PS (1983a) Corner's Rules in Eastern Deciduous Trees: Allometry and Its Implications
for the Adaptive Architecture of Trees. Bull Torrey Bot Club 110:203–212.

White PS (1983b) Evidance that temperate east north american evergreen woody plants
follow Corner's rule. New Phytol 95:139–145.

Whitman T, Aarssen LW (2010) The leaf size/number trade-off in herbaceous angiosperms. J
Plant Ecol 3:49–58.

Williamson GB, Wiemann MC (2010) Measuring wood specific gravity…Correctly. Am J
Bot 97:519–524.

Wright IJ, Dong N, Maire V, Prentice IC, Westoby M, Díaz S, Gallagher RV, Jacobs BF,
Kooyman R, Law EA, Leishman MR, Niinemets Ü, Reich PB, Sack L, Villar R,
Wang H, Wilf P (2017) Global climatic drivers of leaf size. Science 357:917–921.

Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J,
Chapin T, Cornelissen JHC, Diemer M, Flexas J, Garnier E, Groom PK, Gulias J,
Hikosaka K, Lamont BB, Lee T, Lee W, Lusk C, Midgley JJ, Navas M-L, Niinemets
Ü, Oleksyn J, Osada N, Poorter H, Poot P, Prior L, Pyankov VI, Roumet C, Thomas
SC, Tjoelker MG, Veneklaas EJ, Villar R (2004) The worldwide leaf economics
spectrum. Nature 428:821–827.

Yan E-R, Wang X-H, Chang SX, He F (2013) Scaling relationships among twig size, leaf size
and leafing intensity in a successional series of subtropical forests. Tree Physiol
33:609–617.

Yang D, Li G, Sun S (2008) The Generality of Leaf Size versus Number Trade-off in
Temperate Woody Species. Ann Bot 102:623–629.

Zalamea P-C, Heuret P, Sarmiento C, Rodríguez M, Berthouly A, Guitet S, Nicolini E,
Delnatte C, Barthélémy D, Stevenson PR (2012) The Genus Cecropia: A Biological
Clock to Estimate the Age of Recently Disturbed Areas in the Neotropics. PLoS ONE
7:e42643.

Zalamea P-C, Stevenson PR, Madriñán S, Aubert P-M, Heuret P (2008) Growth pattern and
age determination for Cecropia sciadophylla (Urticaceae). Am J Bot 95:263–271.

Zanne AE, Westoby M, Falster DS, Ackerly DD, Loarie SR, Arnold SEJ, Coomes DA (2010)
Angiosperm wood structure: Global patterns in vessel anatomy and their relation to
wood density and potential conductivity. Am J Bot 97:207–215.

FIGURE LEGENDS

Fig. 1. Habits, morphological and petiole-anatomical aspects of *C. obtusa* (Urticaceae). (A)

Habits of *C. obtusa*. (B) Leaf of *C. obtusa*. (C) Petiole cross-sectional anatomy of *C. obtusa*

in the middle of the petiole. (D) Close-up of petiole constitutive tissues: pith (Pith),

sclerenchyma (Scle), interfascicular parenchyma (Pint), primary xylem (XI), secondary xylem

(XII), total phloem (Phlo), primary phloem (Phl), secondary phloem (PhII), cortical

parenchyma (Pcor), collenchyma (Coll) and epidermis (Epi). Simple arrows indicate tylosed

vessels. (E) Tissues and corresponding layer masks studied. (F) "Wavy" cambium with a sub-

bicyclic array of vascular bundles. Arrows represent laticiferous canals. (G) Cambial

discontinuities, island-like vascular bundles and a bicyclic array of vascular bundles.

Fig. 2. Sampling and measurement design. (a) petiole length (L_{pet}); (b) the number of lobes

(N_{lobe}), here there are 7 lobes; (c) the length of the main lobe (LM_{lobe}); (d) The lamina mass

area (LMA) is measured based on three tablets if a circular iron punch; (e) in the P-leaves,

one segment is used to measurable Se petiole specific gravity (PSG); (f) in the P-leaves, one

segment is used for the anatomical study from which is notably derived the petiole cross-

sectional area (A_{pet}); (g) the lamina surface area (A_{lamina}).

Fig. 3. (A) Relationship between the undamaged lamina area (cm²) and the length of the main

lobe (cm) based on a non-linear least square regression ($P < 0.001$; $n = 888$). An estimated

lamina area can be derived from the length of the main lobe ($y = 0.90672 * x^{2.10950}$). Dashed

lines represent the confidence interval at 95%. (B) Comparison between undamaged and

measured lamina area (filled circles, dashed line; $P < 0.001$; $n = 892$) and estimated lamina

area (empty circles, solid line; $P < 0.001$; $n = 1317$) corresponding to the petiole cross-

sectional area. Curves are derived from the SMA regression on a log-log scale.

Fig. 4. Relationship between the ratio of the number of lobes to undamaged lamina area and

the petiole cross-sectional area ($P < 0.001$). The curve is derived from the SMA regression on

a log-log scale. Inset shows data on a log-log scale.

Fig. 5. (A) Relationship between the cross-sectional area of the different petiole tissues and

the total petiole cross-sectional area on a log-log scale ($P < 0.001$). (B) Relationship between

the proportions of the different tissues and the petiole cross-sectional area ($P < 0.01$). Curves

are derived from the SMA regression on a log-log scale. (C) Boxplots of the relative

proportions of the different tissues according to the petiole cross-sectional area. Pink: pith, brown: sclerenchyma, cyan: interfascicular parenchyma, blue: primary xylem, red: secondary xylem, green: phloem, orange: cortical parenchyma, purple: collenchyma.

Fig. 6. Relationships between the petiole cross-sectional area and (A) the xylem surface area ($P < 0.001$), (B) the mean hydraulic diameter ($P < 0.001$), (C) the vessel frequency ($P < 0.001$), and (D) the ratio of mean vessel area to frequency ($P < 0.001$). Curves are derived from the SMA regression on a log-log scale. Insets show data on a log-log scale. Black: data relative the total xylem. Blue: data relative to the primary xylem. Red: data relative to the secondary xylem.

849 Fig. 7. Relationships between the petiole cross-sectional area and (A) theoretical hydraulic conductivity ($P < 0.001$), (B) xylem-specific hydraulic conductivity ($P < 0.001$) and (C) leaf-specific hydraulic conductivity ($P < 0.001$). Curves are derived from the SMA regression on a log-log scale. Insets show data on a log-log scale. Black: data relative to the total xylem. Blue: data relative to the primary xylem. Red: data relative to the secondary xylem.

856 TABLES

857 TABLE 1. List of the morphological, anatomical and hydraulic traits analysed.

Trait	Abbreviation	Unit	Formula	Reference
Ellipse surface area			$A = \pi * d_1/2 * d_2/2$ (d = diameter)	
Petiole cross-sectional area	A_{pet}	mm²		
Petiole length	L_{pet}	mm²		
Main lobe length	LM_{lobe}	cm		
Lamina area	A_{lamina}	cm²		
Number of lobes	N_{lobe}	n		
Petiole dry mass	M_{pet}	g		
Lamina dry mass	M_{lamina}	g		
Lamina mass area	LMA	g mm^2		
Petiole specific gravity	PSG	g mm^3		
Number of vessels	N_{vessel}	n		
Mean hydraulic diameter	Dmh	μm	$Dmh = \left(\dfrac{\sum D_H^4}{N}\right)^{1/4}$	Tyree & Zimmermann, 2002 (Steppe and Lemeur 2007) Scholz *et al.*, 2013
Vessel frequency	f_{vessel}	n mm^2	$f_{vessel} = \dfrac{N_{vessel}}{Xylem\ surface\ area}$	
Lumen fraction	F	%	$F = Mean\ vessel\ area * f_{vessel}$	Zanne *et al.*, 2010
Mean vessel area to density ratio	S	μm²/μm²	$S = \dfrac{Mean\ vessel\ area}{fvessel}$	Zanne *et al.*, 2010
Ellipse conductivity	$K_{ellipse}$	kg s^{-1} MPa^{-1} m	$K_{ellipse} = \dfrac{\pi}{64\eta}\dfrac{d_1^3 d_2^3}{d_1^2 + d_2^2}$ for η = 1.002*10^{-9} MPa s^{-1} at 20°C	Tyree & Zimmermann, 2002 Steppe & Lemeur, 2007 Scholz *et al.*, 2013
Theoretical hydraulic conductivity	K_{th}	kg s^{-1} MPa^{-1} m	$K_{th} = \sum K_{ellipse}$	Tyree & Zimmermann, 2002 (Scholz et al. 2013)
Xylem-specific hydraulic conductivity	K_s	kg s^{-1} MPa^{-1} m^{-1}	$K_s = \dfrac{K_{th}}{Xylem\ surface\ area}$	Tyree & Zimmermann, 2002

860 TABLE 2. Correlation matrix for leaf-level morphological traits based on SMA regressions
861 on a log-log scale.

	A_{pet}	L_{pet}	A_{lamina}	N_{lobe}	M_{pet}	M_{lamina}	LMA	PSG
A_{pet}	-	a 0.66***	a 1.28***	a 0.19***	a 1.47***	a 1.35***	a 0.34***	-0.26
L_{pet}	0.734	-	a 1.90***	a 0.29***	a 2.26***	a 2.03***	-0.51	a -0.34*
A_{lamina}	0.880	0.837	-	a 0.15***	a 1.15***	a 1.03***	-0.28	a -0.22**
N_{lobe}	0.186	0.160	0.139	-	a 7.66***	a 7.03***	a 1.79***	-1.56
M_{pet}	0.934	0.829	0.895	0.199	-	a 0.91***	a 0.23***	-0.16
M_{leaf}	0.935	0.776	0.924	0.173	0.962	-	a 0.26***	-0.19
LMA	0.063	<0.001	<0.001	0.028	0.051	0.068	-	a 0.65***
PSG	<0.001	0.074	0.166	0.025	0.001	<0.001	0.334	-

x-axis

Bold values are for significant correlations ($P < 0.05$).
*: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$

SUPPLEMENTARY TABLES

SUPPLEMENTARY TABLE 1. Correlation matrix for tissue surface area and proportion
traits based on SMA regressions on a log-log scale. Allometric exponents (slopes) are
presented.

SUPPLEMENTARY TABLE 2. Correlation matrix between morphological, anatomical and
hydraulic traits based on SMA regressions on a log-log scale. Allometric exponent (slopes)
are presented.
870
871

A

B

C

1 mm

D

Epi

Coll

Pcor

Phlo

Phl

PhII

XII

XI

Pint

Scle

Pith

250 µm

E

F

1 mm

G

1 mm

	Pith		XI		XII		Pcor
	Scle		Pint		Phlo		Coll

(d) **LAM**$_{lam}$

LM$_{lobe}$

(c)

(g) **A**$_{lamina}$

7 (b) **N**$_{lobe}$

(e) **PSG**

(f) **Anatomy**

(a) **L**$_{pet}$ + **A**$_{pet}$

8mm

5cm