Sensory neurons sprouting is dependent on osteoclast-derived extracellular vesicles

involving the activation of epidermal growth factor receptors

Estrela Neto[1, 2], Luís Leitão[1, 2, 3], José Mateus[1, 2], Daniela M. Sousa[1, 2], Francisco Conceição[1, 2, 3], Cecília J.

Alves[1, 2], Richard O. C. Oreffo[5], Jonathan West[6], Paulo Aguiar[1, 2] and Meriem Lamghari[1, 2, *]

[1] i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen 280, 4200-

135 Porto, Portugal

[2] INEB - Instituto de Engenharia Biomédica, Universidade do Porto, Rua Alfredo Allen 280, 4200-135

Porto, Portugal

[3] ICBAS - Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, Rua de Jorge Viterbo

Ferreira n.º 228, 4050-313 Porto, Portugal

[4] Centre for Human Development, Stem Cells and Regeneration, Human Development and Health,

Tremona Rd, Southampton SO16 6YD, United Kingdom

[5] Institute for Life Sciences and Cancer Sciences, University of Southampton, University of Southampton,

Highfield Campus, Southampton SO17 1BJ, United Kingdom

*Correspondence: Meriem Lamghari, lamghari@ineb.up.pt

Significance Statement: Sensory nerve fibers sprouting in bone pathologies is highly associated with pain.

Thus, understanding the mechanisms behind sensory nerves ingrowth, sprouting and electrical activity,

within the bone compartment, is essential for improving the strategies to overcome pain in bone disorders.

We provide a new mechanism on the sensory nerves sprouting, indicating that the effect is dependent on

the extracellular vesicles (EV) released by osteoclasts, through the epidermal growth factor receptor family

targeting, by integrin independent pathways. We show different electrophysiology patterns being triggered

in the presence of osteoclasts secretome and the abolishment of sensory neurons firing rate in EV-depleted

conditions. Overall, our results elucidate novel mechanisms on the peripheral nerves sprouting, essential

for pursuing new targets for bone pain therapies.

Abstract

The patterning of peripheral innervation is accomplished through the tissue expression, in specific space

and timeframe, of attractive or repulsive axonal guidance cues. At the bone microenvironment, neurotrophic

factors such as nerve growth factor, brain-derived neurotrophic factor, vascular endothelial growth factor,

netrin-1 and others were described to regulate the nerve ingrowth towards the bone compartment, by acting

directly on receptors expressed at the nerve terminals. Interestingly, besides the gradient of soluble factors,

neurons were described to be responsive to extracellular vesicles (EV) derived from myelinating cells and

mesenchymal stem cells.

Here we provide evidence on a new mechanism by which peripheral innervation can be coordinated. We

show that sensory nerves outgrowth and electric signal propagation are dependent on the EV secreted by

osteoclasts, the bone resorbing cells. Furthermore, we demonstrate that the axonal sprouting is achieved

through the activation of epidermal-growth factor receptor (EGFR) family signaling pathway. We proved

that the EV-depleted osteoclast secretome leads to a significant decrease of neurons firing rate and axonal

sprouting, concomitant with a decrease of EGFR/ErbB2 activation levels. Excitingly, the proteomic

analysis of the osteoclast-derived EV cargo shows a high correlation with synaptic components reinforcing

the role on sensory neurons/osteoclast crosstalk.

Our findings that osteoclast-derived EV hold effect in axonal outgrowth, contributing actively to the

dynamics of the sensory neurons sprouting and electrophysiology, is a step toward unraveling target

mechanisms to control electrical signal propagation and nerve fibers sprouting and consequently open new

avenues for the development of innovative therapies to control bone pain.

Keywords: sensory neurons sprouting; extracellular vesicles; epidermal growth factor receptor

(EGFR/ErbB2) signaling; electrophysiology; osteoclast secretome.

Introduction

Nerve fibers present in the bone microenvironment have been directly implicated in bone homeostasis and regeneration (Chenu, 2004; Chenu and Marenzana, 2005; Elefteriou, 2008; Elefteriou et al., 2014; Patel and Elefteriou, 2007). Anatomical mapping of innervation during skeletal development shows co-localization of nerve fibers with a rich capillary network in areas with high osteogenic activity (Gomez et al., 2005; Hukkanen et al., 1993; Li et al., 2007; Sisask et al., 2013; Tomlinson et al., 2016) and the pattern of neuronal dynamics observed during bone development is typically repeated (Hukkanen et al., 1993; Li et al., 2001; Madsen et al., 1998; Nordsletten et al., 1994), highlighting the critical role of neuronal regulation. Several denervation studies have revealed an impairment on fracture healing, emphasizing the impact of the peripheral nervous system on bone turnover (Bove et al., 2009; Cherruau et al., 1999; Edoff et al., 1997; Hill and Elde, 1991; Song et al., 2012). In line with these observations, sensory innervation has attracted significant interest concerning bone formation. Beyond their role in bone nociception, evidence demonstrates that the lack of sensory nerves significantly impairs bone mass accrual (Chen et al., 2019; Fukuda et al., 2013; Hayashi et al., 2012), indicating a key role in bone homeostasis. In the opposite way, undifferentiated bone marrow stromal cells are acknowledged to hold a neurotrophic potential. Both the bone marrow cells and/or their secretome are widely used to improve nerve regeneration (Brohlin et al., 2012; Chen et al., 2002; Gu et al., 2010; Han et al., 2016; Lu et al., 2005; Neuhuber et al., 2005; Novikova et al., 2011; Wilkins et al., 2009), mainly through the secretion of neurotrophic factors such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), vascular endothelial growth factor (VEGF), platelet-derived growth factor (PDGF) or epidermal growth factor (EGF) (Han et al., 2016). These growth factors are known to activate their respective receptors in axonal terminals, ultimately promoting axonal regeneration, elongation and guidance (Chao, 2003).

Interestingly, in bone pathological conditions such as bone cancer, ectopic sprouting of nerve fibers in the bone marrow, mineralized bone, and periosteum is detected (Bloom et al., 2011; Chartier et al., 2014; Ghilardi et al., 2010; Jimenez-Andrade et al., 2010). Osteoclasts, the bone resorbing cells, play a major role in the activation of nociceptors. Matrix resorption in the bone compartment results in active proton secretion into the extracellular space and a decrease in the pH milieu (Mantyh, 2014; Middlemiss et al., 2011; Yoneda et al., 2011), stimulating the acid-sensing ions channels in nerve endings and causing pain. Notably, clinical data showed that osteoclast inhibitors significantly reduced cancer-associated bone pain (Cleeland et al.,

2013; Kohno et al., 2005). While such studies provide support for the role of osteoclasts in the activation

of nociceptors in diseased bone (Yoneda et al., 2015b), recent reports start to reveal how osteoclasts

contribute to the changes in the innervation pattern in normal or diseased bone. Zhu *et al*. have shown that

osteoclasts can actively promote axonal growth on subchondral bone, through netrin-1 signaling, in an

osteoarthritis model (Zhu et al., 2019). Moreover, osteoclast-derived netrin-1 was also linked to an increase

in the sensory innervation of porous endplates of intervertebral disc, causing an low back pain (Ni et al.,

2019). Therefore, deciphering the mechanisms by which peripheral innervation can be controlled by the

osteoclasts constitute a major advantage for developing new strategies to tackle bone pain.

The current study has examined the effect of osteoclasts secretome on the growth and electrical signal

propagation on sensory nerve fibers. We demonstrate new mechanisms by which peripheral innervation is

controlled, namely indicating that the effects are dependent on the extracellular vesicles (EV) released by

osteoclasts, through the epidermal growth factor receptor family targeting, by integrin independent

pathways.

Materials and methods

All animal procedures were approved by the IBMC/INEB ethics committee and by the Portuguese Agency

for Animal Welfare (*Direção-Geral de Alimentação e Veterinária*) in accordance with the EU Directive

(2010/63/EU) and Portuguese law (DL 113/2013). Mice were housed at 22°C with a 12 h light/dark cycle

with *ad libitum* access to water and food. Adult C57Bl/6 male mice (6-8 weeks-old) were sacrificed in a

carbon dioxide chamber to obtain bone marrow cells, while pregnant females were sacrificed by cervical

dislocation.

Cell culture

Bone marrow cells cultures

Bone marrow stromal cells (BMSC) were isolated from tibiae and femur by flushing the bone marrow with

α-MEM (Gibco, Thermo Fisher Scientific, Waltham, MA, USA) containing 10% (v/v) heat-inactivated (30

min at 56°C) foetal bovine serum (FBS, Gibco, Thermo Fisher Scientific) and 1% (v/v)

penicillin/streptomycin (P/S, Gibco, Thermo Fisher Scientific). Non-adherent cells were removed after 3

days, and fresh medium was added. Cells were expanded from the colony-forming units for one week.

Afterward, the medium conditioned for 3 days was collected, centrifuged at 140 g, 4°C, 5 min, and stored

at -80°C. The conditioned medium was divided into small aliquots (500 µL - 1 mL) before freezing to avoid

repeated freeze/thaw cycles.

Osteoclast cultures

Bone marrow cells were isolated from tibiae and femur by flushing the bone marrow with α-MEM

containing 10% (v/v) FBS and 1% (v/v) P/S. To generate primary osteoclast precursors, the bone marrow

mononuclear cell suspension was treated with red blood cells lysis buffer (ACK lysing buffer, #A1049201,

Gibco, Thermo Fisher Scientific) for 1 min at room temperature (RT) and, after centrifugation, cells were

plated in 10 cm diameter Petri dishes with 10 ng/mL macrophage colony-stimulating factor (M-CSF,

PeproTech, London, UK) for 24 h. Afterward the M-CSF concentration was increased to 30 ng/mL for an

additional 3 days (pre-osteoclasts). Conditioned medium from osteoclasts precursors was collected,

centrifuged at 140 g, 4°C, 5 min, and stored at -80°C. Adherent cells were then detached with a cell scrapper

and seeded at a density of 5 x 10^4 cells/cm^2 in the presence of 30 ng/mL M-CSF and 100 ng/mL receptor

activator of nuclear factor kappa-B ligand (RANKL, PeproTech) (Bartell et al., 2014; Sousa et al., 2016).

Cells were also seeded on top of bone slices (boneslices.com, Denmark) to collect the secretome from active

resorbing osteoclasts. Cells were maintained at 37°C in a humidified atmosphere of 5% CO_2 for 3-4 days

and then osteoclast conditioned medium was collected and stored at -80°C. Evaluation of the osteoclast

differentiation in supplemental data (Figure S1).

Dorsal root ganglia (DRG) culture

Embryonic DRG were obtained from 16 to 18 days-old (E16-18) murine embryos. Embryos were harvested

and maintained on ice-cold Hank's balanced salt solution (HBSS, Invitrogen). Embryonic ganglia were

accessed through the dorsal side of the embryo after spinal cord removal. The meninges were cleaned from

the isolated DRG, and the roots were cut. The ganglia were kept in cold HBSS until seeding. DRG were

isolated under the stereoscopic magnifier and seeded into the lower wells of a 15-well µ-Slide Angiogenesis

plate (#81506, Ibidi, Martinsried, Germany). Fibrin hydrogels were formed in the plates by applying equal

volumes of a solution of plasminogen-free fibrinogen, pooled from human plasma, and a thrombin solution

containing $CaCl_2$ and aprotinin (final concentration of fibrin components: 6 mg/mL fibrinogen (Sigma-

Aldrich); 2 NIH U/mL thrombin from human plasma (Sigma-Aldrich); 2.5 mM $CaCl_2$ (Sigma-Aldrich); 10

µg/mL aprotinin (Sigma-Aldrich)). Before being used in the preparation of fibrin gels, fibrinogen was

dissolved in ultrapure water, dialyzed against tris-buffered saline (TBS, pH 7.4), sterile-filtered and diluted

to 12 mg/mL with sterile TBS. The fibrin gel was allowed to polymerize for 15 min at 37°C in a 5% CO_2

humidified incubator, before the addition of culture media. DRG were cultured with neurobasal medium

supplemented with 2% v/v B-27 Serum-Free Supplement® (B-27, Invitrogen), 60 µM 5-fluoro-2'-

deoxyuridine (FDU, Sigma-Aldrich), 25 mM glucose (Sigma-Aldrich), 1 mM pyruvate (Sigma-Aldrich),

50 ng/mL 7S NGF (Calbiochem, Merck Millipore), 2 mM glutamine (BioWittacker, Lonza, Basel,

Switzerland) and 1% P/S. Embryonic DRG explant cultures were left undisturbed for 24 h. Afterward,

neurobasal media was replaced by the conditioned media collected from BMSC and osteoclast cultures at

different stages of differentiation (200 mg of total protein/well) for an additional three days. Controls with

neurobasal supplemented with NGF and α-MEM supplemented with M-CSF and RANKL (no contact with

cells) were performed.

Quantification of axonal growth

Axonal outgrowth was quantified after 72 h of treatment with conditioned medium and controls. The

embryonic DRG samples were fixed with 2% PFA in PBS for 10 min followed by 10 min at 37°C with 4%

of PFA in PBS with 4% sucrose. Ganglia were permeabilized with 0.25% (v/v) Triton X-100 in PBS and

incubated, for 30 min at RT, with blocking solution composed of 5% v/v normal goat serum (Invitrogen)

and 5% v/v FBS in PBS. DRG cultures were incubated with an antibody directed against the neuronal-

specific marker βIII tubulin (Promega, Madison, WI, United States) diluted 1:2000 in blocking solution,

overnight at 4°C. Afterward, cells were washed and incubated for 1 h at RT with the secondary antibody

(Alexafluor 488/568, Invitrogen). Images were captured either with a confocal laser scanning microscope

(CLSM) Leica SP2 AOBS SE equipped with LCS 2.61 software (Leica Microsystems, Wetzlar, Germany)

at the i3S Advanced Light Microscopy Unit or with an IN Cell Analyzer 2000 equipped with IN Cell

Investigator software (GE Healthcare, Little Chalfont, UK) at the i3S BioSciences Screening Unit. Radial

outgrowth, defined as the area between the ganglion edge and the outgrowth front, was determined. The

outgrowth area was computed according to Bessa *et al.* (Bessa et al., 2013).

Quantification of secretome-neurotrophins by enzyme-linked immunosorbent assay (ELISA)

Conditioned media from osteoclasts and BMSC were concentrated 10 times by centrifugation at 4000 g for

45 min, using 3 kDa MW cut-off filter units (Merck Millipore). In order to detect and quantify the amount

of NGF, BDNF, neurotrophin-3 (NT-3) and NT4/5 in the conditioned medium, a multi-neurotrophin rapid

screening ELISA kit (#BEK-2231, Tebu-bio, France) was used according to the manufacturers' protocol.

The concentration of netrin-1 in the concentrated conditioned media was measured using the ELISA

development kit (EKC37454, Biomatik), also according to the manufacturer's instructions.

Analysis of mature osteoclast total secretome by information dependent acquisition (IDA) / liquid

chromatography – mass spectrometry (LC-MS)

Protein identification was performed by nanoLC-MS/MS. This equipment comprises an Ultimate 3000

liquid chromatography system coupled to a Q-Exactive Hybrid Quadrupole-Orbitrap mass spectrometer

(Thermo Scientific, Bremen, Germany). Samples were loaded onto a trapping cartridge (Acclaim PepMap

C18 100Å, 5 mm x 300 µm i.d., 160454, Thermo Scientific) in a mobile phase of 2% ACN, 0.1% FA at 10

µL/min. After 3 min loading, the trap column was switched in-line to a 50 cm by 75µm ID EASY-Spray

column (ES803, PepMap RSLC, C18, 2 µm, Thermo Scientific, Bremen, Germany) at 300 nL/min.

Separation involved mixing A: 0.1% FA, and B: 80% ACN, with the following gradient: 5 min (2.5% B to

10% B), 120 min (10% B to 30% B), 20 min (30% B to 50% B), 5 min (50% B to 99% B) and 10 min (hold

99% B). Subsequently, the column was equilibrated with 2.5% B for 17 min. Data acquisition was

controlled by Xcalibur 4.0 and Tune 2.9 software (Thermo Scientific, Bremen, Germany).

The mass spectrometer was operated in data-dependent (dd) positive acquisition mode alternating between

a full scan (m/z 380-1580) and subsequent HCD MS/MS of the 10 most intense peaks from full scan

(normalized collision energy of 27%). The ESI spray voltage was 1.9 kV. Global settings: use lock masses

best (m/z 445.12003), lock mass injection Full MS, full-width half-maxima (FWHM) 15s. Full scan

settings: 70k resolution (m/z 200), AGC target 3e6, maximum injection time 120 ms. dd settings: minimum

AGC target 8e3, intensity threshold 7.3e4, charge exclusion: unassigned, 1, 8, >8, peptide match preferred,

exclude isotopes on, dynamic exclusion 45 s. MS2 settings: microscans 1, resolution 35k (m/z 200), AGC

target 2e5, maximum injection time 110 ms, isolation window 2.0 m/z, isolation offset 0.0 m/z, spectrum

data type profile.

The raw data was processed using the Proteome Discoverer software (Thermo Scientific) and compared

against the UniProt database for *Mus musculus* taxonomic selection. The Sequest HT search engine was

used to identify tryptic peptides. The ion mass tolerance was 10 ppm for precursor ions and 0.02 Da for

fragment ions. Maximum allowed missing cleavage sites was set to 2. Cysteine carbamidomethylation was

defined as constant modification. Methionine oxidation and protein N-terminus acetylation were defined

as variable modifications. Peptide confidence was set to high. The processing node Percolator was enabled

with the following settings: maximum delta Cn 0.05; decoy database search target FDR 1%, validation

based on q-value.

Network analysis was performed using the Cytoscape software (version v3.7.1) (Cytoscape Consortium,

San Diego, CA, US)) with the plugins ClueGO (version v2.5.4) and CluePedia (version v1.5.4). We used

ClueGO's default settings: merge redundant groups with >50.0% overlap; the minimum GO level used was

0 and the maximum GO level was 2; statistical test used was "Enrichment/Depletion (Two-sided

hypergeometric test)", Bonferroni step down p-value correction; Kappa Score Threshold was 0.5; and

number of genes was set at 3 with a minimum percentage at 4.0.

Analysis of Phospho-Receptor Tyrosine Kinase (RTK) activation

A proteome profiler mouse phospho-RTK array kit (#ARY014, R&D system, Minneapolis, MN, USA) was

used to quantify the phosphorylation level of 39 RTKs. After 72 h of treatment with conditioned medium

and controls, protein lysate of DRG was quantified and analyzed. The protein content (from 6-10 DRG) of

the same condition, from each independent experiment, was pooled together. According to the

manufacturer's instructions, for the array analysis, the same amount of protein was added to each

membrane. Each array membrane was exposed to X-ray film using a chemiluminescence detection system

(Amersham, GE Healthcare). The film was scanned using Molecular Imager GS800 calibrated densitometer

(Bio-Rad, Hercules, CA, USA) and pixel density was quantified using Quantity One 1-D Analysis

Software, v 4.6 (Bio-Rad). The results were presented as the mean spot intensity which corresponds to the

mean of the two spots for each receptor within the same membrane array.

Pharmacological inhibition of epidermal growth factor receptor (EGFR) and ErbB2

Embryonic DRG were cultured in 15-well slides for 24 h as previously described. Erlotinib, an EGFR

inhibitor, was added to the conditioned medium at 10 nM, 100 nM, 1 µM, 10 µM and 100 µM and tested

on DRG cultures during 72 h. Trastuzumab (Herceptin®), a monoclonal antibody against the ErbB2

receptor, was kindly donated by the Instituto Português de Oncologia do Porto (IPO, Porto, Portugal).

Trastuzumab was tested in DRG cultures at a concentration of 10 μg/mL in conditioned medium.

Afterward, axonal outgrowth was measured as previously described.

β1 integrin subunit function blocking

To assess the contribution of the β1 integrin subunit on the phosphorylation of EGFR and osteoclast-

induced axonal sprouting, stimulation of DRG with osteoclast secretome was conducted in the absence or

presence of β1 integrin function-blocking antibodies, as previously described in (J. Silva et al., 2017). For

this purpose, monoclonal antibodies against β1 (clone Ha2/5, BD Pharmigen; 10 μg/mL) integrin subunits

were added to the osteoclast secretome and incubated with DRG at 37°C during 72 h. To evaluate the

contribution of non-specific antibody interactions, incubation with isotype-matched control β1 antibodies

(hamster IgM λ1, Pharmigen; 10 μg/mL) was also performed. At the end of this period, the wells were

washed and fixed for immunocytochemistry or the protein collected for RTK phosphorylation array

experiments.

Extracellular vesicles (EV) depletion from the osteoclast secretome and characterization

Osteoclasts were isolated and differentiated as described in previous sections. After seeding on 48 well

plates, cells were cultured with 10% EV-depleted FBS. 24h prior media collection, FBS was reduced to

1%. All steps for the EV depletion were conducted under sterile conditions and in line with the published

Minimal information for studies of extracellular vesicles 2018 guidelines (Théry et al., 2018) and as

described in (Théry et al., 2006). Briefly, the secretome was collected, centrifuged 1 000g 10 min to clear

the cell debris, 2 000g for 10 min, followed by 10 000g for 30 min. The supernatant was then centrifuged

at 100 000g using 70Ti rotor (Beckman Coulter Genomics), for 120 min. The pellet containing exosomes

was then resuspended in PBS and stored at -80°C. All centrifugation steps were performed at 4°C (Tassew

et al., 2017). The supernatant was stored -80°C to perform the experiments comprising the EV-depleted

osteoclast secretome.

Western blot (WB) analysis was performed to the protein content from the EV enriched fraction. Protein

was quantified by DC Protein Assay kit (Bio-Rad). The same amount of protein (15 μg) of the EV enriched

fraction and the corresponding supernatant were prepared in non-reducing loading buffer, denatured at 95°C

for 5 min, and loaded in 10% polyacrylamide SDS-PAGE gels. Resolved proteins were wet-transferred to

nitrocellulose membranes, and membranes blocked with non-fat dry milk 5% solution. Membranes were

probed overnight at 4°C with hamster anti-mouse CD81 antibody, clone Eat2 (MCA1846GA, Bio-Rad).

Membranes were probed with HRP-conjugated secondary antibody (GE Healthcare), incubated with ECL

substrate, and chemiluminescence signal was detected with autoradiographic films (all from GE

Healthcare). Films were scanned on a GS-800 calibrated imaging densitometer (Bio-Rad).

The osteoclast-derived EV enriched fraction was further characterized by nanoparticle tracking analysis

(NTA) and transmission electron microscopy (TEM), as previously described (A. M. Silva et al., 2017).

Briefly, for size and particle concentration evaluation, EV suspensions were diluted 1:500 in filtered PBS

1x and analyzed by NTA in a NanoSight NS300 device with NTA3.0 software. For the TEM negative

staining, 10 µL of samples were mounted on Formvar/carbon film-coated mesh nickel grids (Electron

Microscopy Sciences, Hatfield, PA, USA) and left standing for 2 min. The liquid in excess was removed

with filter paper, and 10 µL of 1% uranyl acetate were added on to the grids and left standing for 10 s, after

which, liquid in excess was removed with filter paper. Visualization was carried out on a JEOL JEM 1400

TEM at 120 kV (Tokyo, Japan). Images were digitally recorded using a CCD digital camera Orious 1100W

(Tokyo, Japan) at the i3S Scientific Platforms Histology and Electron Microscopy.

Microelectrode–microfluidic cultures and electrophysiology recordings

The microfluidic devices were fabricated by mixing the polydimethylsiloxane (PDMS) elastomer

(Sylgard® 184, DowCorning) with curing agent (10:1, w/w), degassed and cured over the mold at 65°C for

2 h. Custom microElectrode-microFluidic (µEF) devices were prepared as described previously (Lopes et

al., 2018). Briefly, PDMS microfluidic chambers were aligned on top of microelectrode array (MEA) chips

(MultiChannel Systems MCS GmbH, Germany), with 252 recording electrodes (30 µm in diameter and

pitch of 100 µm) organized in a 16×16 grid. Microfluidic chambers had an appropriate microchannel

spacing for compartmentalization and probing of axonal activity. Microfluidic chambers were also adapted

by adding an extra smaller reservoir (Ø 3 mm), which allowed the seeding of the DRG in a central position

to the electrode matrix (Neto et al., 2014). µEF devices were composed of two separate compartments

connected by 16 microchannels of 700 µm length × 9.6 µm height × 10 µm width. Each microchannel was

probed with 7 electrodes, thus, every axon extending to the axonal compartment was electrophysiologically

monitored. After mounting, µEFs were sequentially coated with poly-D-lysine (PDL) (0.01 mg/ml)

(Corning) and laminin (0.01 mg/ml) (Sigma-Aldrich). The unbound laminin was aspirated, and chambers

were refilled with complete neurobasal medium and left to equilibrate at 37 °C. Isolated embryonic DRG

explants were placed and cultured as described before. DRG explants extended axons to the axonal

compartment within the first 5 days *in vitro* (DIV). Then, treatments and recordings were performed at 6-7

DIV. This time point was chosen following preliminary studies that showed adequate electrophysiological

maturation and culture viability at this stage (Heiney et al., 2019).

Besides the positive control composed of complete neurobasal supplemented with NGF, three conditions

with osteoclast secretome were tested. The total mature osteoclast secretome, the EV-depleted osteoclast

secretome and secretome from osteoclasts cultured in mineralized substrates.

Recordings at a sampling rate of 20 kHz were performed using a MEA2100 recording system (MCS GmbH,

Germany). In every recording session, the temperature was maintained at 37ºC by an external temperature

controller. After removing the cultures from the incubator, recordings only started after 5 minutes of

habituation to avoid an effect due to mechanical perturbation. Then, a baseline recording (5 minutes) was

obtained. Afterward, the medium from the axonal compartment was gently aspirated and replaced by 100

µl of treatment medium. The larger volume present on the somal compartment maintained a hydrodynamic

pressure difference, inhibiting any flow from the axonal to the somal compartment.

Post-treatment (timepoint 0h) recordings (20-30 minutes) were started as soon as the baseline stabilized

following liquid flow perturbation (less than 1 minute). Cultures were returned to the incubator until 3 hours

after treatment, when a post-treatment recording (timepoint 3h) of 5-10 minutes was performed. The day

after, a final post-treatment recording (timepoint 24h) was performed for 5-10 minutes.

Raw signals were high-pass filtered (200 Hz) and spikes were detected by a threshold set to $5\times$ SD of the

electrode noise. Spike data analysis was carried out in MATLAB R2018a (The MathWorks Inc., USA)

using custom scripts. The mean firing rate (MFR) of each microchannel was calculated by averaging the

MFR of the 5 inner electrodes (typically electrode rows 10-14), due to their superior signal-to-noise ratio.

Microchannels with an MFR of at least 0.1 Hz in a given time point at 6 DIV were considered as active and

included in the analysis.

Statistical analysis

All experiments were run in triplicate and repeated at least three times. Data analysis was performed using

GraphPad Prism 8.20 for Windows (GraphPad Software, San Diego CA, USA). Statistical differences

between groups were calculated using one-way analysis of variance, more precisely, the non-parametric

Kruskal-Wallis test followed by Dunns post test for multiple comparisons. The non-parametric Mann-

Whitney t-test was used to identify statistical differences when only two groups where being compared.

Differences between groups were considered statistically significant when $*0.01 < p < 0.05$, $** 0.001 < p$

< 0.01, $*** p < 0.001$, $**** p < 0.0001$.

Results

Sensory neurons outgrowth under osteoclasts effect is neurotrophin-independent

To understand the response of peripheral sensory nerves to the cells present in the bone compartment we

assessed the effect of the secretome from osteoclasts lineage, at different stages of differentiation, in axonal

growth and sprouting of dorsal root ganglia (DRG). We compared these effects with the neurotrophic effect

of bone marrow stromal cells (BMSC).

Organotypic explants of embryonic DRG cultures were exposed to the secretome derived from BMSC,

osteoclast precursors (pre-OC) and mature osteoclasts (OC) populations. The mature osteoclast secretome

demonstrated an approximate 3-fold stronger influence on sensory neurons growth than the secretome from

BMSC (Figure 1A-B), previously demonstrated to have a positive neurotrophic potential (Gu et al., 2012,

2010). The DRG treated with osteoclastic lineage conditioned media, independently of the osteoclasts

maturation status (pre-OC and mature OC), showed significantly higher axonal network when compared to

the neurobasal-nerve growth factor (neurobasal (+NGF)) positive control (Figure 1A-B). Explant DRG

cultures were also treated with osteoclast differentiation medium (OCm) to rule out the effect of receptor

activator of nuclear factor kappa-B ligand (RANKL) and macrophage colony-stimulating factor (M-CSF)

present in the medium and known to modulate axonal outgrowth (Gutierrez et al., 2013; Imai and Kohsaka,

2002). No significant differences were observed when compared to neurobasal control media.

BMSC are known to promote axon growth through the secretion of neurotrophins (Cantinieaux et al., 2013;

Gu et al., 2012, 2010) and osteoclasts where described to induce nerve outgrowth through netrin-1 action

(Ni et al., 2019; Zhu et al., 2019). To assess if the osteoclast secretome observed effect follow similar

mechanisms, we measured in the conditioned medium the levels of NGF, BDNF, NT-3, NT-4/5 and netrin-

1, by enzyme-linked immunosorbent assay (ELISA). NGF, BDNF, and NT-3 were not detected in the

mature osteoclast secretome, despite our observation of higher neurite outgrowth on embryonic DRG

explant cultures. NT-4/5 was detected in very small concentrations for the three different secretomes, while

NGF and BDNF were detected in the BMSC secretome (Figure 1C). Netrin-1 was not detected in the

different secretomes (data not shown).

Figure 1: **Sensory neurons axonal outgrowth under the effect of bone cells secretome. A.** Representative images of DRG outgrowth after treatment (stained for βIII-tubulin, scale bar 500 µm). Neurobasal supplemented with nerve growth factor (NB); bone marrow stromal cells conditioned medium (BMSC); Pre-osteoclast conditioned media (Pre-OC), and mature osteoclast conditioned media (OC) was collected and added to embryonic DRG culture. **B.** Automatic axonal outgrowth quantification of DRG. Data represented as a violin plot. Non-conditioned osteoclast medium (OCm) represented by the dashed line. Statistical analysis was performed *vs.* controls – NB and OCm (*) and between groups (#). **C.** Analysis of secretome neurotrophins. Quantification of NGF, BDNF, NT-3, and NT-4/5 by ELISA on BMSC, Pre-OC and mature OC conditioned media. Data represented as mean±SD.

Epidermal growth factor receptor family signaling pathway is involved in the axonal sprouting of DRG organotypic cultures under osteoclasts secretome stimulation

To understand the mechanisms activated in the context of axonal outgrowth under osteoclasts stimulation, we determined the phosphorylation level (and therefore, activation level) of receptors tyrosine kinase (RTK) which have been implicated in neuronal development, growth, survival and axonal regeneration (Vigneswara et al., 2012).

Epidermal growth factor receptor (EGFR), ErbB2, and platelet-derived growth factor receptor alpha (PDGFRα) displayed higher activation levels (Figure 2). Interestingly, under osteoclast stimulation, there was increased activation of the ErbB2 receptor in DRG neurons when compared to treatment with BMSC (Figure 2B). ErbB2, an orphan receptor with no characterized ligand, can be activated by spontaneous homodimer formation or by heterodimerization with another ligand-bound or epidermal growth factor (EGF) family transactivated receptor (Macdonald-Obermann et al., 2013; Olayioye, 2000). The high levels of EGFR phosphorylation suggest heterodimerization of EGFR/ErbB2 leading to pathway activation. Low levels of TrkA, TrkB and TrkC phosphorylation were observed further confirming the low contribution of NGF, BDNF, NT-3 and NT-4/5 neurotrophins (Figure 2A). In the context of nerve repair, these results are in agreement with the literature where ErbB receptors expression was shown to be increased in DRG upon

lesion (Mizobuchi and Kanzaki, 2013). Therefore, osteoclasts might promote axonal outgrowth through

EGFR family signaling, described to be involved in neuronal repair.

Figure 2: Analysis of activated receptor tyrosine kinases (RTK) in DRG cultures. A. Phosphorylation
levels of RTK on DRG after treatment with with different secretomes – BMSC; Pre-OC and mature OC.
Heatmap representing the relative spot intensity for the activated receptors calculated from the pixel density
in the X-ray films. **B.** X-ray films of the arrays performed. The graphic represents the mean spot intensity
of the most phosphorylated receptors in embryonic DRG treated with different secretomes – BMSC; Pre-
OC and mature OC, showing the major activation of two different families: epidermal-growth factor
receptors (EGFR), platelet-derived growth factor (PDGF) receptor alpha. For array analysis, protein lysate
from a 6–10 DRG in each condition, from three independent experiments (n = 3), was pooled. One array
was used per condition.

To understand the contribution of EGFR and ErbB2 signaling in osteoclast-induced axonal outgrowth,

receptor-mediated inhibition using pharmacological blockers was performed (Figure 3).

Erlotinib is an EGFR inhibitor that reversibly binds to the intracellular tyrosine kinase domain of the

receptor, blocking EGFR over ErbB2. At high concentrations it inhibits both receptor kinases (Hynes and

Lane, 2005; Macdonald-Obermann et al., 2013; Wood et al., 2004). Our results show that the neurotrophic

effect of osteoclasts was reduced in the presence of the high Erlotinib concentration (Figure 3A-B), without

compromising the cell viability and metabolic activity (supplemental data Figure S2), suggesting that both

EGFR/EGFR homodimers and EGFR/ErbB2 heterodimers might contribute to the osteoclast-mediated

effect in axonal outgrowth.

To unravel the input of ErbB2 in osteoclast neurotrophic effect the monoclonal antibody Trastuzumab (also

known as Herceptin®) was used against the extracellular domain (Nielsen et al., 2013). Upon treatment

with Trastuzumab, no differences in the outgrowth area of the sensory neurons were observed (Figure 3C-

D), indicating that sole inhibition of ErbB2 signaling fails to circumvent the osteoclast secretome stimuli.

These results show that osteoclast-mediated axonal outgrowth required both EGFR- and ErbB2 receptor-

dependent pathways.

Figure 3: Pharmacological inhibition of EGFR and ErbB2. A. Quantification of axonal outgrowth of
sensory neurons blocked with epidermal growth factor receptor (EGFR) inhibitor – Erlotinib. **B.**
Representative images of DRG treated with different concentrations of Erlotinib (βIII tubulin in green and
nuclei in blue, scale bar 500 µm). **C.** Quantification of axonal outgrowth of sensory neurons blocked with
ErbB2 receptor inhibitor – Trastuzumab. **D.** Representative images of DRG treated with different
concentrations of Trastuzumab (βIII tubulin in green and nuclei in blue, scale bar 500 µm). Data represented
as box & whiskers (centerline is plotted at the median. Whiskers represent minimum to maximum range)
*p≤0.05.

EGFR phosphorylation is independent of β1 integrin subunit activation

Integrin-mediated adhesion can enhance signaling pathways by direct phosphorylation of growth factor receptors. β1 integrin cytoplasmic domain was demonstrated to trigger integrin-dependent EGFR phosphorylation (Moro et al., 2002). Therefore, we decided to explore further if β1 integrin activation could contribute to EGFR family phosphorylation and synergistically potentiate axonal growth.

An antibody-based β1 integrin subunit blocker and the respective isotype were added to the osteoclast secretome, and axonal sprouting was measured. No differences in the axonal outgrowth was observed when the anti-β1 integrin isotype was added to the osteoclast secretome (supplemental Figure S3). Regarding the functional blocker, the treatment induced a significant decrease in the axonal outgrowth area when compared to the control (Figure 4A-B). Nevertheless, no alterations of EGFR family phosphorylation were observed when the activity of β1 integrin was reduced (Figure 4C). These results were further supported by immunocytochemistry where phosphorylated EGFR was detected independently on the condition analyzed (Figure 4D).

Figure 4: Effect of β1 integrin blocker in axonal sprouting and EGFR phosphorylation. A.
Quantification of axonal sprouting of DRG stimulated with osteoclast secretome and osteoclast secretome
with the functional anti-β1 integrin. Data represented as box & whiskers (median, whiskers represent
minimum to maximum range), ****p≤0.0001. **B.** Representative images of DRG treated with osteoclast
secretome and osteoclast secretome with the functional anti-β1 integrin. Staining for βIII tubulin in green
and nuclei in blue, scale bar 500 µm. **C.** Levels of EGFR family phosphorylation assessed by RTK array.
Values normalized by the neurobasal control. **D.** Representative images of EGFR phosphorylation by
immunocytochemistry after treatment with NGF-supplemented neurobasal, osteoclast secretome,
osteoclast secretome with the functional anti-β1 integrin and osteoclast secretome with anti-β1 integrin

423   isotype. Phosphorylated EGFR (pEGFR) in red, βIII tubulin in green and nuclei in blue (left column); zoom

424   in images of the isolated channel for the pEGFR (right column). Scale bar 100 μm.

426   Soluble EGFR/ErbB2 activators were undetected in the osteoclast-derived secretome

EGF and TGFα ligands, that preferentially bind to the EGFR homodimers and EGFR/ErbB2 heterodimers

(Hynes and Lane, 2005), activating the signaling cascade, were not detected in the osteoclasts secretome

by ELISA (data not shown). Therefore, to understand which ligands were present in the osteoclast

secretome that could directly or indirectly activate the signaling cascade, we proceed to a higher sensitive

technique and performed proteomic analysis by liquid chromatography-tandem mass spectrometry (LC-

MS/MS) of osteoclast serum-free secretome collected at two different time points: 6 h and 24 h.

Over four hundred proteins were identified in the osteoclast serum-free secretome. The complete set of

identified proteins is available as supplemental data (supplemental Table S1). Protein identification

revealed oscillations in the relative concentration of the soluble factors throughout the conditioning period.

As expected, the higher amount of the identified proteins at 24 h were increased in relative concentration

when compared to 6 h.

Interestingly, the identified proteins were not only assigned to the extracellular region (extracellular matrix

proteins, secreted soluble factors, extracellular vesicles-associated proteins, etc.) but also to synaptic part

(synaptic structural proteins, downstream effectors, cytoskeletal components) in equal percentage,

according to Gene Ontology cellular component terms (ClueGO v2.5.4 + CluePedia (v1.5.4) from the

software Cytoscape (v3.7.1) (Figure 5A). By restricting the analysis to pathways with a higher significance

(p<0.01) and increasing the specificity level of the cellular component terms, we observed that the majority

of the proteins remained associated to the initial detected clusters (Figure 5B and supplemental table S2).

Importantly, neither EGF nor TGFα were identified among the secreted proteins, which is in accordance

with a previous study reporting briefly the analysis of the osteoclast secretome where no EGFR ligands

were detected (Rody et al., 2017). Nevertheless, our thorough proteomic analysis of the secretome revealed

that the osteoclasts secrete molecules to interact with neuronal and non-neuronal cells to directly influence

neuronal processes. Dihydropyrimidinase-related protein 2 (Dpysl2), also known as collapsin response

mediator protein 2 (Crmp2), here found linked to the synaptic cluster. Moreover, in the extracellular region

cluster, we detected the protein delta homolog 1 (Dlk1) and Serpin F1 (SerpinF1), also known as pigment

epithelium-derived factor (PEDF). Still, none of the identified proteins are described to contribute directly

or indirectly to the EGFR/ErbB2 signaling cascade.

Figure 5: Cytoscape network of the main cellular components associated to osteoclasts-secreted
proteins at 6 h and 24 h. A. Network analysis using the plugin ClueGO (v2.5.4) + CluePedia (v1.5.4) from

the software Cytoscape (v3.7.1) to analyze the cellular components represented by the identified proteins
from osteoclast secretome. Gene ontology (GO) term fusion was applied. Pathways with p value lower than
0.05 were considered. Tree interval was applied from 0 to 1 level. Genes from the different cellular
components are shown in red. **B.** Protein network (cerebral view layout) representing the most significant
clusters associated with the identified proteins. Gene ontology (GO) term fusion was applied. Pathways
with p value lower than 0.01 were considered. Tree interval was applied from 1 to 2 level. The size of the
nodes is indicative of the number of associated proteins. The increase in red color gradient represents higher
statistical significance.

Extracellular vesicles secreted by osteoclasts carry mediators of the EGFR/ErbB2-dependent axonal sprouting

Neurotrophic proteins, are generally believed to be released from cells in a soluble form and act locally or

move through the extracellular milieu, which lead us to perform the proteomic profiling in the total

secretome. However, it is increasingly appreciated that growth factors can be released from cells in or on

the surface of extracellular vesicles (EV)/exosomes (Jaiswal and Sedger, 2019). We, therefore,

hypothesized that osteoclast-derived EV could play a crucial role in the axonal outgrowth. To test this,

DRG were exposed to EV-depleted osteoclast secretome and the axonal sprouting was measured. The EV

enriched fraction was characterized by NTA, WB and by TEM. The analysis of the size and concentration

of the vesicles showed a normal distribution with a mean size of 141.8 ± 2.7 nm and a concentration of 4.90

x 10^{11} particles/ml (supplemental Figure S4). The analyzed fraction stained positive for the CD81 EV-

specific marker (Figure 6A) and by negative staining for TEM, it was possible to visualize the EV present

in the OC-derived secretome (Figure 6B, white arrowheads), ranging from 40 to 200 nm.

A significant decrease in axonal sprouting was observed in the absence of EV (Figure 6C-D). This suggests

that the EV cargo plays a role in the neurotrophic potential of the osteoclast secretome. To understand if

EV depletion modulates the EGFR/ErbB2 phosphorylation levels, that we previously demonstrated to be

involved in the osteoclast-induced axonal outgrowth, we evaluated the phosphorylation state of the EGFR

family. Remarkably, a substantial decrease in EGFR family phosphorylation was observed in the absence

of osteoclast-derived EV (Figure 6E), confirming that the EGFR family activator is present in the EV cargo

and reinforcing the contribution of this signaling pathway to the osteoclast-mediated axonal growth.

Therefore, the EV enriched fraction was further analyzed by LC-MS/MS for the identification of potential

proteins that can directly or indirectly regulate the EGFR/ErbB2 signaling cascade. Over five hundred

proteins were identified (supplemental table S3). New clusters were detected, when comparing to the

previous analysis performed for the whole secretome. Highly significant clusters containing proteins related

to DNA binding, cell-substrate interactions, proteic complexes and membrane-associated proteins were

revealed (supplemental table S4). Very interestingly, the nodes with higher protein contribution remained

the synaptic part (27%) and extracellular region (15%). In addition, according to the proteomic analysis,

SH3 domain-binding glutamic acid-rich-like protein (SH3BGRL) previously described to interact with the

EGFR signalling cascade (Schulze et al., 2005; Taguchi et al., 2011) and promote the dowstream activation

of c-Src (Wang et al., 2016) was detected. The proteins present in the nodes with SH3BGRL were closely

related to several biological processes including developmental growth with high contribution to positive

regulation of cell growth (Figure 6F and supplemental table S5).

Figure 6: Dorsal root ganglia (DRG) axonal network growth is dependent on the osteoclasts
extracellular vesicles (EV). A. Characterization of osteoclasts-derived EV by Western blot using CD81
membrane marker (EV *vs.* Supernatant (SN) EV depleted). **B.** Transmission electron microscopy of
osteoclasts-derived EV (white arrows) by negative staining (100 000x). **C.** Representative images of DRG
treated with osteoclast secretome and EV-depleted osteoclast secretome. Staining for βIII tubulin in green
and nuclei in blue, scale bar 500 µm. **D.** Quantification of axonal sprouting of DRG stimulated with
osteoclast secretome vs. osteoclast secretome lacking the EV. Data represented as box & whiskers (median,
whiskers represent minimum to maximum range), *p≤0.05. **E.** Levels of RTK phosphorylation assessed by
RTK array. Values normalized to the neurobasal control. **F.** Network analysis of osteoclast-derived EV
enriched fraction according to the cellular components. Protein network (edge-weighted spring embedded
layout) representing the most significant clusters associated with the identified proteins. Gene ontology
(GO) term fusion was applied. Pathways with p value lower than 0.01 were considered. Tree interval was
applied from 1 to 2 level. The size of the nodes is indicative of the number of associated proteins. The
increase in red color gradient represents higher statistical significance. High resolution networks provided
as supplemental material.

Electrical activity induced by osteoclast secretome exposure is eradicated upon EV depletion

To unravel the electrophysiological implications of the axonal exposure to the osteoclast secretome, a

combination of substrate-integrated microelectrode arrays (MEAs) with custom-made microfluidic

chambers (Lopes et al., 2018) were used. MEAs enable non-invasive, thus repeatable, recordings of

extracellular action potentials. Although recordings of random DRG cultures have been previously

demonstrated (Black et al., 2019), these are not adapted to the study of peripheral innervation. Here, we

employed microElectrode-microFluidic (µEF) devices (Figure 7A), which allowed us to monitor axonal

activity with great fidelity (Figure 7B) (Heiney et al., 2019; Lopes et al., 2018).

Unlike central nervous system neurons in culture (e.g. hippocampal neurons), DRG did not fire in bursts

but rather exhibited sporadic spontaneous spiking at baseline. Under normal conditions, this relatively low

level of spontaneous activity also occurs *in vivo* (Weng et al., 2012). Regardless of the explant position

(outside or inside the array area), in most cases, we could only detect activity within the microchannels

(Figure 7D). Still, this allowed us to directly compare baseline and post-treatment levels of axonal activity,

as most axons within the microchannels are expected to have extended to the axonal compartment.

DRG terminals were exposed to three conditions comprising osteoclast secretome: i) mature, non-resorbing.

osteoclasts (CM), ii) resorbing osteoclasts on mineralized substrates (bone slices – BS) and iii) EV-depleted

osteoclasts secretome and compared to NGF-containing neurobasal control (NB).

Electrophysiology recordings show that the DRG exposed to osteoclasts secretome present an increased

mean firing rate (MFR) when compared to the baseline recording immediately before treatment (Figure

7D-F). While the control treatment, supplemented with NGF, did not produce a significant effect in the

MFR, the DRG treated with secretome from resorbing osteoblasts (BS) presented a highly significative

effect. Previous reports have shown that medium acidification and the molecules released by osteoclasts

and matrix itself, upon matrix degradation, contribute to the activation of nociceptors on the nerve terminals

(Hiasa et al., 2017; Yoneda et al., 2015a). Here we show that the secretome from mature, non-resorbing,

osteoclasts, also contribute to the electrical activation and signal propagation across sensory neurons.

Excitingly, the effect on the firing rate of the sensory neurons was dramatically reduced upon EV depletion

from the osteoclast secretome (Figure 7D-F). The effect produced by EV-depleted secretome, when

compared to the pre-treatment baseline, was 598% lower than the parallel effect produced by osteoclasts

CM and 1990% lower than the BS secretome (Figure 7F). Therefore, we postulate that the sensory neurons

electrophysiological activity is dependent on the osteoclasts secreted EV, still, the mechanisms that support

this effect deserve further elucidation.

Figure 7: Electrophysiology studies on dorsal root ganglia (DRG) neurons in microfluidic devices stimulated with osteoclasts secretome. A. Phase-contrast microscopy image mosaic of an organotypic dorsal root ganglia (DRG) culture at 7 days *in vitro* (DIV). The whole microelectrode array (MEA) (1.5 × 1.5 mm) active area is shown by a combination of 9 mosaic images (10× objective) from different parts of the culture. A PDMS device composed of 16 microchannels (10 μm width; 700 μm length) is aligned to encompass 7 microelectrodes. Details of axonal morphology can be seen in the somal compartment, microchannels and axonal compartment (scale bar = 200 μm). A schematized version of a microchannel is shown on the right. **B.** Electrophysiological trace of 30 seconds of baseline activity from an electrode (within a microchannel) at 6 DIV and corresponding spike raster plot. Inset shows a single spike. **C.** Schematic diagram of experiments and timeline of recordings. **D.** Representative activity maps of baseline and post-treatment (time 0h) activity for each condition (microchannel area only, electrode rows 9-15). Each pixel corresponds to an electrode and the mean firing rate (MFR) is color-coded. Corresponding raster

plots of 60 seconds of activity are shown below each activity map. Each row corresponds to the spike raster
plot from the central electrode of every microchannel. **E.** Box-and-whiskers plots of the active
microchannels' MFR at baseline and 0h, 3h and 24 post-treatment. Data from 35-61 microchannels from
3-5 independent μEFs. **F.** Before-after plot of every active microchannel after treatment. ns = not
significant, *0.01 < p < 0.05, ** 0.001 < p < 0.01, *** p < 0.001, **** p < 0.0001.

Discussion

We provide evidence on a new mechanism by which peripheral innervation can be coordinated. We show

that sensory nerves outgrowth and electric signal propagation are dependent on the EV secreted by

osteoclasts, the bone resorbing cells. Furthermore, we demonstrate that the axonal sprouting is achieved

through the activation of epidermal-growth factor receptor (EGFR) and ErbB2 receptors, highlighting their

involvement in the crosstalk, independent of β1 integrin function. Furthermore, inhibition of EGFR/EGFR

and EGFR/ErbB2 dimers uncovers the contribution of this signaling cascade to the osteoclast-mediated

axonal outgrowth. We proved that the EV-depleted osteoclast secretome leads to a significant decrease of

neurons firing rate and axonal sprouting, concomitant with a decrease of EGFR/ErbB2 activation levels.

Excitingly, the proteomic analysis of the osteoclast-derived EV cargo shows a high correlation with

synaptic components reinforcing the role on sensory neurons/osteoclast crosstalk.

Different studies point to a parallel between osteoclast activity and sensory neurons outgrowth in

physiological and pathological conditions (Cheung et al., 2009; Clarke, 2008; Gajda et al., 2005; Kumar et

al., 2011; Mediati et al., 2014; Nagae et al., 2006; Ni et al., 2019; Reddy et al., 1999; Riminucci et al., 2003;

Sisask et al., 1996, 2013; Zhu et al., 2019). Several reports have described that during embryonic long bone

development, osteoclasts are found in the process of endochondral ossification, degrading the newly formed

trabecular bone, creating the bone marrow space (Clarke, 2008). In another hand, longitudinal studies on

the innervation process showed nerves appearing in areas with high osteogenic and chondrogenic activity

during embryonic development (Gajda et al., 2005; Sisask et al., 2013, 1996). Thus, the presence of

osteoclasts in stages where innervation occurs suggests a role of these cells in the innervation pattern. Bone

pathologies with increased osteoclast number (Paget's disease (Kumar et al., 2011; Reddy et al., 1999),

fibrous dysplasia (Riminucci et al., 2003), *osteogenesis imperfecta* (Cheung et al., 2009)), and with

increased osteoclast activity (osteoporosis and metastatic bone disease), are frequently associated with

activation of sensory nerves (Mantyh, 2013; Mediati et al., 2014; Nagae et al., 2006). Although the pattern

of innervation in these pathological conditions has only been described in metastatic bone disease, where

exuberant sprouting of sensory nerve fibers has been reported, several studies indicate osteoclasts play a

fundamental role in bone pain (Julius and Basbaum, 2001; Mantyh, 2014; Nagae et al., 2006; Yoneda et

al., 2015b). Recently, Zhu *et al.* have shown that osteoclasts can actively promote axonal growth on

subchondral bone, through netrin-1 signaling, in an osteoarthritis model (Zhu et al., 2019). Moreover,

osteoclasts-derived netrin-1 was also linked to an increase in the sensory innervation of porous endplates

of intervertebral disc (Ni et al., 2019). We were unable to detect NGF, BDNF, NT-3, NT4/5 nor netrin-1 in

the osteoclast secretome by using comparable settings for osteoclast *in vitro* differentiation and netrin-1

quantification (mass spectrometry analysis and ELISA (data not shown)). In light of our results, we

hypothesize that other signaling mechanisms were taking place. Herein we aimed to understand the main

activation mechanisms on DRG explant cultures. Our work highlights new key players in this dynamic

crosstalk. We detected high levels of EGFR and ErbB2 activation on DRG upon osteoclast secretome

stimuli.

EGFR activation is involved in neuron differentiation and survival and has been implicated in responses to

neuronal damage (Mizobuchi and Kanzaki, 2013; Nilsson and Kanje, 2005; Tsai et al., 2010; Xian and

Zhou, 1999). Nevertheless, there is an extensive discussion in the literature concerning the direct

involvement of EGFR activation/inhibition in axonal regeneration (Berry et al., 2011; Douglas et al., 2009;

Liu et al., 2006; Wong and Guillaud, 2004). EGFR phosphorylation has been implicated in signaling

inhibition of axonal growth in central nervous system (Berry et al., 2011; Koprivica et al., 2005; Liu et al.,

2006). Koprivica *et al.* showed that EGFR inhibitors effectively promoted neurite outgrowth from cultured

DRG (Koprivica et al., 2005). In contrast, others reported that EGFR knockdown did not induce neurite

outgrowth of retinal ganglion cells, and inhibitors of EGFR were shown to enhance neurite outgrowth in

the absence of the receptor (Berry et al., 2011). We and others previously demonstrated an increased

expression of the EGFR family in DRG after lesion (Mizobuchi and Kanzaki, 2013; Neto et al., 2017; Xian

and Zhou, 1999). Our data indicate that EGFR signaling has a role in axonal outgrowth promoted by

osteoclast secretome. Upon EGFR inhibition with Erlotinib, described to inhibit both EGFR and ErbB2

receptor kinases at high concentrations (Macdonald-Obermann et al., 2013; Wood et al., 2004), we

observed a significant reduction in the axonal outgrowth area. These findings are consistent with prior

observations that the phosphorylation of EGFR enhances neurite outgrowth (Evangelopoulos et al., 2009;

Goldshmit et al., 2004a, 2004b; Nilsson and Kanje, 2005; Tsai et al., 2010; Xu et al., 2014). The activation

of EGFR in an integrin-dependent and EGFR ligand-independent manner has been described (Bill et al.,

2004), where β1 integrin activation can enhance direct phosphorylation of EGFR (Balanis et al., 2011; Bill

et al., 2004; Moro et al., 2002; Morozevich et al., 2012). We detected a significant decrease in axonal

sprouting upon β1 integrin blocking, yet the EGFR family phosphorylation was not impaired, suggesting

that activation of EGFR signaling cascade is not dependent on the cell-matrix or cell-cell mediated

interactions.

Neurons have been described to be highly responsive to EV and exosomes derived from myelinating cells

(oligodendrocytes and Schwann cells) (Budnik et al., 2016; Frühbeis et al., 2013; Lopez-Verrilli et al.,

2013). Microglial-derived EV were also reported to promote synaptic refinement and instructing neurons

upon inflammatory stimuli (Paolicelli et al., 2019). More recently, neurons were shown to be reactive to

EV derived from other cellular populations, as seen for the response of cortical neurons to mesenchymal

stem cells-derived EV (Xin et al., 2012; Zhang et al., 2017) and neuroblastoma cell line to EV from

adipocyte derived Schwann cell-like (Ching et al., 2018). Osteoclasts secrete EV at the bone

microenvironment in both physiological and pathological conditions (Yuan et al., 2018), which have been

associated as key players underlying the osteoclast-osteoblast communication (Collison, 2017; Huynh et

al., 2016a), tumor proliferation, angiogenesis, and tumor cell survival (Huynh et al., 2016b; Lee et al., 2007;

Rossi et al., 2018).

Here, we firstly described that sensory neurons are responsive to osteoclast-derived EV. Our results

strengthen the importance on the communication between sensory terminals and osteoclasts and unravels

new mechanism underlying the interplay. We indicate that the EV signaling mechanism underlie sensory

processing and nociception signal propagation, largely unknow in the literature (Fowler, 2019).

Remarkably, EV-depleted osteoclast secretome produced not only a significant decrease in axonal growth,

but also a striking reduction in EGFR family phosphorylation. Once again, the proteome of EV cargo

revealed no presence of EGF or TGFα, which is in accordance with recent data reporting the proteomic

profile of the osteoclasts secreted vesicles upon different substrates (Rody et al., 2019). Importantly, the

proteomic screening of the EV cargo, revealed the presence of SH3BGRL, proved to bind to the

intracellular EGFR domain (Chiang et al., 2015; Schulze et al., 2005; Taguchi et al., 2011). More

interestingly, phosphotyrosine interactome of the ErbB-receptor kinase family showed that ErbB2 has few

interaction partners being SH3BGRL one of them on residue Y0923 (Schulze et al., 2005). Noteworthy,

SH3BGRL promote the dowstream activation of c-Src (Wang et al., 2016), which is central to multiple

signal transduction pathways including axonal elongation, adhesion, growth, and cell survival (Irby and

Yeatman, 2000; Robles et al., 2005). The main mechanism underlying the EV interaction with sensory

neurons require further studies, still it has been reported that mesenchymal stem cells-derived EV are

capable of entering cortical neurons both through cell body and axonal terminals (Zhang et al., 2017).

We also show that different electrophysiological patterns are revealed depending on the osteoclast

secretome tested. The greater effect observed in the increasing of the MFR was related to the secretome

from osteoclasts cultured on mineralized substrates, as expected. Osteoclasts were shown to increase the

expression of sodium channel Na_v 1.8 in sensory nerve endings in the endplates leading to spinal

hypersensitivity (Ni et al., 2019). Protons released by the osteoclasts upon matrix degradation are known

to stimulate the acid-sensing ions channels contributing to the activation of nociceptors on the nerve

terminals (Hiasa et al., 2017; Yoneda et al., 2015a). We demonstrate that, not only the products released

through matrix degradation process are mediating the sensory fibers activation, but also the products

secreted by osteoclasts in non-degrading scenario. The EV-depletion resulted in a major decrease on the

MFR reflecting a possible pathway to be addressed to understand the nerves sensitization on bone

microenvironment.

In conclusion, we provide a new mechanism on the sensory nerves sprouting, indicating that the effect is

dependent on the extracellular vesicles (EV) released by osteoclasts, through the epidermal growth factor

receptor family targeting, by integrin independent pathways. We show different electrophysiology patterns

being triggered in the presence of osteoclasts secretome and the abolishment of sensory neurons firing rate

in EV-depleted conditions. Clearly, besides the activation of nociceptors through microenvironment

acidification or soluble molecules, osteoclasts secreted vesicles can be implicated in cancer-associated bone

pain through the control of nerve sprouting and electrical signal propagation, consequently remodeling the

pathological innervation pattern and activity. Furthermore, studies aiming to manipulate osteoclast-derived

EV cargo can be employed to further dissect the bone-peripheral nervous system communication. This is a

subject that deserves further investigation to i) control/reduce the formation of neuroma-like structures and

ii) specify new candidates as potential targets for bone pain therapies.

Author contributions

Conceptualization, Estrela Neto, Meriem Lamghari and Paulo Aguiar; Methodology, Estrela Neto, Luís Leitão, José Mateus, Daniela Monteiro Sousa and Francisco Conceição; Investigation, Estrela Neto, Luís Leitão, José Mateus, Daniela Monteiro Sousa and Cecília Juliana Alves; Writing – Original Draft, Estrela Neto, José Mateus and Meriem Lamghari; Writing – Review & Editing, all authors; Funding Acquisition, Meriem Lamghari and Paulo Aguiar; Supervision, Richard Oreffo, Jonathan West and Meriem Lamghari.

Acknowledgments

The authors would like to acknowledge to João H. Morais-Cabral from i3S for the fruitful scientific discussions and input for this work. To Susana Santos, Andreia Silva and José Teixeira for the help with EV isolation protocol. To Maria José Oliveira, José Carlos Machado and Ana Paula Pêgo groups from i3S for the phospho-EGFR antibody, Erlotinib and anti-β1 integrin antibodies, respectively. The microfluidic design adapted to the commercial microelectrode arrays used in this study were fabricated at INESC - Microsystems and Nanotechnologies, Portugal, under the supervision of João Pedro Conde and Virginia Chu.

This work was financed by FEDER - Fundo Europeu de Desenvolvimento Regional funds through the COMPETE 2020 - Operacional Programme for Competitiveness and Internationalization (POCI), Portugal 2020, and by Portuguese funds through FCT/MCTES in the framework of the project SproutOC: POCI-01-0145-FEDER-030158, PTDC/MED-PAT/30158/2017. LL, JM and FC are recipients of Ph.D. fellowships (SFRH/BD/109686/2015, PD/BD/135491/2018 and SFRH/BD/128771/2017, respectively). DMS is recipient of Post-Doc fellowship (SFRH/BPD/115341/2016). The authors acknowledge the support of the i3S Scientific Platforms BioSciences Screening, Bioimaging, and Histology and Electron Microscopy, members of the national infrastructure PPBI - Portuguese Platform of Bioimaging. The mass spectrometry technique was performed at the Proteomics i3S Scientific Platform with the assistance of Hugo Osório.

The authors declare no competing interests.

References

Balanis, N., Yoshigi, M., Wendt, M.K., Schiemann, W.P., Carlin, C.R., 2011. β3 integrin-EGF receptor cross-talk activates p190RhoGAP in mouse mammary gland epithelial cells. Mol. Biol. Cell 22, 4288–301.

Bartell, S.M., Kim, H.-N., Ambrogini, E., Han, L., Iyer, S., Serra Ucer, S., Rabinovitch, P., Jilka, R.L., Weinstein, R.S., Zhao, H., et al., 2014. FoxO proteins restrain osteoclastogenesis and bone resorption by attenuating H2O2 accumulation. Nat. Commun. 5, 3773.

Berry, M., Ahmed, Z., Douglas, M.R., Logan, A., 2011. Epidermal growth factor receptor antagonists and CNS axon regeneration: Mechanisms and controversies. Brain Res. Bull. 84, 289–299.

Bessa, S., Quelhas, P., Amaral, I.F., 2013. Automatic Quantification of Cell Outgrowth from Neurospheres. In: Sanches, J.M., Micó, L., Cardoso, J.S. (Eds.), Pattern Recognition and Image Analysis: 6th Iberian Conference, IbPRIA 2013, Funchal, Madeira, Portugal, June 5-7, 2013. Proceedings. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 141–148.

Bill, H.M., Knudsen, B., Moores, S.L., Muthuswamy, S.K., Rao, V.R., Brugge, J.S., Miranti, C.K., 2004. Epidermal Growth Factor Receptor-Dependent Regulation of Integrin-Mediated Signaling and Cell Cycle Entry in Epithelial Cells. Mol. Cell. Biol. 24, 8586–8599.

Black, B.J., Atmaramani, R., Plagens, S., Campbell, Z.T., Dussor, G., Price, T.J., Pancrazio, J.J., 2019. Emerging neurotechnology for antinoceptive mechanisms and therapeutics discovery. Biosens. Bioelectron. 126, 679–689.

Bloom, A.P., Jimenez-Andrade, J.M., Taylor, R.N., Castañeda-Corral, G., Kaczmarska, M.J., Freeman, K.T., Coughlin, K.A., Ghilardi, J.R., Kuskowski, M.A., Mantyh, P.W., 2011. Breast cancer-induced bone remodeling, skeletal pain, and sprouting of sensory nerve fibers. J. Pain 12, 698–711.

Bove, S.E., Flatters, S.J.L., Inglis, J.J., Mantyh, P.W., 2009. New advances in musculoskeletal pain. Brain Res. Rev. 60, 187–201.

Brohlin, M., Kingham, P.J., Novikova, L.N., Novikov, L.N., Wiberg, M., 2012. Aging Effect on Neurotrophic Activity of Human Mesenchymal Stem Cells. PLoS One 7, 67–75.

Budnik, V., Ruiz-Cañada, C., Wendler, F., 2016. Extracellular vesicles round off communication in the nervous system. Nat. Rev. Neurosci. 17, 160–72.

Cantinieaux, D., Quertainmont, R., Blacher, S., Rossi, L., Wanet, T., Noël, A., Brook, G., Schoenen, J., Franzen, R., 2013. Conditioned Medium from Bone Marrow-Derived Mesenchymal Stem Cells Improves Recovery after Spinal Cord Injury in Rats: An Original Strategy to Avoid Cell Transplantation. PLoS One 8.

Chao, M. V, 2003. Neurotrophins and their receptors: A convergence point for many signalling pathways. Nat. Rev. Neurosci. 4, 299–309.

Chartier, S.R., Thompson, M.L., Longo, G., Fealk, M.N., Majuta, L.A., Mantyh, P.W., 2014. Exuberant sprouting of sensory and sympathetic nerve fibers in nonhealed bone fractures and the generation and maintenance of chronic skeletal pain. Pain 155, 2323–2336.

Chen, Hao, Hu, B., Lv, X., Zhu, S., Zhen, G., Wan, M., Jain, A., Gao, B., Chai, Y., Yang, M., et al., 2019. Prostaglandin E2 mediates sensory nerve regulation of bone homeostasis. Nat. Commun. 10.

Chen, X., Katakowski, M., Li, Y., Lu, D., Wang, L., Zhang, L., Chen, J., Xu, Y., Gautam, S., Mahmood, A., et al., 2002. Human bone marrow stromal cell cultures conditioned by traumatic brain tissue extracts: Growth factor production. J. Neurosci. Res. 69, 687–691.

Chenu, C., 2004. Role of innervation in the control of bone remodeling. J. Musculoskelet. Neuronal Interact. 4, 132–134.

Chenu, C., Marenzana, M., 2005. Sympathetic nervous system and bone remodeling. Jt. Bone Spine 72, 481–483.

Cherruau, M., Facchinetti, P., Baroukh, B., Saffar, J.L., 1999. Chemical sympathectomy impairs bone resorption in rats: a role for the sympathetic system on bone metabolism. Bone 25, 545–551.

Cheung, M.S., Glorieux, F.H., Rauch, F., 2009. Large osteoclasts in pediatric osteogenesis imperfecta patients receiving intravenous pamidronate. J. Bone Miner. Res. 24, 669–674.

Chiang, C.-Y., Pan, C.-C., Chang, H.-Y., Lai, M.-D., Tzai, T.-S., Tsai, Y.-S., Ling, P., Liu, H.-S., Lee, B.-F., Cheng, H.-L., et al., 2015. SH3BGRL3 Protein as a Potential Prognostic Biomarker for Urothelial Carcinoma: A Novel Binding Partner of Epidermal Growth Factor Receptor. Clin. Cancer Res. 21, 5601–5611.

Ching, R.C., Wiberg, M., Kingham, P.J., 2018. Schwann cell-like differentiated adipose stem cells promote neurite outgrowth via secreted exosomes and RNA transfer. Stem Cell Res. Ther. 9, 266.

Clarke, B., 2008. Normal Bone Anatomy and Physiology. Clin. J. Am. Soc. Nephrol. 3, S131–S139.

Cleeland, C.S., Body, J.-J., Stopeck, A., von Moos, R., Fallowfield, L., Mathias, S.D., Patrick, D.L., Clemons, M., Tonkin, K., Masuda, N., et al., 2013. Pain outcomes in patients with advanced breast

cancer and bone metastases. Cancer 119, 832–838.

Collison, J., 2017. Bone: Extracellular vesicles in bone cell crosstalk. Nat. Rev. Rheumatol. 14, 2–3.

Douglas, M.R., Morrison, K.C., Jacques, S.J., Leadbeater, W.E., Gonzalez, A.M., Berry, M., Logan, A.,
Ahmed, Z., 2009. Off-target effects of epidermal growth factor receptor antagonists mediate retinal
ganglion cell disinhibited axon growth. Brain 132, 3102–3121.

Edoff, K., Hellman, J., Persliden, J., Hildebrand, C., 1997. The developmental skeletal growth in the rat
foot is reduced after denervation. Anat. Embryol. (Berl). 195, 531–8.

Elefteriou, F., 2008. Regulation of bone remodeling by the central and peripheral nervous system. Arch
Biochem Biophys 473, 231–236.

Elefteriou, F., Campbell, P., Ma, Y., 2014. Control of bone remodeling by the peripheral sympathetic
nervous system. Calcif Tissue Int 94, 140–151.

Evangelopoulos, M.E., Weis, J., Krüttgen, A., 2009. Mevastatin-induced neurite outgrowth of
neuroblastoma cells via activation of EGFR. J. Neurosci. Res. 87, 2138–2144.

Fowler, C.D., 2019. NeuroEVs: Characterizing Extracellular Vesicles Generated in the Neural Domain. J.
Neurosci. 39, 9262–9268.

Frühbeis, C., Fröhlich, D., Kuo, W.P., Amphornrat, J., Thilemann, S., Saab, A.S., Kirchhoff, F., Möbius,
W., Goebbels, S., Nave, K.-A., et al., 2013. Neurotransmitter-triggered transfer of exosomes mediates
oligodendrocyte-neuron communication. PLoS Biol. 11, e1001604.

Fukuda, T., Takeda, S., Xu, R., Ochi, H., Sunamura, S., Sato, T., Shibata, S., Yoshida, Y., Gu, Z., Kimura,
A., et al., 2013. Sema3A regulates bone-mass accrual through sensory innervations. Nature 497, 490–
493.

Gajda, M., Litwin, J. a, Cichocki, T., Timmermans, J.P., Adriaensen, D., Timmermanns, J.P., Adriaensen,
D., 2005. Development of sensory innervation in rat tibia: co-localization of CGRP and substance P
with growth-associated protein 43 (GAP-43). J. Anat. 207, 135–144.

Ghilardi, J.R., Freeman, K.T., Jimenez-Andrade, J.M., Mantyh, W.G., Bloom, A.P., Kuskowski, M. a,
Mantyh, P.W., 2010. Administration of a tropomyosin receptor kinase inhibitor attenuates sarcoma-
induced nerve sprouting, neuroma formation and bone cancer pain. Mol. Pain 6, 87.

Goldshmit, Y., Greenhalgh, C.J., Turnley, A.M., 2004a. Suppressor of cytokine signalling-2 and epidermal
growth factor regulate neurite outgrowth of cortical neurons. Eur. J. Neurosci. 20, 2260–2266.

Goldshmit, Y., Walters, C.E., Scott, H.J., Greenhalgh, C.J., Turnley, A.M., 2004b. SOCS2 induces neurite
outgrowth by regulation of epidermal growth factor receptor activation. J. Biol. Chem. 279, 16349–
16355.

Gomez, C., Burt-Pichat, B., Mallein-Gerin, F., Merle, B., Delmas, P.D., Skerry, T.M., Vico, L., Malaval,
L., Chenu, C., 2005. Expression of Semaphorin-3A and its receptors in endochondral ossification:
potential role in skeletal development and innervation. Dev Dyn 234, 393–403.

Gu, W., Zhang, F., Xue, Q., Ma, Z., Lu, P., Yu, B., 2012. Bone mesenchymal stromal cells stimulate neurite
outgrowth of spinal neurons by secreting neurotrophic factors. Neurol. Res. 34, 172–80.

Gu, Y., Wang, J., Ding, F., Hu, N., Wang, Y., Gu, X., 2010. Neurotrophic actions of bone marrow stromal
cells on primary culture of dorsal root ganglion tissues and neurons. J Mol Neurosci 40, 332–341.

Gutierrez, H., Kisiswa, L., O'Keeffe, G.W., Smithen, M.J., Wyatt, S., Davies, A.M., 2013. Regulation of
neurite growth by tumour necrosis superfamily member RANKL. Open Biol. 3, 120150.

Han, S., Wang, B., Li, X., Xiao, Z., Han, J., Zhao, Y., Fang, Y., Yin, Y., Chen, B., Dai, J., 2016. Bone
marrow-derived mesenchymal stem cells in three-dimensional culture promote neuronal regeneration
by neurotrophic protection and immunomodulation. J. Biomed. Mater. Res. Part A 104, 1759–1769.

Hayashi, M., Nakashima, T., Taniguchi, M., Kodama, T., Kumanogoh, A., Takayanagi, H., 2012.
Osteoprotection by semaphorin 3A. Nature 485, 69–74.

Heiney, K., Mateus, J.C., Lopes, C.D.F., Neto, E., Lamghari, M., Aguiar, P., 2019. µSpikeHunter: An
advanced computational tool for the analysis of neuronal communication and action potential
propagation in microfluidic platforms. Sci. Rep. 9, 5777.

Hiasa, M., Okui, T., Allette, Y.M., Ripsch, M.S., Sun-Wada, G.-H., Wakabayashi, H., Roodman, G.D.,
White, F.A., Yoneda, T., 2017. Bone Pain Induced by Multiple Myeloma Is Reduced by Targeting
V-ATPase and ASIC3. Cancer Res. 77, 1283–1295.

Hill, E.L., Elde, R., 1991. Distribution of CGRP-, VIP-, D beta H-, SP-, and NPY-immunoreactive nerves
in the periosteum of the rat. Cell Tissue Res. 264, 469–480.

Hukkanen, M., Konttinen, Y.T.T., Santavirta, S., Paavolainen, P., Gu, X.-H.H., Terenghi, G., Polak,
J.M.M., 1993. Rapid proliferation of calcitonin gene-related peptide-immunoreactive nerves during
healing of rat tibial fracture suggests neural involvement in bone growth and remodelling.
Neuroscience 54, 969–979.

Huynh, N., VonMoss, L., Smith, D., Rahman, I., Felemban, M.F., Zuo, J., Rody, W.J., McHugh, K.P.,
Holliday, L.S., 2016a. Characterization of Regulatory Extracellular Vesicles from Osteoclasts. J.

Dent. Res. 95, 673–679.
Huynh, N., VonMoss, L., Smith, D., Rahman, I., Felemban, M.F.F., Zuo, J., Rody, W.J.J., McHugh, K.P.P.,
Holliday, L.S.S., Holliday, L.S.S., et al., 2016b. Characterization of Regulatory Extracellular
Vesicles from Osteoclasts 95.
Hynes, N.E., Lane, H. a, 2005. ERBB receptors and cancer: the complexity of targeted inhibitors. Nat. Rev.
Cancer 5, 341–54.
Imai, Y., Kohsaka, S., 2002. Intracellular signaling in M-CSF-induced microglia activation: Role of Iba1.
Glia 40, 164–174.
Irby, R.B., Yeatman, T.J., 2000. Role of Src expression and activation in human cancer. Oncogene 19,
5636–5642.
Jaiswal, R., Sedger, L.M., 2019. Intercellular Vesicular Transfer by Exosomes, Microparticles and
Oncosomes - Implications for Cancer Biology and Treatments. Front. Oncol. 9, 125.
Jimenez-Andrade, J.M., Bloom, A.P., Stake, J.I., Mantyh, W.G., Taylor, R.N., Freeman, K.T., Ghilardi,
J.R., Kuskowski, M.A., Mantyh, P.W., 2010. Pathological Sprouting of Adult Nociceptors in Chronic
Prostate Cancer-Induced Bone Pain. J. Neurosci. 30, 14649–14656.
Julius, D., Basbaum, A.I., 2001. Molecular mechanisms of nociception. Nature 413, 203–210.
Kohno, N., Aogi, K., Minami, H., Nakamura, S., Asaga, T., Iino, Y., Watanabe, T., Goessl, C., Ohashi, Y.,
Takashima, S., 2005. Zoledronic Acid Significantly Reduces Skeletal Complications Compared With
Placebo in Japanese Women With Bone Metastases From Breast Cancer: A Randomized, Placebo-
Controlled Trial. J. Clin. Oncol. 23, 3314–3321.
Koprivica, V., Cho, K.-S.S., Park, J.B., Yiu, G., Atwal, J., Gore, B., Kim, J. a, Lin, E., Tessier-Lavigne,
M., Chen, D.F., et al., 2005. EGFR Activation Mediates Inhibition of Axon Regeneration by Myelin
and Chondroitin Sulfate Proteoglycans. Science (80-.). 310, 106–110.
Kumar, K.R., Liang, C., Needham, M., Burke, D., Sue, C.M., Ng, K., 2011. Axonal hyperpolarization in
inclusion-body myopathy, paget disease of the bone, and frontotemporal dementia (IBMPFD).
Muscle Nerve 44, 191–196.
Lee, D.Y., Deng, Z., Wang, C.-H., Yang, B.B., 2007. MicroRNA-378 promotes cell survival, tumor growth,
and angiogenesis by targeting SuFu and Fus-1 expression. Proc. Natl. Acad. Sci. 104, 20350–20355.
Li, J., Ahmad, T., Spetea, M., Ahmed, M., Kreicbergs, A., 2001. Bone reinnervation after fracture: a study
in the rat. J Bone Min. Res 16, 1505–1510.
Li, J., Kreicbergs, A., Bergstrom, J., Stark, A., Ahmed, M., 2007. Site-specific CGRP innervation coincides
with bone formation during fracture healing and modeling: A study in rat angulated tibia. J Orthop
Res 25, 1204–1212.
Liu, B., Chen, H., Johns, T.G., Neufeld, A.H., 2006. Epidermal Growth Factor Receptor Activation: An
Upstream Signal for Transition of Quiescent Astrocytes into Reactive Astrocytes after Neural Injury.
J. Neurosci. 26, 7532–7540.
Lopes, C.D.F., Mateus, J.C., Aguiar, P., 2018. Interfacing Microfluidics with Microelectrode Arrays for
Studying Neuronal Communication and Axonal Signal Propagation. J. Vis. Exp. 1–8.
Lopez-Verrilli, M.A., Picou, F., Court, F.A., 2013. Schwann cell-derived exosomes enhance axonal
regeneration in the peripheral nervous system. Glia 61, 1795–1806.
Lu, P., Jones, L.L., Tuszynski, M.H., 2005. BDNF-expressing marrow stromal cells support extensive
axonal growth at sites of spinal cord injury. Exp. Neurol. 191, 344–360.
Macdonald-Obermann, J.L., Adak, S., Landgraf, R., Piwnica-Worms, D., Pike, L.J., 2013. Dynamic
analysis of the epidermal growth factor (EGF) receptor-ErbB2-ErbB3 protein network by luciferase
fragment complementation imaging. J. Biol. Chem. 288, 30773–30784.
Madsen, J.E.E., Hukkanen, M., Aune, A.K.K., Basran, I., Møller, J.F., Polak, J.M.M., Nordsletten, L.,
Moller, J.F., Polak, J.M.M., Nordsletten, L., 1998. Fracture healing and callus innervation after
peripheral nerve resection in rats. Clin Orthop Relat Res 230–240.
Mantyh, P., 2013. Bone cancer pain: Causes, consequences, and therapeutic opportunities. Pain 154, S54–
S62.
Mantyh, P.W., 2014. The neurobiology of skeletal pain. Eur. J. Neurosci. 39, 508–519.
Mediati, R.D., Vellucci, R., Dodaro, L., 2014. Pathogenesis and clinical aspects of pain in patients with
osteoporosis. Clin. Cases Miner. Bone Metab. 11, 169–72.
Middlemiss, T., Laird, B.J.A., Fallon, M.T., 2011. Mechanisms of Cancer-induced Bone Pain. Clin. Oncol.
23, 387–392.
Mizobuchi, S., Kanzaki, H., 2013. Spinal nerve injury causes upregulation of ErbB2 and ErbB3 receptors
in rat dorsal root ganglia. J. Pain Res. 6, 87–94.
Moro, L., Dolce, L., Cabodi, S., Bergatto, E., Erba, E.B., Smeriglio, M., Turco, E., Retta, S.F., Giuffrida,
M.G., Venturino, M., et al., 2002. Integrin-induced epidermal growth factor (EGF) receptor activation
requires c-Src and p130Cas and leads to phosphorylation of specific EGF receptor tyrosines. J. Biol.

880   Chem. 277, 9405–9414.

881 Morozevich, G.E., Kozlova, N.I., Ushakova, N.A., Preobrazhenskaya, M.E., Berman, A.E., 2012. Integrin
882   α5β1 simultaneously controls EGFR-dependent proliferation and Akt-dependent pro-survival
883   signaling in epidermoid carcinoma cells. Aging (Albany. NY). 4, 368–74.

884 Nagae, M., Hiraga, T., Wakabayashi, H., Wang, L., Iwata, K., Yoneda, T., Clerck, L.S. De, 2006.
885   Osteoclasts play a part in pain due to the inflammation adjacent to bone. Bone 39, 1107–15.

886 Neto, E., Alves, C.J., Leitão, L., Sousa, D.M., Alencastre, I.S., Conceição, F., Lamghari, M., 2017. Axonal
887   outgrowth, neuropeptides expression and receptors tyrosine kinase phosphorylation in 3D
888   organotypic cultures of adult dorsal root ganglia. PLoS One 12, e0181612.

889 Neto, E., Alves, C.J., Sousa, D.M., Alencastre, I.S., Lourenço, A.H., Leitão, L., Ryu, H.R., Jeon, N.L.,
890   Fernandes, R., Aguiar, P., et al., 2014. Sensory neurons and osteoblasts: close partners in a
891   microfluidic platform. Integr. Biol. 6, 586.

892 Neuhuber, B., Timothy Himes, B., Shumsky, J.S., Gallo, G., Fischer, I., 2005. Axon growth and recovery
893   of function supported by human bone marrow stromal cells in the injured spinal cord exhibit donor
894   variations. Brain Res 1035, 73–85.

895 Ni, S., Ling, Z., Wang, X., Cao, Y., Wu, T., Deng, R., Crane, J.L., Skolask, R., Demehril, S., Zhen, G., et
896   al., 2019. Sensory innervation in porous endplates by Netrin-1 from osteoclasts mediates PGE2-
897   induced spinal hypersensitivity in mice. Nat. Commun. 10.

898 Nielsen, D.L., Kümler, I., Palshof, J.A.E., Andersson, M., 2013. Efficacy of HER2-targeted therapy in
899   metastatic breast cancer. Monoclonal antibodies and tyrosine kinase inhibitors. The Breast 22, 1–12.

900 Nilsson, A., Kanje, M., 2005. Amphiregulin acts as an autocrine survival factor for adult sensory neurons.
901   Neuroreport 16, 213–8.

902 Nordsletten, L., Madsen, J.E., Almaas, R., Rootwelt, T., Halse, J., Konttinen, Y.T., Hukkanen, M.,
903   Santavirta, S., 1994. The neuronal regulation of fracture healing. Effects of sciatic nerve resection in
904   rat tibia. Acta Orthop. Scand. 65, 299–304.

905 Novikova, L.N., Brohlin, M., Kingham, P.J., Novikov, L.N., Wiberg, M., 2011. Neuroprotective and
906   growth-promoting effects of bone marrow stromal cells after cervical spinal cord injury in adult rats.
907   Cytotherapy 13, 873–887.

908 Olayioye, M.A., 2000. The ErbB signaling network: receptor heterodimerization in development and
909   cancer. EMBO J. 19, 3159–3167.

910 Paolicelli, R.C., Bergamini, G., Rajendran, L., 2019. Cell-to-cell Communication by Extracellular Vesicles:
911   Focus on Microglia. Neuroscience 405, 148–157.

912 Patel, M.S., Elefteriou, F., 2007. The new field of neuroskeletal biology. Calcif Tissue Int 80, 337–347.

913 Reddy, S. V, Menaa, C., Singer, F.R., Demulder, A., Roodman, G.D., 1999. Cell biology of Paget's disease.
914   J. Bone Miner. Res. 14 Suppl 2, 3–8.

915 Riminucci, M., Kuznetsov, S.A., Cherman, N., Corsi, A., Bianco, P., Gehron Robey, P., 2003.
916   Osteoclastogenesis in fibrous dysplasia of bone: In situ and in vitro analysis of IL-6 expression. Bone
917   33, 434–442.

918 Robles, E., Woo, S., Gomez, T.M., 2005. Src-dependent tyrosine phosphorylation at the tips of growth cone
919   filopodia promotes extension. J. Neurosci. 25, 7669–81.

920 Rody, W.J., Chamberlain, C.A., Emory-Carter, A.K., McHugh, K.P., Wallet, S.M., Spicer, V., Krokhin,
921   O., Holliday, L.S., Holliday, L.S., 2019. The proteome of extracellular vesicles released by clastic
922   cells differs based on their substrate. PLoS One 14, e0219602.

923 Rody, W.J., Krokhin, O., Spicer, V., Chamberlain, C.A., Chamberlain, M., McHugh, K.P., Wallet, S.M.,
924   Emory, A.K., Crull, J.D., Holliday, L.S., 2017. The use of cell culture platforms to identify novel
925   markers of bone and dentin resorption. Orthod. Craniofac. Res. 20, 89–94.

926 Rossi, M., Battafarano, G., D'Agostini, M., Del Fattore, A., 2018. The Role of Extracellular Vesicles in
927   Bone Metastasis. Int. J. Mol. Sci. 19.

928 Schulze, W.X., Deng, L., Mann, M., 2005. Phosphotyrosine interactome of the ErbB-receptor kinase
929   family. Mol. Syst. Biol. 1, 2005.0008.

930 Silva, A.M., Almeida, M.I., Teixeira, J.H., Maia, A.F., Calin, G.A., Barbosa, M.A., Santos, S.G., 2017.
931   Dendritic Cell-derived Extracellular Vesicles mediate Mesenchymal Stem/Stromal Cell recruitment.
932   Sci. Rep. 7, 1667.

933 Silva, J., Bento, A.R., Barros, D., Laundos, T.L., Sousa, S.R., Quelhas, P., Sousa, M.M., Pêgo, A.P.,
934   Amaral, I.F., 2017. Fibrin functionalization with synthetic adhesive ligands interacting with α6β1
935   integrin receptor enhance neurite outgrowth of embryonic stem cell-derived neural stem/progenitors.
936   Acta Biomater. 59, 243–256.

937 Sisask, G., Bjurholm, A., Ahmed, M., Kreicbergs, A., 1996. The development of autonomic innervation in
938   bone and joints of the rat. J. Auton. Nerv. Syst. 59, 27–33.

Sisask, G., Silfverswärd, C.-J., Bjurholm, A., Nilsson, O., 2013. Ontogeny of sensory and autonomic nerves

in the developing mouse skeleton. Auton. Neurosci. 177, 237–243.

Song, D., Jiang, X., Zhu, S., Li, W., Khadka, A., Hu, J., 2012. Denervation impairs bone regeneration
during distraction osteogenesis in rabbit tibia lengthening. Acta Orthop 83, 406–410.

Sousa, D.M., Conceição, F., Silva, D.I., Leitão, L., Neto, E., Alves, C.J., Alencastre, I.S., Herzog, H.,
Aguiar, P., Lamghari, M., 2016. Ablation of Y1 receptor impairs osteoclast bone-resorbing activity.
Sci. Rep. 6, 33470.

Taguchi, A., Politi, K., Pitteri, S.J., Lockwood, W.W., Faça, V.M., Kelly-Spratt, K., Wong, C.-H., Zhang,
Q., Chin, A., Park, K.-S., et al., 2011. Lung Cancer Signatures in Plasma Based on Proteome Profiling
of Mouse Tumor Models. Cancer Cell 20, 289–299.

Tassew, N.G., Charish, J., Shabanzadeh, A.P., Luga, V., Harada, H., Farhani, N., D'Onofrio, P., Choi, B.,
Ellabban, A., Nickerson, P.E.B., et al., 2017. Exosomes Mediate Mobilization of Autocrine Wnt10b
to Promote Axonal Regeneration in the Injured CNS. Cell Rep. 20, 99–111.

Théry, C., Amigorena, S., Raposo, G., Clayton, A., 2006. Isolation and Characterization of Exosomes from
Cell Culture Supernatants and Biological Fluids. Curr. Protoc. Cell Biol. 30, 3.22.1-3.22.29.

Théry, C., Witwer, K.W., Aikawa, E., Alcaraz, M.J., Anderson, J.D., Andriantsitohaina, R., Antoniou, A.,
Arab, T., Archer, F., Atkin-Smith, G.K., et al., 2018. Minimal information for studies of extracellular
vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular
Vesicles and update of the MISEV2014 guidelines. J. Extracell. vesicles 7, 1535750.

Tomlinson, R.E., Li, Zhi, Zhang, Q., Goh, B.C., Li, Zhu, Thorek, D.L.J., Rajbhandari, L., Brushart, T.M.,
Minichiello, L., Zhou, F., et al., 2016. NGF-TrkA Signaling by Sensory Nerves Coordinates the
Vascularization and Ossification of Developing Endochondral Bone. Cell Rep. 1–13.

Tsai, N.-P., Tsui, Y.-C., Pintar, J.E., Loh, H.H., Wei, L.-N., 2010. Kappa opioid receptor contributes to
EGF-stimulated neurite extension in development. Proc. Natl. Acad. Sci. U. S. A. 107, 3216–21.

Vigneswara, V., Kundi, S., Ahmed, Z., 2012. Receptor tyrosine kinases: molecular switches regulating
CNS axon regeneration. J. Signal Transduct. 2012, 361721.

Wang, H., Liu, B., Al-Aidaroos, A.Q.O., Shi, H., Li, L., Guo, K., Li, J., Tan, B.C.P., Loo, J.M., Tang, J.P.,
et al., 2016. Dual-faced SH3BGRL: oncogenic in mice, tumor suppressive in humans. Oncogene 35,
3303–13.

Weng, X., Smith, T., Sathish, J., Djouhri, L., 2012. Chronic inflammatory pain is associated with increased
excitability and hyperpolarization-activated current (Ih) in C- but not Aδ-nociceptors. Pain 153, 900–
914.

Wilkins, A., Kemp, K., Ginty, M., Hares, K., Mallam, E., Scolding, N., 2009. Human bone marrow-derived
mesenchymal stem cells secrete brain-derived neurotrophic factor which promotes neuronal survival
in vitro. Stem Cell Res. 3, 63–70.

Wong, R.W.C., Guillaud, L., 2004. The role of epidermal growth factor and its receptors in mammalian
CNS. Cytokine Growth Factor Rev. 15, 147–156.

Wood, E.R., Truesdale, A.T., McDonald, O.B., Yuan, D., Hassell, A., Dickerson, S.H., Ellis, B., Pennisi,
C., Horne, E., Lackey, K., et al., 2004. A Unique Structure for Epidermal Growth Factor Receptor
Bound to GW572016 (Lapatinib). Cancer Res. 64, 6652–6659.

Xian, C.J., Zhou, X.-F.F., 1999. Neuronal-glial differential expression of TGF-alpha and its receptor in the
dorsal root ganglia in response to sciatic nerve lesion. Exp. Neurol. 157, 317–326.

Xin, H., Li, Y., Buller, B., Katakowski, M., Zhang, Y., Wang, X., Shang, X., Zhang, Z.G., Chopp, M.,
2012. Exosome-Mediated Transfer of miR-133b from Multipotent Mesenchymal Stromal Cells to
Neural Cells Contributes to Neurite Outgrowth. Stem Cells 30, 1556–1564.

Xu, M., Zhou, H., Hu, C., Liang, Y., Hu, L., Chen, D., 2014. The mechanisms of EGFR in the regulation
of axon regeneration 101–105.

Yoneda, T., Hata, K., Nakanishi, M., Nagae, M., Nagayama, T., Wakabayashi, H., Nishisho, T., Sakurai,
T., Hiraga, T., 2011. Involvement of acidic microenvironment in the pathophysiology of cancer-
associated bone pain. Bone 48, 100–105.

Yoneda, T., Hiasa, M., Nagata, Y., Okui, T., White, F., 2015a. Contribution of acidic extracellular
microenvironment of cancer-colonized bone to bone pain. Biochim. Biophys. Acta - Biomembr.
1848, 2677–2684.

Yoneda, T., Hiasa, M., Nagata, Y., Okui, T., White, F.A., 2015b. Acidic microenvironment and bone pain
in cancer-colonized bone. Bonekey Rep. 4, 690.

Yuan, F.-L., Wu, Q.-Y., Miao, Z.-N., Xu, M.-H., Xu, R.-S., Jiang, D.-L., Ye, J.-X., Chen, F.-H., Zhao, M.-
D., Wang, H.-J., et al., 2018. Osteoclast-Derived Extracellular Vesicles: Novel Regulators of
Osteoclastogenesis and Osteoclast-Osteoblasts Communication in Bone Remodeling. Front. Physiol.
9, 628.

Zhang, Y., Chopp, M., Liu, X.S., Katakowski, M., Wang, X., Tian, X., Wu, D., Zhang, Z.G., 2017.
Exosomes Derived from Mesenchymal Stromal Cells Promote Axonal Growth of Cortical Neurons.

Mol. Neurobiol. 54, 2659–2673.

Zhu, S., Zhu, J., Zhen, G., Hu, Y., An, S., Li, Y., Zheng, Q., Chen, Z., Yang, Y., Wan, M., et al., 2019. Subchondral bone osteoclasts induce sensory innervation and osteoarthritis pain. J. Clin. Invest. 129, 1076–1093.