The origin of the odorant receptor gene family in insects

Philipp Brand[1,*], Hugh M. Robertson[2,*], Wei Lin[3], Ratnasri Pothula[4], William E.

Klingeman[5], Juan Luis Jurat-Fuentes[4], and Brian R. Johnson[3]

[1] Department of Evolution and Ecology, Center for Population Biology, University of

California, Davis, Davis, CA 95616, USA

[2] Department of Entomology, University of Illinois at Urbana-Champaign, Urbana, IL

61801, USA

[3] Department of Entomology and Nematology, University of California, Davis, Davis,

CA 95616, USA

[4] Department of Entomology and Plant Pathology, University of Tennessee, Knoxville

TN 37996, USA

[5] Department of Plant Sciences, University of Tennessee, Knoxville TN 37996, USA

* Shared first authorship

Corresponding authors: Philipp Brand, pbrand@ucdavis.edu, and Hugh Robertson,

hughrobe@uiuc.edu

Abstract

The sense of smell enables the detection and discrimination of airborne chemicals via chemosensory receptors that have evolved independently multiple times throughout the tree of life. In insects, the odorant receptor (OR) gene family is the major chemosensory gene family involved in olfaction and its origin has been hypothesized to coincide with the evolution of a terrestrial lifestyle in hexapods. Missbach et al. (2014) challenged this view and suggested that ORs evolved with an ancestral OR co-receptor (Orco) after the origin of terrestriality, hypothesizing that the OR/Orco system is an adaptation to winged flight in insects instead. Building upon this work, we investigated the genomes of basal hexapod and insect lineages including Collembola, Diplura, Archaeognatha, Zygentoma, Odonata, and Ephemeroptera in an effort to identify the origin of the insect OR gene family. While absent from all non-insect hexapod lineages analyzed, ORs are present in all insect genomes. Orco is absent only in the most ancient insect lineage Archaeognatha. A fully functional OR/Orco system was present in our newly generated genome data of the Zygentoma *Thermobia domestica*. We suggest that ORs did evolve as adaptation to a terrestrial lifestyle outside high-humidity habitats, and not winged flight, representing a key evolutionary novelty in the ancestor of all insects. The OR family is therefore the first known molecular synapomorphy for the Class Insecta.

Introduction

From bacteria to mammals, living organisms of all levels of complexity have evolved chemosensory receptors to detect and discriminate chemicals in the environment (Wuichet and Zhulin, 2010; Hansson and Stensmyr, 2011). The largest metazoan gene families for example encode tens to hundreds of odorant receptors (ORs) that interact with volatile chemicals at the sensory periphery underlying the sense of smell (Sánchez-Gracia et al., 2009; Niimura et al., 2014). OR gene families have evolved multiple times throughout the metazoans, including independent origins in vertebrates, nematodes, and insects (Hansson and Stensmyr, 2011). In insects, the OR gene family evolved from within the ancestral gustatory receptor (GR) gene family (Scott et al., 2001; Robertson et

al., 2003) that extends back to ancient metazoan lineages (Robertson, 2015; Saina et al., 2015; Eyun et al., 2017). ORs are absent from non-insect arthropod genomes (Peñalva-Arana et al., 2009; Almeida et al., 2015; Gulia-Nuss et al., 2016; Ngoc et al., 2016; Eyun et al., 2017), and have been hypothesized to have evolved concomitant with the evolution of terrestriality in hexapods (Robertson et al., 2003).

The lack of molecular resources for early branching hexapod and insect lineages has prevented the precise dating of the origin of insect ORs. Only recently, whole-genome sequencing efforts suggested that ORs are absent in non-insect hexapods such as Collembola (Wu et al., 2017) but present in early branching pterygote insects such as damselflies (Odonata; Ioannidis et al., 2017). Efforts to understand more precisely the origin of the OR gene family within hexapods were greatly advanced by the findings of Missbach et al. (2014) who sequenced transcriptomes of the chemosensory organs of two apterygote insects, the bristletail *Lepismachilis y-signata* (Archaeognatha) and the firebrat *Thermobia domestica* (Zygentoma). They identified three ORs in the firebrat, which they named TdomOrco1-3, with apparent similarity to the neopteran odorant receptor co-receptor (Orco; Vosshall and Hansson, 2011). Orco is a highly conserved single-copy gene present in all other insects studied to date and encodes a protein that is a partner with each of the other "specific" ORs (Benton et al., 2006) which is required for OR-based olfaction in insects (Larsson et al., 2004). In contrast, Missbach et al. (2014) could not find ORs or Orco relatives in their bristletail transcriptome, instead finding only members of the ionotropic receptor (IR) gene family. Given evidence that IRs serve olfactory roles in terrestrial crustaceans and insects (Rytz et al., 2013; Groh-Lunow et al., 2015; Rimal and Lee, 2018), they argued that olfaction in early-branching terrestrial hexapods and apterygote insects is entirely IR-dependent, with Orco evolving as ancestral OR from the GR lineage between the Archaeognatha and Zygentoma. Based on these findings, Missbach et al. (2014) suggested that the Orco/OR system evolved together with flight in pterygote insects and left off with the observation that "the existence of three Orco types remains mysterious".

Recently, phylogenetic analysis of the OR gene family of the damselfly *Calopteryx*

splendens suggested that at least one of the three Orco-like ORs from *T. domestica*,

TdomOrco3, might be a specific OR instead of an Orco (Ioannidis et al., 2017). If this is

correct, then the entire Orco/OR system evolved before winged insects, which would

explain the "mystery" of three apparent Orco types in Zygentoma. In an effort to identify

the origin of the insect OR gene family and the Orco/OR system, we investigated the

genome sequences of species belonging to multiple ancient terrestrial hexapod and insect

orders, including Collembola (springtails), Diplura (two-pronged bristletails),

Archaeognatha (jumping bristletails), Zygentoma (silverfish and firebrats), Odonata

(damselflies and dragonflies), and Ephemeroptera (mayflies).

Results and Discussion

ORs were present in the ancestor of insects

We detected no ORs in two non-insect hexapod lineages, Collembola and Diplura (a

genome sequence is not available for the third lineage, the Protura), despite extensive

annotation efforts. In contrast, we identified genes with similarity to known insect ORs in

all other genomes investigated (Figure 1; details in Supplemental Material, Tables S2 and

S3). These included one species of each of the earliest branching insect lineages, the

Archeognatha, Zygentoma, Ephemeroptera, and Odonata (Misof et al., 20014).

Accordingly, ORs were likely present in the ancestor of all insects but absent from all

non-insect hexapod lineages. This suggests that the origin of the OR gene family

coincided with the evolution of insects. Thus, our analysis does not support the

hypothesis that ORs evolved with the evolution of winged flight in insects (Missbach et

al., 2014) but is compatible with the hypothesis that they evolved with terrestriality in

insects (Robertson et al. 2003). Terrestrial hexapod lineages without ORs (*i.e.* Diplura

and Collembola) are confined to highly humid environments including moist soil and

leaf-litter. This is analogous to terrestrial crustaceans, which evolved OR-independent

modified sensory systems allowing the detection of airborne odors on land (Stensmyr et

al., 2005; Hansson et al., 2010). While terrestrial crustaceans employ the same olfactory

gene-families as their marine ancestors in combination with anatomical adaptations to

terrestrial olfaction, their sense of smell is highly dependent on humidity (Krång et al.,

2012). Although the molecular and physiological basis of dipluran and collembolan

olfaction is unexplored, they are known to have a highly sensitive sense of smell used in

pheromone communication and foraging behavior (Verhoef et al., 1977; Purrington et al.,

1991; Staaden et al., 2011). It is thus possible that ORs evolved as an adaptation to a

terrestrial lifestyle outside ancestrally humid environments in insects.

Figure 1 Origin of the insect odorant receptor gene family. The number of ORs and OR co-receptors (Orcos) for all
ancient insect and non-insect orders analyzed was mapped on the hexapod phylogeny sensu Misof et al. (2014). ORs
are present in all insects but absent from non-insect hexapod genomes, and thus likely represent a molecular
synapomorphy for the Clade Insecta. Orco is present in all but Archaeognatha, the earliest branching insect order. This
suggests two scenarios including either the loss of Orco in Archaeognatha or an Orco origin following the evolution of
ORs (as indicated). The OR gene family likely evolved as an adaptation to a terrestrial lifestyle outside ancestrally
humid environments in insects. *: The genomes of all neopteran insects analyzed to date encode ORs, ranging from 10
ORs in head lice (Kirkness et al., 2010) to more than 300 ORs in ants (Smith et al., 2011a; b).

The *Thermobia domestica* genome harbors a full Orco/OR gene family repertoire

With the exception of the Zygentoma, all lineages analyzed had genome data either

published (Faddeeva-Vakhrusheva et al., 2016; 2017; Wu et al., 2017) or available from

the i5k Pilot Project from the Baylor College of Medicine at the i5k NAL Workspace

(Poelchau et al., 2015). To complete taxon sampling of early branching hexapod and

insect lineages, we produced a draft genome assembly for *T. domestica* (Supplemental

Material; Figure S1; Table S1). This enabled direct comparison to Missbach et al. (2014)

and revealed that the *T. domestica* genome encodes far more than the three Orco-like OR

proteins. Our manual annotation revealed 43 ORs encoded by 31 genes including the

three previously identified genes (TdomOrco1-3; Missbach et al., 2014). Four genes are

modeled as exhibiting alternative splicing leading to the additional protein isoforms

(Supplemental Material). We used the antennal transcriptome of Missbach et al. (2014)

for support of intron-exon boundaries, however only a few transcriptome reads mapped

to the "specific" OR genes (Table S2), indicating that the RNA-seq analysis of Missbach

et al. (2014) did not sequence to a sufficient depth to reconstruct these low-expressed

transcripts.

Phylogenetic analyses of all ORs we annotated in the bristletail *Machilis hrabei* (5 ORs),

the dragonfly *Ladona fulva* (4 ORs), and the mayfly *Ephemera danica* (47 ORs), as well

as the previously annotated damselfly *C. splendens* (6 ORs; Ioannidis et al., 2017)

revealed that one of the *T. domestica* ORs (TdomOrco2) clustered confidently with the

Orco lineage in pterygote insects (Figure 2; Figure S2). We believe this is the sole Orco

relative because it shares unique features with the pterygote Orco proteins, such as a

TKKQ motif in the expanded intracellular loop 2 (positions 327-330 in DmelOrco), and

so we simply rename it TdomOrco. TdomOr1-8 represent a set of Orco-like proteins that

share a common gene structure with TdomOrco, with introns in phases 0-2-0-0-0. These

last four introns are present in all the other TdomOr genes, as well as those of the

bristletail, Odonata, and mayfly, and correspond to the four introns identified by

Robertson et al. (2003) as being ancestral to the OR family. The first phase-0 intron of

Orco and Or1-8 is the only additional intron shared by most pterygote Orco genes.

With the exception of the bristletail *M. hrabei*, all insect genomes analyzed have both

single genes with high similarity to Orco and multiple genes with similarity to specific

ORs. The *M. hrabei* genome did not encode an Orco, but instead contains 5 ORs of high

similarity that form a highly supported clade in the gene phylogeny (Figure 2). We also

could not find an Orco in the deep RNAseq transcriptome Missbach et al. (2014)

generated for their bristletail, *L. y-signata*. This finding leaves open two possibilities.

First, bristletails might have lost their Orco gene. Second, the OR family might have

originated with a few specific ORs like those of *M. hrabei*, with the Orco lineage

evolving between Archaegnatha and Zygentoma. Phylogenetic analysis using various sets

of GRs from other insects, arthropods, and animals as outgroup to root the OR family tree

does not resolve this question confidently (data not shown). In any case, these five

specific ORs in *M. hrabei*, at least one of which is present in *L. y-signata*, must function

in the absence of Orco, perhaps alone or as dimers.

Finally, we note that while insects are defined by morphological and developmental

synapomorphies (shared derived characters that are unique to a taxon), to the best of our

knowledge presence of the OR gene family is the first molecular synapomorphy for the

Class Insecta.

Figure 2 Odorant receptor (OR) gene family phylogeny including representatives of all apterygote and
paleopteran insect orders. The Maximum Likelihood tree demonstrates monophyly of the single-copy insect Orco
with high bootstrap support. The *M. hrabei* genome lacks Orco but encodes five highly similar ORs clustering in a
single highly-supported clade. *T. domestica* has a fully developed functional OR/Orco system. The red arrowhead
indicates the location of the three *T. domestica* ORs identified by Missbach et al. (2014).

Materials and Methods

A detailed Materials and Methods section can be found in the Supplementary Material.

Sequencing and assembly of the *Thermobia domestica* genome

Sequencing and assembly of the *T. domestica* genome followed Brand et al. (2018). Briefly, we sequenced the DNA extracted from a single *T. domestica* individual for assembly with the DISCOVAR v1 pipeline (Weisenfeld et al., 2014). Quality assessment of the resulting assembly was based on standard N statistics, k-mer distribution analysis, and the BUSCO v2 pipeline (Simão et al., 2015) as previously described (Brand et al., 2017).

Odorant receptor annotation

The genomes of the dipluran *Catajapyx aquinolaris,* the collembolans *Holacanthella duospinosa, Orchesella cincta,* and *Folsomia candida*, the firebrat *Thermobia domestica*, the bristletail *Machilis hrabei,* the dragonfly *Ladona fulva*, and the mayfly *Ephemera danica* were used for manual OR gene annotation (detailed in Supplementary Material).

Odorant receptor gene family analysis

OR protein alignments were produced with CLUSTALX v2 (Larkin et al., 2007) and trimmed using TrimAl (Capella-Gutiérrez et al., 2009). The resulting alignment was used for gene tree inference using RaxML (Stamatakis et al., 2005) under the JTT + G substitution model with 20 independent ML searches and 1000 bootstrap replicates as previously described (Brand and Ramírez, 2017).

Acknowledgements

We thank Stephen Richards for permission to examine unpublished genome sequences from the i5k pilot project, and Kimberly Walden for assistance with BLAST searches.

This work was funded by a National Science Foundation grant, IOS-1456678, to Brian

Johnson and Juan Luis Jurat-Fuentes and a Hatch grant to Brian Johnson (CA-D-ENM

2161-H).

References

Almeida FC, Sánchez-Gracia A, Walden KKO, Robertson HM, Rozas J. 2015. Positive
selection in extra cellular domains in the diversification of *Strigamia maritima*
chemoreceptors. Front Ecol Evol 3:1669.

Benton R, Sachse S, Michnick SW, Vosshall LB. 2006. Atypical membrane topology and
heteromeric function of *Drosophila* odorant receptors in vivo. PLoS Biol 4:e20.

Brand P, Lin W, Johnson BR. 2018. The Draft Genome of the Invasive Walking Stick,
Medauroidea extradentata, Reveals Extensive Lineage-Specific Gene Family
Expansions of Cell Wall Degrading Enzymes in Phasmatodea. G3: Genes, Genomes,
Genetics:g3.200204.2018.

Brand P, Ramírez SR. 2017. The Evolutionary Dynamics of the Odorant Receptor Gene
Family in Corbiculate Bees. Genome Biol Evol 9:2023–2036.

Brand P, Saleh N, Pan H, Li C, Kapheim KM, Ramírez SR. 2017. The Nuclear and
Mitochondrial Genomes of the Facultatively Eusocial Orchid Bee *Euglossa dilemma*.
G3: Genes, Genomes, Genetics 7:2891–2898.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–
1973.

Eyun S-I, Soh HY, Posavi M, Munro JB, Hughes DST, Murali SC, Qu J, Dugan S, Lee
SL, Chao H, Dinh H, Han Y, Doddapaneni H, Worley KC, Muzny DM, Park E-O,
Silva JC, Gibbs RA, Richards S, Lee CE. 2017. Evolutionary History of
Chemosensory-Related Gene Families across the Arthropoda. Mol Biol Evol
34:1838–1862.

Faddeeva-Vakhrusheva A, Derks MFL, Anvar SY, Agamennone V, Suring W, Smit S,
van Straalen NM, Roelofs D. 2016. Gene Family Evolution Reflects Adaptation to
Soil Environmental Stressors in the Genome of the Collembolan *Orchesella cincta*.
Genome Biol Evol 8:2106–2117.

Faddeeva-Vakhrusheva A, Kraaijeveld K, Derks MFL, Anvar SY, Agamennone V,
Suring W, Kampfraath AA, Ellers J, Le Ngoc G, van Gestel CAM, Marien J, Smit S,
van Straalen NM, Roelofs D. 2017. Coping with living in the soil: the genome of the
parthenogenetic springtail *Folsomia candida*. BMC Genomics 18.

Groh-Lunow KC, Getahun MN, Grosse-Wilde E, Hansson BS. 2015. Expression of

ionotropic receptors in terrestrial hermit crab's olfactory sensory neurons. Front Cell
Neurosci 8:198.

Gulia-Nuss M, Nuss AB, Meyer JM, Sonenshine DE, Roe RM, Waterhouse RM, Sattelle
DB, la Fuente de J, Ribeiro JM, Megy K, Thimmapuram J, Miller JR, Walenz BP,
Koren S, Hostetler JB, Thiagarajan M, Joardar VS, Hannick LI, Bidwell S,
Hammond MP, Young S, Zeng Q, Abrudan JL, Almeida FC, Ayllon N, Bhide K,
Bissinger BW, Bonzon-Kulichenko E, Buckingham SD, Caffrey DR, Caimano MJ,
Croset V, Driscoll T, Gilbert D, Gillespie JJ, Giraldo-Calderon GI, Grabowski JM,
Jiang D, Khalil SMS, Kim D, Kocan KM, Koci J, Kuhn RJ, Kurtti TJ, Lees K, Lang
EG, Kennedy RC, Kwon H, Perera R, Qi Y, Radolf JD, Sakamoto JM, Sánchez-
Gracia A, Severo MS, Silverman N, Simo L, Tojo M, Tornador C, Van Zee JP,
Vazquez J, Vieira FG, Villar M, Wespiser AR, Yang Y, Zhu J, Arensburger P,
Pietrantonio PV, Barker SC, Shao R, Zdobnov EM, Hauser F, Grimmelikhuijzen
CJP, Park Y, Rozas J, Benton R, Pedra JHF, Nelson DR, Unger MF, Tubio JMC, Tu
Z, Robertson HM, Shumway M, Sutton G, Wortman JR, Lawson D, Wikel SK, Nene
VM, Fraser CM, Collins FH, Birren B, Nelson KE, Caler E, Hill CA. 2016. Genomic
insights into the *Ixodes scapularis* tick vector of Lyme disease. Nature
Communications 7.

Hansson BS, Harzsch S, Knaden M, Stensmyr M. 2010. The Neural and Behavioral Basis
of Chemical Communication in Terrestrial Crustaceans. In: Chemical
Communication in Crustaceans. New York, NY: Springer, New York, NY. p 149–
173.

Hansson BS, Stensmyr MC. 2011. Evolution of insect olfaction. Neuron 72:698–711.

Ioannidis P, Simão FA, Waterhouse RM, Manni M, Seppey M, Robertson HM, Misof B,
Niehuis O, Zdobnov EM. 2017. Genomic features of the damselfly *Calopteryx*
splendens representing a sister clade to most insect orders. Genome Biol
Evol:evx006.

Kirkness EF, Haas BJ, Sun W, Braig HR, Perotti MA, Clark JM, Lee SH, Robertson HM,
Kennedy RC, Elhaik E, Gerlach D, Kriventseva EV, Elsik CG, Graur D, Hill CA,
Veenstra JA, Walenz B, Tubío JMC, Ribeiro JMC, Rozas J, Johnston JS, Reese JT,
Popadic A, Tojo M, Raoult D, Reed DL, Tomoyasu Y, Kraus E, Krause E, Mittapalli
O, Margam VM, Li H-M, Meyer JM, Johnson RM, Romero-Severson J, VanZee JP,
Alvarez-Ponce D, Vieira FG, Aguadé M, Guirao-Rico S, Anzola JM, Yoon KS,
Strycharz JP, Unger MF, Christley S, Lobo NF, Seufferheld MJ, Wang N, Dasch GA,
Struchiner CJ, Madey G, Hannick LI, Bidwell S, Joardar V, Caler E, Shao R, Barker
SC, Cameron S, Bruggner RV, Regier A, Johnson J, Viswanathan L, Utterback TR,
Sutton GG, Lawson D, Waterhouse RM, Venter JC, Strausberg RL, Berenbaum MR,
Collins FH, Zdobnov EM, Pittendrigh BR. 2010. Genome sequences of the human
body louse and its primary endosymbiont provide insights into the permanent
parasitic lifestyle. Proc Natl Acad Sci USA 107:12168–12173.

Krång A-S, Knaden M, Steck K, Hansson BS. 2012. Transition from sea to land:

olfactory function and constraints in the terrestrial hermit crab *Coenobita clypeatus*.
Proc Biol Sci 279:3510–3519.

Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,
Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG.
2007. Clustal W and Clustal X version 2.0. Bioinformatics 23:2947–2948. Available
from: https://academic.oup.com/bioinformatics/article-
lookup/doi/10.1093/bioinformatics/btm404

Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB. 2004.
Or83b encodes a broadly expressed odorant receptor essential for *Drosophila*
olfaction. Neuron 43:703–714.

Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J,
Flouri T, Beutel RG, Niehuis O, Petersen M, Izquierdo-Carrasco F, Wappler T, Rust
J, Aberer AJ, Aspöck U, Aspöck H, Bartel D, Blanke A, Berger S, Böhm A, Buckley
TR, Calcott B, Chen J, Friedrich F, Fukui M, Fujita M, Greve C, Grobe P, Gu S,
Huang Y, Jermiin LS, Kawahara AY, Krogmann L, Kubiak M, Lanfear R, Letsch H,
Li Y, Li Z, Li J, Lu H, Machida R, Mashimo Y, Kapli P, McKenna DD, Meng G,
Nakagaki Y, Navarrete-Heredia JL, Ott M, Ou Y, Pass G, Podsiadlowski L, Pohl H,
Reumont von BM, Schütte K, Sekiya K, Shimizu S, Slipinski A, Stamatakis A, Song
W, Su X, Szucsich NU, Tan M, Tan X, Tang M, Tang J, Timelthaler G, Tomizuka S,
Trautwein M, Tong X, Uchifune T, Walzl MG, Wiegmann BM, Wilbrandt J, Wipfler
B, Wong TKF, Wu Q, Wu G, Xie Y, Yang S, Yang Q, Yeates DK, Yoshizawa K,
Zhang Q, Zhang R, Zhang W, Zhang Y, Zhao J, Zhou C, Zhou L, Ziesmann T, Zou
S, Li Y, Xu X, Zhang Y, Yang H, Wang J, et al. 2014. Phylogenomics resolves the
timing and pattern of insect evolution. Science 346:763–767.

Missbach C, Dweck HK, Vogel H, Vilcinskas A, Stensmyr MC, Hansson BS, Grosse-
Wilde E. 2014. Evolution of insect olfactory receptors. Elife 3:e02115.

Ngoc PCT, Greenhalgh R, Dermauw W, Rombauts S, Bajda S, Zhurov V, Grbic M, Van
de Peer Y, Van Leeuwen T, Rouze P, Clark RM. 2016. Complex Evolutionary
Dynamics of Massively Expanded Chemosensory Receptor Families in an Extreme
Generalist Chelicerate Herbivore. Genome Biol Evol 8:3323–3339.

Niimura Y, Matsui A, Touhara K. 2014. Extreme expansion of the olfactory receptor
gene repertoire in African elephants and evolutionary dynamics of orthologous gene
groups in 13 placental mammals. Genome Res 24:1485–1496.

Peñalva-Arana DC, Lynch M, Robertson HM. 2009. The chemoreceptor genes of the
waterflea Daphnia pulex: many Grs but no Ors. BMC Evol Biol 9:79.

Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee C-Y, Lin H, Lin J-W,
Hackett K. 2015. The i5k Workspace@NAL--enabling genomic data access,
visualization and curation of arthropod genomes. Nucleic Acids Res 43:D714–9.

Purrington FF, Kendall PA, Bater JE, Stinner BR. 1991. Alarm Pheromone in a

Gregarious Poduromorph-Collembolan (Collembola, Hypogastruridae). Great Lakes
Entomologist 24:75–78.

Rimal S, Lee Y. 2018. The multidimensional ionotropic receptors of *Drosophila*
melanogaster. Insect Mol Biol 27:1–7.

Robertson HM, Warr CG, Carlson JR. 2003. Molecular evolution of the insect
chemoreceptor gene superfamily in *Drosophila melanogaster*. Proc Natl Acad Sci
USA 100.

Robertson HM. 2015. The Insect Chemoreceptor Superfamily Is Ancient in Animals.
Chem Senses:bjv046.

Rytz R, Croset V, Benton R. 2013. Ionotropic receptors (IRs): chemosensory ionotropic
glutamate receptors in *Drosophila* and beyond. Insect Biochem Mol Biol 43.

Saina M, Busengdal H, Sinigaglia C, Petrone L, Oliveri P, Rentzsch F, Benton R. 2015.
A cnidarian homologue of an insect gustatory receptor functions in developmental
body patterning. Nature Communications 6.

Sánchez-Gracia A, Vieira FG, Rozas J. 2009. Molecular evolution of the major
chemosensory gene families in insects. Heredity 103:208–216.

Scott K, Brady R, Cravchik A, Morozov P, Rzhetsky A, Zuker C, Axel R. 2001. A
chemosensory gene family encoding candidate gustatory and olfactory receptors in
Drosophila. Cell 104:661–673.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31:3210–3212.

Smith CD, Zimin A, Holt C, Abouheif E, Benton R, Cash E, Croset V, Currie CR, Elhaik
E, Elsik CG, Favé M-J, Fernandes V, Gadau J, Gibson JD, Graur D, Grubbs KJ,
Hagen DE, Helmkampf M, Holley J-A, Hu H, Viniegra ASI, Johnson BR, Johnson
RM, Khila A, Kim JW, Laird J, Mathis KA, Moeller JA, Munoz-Torres MC, Murphy
MC, Nakamura R, Nigam S, Overson RP, Placek JE, Rajakumar R, Reese JT,
Robertson HM, Smith CR, Suarez AV, Suen G, Suhr EL, Tao S, Torres CW, van
Wilgenburg E, Viljakainen L, Walden KKO, Wild AL, Yandell M, Yorke JA,
Tsutsui ND. 2011a. Draft genome of the globally widespread and invasive Argentine
ant (*Linepithema humile*). Proc Natl Acad Sci USA 108:5673–5678.

Smith CR, Smith CD, Robertson HM, Helmkampf M, Zimin A, Yandell M, Holt C, Hu
H, Abouheif E, Benton R, Cash E, Croset V, Currie CR, Elhaik E, Elsik CG, Favé M-
J, Fernandes V, Gibson JD, Graur D, Gronenberg W, Grubbs KJ, Hagen DE,
Viniegra ASI, Johnson BR, Johnson RM, Khila A, Kim JW, Mathis KA, Munoz-
Torres MC, Murphy MC, Mustard JA, Nakamura R, Niehuis O, Nigam S, Overson
RP, Placek JE, Rajakumar R, Reese JT, Suen G, Tao S, Torres CW, Tsutsui ND,
Viljakainen L, Wolschin F, Gadau J. 2011b. Draft genome of the red harvester ant

384   *Pogonomyrmex barbatus*. Proc Natl Acad Sci USA 108:5667–5672.

385 Staaden S, Milcu A, Rohlfs M, Scheu S. 2011. Olfactory cues associated with fungal
386   grazing intensity and secondary metabolite pathway modulate Collembola foraging
387   behaviour. Soil Biology & Biochemistry 43:1411–1416.

388 Stamatakis A, Ludwig T, Meier H. 2005. RAxML-III: a fast program for maximum
389   likelihood-based inference of large phylogenetic trees. Bioinformatics 21:456–463.

390 Stensmyr MC, Erland S, Hallberg E, Wallén R, Greenaway P, Hansson BS. 2005. Insect-
391   like olfactory adaptations in the terrestrial giant robber crab. Current Biology
392   15:116–121.

393 Verhoef HA, Nagelkerke CJ, Joosse E. 1977. Aggregation Pheromones in Collembola.
394   Journal of Insect Physiology 23:1009–1013.

395 Vosshall LB, Hansson BS. 2011. A unified nomenclature system for the insect olfactory
396   coreceptor. Chem Senses 36:497–498.

397 Weisenfeld NI, Yin S, Sharpe T, Lau B, Hegarty R, Holmes L, Sogoloff B, Tabbaa D,
398   Williams L, Russ C, Nusbaum C, Lander ES, MacCallum I, Jaffe DB. 2014.
399   Comprehensive variation discovery in single human genomes. Nat Genet 46:1350–
400   1355.

401 Wu C, Jordan MD, Newcomb RD, Gemmell NJ, Bank S, Meusemann K, Dearden PK,
402   Duncan EJ, Grosser S, Rutherford K, Gardner PP, Crowhurst RN, Steinwender B,
403   Tooman LK, Stevens MI, Buckley TR. 2017. Analysis of the genome of the New
404   Zealand giant collembolan (*Holacanthella duospinosa*) sheds light on hexapod
405   evolution. BMC Genomics 18.

406 Wuichet K, Zhulin IB. 2010. Origins and diversification of a complex signal transduction
407   system in prokaryotes. Sci Signal 3:ra50–ra50.

408