TrackSig: reconstructing evolutionary trajectories of mutation signature exposure

Yulia Rubanova[1]**, Roujia Li**[2]**, Jeff Wintersinger**[1]**, Nil Sahin**[4]**, Amit Deshwar**[3]**, PCAWG-11 working group**[5]**, and Quaid Morris**[1,2,3,4,*]

[1]Department of Computer Science, University of Toronto, Toronto, Canada
[2]Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, Canada
[3]Department of Electrical and Computer Engineering, University of Toronto, Toronto, Canada
[4]Department of Molecular Genetics, University of Toronto, Toronto, Canada
[5]Various affiliations
[*]quaid.morris@utoronto.ca

ABSTRACT

We present a new method, TrackSig, to estimate the evolutionary trajectories of signatures of somatic mutational processes. TrackSig infers an approximate order in which the somatic mutations accumulated, and then fits the signature exposures as a function of the ordering. Using simulations, we assess TrackSig's reconstruction accuracy. We find 1.9% median exposure error on simulations with one to three change-points. The size and the direction of the signature change is consistent in 87% and 95% of cases respectively. There were an average of 0.02 missed change-points and 0.37 false positive change-points per sample. The code is available at https://github.com/YuliaRubanova/TrackSig.

1 Introduction

Somatic mutations accumulate throughout our lifetime, arising from external sources or from processes intrinsic to the cell[1,2]. Some sources generate characteristic patterns of mutations. For example, smoking is associated with G to T mutations; UV radiation is associated with C to T mutations[3–5]. Some processes provide a constant source of mutations[6] while others are sporadic[7].

Using mutational signature analysis, one can estimate the contribution of different mutation processes to the collection of somatic mutations present in a sample. In this type of analysis, single nucleotide variants (SNVs) are classified 96 types based on the type of substitution and tri-nucleotide context (e.g., ACG to a ATG)[2]. Mutational signatures across the 96 types were initially derived by non-negative matrix factorization[2]. Many of the signatures are strongly associated with known mutational processes including smoking[7], non-homologous double strand break repair[2], and ionizing radiation[8]. The exposures of some signatures are correlated with patient age[6] and suggesting their use as a molecular clock[9]. Because signature analysis can identifies the DNA damage repair pathways absent in the cancer, and can predict prognosis[10] or guide treatment choice[11].

Formally, a *mutational signature* is a probability distribution over these 96 types, where each element is a probability of generating mutations from the corresponding type[12]. Each signature is assigned an *exposure* which represents the proportion of mutations that the signature generates. These can be computed for pre-defined signatures from the total mutational spectrum of a sample by constrained regression[13,14].

Mutational sources can change over time. Mutations caused by carcinogen exposure stop accumulating when the exposure ends[7]. Mutations associated with defective DNA damage repair, such as BRCA1 loss[1,2] will begin to accumulate after that loss. Recent analyses have reported modest changes in signature exposures between clonal and subclonal populations[9,15] based on groups of mutations identified by clustering their variant allele frequencies (VAFs). However, the accuracy of these methods relies heavily on the sensitivity

and precision of this clustering, which is low for typical whole genome sequencing coverage[16].

In this paper we introduce TrackSig, a new method to reconstruct signature exposures across time without VAF clustering. We use VAF to approximately order mutations based on their prevalence within the cancer cell population and then track changes in signature exposure consistent with this ordering.

Using TrackSig, we have previously demonstrated that signature exposures change often during the lifetime of a cancer and that these changes can help identify new subclonal lineages[17]. Here, we use bootstrap analyses and realistic simulations to help assess the accuracy of those reconstructions.

2 Methods

TrackSig has two stages. First, we sort single nucleotide variants (SNVs) by an estimated order of their accumulation. We compute this estimate using their variant allele frequencies (VAFs) and a copy number aberration (CNA) reconstruction of the samples. Next, we infer a trajectory of the mutational signature exposures over the estimated ordering of the SNVs. We estimated exposure trajectory for each signature is a piecewise constant function of the SNV ordering with a small number of change-points. These stages are described in detail below.

TrackSig is designed to be applied to VAF frequency data from a single, heterogeneous tumour sample. However, if an ordering of mutations is available through another sources, for example, a reconstruction of the cancer phylogeny, then this ordering can be used directly and the first stage of TrackSig can be omitted.

2.1 Estimating the order of acquisition of SNVs

We assume SNVs to be persistent and cumulative, meaning that mutations cannot be reverted to the reference state and no position is mutated twice. This is known as the *infinite sites assumption*[1,18].

Under this assumption, SNVs acquired earlier in the evolution of a cancer will generally be more prevalent in the population of tumour cells. In TrackSig, we sort mutations according to decreasing cancer cell fraction (CCF) and use this as an estimate the order of acquisition of mutations.

2.1.1 Estimating cancer cell fraction

Estimating a SNV's CCF requires both an estimate of its VAF and an estimate of the average number of alleles per cell at the locus where the SNV occurs. In TrackSig, we derive this estimate from a CNA reconstruction provided with the VAF inputs.

To account for uncertainty in a SNV's VAF due to the finite sampling, we model the posterior distribution over its VAF using a Beta distribution:

$$\text{VAF} \sim \text{Beta}(n_{\text{var}}, n_{\text{ref}}), \tag{1}$$

where n_{var} is the number of reads carrying a variant, and n_{ref} is the number of reference reads. To simplify the algorithm, and the subsequent sorting step, we sample an estimate of VAF_i (VAF of SNV i) from this distribution. This gives us a single sampled ordering. With a large number of SNVs, we expect little variability in the estimated exposure trajectory due to uncertainty in the VAFs of individual SNVs. With a smaller number of SNVs, multiple orderings can be sampled and the trajectories combined.

If no CNA reconstruction is available, TrackSig assumes that each SNV is in a region of normal copy number and TrackSig estimates CCFs in autosomal regions by setting:

$$\text{CCF}_i = \frac{2 * \text{VAF}_i}{\text{Purity}}, \tag{2}$$

where Purity is the purity (i.e. proportion of cancerous cells) of the sample. If purity is not provided, TrackSig assumes Purity $= 1$.

If a CNA reconstruction is available, TrackSig uses it when converting from VAF to CCF. In regions of subclonal CNAs, making this conversion requires a phylogenetic reconstruction[16,19]. As such, we filter

SNVs in these regions out when ordering SNVs in order to avoid this time consuming operation. However, TrackSig can make use of orderings of SNVs in regions of subclonal CNAs if provided by a phylogeny-aware method[16, 20]. Also, TrackSig assumes there is a maximum of one variant allele per cell, and thus estimates CCF by setting:

$$\mathrm{CCF}_i = \frac{(2 + \mathrm{Purity} * (\mathrm{CN}_i - 2))}{\mathrm{Purity}} * \mathrm{VAF}_i, \tag{3}$$

where Purity is the purity of the sample, and CN_i is the clonal copy number of the locus. If the clonal CNA increases the number of variant alleles per cell, this will lead to CCFs larger than one. As such, these cases are easily detected and corrected.

TrackSig sorts SNVs in order of decreasing estimated CCF and use the rank of the SNV in this list as a "pseudo-time" estimate of its time of appearance. Note that this estimate will have a non-linear relationship to real time, if the overall mutation rate can varies during the tumour's development. If some of the SNVs can be interpreted as clock mutations, as SNV's rank can be converted into an estimate of real time[9].

2.1.2 Smoothing

To derive a smoothed estimate of the exposure trajectory, TrackSig converts the SNV ordering into a set of time points with overlapping subsets of the SNVs.

TrackSig first partitions the ordered mutations into bins of 100 mutations and interpret each bin as one time point. It then merges bins of 100 mutations into bins of 400 in a sliding window manner: the first time point includes first 400 mutations of the sorted set, the second time point contains mutations 100-500 and so on. The *timeline* of the cancer is the collection of the time points. TrackSig reports signature exposure trajectories as a function of points in the timeline. Note that it does not use any information about subclones when partitioning the SNVs and that it is only using CCFs for the SNV from a single sample.

2.2 Computing exposures to mutational signatures

To estimate exposure trajectories, TrackSig partitions the timelines into sets containing one or more time points. Within each of these sets, it estimates signature exposures using mixture of discrete distributions. Full details of the model are provided in the appendix A. In brief, TrackSig models each signature as a discrete distribution over the 96 types and it treats the mutation count vector over the 96 types as a set of independently and identically distributed samples from a mixture of the discrete distributions corresponding to each signature. The mixing coefficients of these distributions are interpreted as their exposures for the mixture model that produced the set of mutations. TrackSig fits these exposures using the Expectation-Maximization algorithm[21].

2.3 Detecting change-points

TrackSig identifies change-points in the timeline where there are discernible differences in the exposure of mutations in the time points before and after the change-points. Specifically, the change-points delineate the partitions of the timeline into sets of mutations with approximately constant exposures. TrackSig fits exposures for this set, as described above. This procedure generates piecewise constant exposure trajectories for each signature and selects change-points using a greedy algorithm.

The greedy algorithm is as follows. At each stage of the algorithm, given the existing partition of the time points, TrackSig considers adding a new change-point between any adjacent pair of time points where there isn't already one. Each such potential change-point will split one of the existing groups of time points into two sets (i.e., before and after the change-point). To score a potential new change-point, it refits the exposures separately to the two new mutation sets associated with the new split of the time points in the old partition. It computes the difference between the sum of the log likelihoods of the two new sets and the log likelihood of the initial partition. This difference is computed for all possible new change-point locations, and selects the one that gives the largest increase in likelihood. If increased likelihood exceeds the complexity

penalty associated with adding a new partition from the Bayesian Information Criteria (BIC), then the new change-point is kept and the algorithm continues. If not, it stops.

To compute the BIC we note that the piecewise linear model has $(\text{\# changepoints} + 1) \cdot (\text{\# signatures} - 1)$ parameters. Change-points split the timeline into $(\text{\# changepoints} + 1)$ sections. In each section, TrackSig fits the signature exposures, which have to sum to one. Therefore there are $(\text{\# signatures} - 1)$ free parameters per section, or $(\text{\# changepoints} + 1) \cdot (\text{\# signatures} - 1)$ free parameters in total. As such, BIC objective takes the following form:

$$BIC = -2\ln\hat{\mathscr{L}} + (\text{\# changepoints} + 1) \cdot (\text{\# signatures} - 1) \cdot \ln(\text{\# timepoints}) \qquad (4)$$

where $\hat{\mathscr{L}}$ is the likelihood of the current model (see Supplement for details).

Once TrackSig selects the change-points, we can directly compare them to subclone boundaries. Note that the trajectories and the change-points are detected without using any subclonal information, thus permitting independent conclusions about the signature behaviour and its relation to subclones.

2.4 Correcting cancer cell fraction greater than 1

If the number of variant alleles per cell is increased by a clonal copy number change, TrackSig's CCF estimates might be greater than 1. To correct for this, when displaying exposure trajectories, it merges all the time points that have average CCF ≥ 1 into one time point. As such, the first time point can contain more than 400 mutations. To determine a signature exposure at this new time point, TrackSig simply takes an average exposure of all merged time points (those having CCF ≥ 1).

2.5 Bootstrapping to estimating exposure uncertainty

TrackSig estimates uncertainty in the exposure estimates by bootstrapping the mutations and refitting the exposure trajectories. Specifically, it takes the random subset of N mutations by sampling uniformly with replacement from the N unfiltered SNVs in the sample under consideration. Using the pre-assigned CCF estimates, we sort the SNVs in decreasing order, as above, re-partition them into time points and recompute exposure estimates. The trajectories obtained from bootstrapped mutation sets have the same number of time points, however the average CCF for each time point can change. We use these bootstrapped trajectories to compute uncertainty estimates for the sizes of exposure changes.

3 Results

TrackSig was applied to 2,486 whole-genome sequencing samples of 39 cancer types as part of the Pancancer Analysis of Whole Genomes (PCAWG) group. The results of these analyses are described elsewhere[17]. Here we describe the simulations establishing TrackSig's performance characteristics and provide some methodological details of TrackSig's use in PCAWG.

3.1 Choice of mutation signatures

Following *Alexandrov et al.*[2], we classify mutations into 96 types based on their three-nucleotide context. Point mutations fall into 6 different mutation types (i.e., C -> [AGT] and T -> [ACG]) excluding complementary pairs. There are 16 (4*4) possible combinations of the 5' and 3' nucleotides. Thus, SNVs are separated into 96 ($K = 16 * 6 = 96$) types.

Within the context of PCAWG, we use the set of 48 signatures developed by PCAWG-7. The first 30 of those signatures are slightly modified versions of original signatures defined by *Alexandrov et al.*[2,12] and have the same numbering and interpretation. The original 30 signatures are described at COSMIC[1]. Signature analysis methods, including TrackSig, fit exposures for only a subset of the signatures. These signatures are called the *active* signatures. The exposures for the non-active signatures are clamped to zero. For example,

---

[1]http://cancer.sanger.ac.uk/cosmic/signatures

(a) Breast cancer

(b) Chronic lymphocytic leukemia

Figure 1. Exposure trajectories for two samples. Each plot is constructed from VAF data from one tumour sample. Each line is an exposure trajectory that depicts inferred exposures for a single signature (y-axis) as a function of decreasing CCF (x-axis). Signatures are indicated by colour. The thin lines are trajectories from each of 30 bootstrap runs. The bold line depicts the mean exposures across bootstraps. The vertical lines indicate time points in the original dataset, and are placed at the average CCF of their associated mutations. As such, the distance between time points in the plot is inversely proportional to the density of the mutations near that CCF. Changes in exposure trajectories are not necessarily aligned with vertical bars because mean CCFs of time points change across bootstraps. Frequency of changepoints between two vertical bars is indicated by shade, the darker shades indicate higher density of change-points. Subclonal boundaries are shown in red vertical lines. Subclonal information is not used in trajectory calculation and is only shown for comparison. **(a) Breast cancer sample** In clonal signatures remain constant with dominating signature 3 (associated with BRCA1 mutations). In the subclone exposure to signature 3 decreases and is replaced by SNVs associated with APOBEC/AID (signatures 2 and 13). **(b) Chronic lymphocytic leukemia sample** Signature 9 (somatic hypermutation) dominates during clonal expansion and drops from 55% exposure to almost zero in the subclone. Signature 5 compensates for this change.

signature 7 is associated with ultraviolet light has been detected almost exclusively in skin cancers[2]. As such, it is only assigned active status in skin cancers. In our analysis, we use the active signatures reported by PCAWG-7. For analyses based on COSMIC signatures, one can use active signatures per cancer type is provided on COSMIC website. TrackSig can also be used to automatically select active signatures, as described in a later section.

3.2 Signatures with most changing exposures
In this section we analyze the variation of signature exposures on PCAWG data across time and across samples. We compute the maximum change of the signatures in each sample, which is simply the difference between maximum and minimum exposure of the signature. To assess whether a signature change is statistically significant, we perform the following procedure. We permute the mutations in each sample and run the trajectory estimation on the permuted set. Since permuted mutations are not sorted in time, we expect no

change in the exposure trajectories over time. The maximum exposure change that we observe on permuted set of mutations does not exceed 6% in any sample. Therefore, we will consider signature changes below 6% to be insignificant (Fig. 2).

(a)   **(b)**

Figure 2. Distribution of maximum signature changes across 2486 PCAWG samples. The red line shows the threshold of 6%, below which we consider changes to be insignificant. **(a)** Changes on random orders of mutations where we don't expect to see change in exposures. **(b)** Exposure changes in TrackSig trajectories across all samples (on mutations sorted by CCF). Frequency axis shows the number of samples where we observe the certain exposure change.

As shown by Figure 3, samples typically have only two or three signatures with high exposures. These signatures are usually the most variable (up to 80% max change, 11% on average). Other signature have low exposure and remain constant. On average 3.6% of overall exposure is explained by low-exposure signatures (with exposure <5%), most likely reflecting the level of noise in the data.

Figure 3. Left: Mean signature exposures ranked from the largest to the smallest within each sample. Only the top 5 signatures with the highest exposures in a sample are shown. **Right:** Maximum changes of signature exposures for the corresponding signatures on the left plot. The changes below 6% are omitted as insignificant.

3.3 Trends in signature change per cancer type

The majority of samples have a signature change: 69.6% of samples have a max change >6% in at least one signature; 44.6% of samples have change >10%. However, the number of signature changes correlates to some extent with the number of mutations in the sample. Out of samples with less than 10 timepoints (tp) only 29.7% of samples have a change >6% compared to 79.3% across the rest of the samples (see distribution on fig. 4).

Figure 4. Proportion of tumours that have a significant change greater than 6% exposure depending on the number of time points in a sample. Each bar corresponds to the range of number of time points in a sample.

3.4 Bootstrapping

We assess the variability in exposure trajectories by performing bootstrap on the PCAWG data. We sample mutations with replacement from the original set and re-calculate their exposures and change-points. We perform 30 bootstrap runs for each sample. Fig. 1 shows examples of bootstrapped trajectories from two samples (breast cancer and leukemia).

Signature trajectories calculated on bootstrap data are stable. The mean standard deviation of exposure values calculated at each time point is 2.7%. We also evaluate the consistency of signature changes across the entire exposure trajectory: size of signature change and location of the change-point. The mean standard deviation of the signature change is 4.2% across the bootstraps. This standard deviation does not exceed 5% in 74.8% of samples (does not exceed 10% in 99.5% of samples, fig. B.1).

In TrackSig the number of change-points is calculated during exposure fitting does vary across bootstrap samples. We observe 1.04 standard deviation in the number of change-points. To assess the variability in the location of the change-points, we matched nearby change-points between bootstrap samples and measured their average distance in CCF. Because the number of change-points can change between samples, as a reference, we randomly choose one of the samples that has a number of change-points equal to the median number of change-points among all samples. Then, in all other bootstrap runs, we match each change-point to the closest run in the reference. We found that location of the change-points is consistent across bootstraps: on average, change-points are located 0.046 CCF apart from the closest reference change-point.

3.5 Simulations

To test TrackSig's ability to reconstruct the exposure trajectories, we generate a set of simulated samples with known ground truth. Simulations have 50 time points (average number of time points in PCAWG samples). Each simulation has four active signatures. Two of those signatures are 1 and 5, which are nearly always active in the PCAWG samples. For the remaining two signatures, we test all possible combinations of the other 46 signatures. Thus, we have 1035 (= 46 choose 2) different signature combinations.

We generate simulations with 0 to 3 change-points that are placed randomly on the timeline. For each segment on the timeline, we sample signature exposures from a symmetric Dirichlet distribution with all parameters $\alpha_i = 1$, in other words, all exposure vectors are equally likely. Finally, we sample 100 mutations from the discrete distribution derived using the sampled exposures as mixing coefficients for the four signatures.

The simulations mimic the input from the real data (100 mutations per time point). As a part of TrackSig (section 2.1.2), we replace the initial bins of 100 in each time point with smoothed bins of 400. To account for the smoothing in TrackSig, we smooth the exposures in ground truth by taking the mean exposures across four time points in a sliding-window manner.

Next, we run TrackSig on the simulated data and compare the reconstructed exposure trajectories to the ground truth. We remove change-points with small change, that is, where exposures of all signatures change by less than 6% in reconstructed trajectories. This threshold is derived in section 3.2 from permutation analysis.

We computed the absolute difference between predicted exposures and the ground truth at each time point and take the median across all time points and all four signatures. We called this the median per simulation difference. On the simulations with no change-points, the median of these median per simulation differences is 1%. On simulations with 1 to 3 change-points, this median increases slightly to 1.9%. The cumulation distribution of the median per simulation differences is shown in fig. 5.

For the PCAWG data, we report the maximum exposure change (MEC) across exposure trajectory[17]. The maximum change is the difference between maximum and minimum exposure across all time points in a sample. We also report the direction of change (down if maximum occurs before minimum and up otherwise). Here, we evaluate TrackSig's accuracy in these estimates on the simulated data. The MEC discrepancies between the estimated and ground-truth trajectories is less than 6% in 87% of cases across all signatures in all simulations (fig. 6).

To compare the direction of the exposure change, we divide signatures into three categories: with decreasing exposure, increasing exposure and no exposure change (if max change is less than 6%). The direction of maximum change is consistent in 95% of all signatures across all simulations.

To compute number of false positives and false negatives, we use modified criteria that accounts for the sliding-window smoothing. Specifically, we count a true positive detection if at least one of predicted change-points occur with three time points of an actual one. A false negative is when no predicted change-points are within three time points of an actual change. This criteria is identical to the one we use to evaluate whether a change-point supports a subclonal boundary[17]. We deem a predicted change-point a false positive if it occurs more than three time points away from the closest actual change-point.

Table 1 shows the percentage of simulations where we observe the certain number false positives. On average, we observe 0.37 false positives per simulation. We detect 0.02 false negatives on average per simulation.

3.6 Choosing active signatures

Only a subset of signatures are active in a particular sample, and this subset is largely determined by a cancer type. For the analyses reported above, we use a set of active signatures provided by PCAWG, which contains a list of active signatures per sample (on average, four per sample). For COSMIC signatures the list of active signature per cancer type is available on the website. However, such data is frequently unavailable. Here we explore different ways to select the active signatures, comparing them to those selected by PCAWG-7. Note that is all the approaches described below it is sufficient to fit the signatures to overall mutation counts without separating mutations into time points. Once active signatures are selected, they can be used to compute the exposure trajectories across time.

TrackSig supports three ways to determine active signatures. The first is to simply use the full set of signatures to fit the data. The second is to use all signatures reported as active in the cancer type under

(a)   (b)

Figure 5. Simulation results. **(a)** Median exposure difference between the reconstructed trajectories and the ground truth. Lines correspond to the simulations with 0, 1, 2 or 3 change-points. The median in computed across all signatures and time points in the sample. **(b)** Number of estimated change-points versus number of true change-points.

Figure 6. Simulation results. Distribution of discrepancies between estimated trajectories and the ground truth in terms of maximum signature change. Maximum signature change is the difference between the maximum and minimum exposure across all time points in a sample.

consideration. The final is to fit exposures using one of the two previous methods, and use as active signatures only those signatures with exposures greater than a threshold (by default, 5%) in the initial fit. We evaluated each strategy by compared the active signatures selected by TrackSig with those reported by PCAWG-7.

For the first strategy, we used all 48 signatures and we found on average, 44.7% of overall exposure assigned by TrackSig is assigned to the active signatures selected by PCAWG-7. Each incorrect signatures gets 1.3% of exposure on average. In other words, the incorrect exposure is widely distributed among the signatures. Therefore, we recommend constraining the number of signatures by one of the approaches described below.

Fitting only the cancer-specific signatures improves the correspondence to 68.7% of the total exposure on

Table 1. Simulation results. False positives and false negatives versus change-points in the ground truth. Each cell shows the percentage of simulations that have certain number of false positives/negatives (normalized within the column). See main text for definition of positive and negative time points. The last row of the table shows the average number of false positives per simulation.

		# true change-points			
		0	1	2	3
# false positive change-points (FP)	0	62.61	70.63	75.65	81.16
	1	22.42	20.48	19.13	16.52
	2	11.88	7.15	4.35	2.03
	3	2.42	1.45	0.68	0.19
	4	0.68	0.29	0.19	0.10
Avg # of FP per simulation		0.56	0.4	0.31	0.22

		0	1.0	0.99	0.97	0.95
# false negative change-points (FP)	1	0.0	0.01	0.03	0.05
	2	0.0	0.0	0.0	0.001
Avg # of FN per simulation		0.0	0.008	0.0319	0.048

Figure 7. Example of the simulation with three change-points **(a)** Reconstructed sample **(b)** Ground truth

average. Using sample-specific sets reduced the initial set of potentially active signatures from 48 down to 12 on average (ranging from 4 signatures in Lower Grade Glioma to 24 signatures in Liver Cancer). In this case, we observe that signature 5 and 40 are the most prevalent among the incorrect signatures, having the average exposure of 14% and 12.6% respectively in the samples where they are supposed to be inactive.

Finally, we fit exposure for all 48 signatures and then re-fit only those with the high exposure (for instance, >5%), we exactly recover the active signatures reported by PCAWG-7.

Fitting either per-cancer or per-sample signatures results in more exposure mass to be on the correct signatures and speeds up the computations. Therefore, we recommend choosing per-cancer or per-sample signatures instead of using exposures from the full set.

4 Summary and Conclusions

TrackSig reconstructs the evolutionary trajectories of mutation signature exposures by sorting point mutations by their inferred CCF and then partitioning this sorted list into groups of mutations with constant signature exposures. TrackSig estimates uncertainty in the location of the change-points using bootstrap. TrackSig is designed to be applied to VAF data on SNVs from a single sample, however, it can be applied to sorted lists of point mutations derived from subclonal reconstruction algorithms.

Change-points often correspond to boundaries between subclones[17]. By reconstructing changes in mutation exposures, TrackSig can potentially help identify DNA damage repair processes disrupted in the cell and, in doing so, help inform treatment[11].

4.1 Relationship to previous work

Previously methods to track changes in the signature exposure have relied on pre-specified groups of mutations, such as those occurring before or after whole genome duplications[7,9,22,23] or those classified as clonal or subclonal[1,9]. In contrast, TrackSig uses the distributions of mutation types to group mutation. We have shown that this leads to more sensitive detection of changes in signature exposure.

The approaches mentioned above are limited to 1) the samples where the certain events have occurred or to 2) the ability of other methods to reconstruct subclonal structure of the tumour. The number of time point bins remains restricted to the number of subclones (only 2.6% of samples have more than 2 subclones). Compared to previous approaches, TrackSig allows to look at the timeline tumour development at greater resolution, where the number of time points increases with the number of mutations.

Another important innovation of TrackSig is the used of CCF as a surrogate for evolutionary timing. Similar ideas have been used in human population genetics, where variant allele frequency to get relative order of mutations along the ancestral lineage[24]. In population genetics, allele frequency is calculated across individuals, while we calculate VAF across cell population within a single sample. In TrackSig we introduce a way to calculate cancer cell fraction (CCF) using VAF and use it as a timing estimate instead of VAF. We further improve the estimates by correcting them for CNAs.

4.2 Applicability to other mutation types

It is straightforward to generalize TrackSig to other types of mutations, so long as these mutations can be approximately ordered by their evolutionary time and each mutation can be assigned to a single, pre-specified type. In this paper, we ordered SNVs by decreasing CCF. This same strategy could be naturally extended to indels for which the infinite sites assumption is also valid. The infinite sites assumption should also be valid for structural variants (SVs) associated with well-defined breakpoints, thus permitting TrackSig to be used to track exposures to recently defined SV signatures[23]. The CCFs of SVs can be estimated using the VAFs of split-reads mapping to their breakpoints[25]. Because they cover larger genomic regions, infinite sites is less valid for CNAs, although it is possible to approximately order clonal CNAs based on the inferred multiplicity of SNVs affected by them[9].

TrackSig also requires a pre-defined set of mutation signatures, each of which is a probability distribution over the mutation types. However, if these signatures are unavailable, they can be defined by non-negative matrix factorization, or Latent Dirichlet Allocation[26], if counts across mutation types are available from multiple cancer samples.

The alternative way to obtain more comprehensive view of tumour development is to recover evolution tree of the tumour and investigate mutations separately within each node of the tree. The root node corresponds to clonal expansion, while each of the child nodes denote the subclones. This structure can be reconstructed using a variety of algorithms including PhyloWGS[16, 20, 27], which builds a subclone tree based on mutation cancer cell fraction and copy number. These methods assign mutations to one of the nodes of subclonal hierarchy, allowing to analyze mutational signatures independently for each subclone.

4.3 Sensitivity to misorderings of the SNVs

TrackSig assumes that ordering SNVs by CCF recovers the order in which they accumulated in the genomes of ancestral cells, thus, our conclusions are sensitive to the correctness of this assumption. With a large number of SNVs, we do not expect large deviation in exposure trajectories due to a small amount of uncertainty in CCF. Indeed, TrackSig's exposure trajectory varied little in bootstrap samples. For this same reason, we do not expect exposure trajectories to be impacted if a small fractions of SNVs violate the infinite sites assumption due to high, regional mutation rates.

However, these trajectories can be impacted by incorrect ordering of a large numbers of SNVs. These can occur in two ways. First, misordering can also occur if a CNA changes the number of SNV allele's per cell. For example, daughter cells can fail to inherit SNVs in their mother cells due to a loss of heterozygosity (LOH). If a CNA reconstruction is available, TrackSig will correct for any detected clonal LOH when ordering SNVs, and will not attempt to order SNVs in regions affected by subclonal CNAs, thereby resolving this difficulty. However, if a CNA reconstruction is not available, or it is inaccurate, the accuracy of the exposure trajectories can suffer.

Second, some of the SNVs could be in separate branches of the genealogy of the cancer cell, in which case SNVs with smaller CCFs do not necessarily occur in cells containing the SNVs with higher CCFs. Note that due to the Pigeonhole Principle[1], this requires that the sum of the CCFs of the two branching lineages be less than 100%, i.e. at least one of the two lineages will have CCFs less than 50%. As such, branching cannot occur before 50% CCF, and if it does, then the SNV ordering will remain correct for the major branched lineage (in other words, the one with the higher CCF). In general, the minor branch contains a relatively small number of SNVs for many cancers[17]. Note, however, that a symptom of branching could be an oscillations in the exposure trajectories, if there is a change in signatures in the major branch that is not reflected in the minor branch. To avoid this issue, we focus on consistent changes in only one direction (either signature going up or going down). Alternatively, the exposure trajectories can be estimated separately for each subclone. We don't conduct such analysis in this paper.

4.4 Accounting for overall mutation rate

The timelines reconstructed by TrackSig are computed with a fixed number of mutations in each bin. If overall rate of generating mutations in tumour was constant, our timeline would correspond to the real time. However, tumour mutation rate often accelerates throughout development[28, 29]. Although the changing rate does not affect our analysis, the estimates of the pseudo-time might not be linear of the real time.

Estimating changes in overall mutation rate is difficult. A possible way to correct for this is to adjust the time line based on exposures of signatures 1 and 5. It was suggested that signatures 1 and 5 operate as cell "clock" as the number of mutations contributed by these signatures is be proportional to the age of the individual[6]. However, it requires additional data, such as tumour samples from the same patient at different time steps and the medical history when these samples were taken. Determining the association between our pseudo-time estimates and real time is left for further investigation.

Our method TrackSig provides further insight how signature profile changes throughout tumour development. We show that through signatures analysis we can detect major events in tumour evolution, notably,

transitions to a new subclone. Mutational signatures provide a unique way to recover tumour evolution path, track activities of mutational processes, adjust the treatment strategy and detect changes in therapy response.

Acknowledgments

We thank Pancancer Analysis of Whole Genomes (PCAWG) network, and in particular the PCAWG Evolution and Heterogeneity working group, for providing data, analysis and valuable input on this project. We would in particular like to highlight Peter Van Loo, Clemency Jolly, Stefan Dentro, David Wedge, Paul Boutros, Lydia Liu, and Moritz Gerstung who provided valuable feedback during the development of the TrackSig methodology. We would like to acknowledge SciNet as part of Compute Canada for providing computational resources. This research was partially supported by an NSERC operating grant to QDM and is part of the University of Toronto's Medicine by Design initiative, which receives funding from the Canada First Research Excellence Fund (CFREF).

References

1. Nik-Zainal, S. *et al.* The Life History of 21 Breast Cancers. *Cell* **149**, 994–1007 (2012). URL http://linkinghub.elsevier.com/retrieve/pii/S0092867412005272. DOI 10.1016/j.cell.2012.04.023.

2. Alexandrov, L. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421 (2013). URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3776390/. DOI 10.1038/nature12477.

3. Hainaut, P. & Pfeifer, G. P. Patterns of p53 G–>T transversions in lung cancers reflect the primary mutagenic signature of DNA-damage by tobacco smoke. *Carcinogenesis* **22**, 367–374 (2001).

4. Pfeifer, G. P., You, Y.-H. & Besaratinia, A. Mutations induced by ultraviolet light. *Mutation Research* **571**, 19–31 (2005). DOI 10.1016/j.mrfmmm.2004.06.057.

5. Pfeifer, G. P. *et al.* Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene* **21**, 7435–7451 (2002). DOI 10.1038/sj.onc.1205803.

6. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nature Genetics* **47**, 1402–1407 (2015). URL http://www.nature.com/ng/journal/v47/n12/full/ng.3441.html. DOI 10.1038/ng.3441.

7. Alexandrov, L. B. *et al.* Mutational signatures associated with tobacco smoking in human cancer. *Science* **354**, 618–622 (2016). URL http://science.sciencemag.org/content/354/6312/618. DOI 10.1126/science.aag0299.

8. Behjati, S. *et al.* Mutational signatures of ionizing radiation in second malignancies. *Nature Communications* **7**, 12605 (2016). URL http://www.nature.com/ncomms/2016/160907/ncomms12605/full/ncomms12605.html. DOI 10.1038/ncomms12605.

9. Gerstung, M. *et al.* The evolutionary history of 2,658 cancers. *bioRxiv* (2017). URL https://www.biorxiv.org/content/early/2017/07/11/161562. DOI 10.1101/161562. For the Evolution and Heterogeneity Working Group of the Pan-Cancer Analysis of Whole Genomes Initiative (PCAWG-11), https://www.biorxiv.org/content/early/2017/07/11/161562.full.pdf.

10. McPherson, A. *et al.* Divergent modes of clonal spread and intraperitoneal mixing in high-grade serous ovarian cancer. *Nature Genetics* **48**, 758–767 (2016). URL http://www.nature.com/ng/journal/v48/n7/full/ng.3573.html. DOI 10.1038/ng.3573.

11. Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on mutational signatures. *Nature Medicine* **23**, 517–525 (2017). DOI 10.1038/nm.4292.

12. Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Campbell, P. J. & Stratton, M. R. Deciphering Signatures of Mutational Processes Operative in Human Cancer. *Cell Reports* **3**, 246–259 (2013). URL http://www.cell.com/cell-reports/abstract/S2211-1247(12)00433-0. DOI 10.1016/j.celrep.2012.12.008.

13. Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C. deconstructSigs: delineating mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma evolution. *Genome Biology* **17**, 31 (2016). URL https://doi.org/10.1186/s13059-016-0893-4. DOI 10.1186/s13059-016-0893-4.

14. Shiraishi, Y., Tremmel, G., Miyano, S. & Stephens, M. A Simple Model-Based Approach to Inferring and Visualizing Cancer Mutation Signatures. *PLOS Genetics* **11**, e1005657 (2015). URL http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1005657. DOI 10.1371/journal.pgen.1005657.

15. Yates, L. R. *et al.* Genomic Evolution of Breast Cancer Metastasis and Relapse. *Cancer Cell* **32**, 169–184.e7 (2017). DOI 10.1016/j.ccell.2017.07.005.

16. Deshwar, A. G. *et al.* PhyloWGS: Reconstructing subclonal composition and evolution from whole-genome sequencing of tumors. *Genome Biology* **16**, 35 (2015). URL http://dx.doi.org/10.1186/s13059-015-0602-8. DOI 10.1186/s13059-015-0602-8.

17. Dentro, S. C. *et al.* Pervasive intra-tumour heterogeneity and subclonal selection across cancer types. *To be submitted* (2017). For the Evolution and Heterogeneity Working Group of the Pan-Cancer Analysis of Whole Genomes Initiative (PCAWG-11).

18. Jiao, W., Vembu, S., Deshwar, A. G., Stein, L. & Morris, Q. Inferring clonal evolution of tumors from single nucleotide somatic mutations. *BMC Bioinformatics* **15**, 35 (2014). URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3922638/. DOI 10.1186/1471-2105-15-35.

19. Roth, A. *et al.* PyClone: statistical inference of clonal population structure in cancer. *Nature Methods* **11**, 396–398 (2014). URL http://www.nature.com/nmeth/journal/v11/n4/full/nmeth.2883.html?foxtrotcallback=true. DOI 10.1038/nmeth.2883.

20. Jiang, Y., Qiu, Y., Minn, A. J. & Zhang, N. R. Assessing intratumor heterogeneity and tracking longitudinal and spatial clonal evolutionary history by next-generation sequencing. *Proceedings of the National Academy of Sciences of the United States of America* **113**, E5528–5537 (2016). DOI 10.1073/pnas.1522203113.

21. Dempster, A. P., Laird, N. M. & Rubin, D. B. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society. Series B (Methodological)* **39**, 1–38 (1977). URL http://www.jstor.org/stable/2984875.

22. Jamal-Hanjani, M. *et al.* Tracking the evolution of non-small-cell lung cancer. *New England Journal of Medicine* **376**, 2109–2121 (2017). DOI 10.1056/NEJMoa1616288. PMID: 28445112.

23. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* **534**, 47–54 (2016). URL http://www.nature.com/nature/journal/v534/n7605/full/nature17676.html. DOI 10.1038/nature17676.

24. Harris, K. & Pritchard, J. K. Rapid evolution of the human mutation spectrum. *eLife* **6** (2017). URL http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5435464/. DOI 10.7554/eLife.24284.

25. Cmero, M. *et al.* Svclone: inferring structural variant cancer cell fraction. *bioRxiv* (2017). URL https://www.biorxiv.org/content/early/2017/08/04/172486. DOI

10.1101/172486. https://www.biorxiv.org/content/early/2017/08/04/172486.full.pdf.

26. Blei, D. M., Ng, A. Y. & Jordan, M. I. Latent dirichlet allocation. *J. Mach. Learn. Res.* **3**, 993–1022 (2003). URL http://dl.acm.org/citation.cfm?id=944919.944937.

27. Satas, G. & Raphael, B. J. Tumor phylogeny inference using tree-constrained importance sampling. *Bioinformatics* **33**, i152–i160 (2017). URL +http://dx.doi.org/10.1093/bioinformatics/btx270. DOI 10.1093/bioinformatics/btx270. /oup/backfile/content_public/journal/bioinformatics/33/14/10.1093_bioinformatics_btx270/4/btx270.pdf.

28. Alberts, B. *Essential Cell Biology*, vol. Unit 5.5 (Garland Science, 2010). URL https://books.google.com/books?id=RAwvAQAAIAAJ.

29. Wodarz, D., Newell, A. & Komarova, N. Passenger mutations can accelerate tumor suppressor gene inactivation in cancer evolution. *bioRxiv* (2017). URL https://www.biorxiv.org/content/early/2017/10/13/202531. DOI 10.1101/202531. https://www.biorxiv.org/content/early/2017/10/13/202531.full.pdf.

30. Blei, D. M. Probabilistic topic models. *Commun. ACM* **55**, 77–84 (2012). URL http://doi.acm.org/10.1145/2133806.2133826. DOI 10.1145/2133806.2133826.

A Computing exposure to mutational signatures

We apply topic modeling[30] to infer signature exposures. Withing the time point, we separate mutation separate into K mutation types. Mutation types relate to vocabulary in topic modeling. The types used in TrackSig are described in section 3.1. Then we use mixture of discrete distributions to infer signature exposures. We describe this model below.

We represent each mutation as a K-dimensional binary vector – "one-hot-encoding" of a mutation type. "One-hot-encoding" of a mutation of type k is a binary vector where k-th component is equal to 1, and other components are zeros. We will denote $\mathbf{x}^{(n)}$ to be the "one-hot-encoding of mutation n. A sample containing N mutations is represented as a $N \times K$ binary matrix $\mathbf{X}$, where each column corresponds one mutation.

$$\mathbf{x}^{(n)} = \begin{bmatrix} 0 \\ 0 \\ ... \\ 1 \\ ... \\ 0 \end{bmatrix} ; \quad \mathbf{x}_k^{(n)} = \begin{cases} 1, & \text{mutation } n \text{ belongs to type } k \\ 0, & \text{otherwise} \end{cases} \tag{5}$$

A mutation process is represented as a distribution over mutation types, known as a "*mutation signature*". We will denote signature multinomials as K-dimensional probability vectors $\boldsymbol{\mu}_i$, where $i = \{1..M\}$ is an index over signatures. Signatures are fixed and are not updated during the training.

We aim to estimate signature *exposures* $\boldsymbol{\pi}$ – the proportion of mutations generated by each signature.

We will use the following notation:

K – number of mutation types
M – number of signatures
N – number of mutations
$\mathbf{x}^{(n)}$ – K-dimensional binary vector of mutation n
$x_k^{(n)}$ – k-th component of vector $\mathbf{x}^{(n)}$
$\boldsymbol{\mu}_i$ – i-th signature (K-dimensional vector)

μ_{ik} – k-th component of vector $\boldsymbol{\mu}_i$
$\boldsymbol{\pi}$ – signature exposures (mixture coefficients, M-dimensional vector)
π_i – i-th component of $\boldsymbol{\pi}$ (signature exposure of signature i)
z_n – signature assignment for mutation n

We represent mutation matrix $\mathbf{X}$ as a mixture of signature multinomials $\boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K$ with mixture coefficients $\boldsymbol{\pi}$:

$$\mathbf{X} \sim \text{Multinomial}(N; \sum_{i=1}^{M} \pi_i \, \boldsymbol{\mu}_i) \tag{6}$$

We denote z_n to be the signature assignment of mutation n. The probabilities of mutation n to be assigned to i-th signature are equal to the mixing coefficients:

$$p(z_n = i | \boldsymbol{\pi}) = \pi_i; \quad i \in \{1..M\} \tag{7}$$

The probability of a mutation n to be generated by signature i is given by:

$$p(\mathbf{x}^{(n)} | z_n = i, \boldsymbol{\pi}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K) = \prod_{k=1}^{K} \mu_{ik}^{x_k^{(n)}}; \quad i \in \{1..M\}; \quad n \in \{1..N\} \tag{8}$$

Then the log likelihood of the collection of mutations in a sample:

$$\log \mathscr{L}(\mathbf{X} | \boldsymbol{\pi}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K) = \sum_{n=1}^{N} \log p(\mathbf{x}^{(n)} | \boldsymbol{\pi}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K) =$$
$$\sum_{n=1}^{N} \log \sum_{i=1}^{M} p(\mathbf{x}^{(n)} | z_n = i, \boldsymbol{\pi}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K) p(z_n = i | \boldsymbol{\pi}) \tag{9}$$

To estimate the exposures, we fit mixing coefficients $\boldsymbol{\pi}$ in each bin using Expectation-Maximization (EM) algorithm[21]. The EM algorithm iterates between updating a posterior distribution over z_n and updating an estimate of the mixing coefficients $\boldsymbol{\pi}$

We start with initializing EM algorithm with uniform mixing coefficients:

$$\pi_i^{(0)} = \frac{1}{M}; \quad i \in \{1..M\} \tag{10}$$

Then, we repeat the following E-step and M-step until the algorithm converges.

In E-step, at the t-th iteration, the posterior probabilities of mutation assignments to signatures are estimated as such:

$$p(z_n = i | \mathbf{x}^{(n)}, \boldsymbol{\pi}^{(t-1)}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K) = \pi_i^{(t-1)} \prod_{k=1}^{K} \mu_{ik}^{x_k^{(n)}}; \quad i \in \{1..M\}; \quad n \in \{1..N\} \tag{11}$$

In M-step we update the estimates of the mixing coefficients:

$$\pi_i^{(t)} = \frac{1}{N} \sum_{n=1}^{N} p(z_i | \mathbf{x}^{(n)}, \boldsymbol{\pi}^{(t-1)}, \boldsymbol{\mu}_1, ..\boldsymbol{\mu}_K); \quad i \in \{1..M\} \tag{12}$$

The algorithm has converged when the value of π is updated by less than 0.001 between iterations. The resulting mixture coefficients as the exposures of the mutational signatures. We show the exposures as percentage for the convenience of interpretation.

B Supplementary figures and tables

Table 2. Simulation results. Predicted change-points versus change-points in the ground truth. Each cell shows the percentage of simulations which have certain number of estimated change-points (normalized within a column). Note that due to smoothing, there might be several predicted change-points that correspond to the same change-point in the ground truth. In this case, predicted change-points have to be located no more than 3 time points away from the ground truth.

		# true change-points			
		0	1	2	3
	0	62.61	0.58	0.00	0.00
	1	22.42	15.27	1.06	0.00
	2	11.88	50.05	9.37	1.06
# predicted	3	2.42	24.25	28.12	4.83
change-	4	0.68	7.73	36.91	19.32
points	5	0.00	1.74	18.65	32.08
	6	0.00	0.39	4.83	27.83
	7	0.00	0.00	0.97	11.30
	>=8	0.00	0.00	0.10	3.57

(a)   (b)

Figure B.1. Discrepancies in signature exposures on bootstrap data. **(a)** Standard deviations of signature exposures at each time point for each signature across bootstraps. **(b)** Standard deviations of exposure *change* at each time point for each signature across bootstraps.