Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

Authors:

Ryan K Waples*, Anders Albrechtsen*,[1], Ida Moltke*,[1]

*Department of Biology, Section for Computational and RNA Biology, University of Copenhagen, 2200 Copenhagen N, Denmark,

[1]Corresponding authors: Department of Biology, University of Copenhagen, Ole Maaloes Vej 5, 2200 Copenhagen N, Denmark. E-mail: albrecht@binf.ku.dk; and ida@binf.ku.dk

Abstract

Knowledge of how individuals are related is important in many areas of research and numerous methods for inferring pairwise relatedness from genetic data have been developed. However, the majority of these methods were not developed for situations where data is limited. Specifically, most methods rely on the availability of population allele frequencies, the relative genomic position of variants, and accurate genotype data. But in studies of non-model organisms or ancient human samples, such data is not always available. Motivated by this, we present a new method for pairwise relatedness inference, which requires neither allele frequency information nor information on genomic position. Furthermore, it can be applied to both genotype data and to low-depth sequencing data where genotypes cannot be accurately called. We evaluate it using data from SNP arrays and low-depth sequencing from a range of human populations and show that it can be used to infer close familial relationships with a similar accuracy as a widely used method that relies on population allele frequencies. Additionally, we show that our method is robust to SNP ascertainment, which is important for application to a diverse range of populations and species.

Introduction

The ability to infer the familial relationship between a pair of individuals from genetic data plays a key role in several research fields. In conservation biology, it is used to design breeding programs that minimize inbreeding (Kardos *et al.* 2015), in archeology it is helpful to understand burial patterns and other cultural traditions (Baca *et al.* 2012; Sikora *et al.* 2017), and in population and disease genetics it is often used to exclude relatives, because many analysis methods within those fields assume all sampled individuals are unrelated and violations of this assumption can lead to wrong conclusions (Balding 2006).

Numerous pairwise relatedness inference methods have been developed, e.g., Thompson (1975), Lee (2003), Purcell *et al.* (2007), Albrechtsen *et al.* (2009), Manichaikul *et al.* (2010), Stevens *et al.* (2011), Korneliussen and Moltke (2015), Conomos *et al.* (2016), Dou *et al.* (2017) and many are available in popular software packages, like PLINK (Purcell *et al.* 2007), and KING (Manichaikul *et al.* 2010). Most of these methods estimate either the three relatedness coefficients k_0, k_1 and k_2, or the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$ for each pair of diploid individuals, where k_0, k_1 and k_2 are the proportions of the genome where a pair of individuals share 0, 1, or 2 alleles identical by descent (IBD) (Thompson 2000). By definition, alleles are IBD when they are identical due to *recent* shared ancestry, but because alleles can also be identical due to older shared ancestry or recurrent mutations, IBD status cannot be directly observed. Therefore, the pairwise relatedness coefficients and the kinship coefficient have to be estimated, which can be done from patterns of observed genetic identity (identity by state; in short IBS) together with population allele frequencies. Once estimated, these relatedness statistics, k_0, k_1, k_2, and the kinship coefficient can be used to infer familial

relationships by comparison to the expectation of the statistics under different familial

relationships (Hill and Weir 2011).

Although inference of relatedness is of wide interest, most existing methods are not

immediately applicable in studies with limited data or genetic resources. First, most

existing methods require that the allele frequencies of the source population are known.

For most studies in modern humans this is not a problem. However, in studies of

ancient humans or other species, accurate estimates of allele frequencies are not always

available, due to limited number of samples.

Second, several existing methods examine loci in sliding windows or use hidden Markov

models to leverage the non-independence of allele sharing along the genome between

relatives, (e.g., Stevens *et al.* 2011; Monroy Kuhn *et al.* 2017). These methods are

powerful, but require information about the genomic position of variable sites, and for

non-model organisms high-quality reference genome assemblies are often not available.

Finally, nearly all existing methods --both frequency-based and others-- require

genotype data. However, in sequencing studies, samples are often only sequenced to low

depth due to cost and technical issues, which makes it infeasible to call genotypes

accurately (Nielsen *et al.* 2012), precluding the use of these methods. There are a few

methods that estimate relatedness from low depth data by utilizing genotype likelihoods

(e.g., Korneliussen and Moltke 2015), or by using imputed genotype dosages (Dou *et al.*

2017). However, these methods function by leveraging access to hundreds or thousands

of samples to estimate allele frequencies or perform accurate imputation from low depth

sequencing data. They are not designed to apply to datasets with a low number of

samples.

Hence, most existing methods to infer close familial relationships are not immediately

applicable in studies where data is limited, including many studies of non-model

organisms and ancient humans. One of the few exceptions is a simple but elegant test

for pairwise relatedness proposed in Lee (2003), that requires neither information on

allele frequencies nor the relative genomic position of the variable sites. This test relies

entirely on the relative frequency of different genotype combinations within the two

individuals and thus only requires genotype data from the two target individuals. While

useful, this test does not provide any means to distinguish between different types of

close familial relationships; it only provides a statistical test for relatedness vs the null

hypothesis of unrelated.

To our knowledge, there are only two published methods that can be used to distinguish

between different types of familial relationships when neither allele frequencies nor

information about the relative genomic position of sites is obtainable. The first of these

two methods consists of plotting the proportion of the genomic sites in which the two

individuals share both alleles IBS (which we will denote IBS2) versus the proportion of

the genomic sites in which they share zero alleles IBS (which we will denote IBS0). This

method was used in Rosenberg (2006), where it was applied to the Human Genome

Diversity Project (HGDP) dataset. In the resulting plot of the HGDP data (figure 1 of

Rosenberg (2006)), pairs of individuals with the same relationship category form

distinct clusters in such a way that it is possible to identify parent-offspring pairs, full

sibling pairs, and to a lesser extent more distant relationships such as half-

siblings/avuncular/grandparent-grandchild and first cousins.

The second existing method is based in part on the KING-robust kinship estimator

(Manichaikul *et al.* 2010), which was developed to overcome biases in kinship estimates

caused by cryptic population and family structure. More specifically, the KING-robust

estimator was developed to be robust to admixture, i.e. violations of the assumption that

the genotypes of the pair of individuals are sampled from a single homogenous

population, but in practice it has been shown to struggle when applied to admixed

samples (Thornton *et al.* 2012; Moltke and Albrechtsen 2014; Conomos *et al.* 2016).

However, while the KING-robust kinship estimator was not developed with the aim of

providing a kinship estimator that works when allele frequencies within a single

population are unknown, it is directly applicable to this scenario as long as samples are

from the same homogenous population. The reason for this is that, like the test

suggested in Lee (2003), it relies only on the genotype combinations within the two

target individuals and does not require knowledge about of allele frequencies.

Importantly, Manichaikul *et al.* (2010) show it is possible to infer if a pair of individuals

are parent-offspring, full sibling, half siblings/avuncular/grandparent-grandchild,

cousins, or unrelated by jointly considering the KING-robust estimator and the fraction

of sites IBS0 with both these statistics estimated from SNP array data without using any

information about allele frequencies. For example, see figure 3A in Manichaikul *et al.*

(2010), a scatterplot of KING-robust kinship vs the fraction of sites IBS0.

However, both the existing methods discussed above have two key limitations. First, like

most other pairwise relatedness estimation methods, they were developed for accurately

called genotypes. For example, the KING software (Manichaikul *et al.* 2010)

implementing the KING-robust kinship estimator requires accurate genotype data as

input, which can be problematic for studies where only moderate or low-depth

sequencing data is available. Second, both methods rely on estimates of the fraction of

the genomic sites in which the pair of individuals share zero alleles identical by state,

IBS0. This can be problematic because this fraction is highly sensitive to the SNP

ascertainment procedure. This means that the results of the methods are platform-

dependent and are likely to differ between different SNP arrays and especially between

SNP array and sequencing datasets. And in turn, this among other things, means that it

can be difficult to distinguish between full siblings and parent-offspring pairs using

KING.

Motivated by the outlined limitations to the existing methods, we present a method for

relationship inference that - unlike most existing methods - relies neither on allele

frequencies nor on information about the relative position of the variant sites, and

which – unlike other frequency-free methods – 1) is applicable even to sequencing data

of so low depth that accurate genotypes cannot be called from it and 2) is robust to

ascertainment bias.

The method is inspired by previous methods; it uses the KING-robust kinship estimator

and a statistic R0, which is similar to the test statistic from the test for relatedness

suggested by Lee (2003). However, the method is new in two important way. First,

besides relying on the two statistics, R0 and KING-robust kinship, it also relies on a

third new statistic, R1. More specifically, the method consists of using two combinations

of these three statistics, R1-R0 and R1-KING-robust kinship, to infer relationships, and

it is this combination of statistics that makes the method robust to ascertainment bias.

In particular, the inclusion of R1 helps distinguish between full siblings and parent

offspring. Secondly, while the new method can straightforwardly be applied to accurate

genotype data, like other similar methods, we also present two ways to estimate the

three statistics directly from sequencing data that takes the uncertainty of genotypes

into account. This is what makes the method applicable to low-depth sequencing data as

well.

In the following we first fully describe the three statistics, R0, R1 and KING-robust

kinship, how they can be estimated and other methodological details. Next, using

publicly available SNP array data, we show that the new method provides similar

accuracy to the commonly used frequency-based method implemented in PLINK,

without utilizing any information on allele frequencies or genomic positions. Then using

sequence data from the 1000 Genomes, we show that the three statistics can be

estimated directly from sequencing data of low depth (~4x), here defined as depth

insufficient for genotype calling. Moreover, we show that the resulting estimates are

useful and that this is not the case for estimates obtained from genotypes called from the

same data. Next, using different subsets of the same data, we show that this new

method, unlike previous similar methods, is robust to SNP ascertainment. Finally, we

discuss some potential applications and limitations of the new method.

Methods and materials

The R0, R1, and KING-kinship statistics

The method for relationship inference we propose consists of first estimating three

statistics called R0, R1 and KING-robust kinship from genetic data, then constructing

two plots: R1 vs R0 and R1 vs KING-robust kinship, and finally interpreting these plots.

To define the three statistics, R0, R1 and the KING-robust kinship, we note that for a

pair of diploid individuals, all diallelic sites fall into one of nine genotype combinations;

all possible combinations of the two individuals carrying 0, 1, or 2 copies of a specific

allele, e.g. the ancestral allele. For a pair of individuals, we can therefore fully

characterize their genome-wide IBS-sharing pattern in nine counts or proportions

denoted: A, B, C, D, E, F, G, H, and I (Figure 1A), similar to a two-dimensional site-

frequency spectrum (SFS) across the two individuals. This is important to note because

R0 and R1 statistics are defined as simple functions of a subset of these nine counts as

shown in Figure 1B and C, and the KING-robust kinship statistic, originally defined by

Manichaikul *et al.* (2010), can also be re-formulated as a function of these 9 values

(Figure 1D).

The new method is motivated by several observations. First, the expected values of A-I

vary depending on the familial relationship. Consequently, so do functions of these

counts, including R0, R1 and KING-robust kinship. Specifically, for two non-inbred

individuals from the same homogenous population the pairs (R1, R0) and (R1, KING-

robust kinship), both – just like (k_0, k_1, k_2) - have the property that the joint expectation

ranges for the five relationship categories parent-offspring (PO), full siblings (FS), half

siblings/avuncular/grandparent-grandchild (HS), first cousins (C1) and unrelated (UR)

do not overlap (Figure 2, for derivations see supplementary text) regardless of the

underlying allele frequency spectrum. Second, this method requires neither population

allele frequencies nor information about the relative position of the genomics sites

analyzed. The only information that is required to estimate the three statistics is genetic

data from the two individuals of interest. Third, we expect these statistics to be robust to

SNP ascertainment because they are ratios computed from sites that are variable within

the two samples and should thus be unaffected by the number of non-variable sites and

because the (unknown) underlying frequency spectrum should only have a limited effect

on these ratios. Finally, while the counts of the nine genotype combinations, A-I, and

thus R0, R1, and the KING-robust kinship estimator can be calculated from genotype

data, they can also be estimated directly from next-generation sequencing (NGS) data

based on the expected number of sites with each genotype combination. This makes the

approach appropriate even when the sequencing depth is too low for accurate genotype

calling (see below for methodological details).

202   Estimation from sequencing data

The nine genotype combinations, A-I, and thus R0, R1, and the KING-robust kinship for

a pair of individual, can be estimated directly from NGS data via the use of genotype

likelihoods calculated from the aligned sequencing reads. By using genotype likelihoods

it is possible to account for the genotype uncertainly inherent to low-depth NGS data.

We used two distinct, but related approaches to estimate these statistics from

sequencing data that both build on this idea.

The first approach, which we denote the IBS-based approach, considers all ten possible

genotypes at a diallelic site for each of the two individuals of interest and consists of a

maximum-likelihood (ML) estimation of the counts of each of the 100 (10 × 10) possible

genotype pairs (for details see supplementary text). To perform the ML estimation we

used an EM algorithm, which we have added to the ANGSD software package under the

name IBS. After obtaining the estimate of the counts of all the 100 possible genotype

pairs, we convert them into estimates of the genotype combinations A-I, by summing

over the counts that correspond to each combination. For example, the genotype pairs:

AA/AA, CC/CC, GG/GG and TT/TT all contribute to cells A or I of Figure 1. Counts

corresponding to genotype pairs with more than two different alleles (e.g. AC/AG) were

discarded. The advantage of this IBS-based approach is that it does not require

specification of a known allele at each site, and can thus be applied to nearly any

sequencing data set, even with low depth, without any prior information on the alleles at

each site.

The second approach, which we denote the SFS approach, consists of performing ML

estimation of the two-dimensional site-frequency spectrum (SFS) (Nielsen *et al.* 2012).

To find the ML estimate of the SFS, we used an expectation-maximization (EM) method

implemented in the ANGSD software package under the name realSFS (Korneliussen *et*

al. 2014). The SFS method requires one allele to be specified for each site e.g., the

ancestral, consensus or reference allele. The model underlying this method assumes that

all genotypes for each site have the possibility of containing this specified allele and up

to one other unspecified allele. To specify the alleles that do exist at each site, we used

the consensus sequences from the highest depth individual (NA19042) and restricted

our analysis to sites where the depth in this individual was at least three.

For both these genotype likelihood-based methods, we analyzed each chromosome

separately due to computational constraints. We used the original `GATK` genotype

likelihood model (Mckenna *et al.* 2010) with independent errors, as implemented in

`ANGSD`, to calculate the likelihood of sequencing data for each possible genotype. We also

tested the `samtools` genotype likelihood model (Li 2011) with a more complicated error

structure, but found that it produced worse results in this case (data not shown). Both

`IBS` and `realSFS` produces the expected counts for each value A-I and the values for

each chromosome were summed over to produce a genome-wide estimate. For example

command lines for each analysis, see supplemental text.

Confidence intervals

All of the above estimation methods treat each site as independent. This assumption

should not affect our expectation of each statistic (Wiuf 2006), but statistical non-

independence (e.g. linkage disequilibrium and IBD) does affect standard estimates of

uncertainty. To quantify the uncertainty, we therefore estimated confidence intervals for

all statistics using a block-jackknife procedure. Confidence intervals were calculated by

leaving each chromosome out (chromosome jackknife), which takes both the IBD and

the LD correlation into account. The weighted block jackknife variance estimator of

Busing *et al.* (1999) was used to estimate the variance from the distribution of estimates

for each statistic. The square root of this variance was interpreted as the standard error

in our estimate.

Application to real datasets

254   To assess the utility of the new method, we applied it to two different publicly available

255   datasets: SNP array data from seven HGDP populations (Rosenberg 2006), and

256 sequencing data from five related individuals from the Luhya in Webuye, Kenya (LWK)
257 population of the 1000 Genomes Project phase 3 (The Genomes Project 2015).

HGDP SNP array

The HGDP SNP array dataset was downloaded from `ftp.cephb.fr/hgdp_supp1` (accessed
Jan. 13, 2017) and followed the quality control steps described in Rosenberg (2006) to
exclude mislabeled and duplicate samples. We selected seven populations from the
HGDP based on the presence of several close familial relationships; five non-African
populations: Surui, Pima, Karitiana, Maya, and Melanesian, and two African
populations: Mbuti Pygmies and Biaka Pygmies.

To ensure a fair comparison to the identity by descent inference method in PLINK, and
by proxy to other commonly used methods, we constructed data sets where these
methods have been shown to perform well. Specifically, we excluded individuals
showing obvious signs of admixture or inbreeding from the selected HGDP populations
as well as genotypes that may be inaccurate. Specifically, we ran `ADMIXTURE` (Alexander
et al. 2009) separately for each of the seven target population also including French,
and Han samples (as well as Yoruban samples the non-African target populations) and
excluded 16 individuals with >5% contribution from more than one ancestry
component. Then we estimated the inbreeding coefficient, F, for each of the remaining
individuals using PLINK and excluded two individuals with F > 0.0625. This left us with
a total of 142 individuals from the seven populations: Surui N=20, Pima = 20, Karitiana
N=21, Maya N=16, Melanesian N=19, Biaka Pygmies N=31 and Mbuti Pygmies N=15.
Finally, for each of the seven population samples we constructed a final set of accurate
genotypes by retaining genotypes from autosomal loci with genotyping rate (>99%),
MAF (>5%), HWE ($p > 10^{-4}$, with midp adjustment).

The R0, R1, and KING-robust kinship statistics for each of the 2902 within-population

pairs of individuals were then calculated from all sites where both individuals had non-

missing genotypes.

1000 Genomes sequencing data

To get sequencing data from several different relationship categories, we selected five

individuals from two families in the Phase 3 1000 Genomes (1000G) Luhya in Webuye,

Kenya (LWK) population: NA19027, NA19042, NA19313, NA19331, NA19334. Across

the five individuals, there is one pair of half-sibs (NA19027 & NA19042), and a separate

trio of related individuals with a pair of full siblings (NA19331 & NA19334), one parent-

offspring relationship (NA19313 & NA19331), and another unspecified second-degree

relationship (NA19313 & NA19334), possibly avuncular (The Genomes Project 2015).

These stated relationships leave six unrelated pairs among the five individuals.

For each pair of the five LWK individuals we estimated the R0, R1 and KING-robust

kinship statistics in five different ways: 1&2) by applying the two different sequencing-

based approaches described above to the 1000G aligned sequence files (~4x bam files),

3) by simple genotype counting based on the phased and high-quality curated genotypes

provided in the hg37 1000G VCF files, 4) genotype counting based on the subset of sites

in 3 that overlap with the Illumina 650Y sites for the HGDP data (to investigate

ascertainment, see below) and 5) by re-calling genotypes from the same 1000G bam files

and then simple counting from the called genotypes. In set 5 genotypes were called in a

basic manner meant to mimic data from a species with a reference genome but few

other genetic resources. Specifically, we used `samtools mpileup` (v1.3.1) to summarize

the reads overlapping each position, and `bcftools call` (v1.3.1) to assign the most

likely genotype at each position. We used mostly default settings; non-default flags to

`samtools` specified skipping indel positions. Non-default flags to `bcftools` specified

using the consensus caller and requesting only variant sites. For all sequence-based

analyses we only considered reads with a minimum phred-scaled quality score of 30 and

bases with minimum phred-scaled quality score of 20 and we restricted our analyses to

regions with a GEM 75mer mappability of 1 (Derrien *et al.* 2012). All of these methods

and filters can be applied to any reference-based sequencing study.

Assessing the effect of SNP ascertainment

To evaluate the effect of SNP ascertainment, we created a subset of the high-quality

genotype data from the five 1000 Genomes individuals. We selected the sites that

overlap with the Illumina 650Y array that was used for the HGDP and estimated our

three relatedness statistics. We compared the results from this subset of HGDP sites to

results for the full genotype dataset and also to the sequence-based analyses. For an

additional comparison, we also performed the same comparison for the methods

presented in Rosenberg (2006) and Manichaikul *et al.* (2010) by constructing

scatterplots by their methods.

Comparison to other methods

To get a categorization of relationships for the HGDP genotype dataset described above

based on a standard, commonly used allele frequency-based method, we first applied

the allele frequency based relatedness estimation program `PLINK` (v1.9) (Chang *et al.*

2015) to the individuals from each population separately to estimate the genome-wide

IBD fractions k_0, k_1, k_2, (denoted Z0, Z1 and Z2 in PLINK (Purcell *et al.* 2007)). Next, we

applied the relationship criteria proposed in Table 1 of Manichaikul *et al.* (2010) to the

relatedness coefficient estimates: the estimated k values were combined into an estimate

of the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$, and a relationship degree was assigned to each

pair of individuals based on comparing the estimated kinship coefficient to the criteria

in the table. Parent-offspring and full siblings were differentiated based on k_2 values.

This provided us with a categorization into five categories: parent offspring (PO), full

siblings (FS), half-siblings / avuncular / grandparent-grandchild (HS), first cousin (C1),

and unrelated (UR). To achieve additional resolution, we further divided the last

category (UR) into two: distantly related (DR) and unrelated (UR). We did this by

simply extending the logic behind the criteria proposed above. Specifically, we set the

kinship threshold between DR and UR to $1/2^{13/2}$, which corresponds to including 2nd to

5th degree relatives in the DR category.

To estimate the statistics for identifying related individuals proposed by Rosenberg

(Rosenberg 2006) and KING (Manichaikul *et al.* 2010) we note that they can all be

calculated directly from the same nine proportions, A-I, as R1, R2 and King-robust

kinship:

KING-kingship = (E − 2(C+G))/ (B+D+H+F+2E)

Fraction IBS0 = (C+G)/(A+B+C+D+E+F+G+H+I)

Fraction IBS2 = (A+E+I)/(A+B+C+D+E+F+G+H+I)

We used these formulas, because they allow for estimation of the statistics directly from

sequence data in the same manner as for R0, R1 and KING-robust kinship. This allowed

us to use sequence data to compare all the methods on a level playing field.

Results

348 To assess to how the new method works in practice, we applied it to real data from

349 different data generation platforms and compared the results to those of other methods.

Application to SNP array data

We first applied the method to SNP array data from the Human Genome Diversity Project (HGDP) to see how the method works on data, on which standard, commonly used allele frequency-based methods like PLINK are known to perform well. More specifically, we applied it to accurately called genotypes from unadmixed and uninbred samples from seven populations originating from the HGDP. This resulted in the R1-R0 and R1-KING-robust kinship plots shown in Figure 3A and B (for population-specific plots see Figures S1 and S2). Because the true relationships for the pairs of individuals are not known, we colored each point in Figure 3A and B not according to the truth, but according to the relationship category inferred using a standard, commonly used allele frequency-based method (Figure 3C, Figure S3, for further details see Methods). Since there are least 15 individuals in each of the seven selected HGDP populations, the allele frequency estimates for these populations should be reasonably accurate even with some relatedness. Thus, although this classification and coloring scheme does not necessarily reveal the true relationships for all pairs (Hill and Weir 2011) it should provide a fair picture of how well our method performs in comparison to a standard, commonly used allele frequency-based method.

In general, points representing pairs of individuals that according to the allele frequency-based method have the same relationship, clearly cluster together on both the R1-R0 plot and the R1-KING-robust kinship plot (Figure 3) and near their expected ranges. More specifically, almost all pairs identified as parent-offspring (PO) by the frequency-based method are easy to identify as such in both the R1-R0 plot and the R1-KING-robust kinship plot, which is not the case when only a single statistic is used (see also Figures S1 and S2). The same is true for full siblings (FS). Points classified as half-siblings/avuncular/grandparent-grandchild (HS) or first cousins (C1) by the frequency-based method have a minimal overlap with each other and with less-related pairs (see

highlighted areas of Figure 3A and B). But notably the few pairs of individuals that are

difficult to classify are the same pairs as those that are edge cases for the allele

frequency-based method. This is apparent in an R1-R0 plot of the HGDP data

constructed excluding pairs that are closer than 0.01 to the kinship coefficient

thresholds that the frequency based method used when categorizing relationships

(Figure S4).

Application to sequencing data

To assess how well the new method works on sequencing data, we applied it to

sequencing data from five low-depth (~4x) human genomes from the 1000 Genomes

project. Among the five selected samples there is a parent-offspring pair, a pair of full

siblings, a pair of half siblings, an unspecified 2^{nd} degree relationship (e.g. avuncular),

and the rest are unrelated. We estimated the R0, R1 and KING-robust kinship for each

pair in several ways. First, by using an IBS-based approach that estimates the

proportion all pairwise combinations of the 10 possible genotypes (Figure 4, "IBS").

Second, we estimated the two-dimensional site frequency spectrum (2D-SFS) of each

pair with a bi-allelic model and calculated R0, KING-robust kinship, and R1 based on

this spectrum (Figure 4, "realSFS"). Both these approaches base their estimates on

genotype likelihoods calculated from the sequencing read data, instead of on called

genotypes, and take the uncertainty of the underlying genotypes that is inherent to low-

depth sequencing data into account. The key difference between them is that the SFS-

based approach requires specification of an allele at each site, potentially excluding

studies with low-depth data from non-model species, whereas the IBS approach has no

such requirement (for methodological details see Methods and materials). The methods

also differ in how they deal with sites with more than two unique alleles. We include

both methods here to compare their performance on real data.

Additionally, we calculated the three statistics from the high-quality phased genotypes

for the same five individuals available from the 1000 Genomes Project Phase 3 (Figure

4, "1000G sites") to see how well the two genotype-likelihood-based approaches applied

to low-depth sequencing data perform compared to direct calculations from high-quality

genotype data for the same samples. In this comparison, results obtained by using the

genotype likelihood-based approaches applied to low-depth sequencing data are close to

those obtained from the high-quality genotypes for all of the pairs (Figure 4).

Chromosome block-jackknife estimates of uncertainty were generally small, and varied

by relationship type, with the pair of full siblings having the most uncertainty in R0, R1,

and KING-robust kinship. Notably, the IBS-based approach provided meaningful

results, even though it did not make use of any external allele information. With the IBS

method, unrelated individuals had a slight decrease in R0 and a slight increase in R1

and kinship, relative to the SFS method, making them appear slightly more related than

expected for unrelated individuals from a homogenous population. However, despite

this bias, it is still possible to correctly categorize of all the pairs to their relationship

category within the set of close familial relationships [PO, FS, HS, or UR], suggesting

that this method can be used when not enough information is available for the realSFS

method.

Finally, we also made R1-R0 plot and the R1-KING-robust kinship plots based on

genotypes that we obtained through a standard genotype calling procedure from the raw

read data. We did this to investigate if the genotype likelihood-based approaches are

necessary or one could just as well use genotypes called from the ~4x data. As expected,

genotype calling had a large negative effect on the outcome; in the resulting R1-R0 and

R1-KING-robust kinship plots the half siblings appear where first cousins are expected

and both the parent-offspring and full sibling pairs appear in the expected range of half

siblings (Figure 4, "called genotypes"). These results demonstrate the potential pitfalls

of basing relationship inferences, including R1-R0 and R1-KING-robust kinship plots,

on genotypes called from low-depth data. Notably, this is also the case for the methods

presented in (Rosenberg 2006) and (Manichaikul *et al.* 2010) (Figure 5A). Hence, for

~4x sequencing data, calling genotypes without external information, such as an

imputation reference panel is not a good alternative to genotype likelihood-based

approaches, which means that software packages like KING cannot be used for this type

of data (as it requires input genotypes).

Assessing the effect of SNP ascertainment scheme

To assess the effect of SNP ascertainment on the new method, and thus the effect of the

data-generation platform or SNP filtering, we applied it to three different subsets of data

from the five 1000 Genomes individuals. The results for each of the three ascertainment

schemes (whole genomes, 1000 Genomes release sites, Illumina 650Y SNP array sites)

are similar (left four panels, Figure 4), showing SNP ascertainment does not have a large

effect.

For comparison, we performed the same assessment of the methods presented in

Rosenberg (Rosenberg 2006) and (Manichaikul *et al.* 2010) by constructing scatterplots

of the same type as those shown in their papers (Figure 5). This shows that both these

other methods are much more affected by ascertainment than the method proposed

here. In particular, the Rosenberg method is affected on both its x-axis (IBS0) and its y-

axis (IBS2), which means that the expected region of the plot for each relationship will

be different for different data sets (Figure 5A). The method presented in Manichaikul *et*

al. 2010 is affected by the SNP ascertainment mainly on its x-axis (IBS0, Figure 5B).

Therefore, the ascertainment mainly affects the ability to distinguish between parent-

offspring and full siblings, since the y-axis, which is only slightly affected by

ascertainment, is the kinship coefficient, which can be used to distinguish between most

close relationships except for parent-offspring and full siblings. The x-axis, IBS0, is

included in part to help make the distinction between PO and FS (Manichaikul *et al.*

2010), but this ability is clearly affected by SNP ascertainment (Figure 5B).

455 Discussion

We have presented a simple new method for inferring if and how two individuals are

related based solely on genetic data from the two target individuals. This new method

has several advantages compared to most existing relatedness inference methods, which

are almost all dependent on additional information. First, it does not require any

information about allele frequencies. Second, it does not require information about the

relative genomic positions of the sites included in the analysis. Third, although the

method can straightforwardly be applied to genotype data (where it boils down to

simple counting, addition, and division), we demonstrate two ways in which it can also

be applied directly to sequencing data via genotype likelihoods. And importantly, we

show that by using these, the method provides useful results when applied to 4x

sequencing data. In contrast, we also show that ours and similar methods like KING fail

to provide useful results if applied to genotypes called from the same ~4x data instead.

This implies that – unlike previous methods - this new method can be used even if all

you have is low-depth sequencing data from a few individuals from a species without a

reference genome. This is not likely to be the typical situation in studies of modern

humans, however it is much more likely to occur in studies of with non-model

organisms or ancient samples. Fourth and finally, we have shown that the new method

is robust to the SNP ascertainment present in common SNP array data sets.

Comparison to similar methods

The new method is based on plotting two statistics, R0 and KING-robust kinship against a third, new statistic R1. The R0 statistic is derived from the test statistic proposed by Lee (2003) to test for relatedness; the only differences are that the numerator and denominator are flipped and that E, the proportion of sites where both individuals are heterozygous, is included in the denominator in the statistic defined by Lee but absent in R0, making R0 similar to the PPC statistic in PLINK. The KING-robust kinship estimator was proposed by Manichaikul *et al.* (2010), here we extend it to estimation directly from sequencing data. Notably, our results suggest that this extension is important for low-depth sequencing data, because estimates based on genotypes called from low-depth sequencing data are very poor (rightmost panels of Figure 4 and 5). And importantly this extension can also be used for similar statistics and thus makes existing methods based on such statistics, like KING-robust kinship, more widely applicable.

However, this extension is not the only contribution of this study. Another key new contribution of this study is the fact that R0 and KING-robust kinship are combined with the new R1 statistic instead of with IBS0 (the proportion of sites where the two individuals share zero alleles IBS) as was previously proposed by Rosenberg (2006) and Manichaikul *et al.* (2010). As we have shown, the fraction of sites that are IBS0 or IBS2 is very sensitive SNP ascertainment, meaning that results are only comparable within each different ascertainment scheme. Across different ascertainment schemes, such as between SNP arrays and sequencing data, the method from Rosenberg (2006), where IBS0 is combined with IBS2, is difficult, if not impossible, to use for relatedness inference because the fraction of sites that are IBS2 and IBS0 varies so wildly. On the other hand, KING-robust kinship is still very useful, but it loses its ability to distinguish between parent-offspring and full siblings, as IBS0 was used for this. Due to this sensitivity to ascertainment samples cannot be analyzed in isolation, and must be placed

in the context of other samples with known relationships and the same ascertainment

scheme. This requirement makes it difficult to apply these previous methods to ancient

humans or other species with limited sample sizes.

In contrast, the ability to identify relatives based on expected values is maintained in the

new method, regardless of ascertainment scheme due to the use of $R1$, instead of IBS0,

which makes the new method robust to SNP ascertainment. Here the $R1$-$R0$ plot is

especially useful, because parent-offspring pairs, tend to have an $R0$ estimate extremely

close to 0, making them particularly easy to identify. The $R1$-KING-robust kinship, on

the other hand, has the appealing aspect that the kinship axis has a biological

interpretation, defined as the probability that two alleles sampled at random from two

individuals are identical by descent. Hence the two plots types, $R1$-$R0$ and $R1$-KING-

robust kinship, each have their advantages.

Additionally, it is worth noticing that $R1$ not only makes it easier to distinguish between

parent-offspring and full siblings it can provide additional information across all

relationship categories; both the new plot types, $R1$-$R0$ and $R1$-KING-robust kinship

provide better separation of several of the different relationship categories than KING-

robust kinship does alone (Figure S2).

Finally, it is worth noticing that the two plots types seem to work better than a range of

other plots constructed from similar ratio statistics that we explored (Figure S5).

Limitations and applications

While the new method provides significant advantages over previous methods in

situations with limited data, it does have some limitations. First, like most other

relatedness inference methods, like PLINK, the proposed method assumes that the

individuals are not inbred and that they originate from the same homogenous,

population. And like many other relationship inference methods, it is not necessarily

robust to violations of these assumptions. Previous studies have shown the effect of

admixture on relatedness inference is complex and can potentially lead to bias in either

direction depending on the circumstances, and this is true even for KING, which was

developed to be robust to admixture (Thornton *et al.* 2012; Moltke and Albrechtsen

2014; Conomos *et al.* 2016; Ramstetter *et al.* 2017). Specific methods have been

developed to correct for admixture when the allele frequencies in the admixing

populations are known (e.g. Thornton *et al.* 2012; Moltke and Albrechtsen 2014), or

enough samples are available (Conomos *et al.* 2016). But since these methods work by

exploiting knowledge about allele frequencies or access to many samples for their

correction; the pairwise R0, R1 and KING-robust kinship statistics cannot be corrected

in a similar manner. However, we note that Lee (2003) showed that the statistic he

proposed for testing for relatedness can also be used to detect if two unrelated samples

are not from the same homogeneous population. If this is the case, the Lee's statistic will

be significantly smaller than 2/3; and equivalently R0 will be significantly above 0.5,

which may be useful when interpreting R0, R1, KING-kinship plots in the presence of

admixture or population structure more generally. Regarding inbreeding, one potential

way to assess if one of the individuals is inbred is to compare heterozygosities across

individuals; non-inbred and non-admixed individuals from the same population should

have similar heterozygosity, so marked heterozygosity differences can be a warning

signal.

A second limitation is that the categorization of different relationship pairs is based on

visual inspection of plots and that in some cases such classification will be difficult

because the actual ranges of the different relationship in the plots are close to each other

and even overlap. However, this is to a large extent due to biological variation caused by

the randomness of recombination (Hill and Weir 2011). This limitation in classification

accuracy is therefore shared with the allele frequency-based inference methods, like the

commonly used PLINK, as illustrated by Figure S5.

Despite these limitations, we believe that the results presented here suggest that the new

method constitutes a helpful new tool for relatedness inference that is especially

advantageous compared to other methods in studies with limited data. Identifying

related individuals is a crucial step in nearly any genetic analysis and removing the

requirements to specify allele frequencies and to have access to accurate genotypes has

the potential allow the identification of relatives even in small studies of non-model

species or ancient samples. These types of studies do not currently have many good

options to address relatedness among their samples. An additional appeal of the IBS-

based approach introduced here, is that because it considers all possible pairwise

genotype categories, it can potentially help identify specific error types that are expected

to occur in DNA from ancient samples, by allowing researchers to specifically identify

transitions. Also, due to its non-reliance on allele frequencies the method is also

potentially useful when relatedness is pervasive among many samples, complicating

allele frequency estimation in the broader population (Mcpeek *et al.* 2004). Hence, the

method will hopefully be useful to many.

Data Accessibility

569 The IBS method is available at:

570 http://www.popgen.dk/angsd/index.php/Genotype_Distribution

571 The data sets used are publicly available.

References

Albrechtsen, A., T. Sand Korneliussen, I. Moltke, T. van Overseem Hansen, F. C. Nielsen *et al.*, 2009 Relatedness mapping and tracts of relatedness for genome-wide data in the presence of linkage disequilibrium. Genet Epidemiol 33**:** 266-274.

Alexander, D. H., J. Novembre and K. Lange, 2009 Fast model-based estimation of ancestry in unrelated individuals. Genome Res 19**:** 1655-1664.

Baca, M., K. Doan, M. Sobczyk, A. Stankovic and P. Weglenski, 2012 Ancient DNA reveals kinship burial patterns of a pre-Columbian Andean community. BMC Genet 13**:** 30.

Balding, D. J., 2006 A tutorial on statistical methods for population association studies. Nat Rev Genet 7**:** 781-791.

Busing, F. M. T. A., E. Meijer and R. Van Der Leeden, 1999 Delete-m jackknife for unequal m. Statistics and Computing 9**:** 3-8.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al.*, 2015 Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4**:** 7.

Conomos, M. P., A. P. Reiner, B. S. Weir and T. A. Thornton, 2016 Model-free Estimation of Recent Genetic Relatedness. Am J Hum Genet 98**:** 127-148.

Derrien, T., J. Estelle, S. Marco Sola, D. G. Knowles, E. Raineri *et al.*, 2012 Fast computation and applications of genome mappability. PLoS One 7**:** e30377.

Dou, J., B. Sun, X. Sim, J. D. Hughes, D. F. Reilly *et al.*, 2017 Estimation of kinship coefficient in structured and admixed populations using sparse sequencing data. PLoS Genet 13**:** e1007021.

Hill, W. G., and B. S. Weir, 2011 Variation in actual relationship as a consequence of Mendelian sampling and linkage. Genet Res (Camb) 93**:** 47-64.

Kardos, M., G. Luikart and F. W. Allendorf, 2015 Measuring individual inbreeding in the age of genomics: marker-based measures are better than pedigrees. Heredity (Edinb) 115**:** 63-72.

Korneliussen, T. S., A. Albrechtsen and R. Nielsen, 2014 ANGSD: Analysis of Next Generation Sequencing Data. BMC Bioinformatics 15**:** 356.

Korneliussen, T. S., and I. Moltke, 2015 NgsRelate: a software tool for estimating pairwise relatedness from next-generation sequencing data. Bioinformatics 31**:** 4009-4011.

Lee, W. C., 2003 Testing the genetic relation between two individuals using a panel of frequency-unknown single nucleotide polymorphisms. Annals of Human Genetics 67**:** 618-619.

Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data.
Bioinformatics 27**:** 2987-2993.

Manichaikul, A., J. C. Mychaleckyj, S. S. Rich, K. Daly, M. Sale *et al.*, 2010 Robust
relationship inference in genome-wide association studies. Bioinformatics 26**:**
2867-2873.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res 20**:** 1297-1303.

McPeek, M. S., X. Wu and C. Ober, 2004 Best linear unbiased allele-frequency
estimation in complex pedigrees. Biometrics 60**:** 359-367.

Moltke, I., and A. Albrechtsen, 2014 RelateAdmix: a software tool for estimating
relatedness between admixed individuals. Bioinformatics 30**:** 1027-1028.

Monroy Kuhn, J. M., M. Jakobsson and T. Günther, 2017 Estimating genetic kin
relationships in prehistoric populations. bioRxiv.

Nielsen, R., T. Korneliussen, A. Albrechtsen, Y. Li and J. Wang, 2012 SNP calling,
genotype calling, and sample allele frequency estimation from New-Generation
Sequencing data. PLoS One 7**:** e37558.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira *et al.*, 2007 PLINK: a
tool set for whole-genome association and population-based linkage analyses. Am
J Hum Genet 81**:** 559-575.

Ramstetter, M. D., T. D. Dyer, D. M. Lehman, J. E. Curran, R. Duggirala *et al.*, 2017
Benchmarking Relatedness Inference Methods with Genome-Wide Data from
Thousands of Relatives. Genetics 207**:** 75-82.

Rosenberg, N. A., 2006 Standardized subsets of the HGDP-CEPH Human Genome
Diversity Cell Line Panel, accounting for atypical and duplicated samples and
pairs of close relatives. Ann Hum Genet 70**:** 841-847.

Sikora, M., A. Seguin-Orlando, V. C. Sousa, A. Albrechtsen, T. Korneliussen *et al.*, 2017
Ancient genomes show social and reproductive behavior of early Upper
Paleolithic foragers. Science 358**:** 659-662.

Stevens, E. L., G. Heckenberg, E. D. Roberson, J. D. Baugher, T. J. Downey *et al.*, 2011
Inference of relationships in population data using identity-by-descent and
identity-by-state. PLoS Genet 7**:** e1002287.

The Genomes Project, C., 2015 A global reference for human genetic variation. Nature
526**:** 68-74.

Thompson, E. A., 1975 The estimation of pairwise relationships. Ann Hum Genet 39**:**
173-188.

Thompson, E. A., 2000 Statistical inference from genetic data on pedigrees, pp. i-169 in
NSF-CBMS regional conference series in probability and statistics. JSTOR.

Thornton, T., H. Tang, T. J. Hoffmann, H. M. Ochs-Balcom, B. J. Caan *et al.*, 2012
Estimating kinship in admixed populations. Am J Hum Genet 91**:** 122-138.

Wiuf, C., 2006 Consistency of estimators of population scaled parameters using
composite likelihood. Journal of Mathematical Biology 53**:** 821-841.

Author Contributions

657   AA, IM, and RKW designed the research. RKW performed research, with input from IM

658   and AA. IM and RKM wrote the paper, with help from AA.

Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

Authors:

Ryan K. Waples
Anders Albrechtsen
Ida Moltke

Figure 1. Overview of the R0, R1, and KING-robust kinship statistics
A Definition of the pairwise genotype categories A - I. Here g_1 and g_2 denotes the genotype for each of the two diploid individuals, 1 and 2, respectively. These genotypes are defined as the number of copies of a certain allele carried by 1 and 2, respectively. We assume diallelic variants such that g_1 and g_2 each has 3 possible values: 0, 1, and 2. For a pair of individuals, there are nine possible genotype combinations. We organize them into a 3x3 matrix and denote them with the letters from A to I. The values A – I can equivalently be either counts or proportions. **B** Definition of the R0 statistic based on the notation illustrated in A. **C** Definition of the R1 statistic based on the notation illustrated in A. **D** Definition of the KING-robust kinship estimator (Manichaikul *et al.* 2010), based on the notation illustrated in A.

Figure 2. Joint expectation ranges for R1-R0 and for R1-KING robust kinship.

A Joint expectation ranges for R0 - R1 for five different relationship categories: Parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1), and unrelated (UR). The expected value for PO is shown as a purple X. The expected value range for monozygotic twins (MZ) is not depicted because it is (R1=0, R0=∞). **B** Joint expectation ranges for KING-robust kinship and R1 jointly for six different relationship categories: Parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1), and unrelated (UR). The expected pair of values for PO is shown as a purple X. The joint expectation ranges shown here are based on expected IBD sharing (i.e. values of k_0, k_1 and k_2) for each relationship category. True IBD sharing for related individuals has some variation due to biological randomness in recombination and is not likely to be exactly equal to their expected values. This can cause derived statistics to fall outside their expected ranges as well. For detail about how the expected ranges were derived see supplementary text.

Figure 3. R1-R0 and R1-KING-robust kinship scatterplots for seven HGDP populations. Each point represents a pair of individuals and is colored according to the relationship category inferred using an allele frequency based approach. Shaded areas and lines show the expected ranges for specific relationship categories, as in Figure 2. Note that in addition to the relationship categories for Figure 2 there is an additional category here representing distantly related pairs (DR). **A** R1-R0 plot for all pairs of individuals within each population. **B** R1-KING-robust kinship plot for all pairs of individuals within each population. **C** Scatterplot of the two relatedness coefficients k_1 and k_2 for all pairs of individuals within each population estimated using the allele frequency-based approach implemented in PLINK. The coloring illustrates how these estimates were used to categorize the pairwise relationships, and thus determine the coloring in panel A, B, D and E. **D** Zoom into the outlined area in A. **E** Zoom into the outlined area in B.

Figure 4. Relatedness plots for all pairs among 5 LWK individuals from the 1000G Project.
A R1-R0 plots for pairs of the same five LWK individuals for five different analysis methods: 1) IBS = reference-free estimation from (~4x) 1000G bam files, 2) realSFS = site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G genotypes from the 1000G project, 4) HGDP sites=1000G genotypes at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes= de novo genotype calling from 1000G bam files. Shaded areas show the joint expected ranges across the two statistics. Points are colored by their true relationship status, as in Figure 3. Thin grey lines show the confidence intervals (+/- 2 SE) with a chromosome jackknife. **B** R1-KING-robust kinship plots for the same data sets, confidence intervals and expected ranges are constructed in the same way. The dashed gray line shows kinship of zero.

Figure 5. Plots of the two other frequency-free methods for different subsets of SNPs. **A** Results from applying the plotting approach from Rosenberg (2006) to pairs of the same five LWK individuals for five different analysis methods: 1) IBS = reference-free estimation from (~4x) 1000G bam files, 2) realSFS = site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G genotypes from the 1000G project, 4) HGDP sites=1000G genotypes at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes= de novo genotype calling from (~4x) 1000G bam files. Pairs are colored by their true relationship status, as in Figure 3. Fraction IBS0/IBS2 are the overall fraction of sites that are IBS0/IBS2, respectively. Thin grey lines centered on each point show confidence intervals (+/- 2 SE) based on a chromosome jackknife. **B** Results from applying the KING-robust based approach to the same pairs of LWK individuals

using the same five different analysis methods as in A. The horizontal black lines show the kinship thresholds used to distinguish unrelated (UR), first-cousins (C1) half siblings (HS), full siblings (FS) and (PO) following (Manichaikul *et al.* 2010) from bottom to top, respectively. Like in A, thin grey lines centered on each point show confidence intervals (+/- 2 SE) based on a chromosome jackknife.