Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

Authors:

Ryan K Waples*, Anders Albrechtsen*[1], Ida Moltke*[1]

*Department of Biology, Section for Computational and RNA Biology, University of Copenhagen, 2200 Copenhagen N, Denmark,

[1]Corresponding authors: Department of Biology, University of Copenhagen, Ole Maaloes Vej 5, 2200 Copenhagen N, Denmark. E-mail: albrecht@binf.ku.dk; and ida@binf.ku.dk

Abstract

Knowledge of how individuals are related is important in many areas of research and numerous methods for inferring pairwise relatedness from genetic data have been developed. However, the majority of these methods were not developed for situations where data is limited. Specifically, most methods rely on the availability of population allele frequencies, the relative genomic position of variants, and accurate genotype data. But in studies of non-model organisms or ancient human samples, such data is not always available. Motivated by this, we present a new method for pairwise relatedness inference, which requires neither allele frequency information nor information on genomic position. Furthermore, it can be applied not only to accurate genotype data but also to low-depth sequencing data from which genotypes cannot be accurately called. We evaluate it using data from a range of human populations and show that it can be used to infer close familial relationships with a similar accuracy as a widely used method that relies on population allele frequencies. Additionally, we show that our method is robust to SNP ascertainment and applicable to low-depth sequencing data generated using different strategies, including resequencing and RADseq, which is important for application to a diverse range of populations and species.

Introduction

The ability to infer the familial relationship between a pair of individuals from genetic data plays a key role in several research fields. In conservation biology, it is used to design breeding programs that minimize inbreeding (Kardos *et al.* 2015), in archeology it is helpful to understand burial patterns and other cultural traditions (Baca *et al.* 2012; Sikora *et al.* 2017), and in population and disease genetics it is often used to exclude relatives, because many analysis methods within those fields assume all analyzed individuals are unrelated and violations of this assumption can lead to wrong conclusions (Balding 2006).

Numerous pairwise relatedness inference methods have been developed, e.g., Thompson (1975), Lee (2003), Purcell *et al.* (2007), Albrechtsen *et al.* (2009), Manichaikul *et al.* (2010), Stevens *et al.* (2011), Korneliussen and Moltke (2015), Conomos *et al.* (2016), Dou *et al.* (2017), and many are available in popular software packages, like PLINK (Purcell *et al.* 2007), and KING (Manichaikul *et al.* 2010). Most of these methods estimate either the three relatedness coefficients k_0, k_1 and k_2, or the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$ for each pair of diploid individuals, where k_0, k_1 and k_2 are the proportions of the genome where a pair of individuals share 0, 1, or 2 alleles identical by descent (IBD) (Thompson 2000). By definition, alleles are IBD when they are identical due to *recent* common ancestry, but because alleles can also be identical due to older common ancestry or recurrent mutations, IBD status cannot be directly observed. Therefore, the pairwise relatedness coefficients and the kinship coefficient have to be estimated, which can be done from patterns of observed genetic identity (identity by state; in short IBS). Once estimated, these relatedness statistics, k_0, k_1, k_2, and the kinship coefficient can be used to infer familial relationships by comparison to the expectation of the statistics for different familial relationships (Hill and Weir 2011).

Although inference of relatedness is of wide interest, most existing methods are not

immediately applicable in studies with limited data or genetic resources. First, most

existing methods require the allele frequencies of the source population. For most

studies in modern humans this is not a problem. However, in studies of ancient humans

or other species, accurate estimates of population allele frequencies are often not

obtainable, because only a low number of samples are available. Second, several existing

methods consider consecutive loci jointly and use sliding windows or hidden Markov

models to leverage the non-independence of allele sharing along the genome between

relatives (Albrechtsen *et al.* 2009; Gusev *et al.* 2009; Stevens *et al.* 2011; Kuhn *et al.*

2018). These methods are powerful but require information about the genomic position

of variable sites, and for non-model organisms high-quality reference genome

assemblies are often not available. Third, other methods avoid allele frequencies, but

rely on access to many samples to provide a necessary context for relationship

classification (e.g., Abecasis *et al.* 2001). Finally, nearly all existing methods --both

frequency-based and others-- require genotype data. However, in sequencing studies,

samples are often only sequenced to low depth due to cost and technical issues. This

makes it infeasible to call genotypes accurately (Nielsen *et al.* 2012), precluding the use

of these methods. There are a few methods that estimate relatedness from low-depth

sequencing data by utilizing genotype likelihoods (e.g., Korneliussen and Moltke 2015),

or by using imputed genotype dosages (Dou *et al.* 2017). However, these methods

function by leveraging access to many samples to estimate allele frequencies or perform

accurate genotype imputation and are therefore not designed to apply to datasets with a

low number of samples.

Hence, most existing methods to infer close familial relationships are not immediately

applicable in studies where data is limited, including many studies of non-model

organisms and ancient humans. One of the few exceptions is a simple but elegant test

for pairwise relatedness proposed in Lee (2003). This test relies entirely on the relative

frequency of different genotype combinations within a pair of individuals and thus only

requires genotype data from the two target individuals. While useful, this test does not

provide any means to distinguish between different types of close familial relationships;

it only provides a statistical test for a pair of individuals of the null hypothesis of them

being unrelated.

There are only a few methods that can be used to distinguish between different types of

familial relationships for a pair of individuals when neither allele frequencies nor

information about the relative genomic position of sites is obtainable. One such method

consists of plotting the proportion of the genomic sites in which the two individuals

share both alleles IBS (which we will denote IBS2) versus the proportion of the genomic

sites in which they share zero alleles IBS (which we will denote IBS0). This method was

used in Rosenberg (2006), where it was applied to the Human Genome Diversity Project

(HGDP) dataset. In the resulting scatter plot of the HGDP data (figure 1 of Rosenberg

(2006)), pairs of individuals with the same relationship category form distinct clusters

so that it is possible to locate parent-offspring pairs, full sibling pairs, and to a lesser

extent more distant relationships such as half-siblings/avuncular/grandparent-

grandchild and first cousins.

Another such method is based in part on the KING-robust kinship estimator

(Manichaikul *et al.* 2010). The KING-robust kinship estimator was developed to be

robust to population structure, but in practice it has been shown to provide biased

kinship estimates when applied to pairs of samples whose four chromosomes are not all

from the same population (Thornton *et al.* 2012; Conomos *et al.* 2016). However, the

KING-robust kinship estimator is directly applicable to samples from the same

homogenous population even when allele frequencies are unknown. The reason for this

is that, like the test suggested in Lee (2003), it relies only on the genotype combinations

within the two target individuals and does not require knowledge about of allele

frequencies. Importantly, Manichaikul *et al.* (2010) show it is possible to infer if a pair

of individuals are parent-offspring, full sibling, half siblings/avuncular/grandparent-

grandchild, first cousins, or unrelated, by jointly considering KING-robust kinship and

the fraction of sites IBS0 using SNP array data without allele frequencies. For example,

see figure 3A in Manichaikul *et al.* (2010); a scatterplot of KING-robust kinship vs the

fraction of sites IBS0.

However, both the methods described above have two important limitations. First, like

most other methods to estimate relatedness, they were developed for genotype data

only. For example, the KING software (Manichaikul *et al.* 2010) implementing the

KING-robust kinship estimator requires genotype data as input, which can be

problematic for studies where only moderate or low-depth sequencing data is available

and calling genotypes is consequently difficult. Second, both methods rely on estimates

of the fraction of sites IBS0, which can be problematic because this fraction, as well as

the fraction of sites IBS1 and IBS2, is highly sensitive to SNP ascertainment. This means

that the results of the methods are platform-dependent and are likely to differ between

different SNP arrays and especially between SNP array and sequencing datasets. And in

turn, this among other things, means that it can be difficult to distinguish between full

siblings and parent-offspring pairs using these methods.

Motivated by the outlined limitations to the existing methods, we present a method for

relationship inference that -unlike most existing methods- relies neither on allele

frequencies nor on information about the relative position of the variant sites, and

which – unlike other frequency-free methods – 1) is applicable even to sequencing data

of so low depth that accurate genotypes cannot be called from it and 2) is robust to SNP

ascertainment bias.

The new method is inspired by previous methods; it uses the KING-robust kinship

estimator and a statistic R0, which is similar to the test statistic from the test for

relatedness suggested by Lee (2003). However, the method is new in two important

ways. First, besides relying on the two statistics, R0 and KING-robust kinship, it also

relies on a third new statistic, R1. More specifically, the method consists of using two

combinations of these three statistics, R1-R0 and R1-KING-robust kinship, to infer

relationships, and it is this combination of statistics that makes the method robust to

ascertainment bias. Secondly, while the new method can straightforwardly be applied to

genotype data like other similar methods, we also present two computational

approaches to estimate the three statistics directly from sequencing data that takes the

uncertainty of genotypes into account, allowing application to low-depth sequencing

data.

In the following we first fully describe the three statistics, R0, R1, and KING-robust

kinship, how they can be estimated and other methodological details. Next, using

simulated and publicly available SNP array data, we show that the new method provides

similar accuracy and precision to the commonly used frequency-based method

implemented in PLINK, when such data is available. Then using sequence data from the

1000 Genomes Project (The Genomes Project 2015), we show that the three statistics

can be estimated directly from sequencing data of low depth (~4x), here defined as

depth insufficient for accurate genotype calling. Moreover, we show that the estimates

obtained in this way are useful for inference of close familial relationships and that this

is not the case for estimates obtained from genotypes called from the same data. Using

different subsets of the same data, we also show that this new method, unlike previous

similar methods, is robust to SNP ascertainment. Finally, we show that the method also

provides useful results when applied to sequencing data down-sampled to approximate

data generated using reduced-representation approaches, e.g. restriction-site associated

DNA sequencing (RADseq) and discuss some potential applications and limitations of
the new method.

Methods and materials

The R0, R1, and KING-kinship statistics

The method for relationship inference we propose consists of estimating three statistics
called R0, R1 and KING-robust kinship from genetic data and interpreting plots of R1 vs
R0 and R1 vs KING-robust kinship.

We define the three statistics, R0, R1, and KING-robust kinship in terms of the genome-
wide IBS sharing pattern of two individuals. At any given diallelic sites, a pair of
individuals will carry one of nine possible genotype combinations; the nine possible
combinations of the two individuals carrying 0, 1, or 2 copies of a specific allele, e.g., the
ancestral allele. We can therefore fully characterize the genome-wide IBS-sharing
pattern of a pair of individuals by nine counts or proportions denoted: A, B, C, D, E, F,
G, H, and I (Figure 1A), similar to a two-dimensional site-frequency spectrum (SFS)
across the two individuals. The R0 and R1 statistics are defined as simple functions of a
subset of these nine counts as shown in Figure 1B and C, and the KING-robust kinship
statistic, originally defined by Manichaikul *et al.* (2010), can also be re-formulated as a
function of these 9 values (Figure 1D).

The new method is motivated by several observations. First, the expected values of A-I
vary depending on the familial relationship between the pair of individuals of interest.
Consequently, so do functions of A-I, including R0, R1, and KING-robust kinship.
Notably, there is no overlap between the joint expectation ranges of [R1, R0] and [R1,
KING-robust kinship] for the four close relationship categories: full siblings (FS), half

siblings/avuncular/grandparent-grandchild (HS), first cousins (C1) and unrelated (UR)

and the range of expected values for parent-offspring (PO) only overlaps with those of

FS in a single point (Figure 2, for derivations see supplementary text). Crucially, this is

true regardless of the underlying allele frequency spectrum and holds for any pair of

non-inbred individuals from the same homogenous population, making [R1, R0] and

[R1, KING-robust kinship] potentially useful for distinguishing between these

relationships. Second, while A-I, and thus R0, R1, and the KING-robust kinship

estimator can be calculated from genotype data, they can also be estimated directly from

next-generation sequencing (NGS) data based on the expected number of sites with each

genotype combination (see below for details). This makes the method appropriate even

when the sequencing depth is too low for accurate genotype calling (see below for

methodological details). Third, regardless of the type of data that is available, R0, R1

and KING-robust kinship can be estimated without the need for population allele

frequencies and information about the relative position of the genomics sites analyzed.

Finally, we expect the three statistics to be robust to SNP ascertainment because they

are ratios computed from sites that are variable within the two samples and should thus

be unaffected by the number of non-variable sites and because the (unknown)

underlying frequency spectrum should only have a limited effect on these ratios.

Estimation from sequencing data

The counts of the nine genotype combinations, A-I, and thus R0, R1, and KING-robust

kinship for a pair of individual, can be estimated directly from NGS data via the use of

genotype likelihoods calculated from the aligned sequencing reads. Genotype

likelihoods provide a means to account for the genotype uncertainly inherent to low-

depth NGS data. We used two distinct, but related, approaches to estimate these

statistics from sequencing data that both build on this idea.

The first approach, which we denote the IBS-based approach, considers all ten possible

genotypes at a diallelic site for each of the two individuals of interest and consists of a

maximum-likelihood (ML) estimation of the counts of each of the 100 (10 × 10) possible

genotype pairs (for details see supplementary text). To perform the ML estimation, we

used an EM algorithm, which we have added to the `ANGSD` software package under the

name `IBS`. After obtaining the estimate of the counts of all the 100 possible genotype

pairs, we convert them into estimates of A-I, by summing over the counts that

correspond to each combination. For example, the genotype pairs: AA/AA, CC/CC,

GG/GG and TT/TT all contribute to cells A or I of Figure 1. Counts corresponding to

genotype pairs with more than two different alleles (e.g. AC/AG) were discarded. The

advantage of this IBS-based approach is that it does not require specification of a known

allele at each site and can thus be applied to nearly any sequencing data set, even with

low-depth, without any prior information on the alleles at each site.

The second approach, which we denote the SFS-based approach, consists of performing

ML estimation of the two-dimensional site-frequency spectrum (SFS) (Nielsen *et al*.

2012). To find the ML estimate of the SFS, we used an expectation-maximization (EM)

method implemented in the `ANGSD` software package under the name `realSFS`

(Korneliussen *et al*. 2014). The SFS-based approach requires one allele to be specified

for each site e.g., the ancestral, consensus, or reference allele. The model underlying this

approach assumes that genotypes for each site have the possibility of containing this

specified allele and up to one other unspecified allele. To specify the alleles that exist at

each site, we used the consensus sequences from the highest depth individual

(NA19042) and restricted our analysis to sites where the depth in this individual was at

least three.

The computational burden of analyzing genome-wide data sets can be significant. For

both the genotype likelihood-based approaches described above, the main limitation is

RAM, as data likelihoods for each site need to be loaded into memory for optimization.

To overcome this limitation, we analyzed each chromosome separately and then

summed over the values for each chromosome to produce a genome-wide estimate. To

calculate the likelihood of sequencing data for each possible genotype, we used the

original GATK genotype likelihood model (Mckenna *et al.* 2010) with independent errors,

as implemented in ANGSD. We also tried to use the samtools genotype likelihood model

(Li 2011) with a more complicated error structure, but found it produced worse results

(data not shown). Both IBS and realSFS produce the expected values for A-I, which we

subsequently use to calculate R0, R1 and KING-robust kinship. For example command

lines for each analysis, see supplemental text.

Confidence intervals

All of the above estimation methods treat each site as independent. This assumption

should not affect our expectation of each statistic (Wiuf 2006), but statistical non-

independence (here due to linkage disequilibrium (LD) and IBD) does affect standard

estimates of uncertainty. To quantify the uncertainty, we therefore estimated confidence

intervals for all statistics using a block-jackknife procedure. Confidence intervals were

estimated by leaving each chromosome out (chromosome jackknife), which takes both

the IBD and the LD correlation into account. The weighted block jackknife variance

estimator of Busing *et al.* (1999) was used to estimate the variance from the distribution

of estimates for each statistic. The square root of this variance was interpreted as the

standard error in our estimate.

Application to simulated data

258 To evaluate the new method and to investigate the effects demography, we simulated

259 genotype data under three different demographic histories: 1) a neutral demography

260 with constant effective population size (N_e), 2) a shrinking demography with a 10x

261 decrease in N_e over the last 100 generations, 3) an expanding demography with a 10x

262 increase in N_e over the last 100 generations. For each of the three scenarios we used the

263 coalescent simulator `msprime` (Kelleher *et al.* 2016) to simulate four haplotypes. These

264 haplotypes were then used to construct pairs of related individuals in five relationship

265 categories: parent offspring (PO), full siblings (FS), half-siblings/avuncular/

266 grandparent-grandchild (HS), first cousin (C1), and unrelated (UR). For unrelated pairs

267 of individuals, the genotype data were constructed by simply splitting the four

268 haplotypes into two pairs. For related pairs of individuals, we constructed the genotypes

269 at each variable site by first sampling whether the individuals shared 0, 1, or 2 alleles

270 IBD according to the expected values of [k0, k1, k2] for the relevant relationship

271 (Supplementary Table S1). Then we used alleles present on the four haplotypes

272 according to the sampled IBD status to construct the genotype at that site. For example,

273 if the individuals were selected to share two alleles IBD at a site, both individuals

274 received the alleles present on the first two haplotypes. The IBD sharing pattern was

275 sampled independently for each SNP and thus LD and biological variation in IBD was

276 not modelled. We concatenated the data from many independent simulations to achieve

277 enough data so the IBD sharing was approximately equal to the expected values.

278 Simulation code is available in the supplemental materials.

Application to real datasets

280 To assess the utility of the new method on more realistic data, we applied it to two

281 different publicly available datasets: SNP array data from seven HGDP populations

282 (Rosenberg 2006), and sequencing data from five related individuals from the Luhya in

283 Webuye, Kenya (LWK) population of the 1000 Genomes Project phase 3 (The Genomes

284 Project 2015).

HGDP SNP array

The HGDP SNP array dataset was accessed Jan. 13, 2017 and we followed the quality control steps described in Rosenberg (2006) to exclude mislabeled and duplicate samples. We selected seven populations from the HGDP based on the presence of several close familial relationships; five non-African populations: Surui, Pima, Karitiana, Maya, and Melanesian, and two African populations: Mbuti Pygmies and Biaka Pygmies.

To ensure a fair comparison to the allele frequency based inference method in PLINK, and by proxy to other commonly used methods, we constructed data sets where these methods have been shown to perform well. Specifically, we excluded individuals showing obvious signs of admixture (n=16) or inbreeding (n=2) from the selected HGDP populations. For details see Supplemental text 2.3.1. This left us with a total of 142 individuals from the seven populations: Surui N=20, Pima = 20, Karitiana N=21, Maya N=16, Melanesian N=19, Biaka Pygmies N=31 and Mbuti Pygmies N=15. For each of these seven populations we constructed a final set of genotypes by retaining genotypes from autosomal loci with genotyping rate >99%, minor allele frequency (MAF) >5%, HWE p-value > 10^{-4}.

The R0, R1, and KING-robust kinship statistics for each of the 2902 within-population pairs of individuals were then calculated from all sites where both individuals had non-missing genotypes.

1000 Genomes sequencing data

To get sequencing data from several different relationship categories, we selected five individuals from two families in the Phase 3 1000 Genomes (1000G) Luhya in Webuye, Kenya (LWK) population: NA19027, NA19042, NA19313, NA19331, NA19334. Across

the five individuals, there is one pair of half-sibs (NA19027 & NA19042), and a separate

trio of related individuals with a pair of full siblings (NA19331 & NA19334), one parent-

offspring relationship (NA19313 & NA19331), and another unspecified second-degree

relationship (NA19313 & NA19334), possibly avuncular (The Genomes Project 2015).

These stated relationships leave six unrelated pairs among the five individuals.

For each pair of the five LWK individuals we estimated the R0, R1, and KING-robust

kinship statistics in five different ways: 1&2) by applying the two different sequencing-

based approaches described above to the 1000G aligned sequence data files (~4x

coverage bam files), 3) by simple genotype counting based on the phased and high-

quality curated genotypes provided in the hg37 1000G VCF files, 4) genotype counting

based on the subset of sites in 3 that overlap with the Illumina 650Y sites for the HGDP

data (to investigate ascertainment, see below) and 5) by calling genotypes from the same

1000G bam files in a basic manner meant to mimic data from a species with a reference

genome but few other genetic resources and then simply counting from the called

genotypes. For set 5, we used `samtools mpileup` (v1.3.1) to summarize the reads

overlapping each position, and `bcftools call` (v1.3.1) to assign the most likely

genotype at each position. We used mostly default settings; non-default flags to

`samtools` specified skipping indel positions. Non-default flags to `bcftools` specified

using the consensus caller. For all sequence-based analyses (1,2 &5) we only considered

reads with a minimum phred-scaled quality score of 30 and bases with minimum phred-

scaled quality score of 20 and we restricted our analyses to genomic regions with a GEM

75mer mappability of 1 (Derrien *et al.* 2012). Notably, all the methods and filters used

here can be applied to any study, including studies with only small contigs, for example

made up of RAD loci, making the results relevant beyond resequencing studies utilizing

well assembled genomes.

Assessing the effect of SNP ascertainment

To evaluate the effect of SNP ascertainment using real data, we created a subset of the

curated genotype data from the five 1000 Genomes individuals. We selected the sites

that overlap with the Illumina 650Y array that was used for the HGDP and estimated

our three relatedness statistics. We compared the results from this subset of HGDP sites

to results for the full genotype dataset and also to the sequence-based analyses. For an

additional comparison, we also performed the same comparison for the methods

presented in Rosenberg (2006) and Manichaikul *et al*. (2010) by constructing

scatterplots by their methods.

We also investigated the effect of SNP ascertainment using the data simulated from a

constant demography (see "Application to simulated data" for details about the

simulations) and compared results for the full dataset to results obtained by including

only sites with a minor allele count >2 out of 40 chromosomes (MAF >5%) in the

analyses.

Assessing the effect of a limited number of sites

To assess the usefulness of this new method on data sets with fewer genomic sites

covered by sequencing reads we constructed reduced size data sets, in a way that

mimicked some aspects of reduced-representation sequencing approaches such as

RADseq. To produce each reduced data set, we selected a specific number of 200bp

windows randomly from the eligible genomic regions and restricted our analysis to sites

falling within them. We used 10k 50k, 100K, and 250k windows, representing ~4x

sequencing coverage on 2M, 10M, 20M, or 50M sites, respectively. All other aspects of

the analyses were the same, expect for that for these data sets, we applied the IBS- and

SFS- approaches to the complete data set, rather than splitting by chromosome as we

358   did for the full dataset. We suggest analyzing the complete data in single run if your

359   computational resources allow it, as we noticed some upward bias in the estimated

360   number of IBS0 sites when the smaller data sets were analyzed separately by

361   chromosome.

Comparison to other methods

To get a categorization of relationships for the HGDP dataset described above based on

a standard, commonly used allele frequency-based method, we first applied the allele

frequency based relatedness estimation program PLINK (v1.9) (Chang *et al.* 2015) to the

individuals from each population separately to estimate the genome-wide IBD fractions

k_0, k_1, k_2. Next, we applied the relationship criteria proposed in Table 1 of Manichaikul *et*

al. (2010) to the relatedness coefficient estimates: the estimated k values were combined

into an estimate of the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$, and a relationship degree was

assigned to each pair of individuals based on comparing the estimated kinship

coefficient to the criteria in the table. Parent-offspring and full siblings were

differentiated based on k_2 values. This provided us with a categorization into five

categories: PO, FS, HS, C1, and UR. To achieve additional resolution, we further divided

the last category (UR) into two: unknown/distantly related (UK-DR) and unrelated

(UN). We did this by simply extending the logic behind the criteria proposed above.

Specifically, we set the kinship threshold between UK-DR and UR to $1/2^{13/2}$, which

corresponds to including 4^{th} to 5^{th} degree relatives in the UK-DR category.

To assess the accuracy and precision of the new method for familial relationship

classification within the HGDP data we examined concordance with the PLINK-based

relationships described above. For this purpose, we assigned a relationship category to

each pair of individuals in two ways: 1) using the statistics R0 and R1 and 2) using a

combination of KING-robust kinship and R0. For the former, we characterized each

possible relationship by a single [R1, R0] point generated from data simulated under a demography with a constant population size over time, detailed in the "Application to simulated data" section and assigned each pair of individuals the relationship of the closest point using a Euclidean distance measure. For the latter, we first used the KING-robust kinship criteria from table 1 of Manichaikul *et al.* (2010) as above. Since this table has overlapping kinship ranges for the PO and FS categories, we used the R0 statistic to distinguish PO from FS relationships: Ignoring rare effects like germline mutations and genotyping errors the expected value for R0 for PO relatives is zero, while for FS the value is above 0; we used an ad hoc cut-off of 0.02.

To estimate the statistics for identifying related individuals proposed by Rosenberg (Rosenberg 2006) and KING (Manichaikul *et al.* 2010), we note that the KING-robust kinship estimator can (as previously described) be calculated directly from the same nine counts, A-I, and so can the fraction of sites IBS0 and IBS2:

KING-robust kinship = $(E - 2(C+G))/ (B+D+H+F+2E)$

Fraction IBS0 = $(C+G)/(A+B+C+D+E+F+G+H+I)$

Fraction IBS2 = $(A+E+I)/(A+B+C+D+E+F+G+H+I)$

We used these formulas in all our comparisons because this allowed us to estimate these statistics not only from genotype data but also directly from sequence data in the same manner as for R0 and R1. However, we note that this is our approach to estimating those statistics, and that existing tools like KING only allow users to estimate the statistics from genotype data.

Results

To assess the performance of the new method, we first applied it to simulated genotype data to ensure that it works on sufficient data and to assess how sensitive it is to the

underlying demographic history of the population the analyzed samples are from. Next,

we applied the new method to real data from different platforms to assess its

performance on more realistic data. Finally, we performed a couple of additional

analyses to access how robust the new method is to SNP ascertainment and to having

data from only a limited number of sites available. Below we describe the results of all

these analyses.

Application to simulated data

We first applied the new method to simulated genotype data from several different

relationship pairs from populations with three different demographic histories: 1)

constant N_e, 2) 10-fold increase in N_e over the past 100 generations, and 3) 10-fold

decrease in N_e over the past 100 generations. We see very similar results across all three

demographic scenarios and in all cases the R1-R0 and R1-KING-robust kinship values

obtained from the simulated data were within the theoretically derived ranges of

expectations (Figure 2). These results demonstrate that the method works if sufficient

high-quality genotype data is available. Furthermore, they demonstrate that the range of

examined population size histories has a limited effect, even though demographic

history affects the allele frequency spectrum. In turn, this suggests that the range of

expected values that are realistic for real data is markedly smaller than the theoretically

derived ranges also shown in the figure, which is useful for classification purposes.

Application to SNP array data

Next, we applied the method to SNP array data from the Human Genome Diversity

Project (HGDP) to see how the method works on real data, for which standard allele

frequency-based methods like PLINK are known to perform well. More specifically, we

applied it to genotype data from unadmixed and non-inbred samples from seven

populations originating from the HGDP. This resulted in the R1-R0 and R1-KING-

robust kinship plots shown in Figure 3A and B (for population-specific plots see

Supplementary Figures S1 and S2). The true relationships for the pairs of individuals are

not known; we instead colored each point in Figure 3A and B according to the

relationship category inferred based on results from the standard, commonly used allele

frequency-based method PLINK (Figure 3C, Supplementary Figure S3). Since there are

least 15 individuals in each of the seven selected HGDP populations, the allele frequency

estimates for these populations should be reasonably accurate even with some

relatedness. Hence, with the large amount of data available in this dataset, the allele

frequency-based method should provide correct inference of most, if not all, pairs closer

than first cousins, but may not be able to fully distinguish first cousins from more

distantly related pairs.

In Figures 3A and B, points from each relationship category clearly cluster together on

both the R1-R0 plot and the R1-KING-robust kinship plot. Moreover, these clusters are

located near both their theoretically derived ranges of expected values and the values

from simulated data. In a similar manner, the k_1-k_2 values for the same pairs of

individuals cluster close to the expected and simulated values of k_1 and k_2 (Figure 3C).

Almost all pairs identified as parent-offspring (PO) by the frequency-based method are

easy to identify as such in both the R1-R0 plot and the R1-KING-robust kinship plot,

which is not the case when only a single statistic is used (see also Supplemental Figures

S1 and S2). The same is true for full siblings (FS). Furthermore, points classified as half-

siblings/avuncular/grandparent-grandchild (HS) or first cousins (C1) by the frequency-

based method have a minimal overlap with each other and with less-related pairs

(Figure 3A and B). The few pairs of individuals that were difficult to classify are the

same pairs as those that are edge cases for the allele frequency-based method. This is

apparent in an R1-R0 plot of the HGDP data constructed excluding pairs that are closer

than 0.01 to the kinship coefficient thresholds that the frequency-based method used

when classifying relationships (Supplementary Figure S4).

To quantify precision and accuracy, we examined the concordance between

classifications based on the new method and the PLINK-based classification. We tried

two simple classification schemes: one based on R1-R0, which uses proximity to the

values we obtained from simulated data from a constant N_e demography, and one based

on KING-robust kinship (for details see Methods and Materials). The results supported

the visual assessment: both classification schemes are highly concordant with the

classifications obtained using the frequency-based method (Supplemental Figure S5).

Mean precision across all relationship categories was 0.90 for the R1-R0 method, vs

0.89 for KING-robust kinship. Mean recall across all relationship categories was 0.88

for R1-R0, vs 0.89 for KING-robust kinship. The relationship categories for which the

method has the lowest precision is the first cousins versus less related pairs, where the

allele frequency-based method is also known to have a hard time making classifications.

For PO, FS and UR alone the mean precision is as high as 0.99 for R1-R0 and 0.96 for

KING-robust-kinship, and the mean recall for these three categories is as high as 0.96

for R1-R0 and 0.99 for KING-robust kinship. Hence the new method provides

comparable performance to a frequency-based method when sufficient genotype data is

available, but without the need for allele frequency information.

Application to sequencing data

To assess how well the new method works on more limited real data, we applied it to

sequencing data from five low-depth (~4x) human genomes from the 1000 Genomes

project. Among the five selected samples there is a parent-offspring pair, a pair of full

siblings, a pair of half siblings, an unspecified 2^{nd} degree relationship (e.g. avuncular),

and the rest are unrelated. We estimated the R0, R1 and KING-robust kinship for each

pair in several ways. First, by using an IBS-based approach that estimates the

proportion all pairwise combinations of the 10 possible genotypes (Figure 4, "IBS").

Second, by using an SFS-based approach where we estimated the two-dimensional site

frequency spectrum (2D-SFS) of each pair with a bi-allelic model and calculated R0,

KING-robust kinship, and R1 based on this spectrum (Figure 4, "realSFS"). Both these

approaches base their estimates on genotype likelihoods calculated from the sequencing

read data, instead of on called genotypes, and take the uncertainty of the underlying

genotypes that is inherent to low-depth sequencing data into account. The key

difference between them is that the SFS-based approach requires specification of an

allele known to exist at each site, whereas the IBS-based approach has no such

requirement, making it more generally applicable. The approaches also differ in how

they deal with sites with more than two unique alleles, either excluding them (IBS-based

approach) or integrating over the two-allele possibilities (SFS-based approach), but

these sites are rare (mean fraction as estimated by IBS: 1.8E-6) so the impact of

discarding them is minimal.

With the values estimated using the SFS-based approach it is possible to visually classify

of all the pairs to their relationship category within the set of close familial relationships

[PO, FS, HS, or UR], or by using one of the classification methods introduced earlier).

Results for the IBS-based approach were similar, but unrelated individuals have a slight

decrease in R0 and a slight increase in R1 and KING-robust kinship, compared to the

SFS-based approach. This makes unrelated individuals appear slightly more related

than expected for unrelated individuals from a homogenous population. However,

despite this bias, it is still possible to correctly classify of all the pairs to their

relationship category, suggesting that the IBS-based approach can be used when not

enough information is available for the SFS-based approach. The chromosome block-

jackknife estimates of uncertainty for the genotype likelihood-based methods were

small, and varied by relationship type, with the pair of full siblings having the most

uncertainty in R0, R1, and KING-robust kinship.

We also calculated the three statistics from the high-quality phased genotypes for the

same five individuals available from the 1000 Genomes Project Phase 3 (Figure 4,

"1000G sites") to see how well the two genotype-likelihood-based approaches applied to

low-depth sequencing data perform compared to direct calculations from high-quality

genotype data for the same samples. In this comparison, results obtained by using the

genotype likelihood-based approaches applied to low-depth sequencing data are close to

those obtained from the high-quality genotypes for all the pairs (Figure 4).

Finally, we also made R1-R0 plot and the R1-KING-robust kinship plots based on

genotypes that we obtained through a standard genotype calling procedure from the raw

read data. We did this to investigate if the genotype likelihood-based approaches are

necessary or one could just as well use genotypes called from the ~4x data. As expected,

genotype calling had a large negative effect on the outcome; in the resulting R1-R0 and

R1-KING-robust kinship plots the half siblings appear within the range of expected

values for first cousins and both the parent-offspring and full sibling pairs appear within

the range of expected values for half siblings (Figure 4, "called genotypes"). These

results demonstrate the pitfalls of basing any relationship inferences, including R1-R0

and R1-KING-robust kinship plots, on genotypes called from low-depth data. Notably,

this is also the case for the methods presented in Rosenberg (2006) and Manichaikul *et*

al. (2010) (Figure 5A). This clearly demonstrate that, with ~4x sequencing data, calling

genotypes without external information, such as an imputation reference panel, is not a

good alternative to a genotype likelihood-based approach. This implies that that

software packages designed to work only on genotype data, such as KING, should not be

used on data like this.

Assessing the effect of SNP ascertainment

To assess the effect of SNP ascertainment, we applied the new method to three different subsets of data from the five 1000 Genomes individuals. The results for each of the three ascertainment schemes (all sites covered by sequencing data, 1000 Genomes release sites, Illumina 650Y SNP array sites) are similar (left four panels, Figure 4), showing SNP ascertainment does not have a large effect.

For comparison, we performed the same assessment of the methods presented in Rosenberg (2006) and Manichaikul *et al.* (2010) by constructing scatterplots of the same type as those shown in their papers (Figure 5). This revealed that both these other methods are much more affected by ascertainment than the method proposed here. In particular, the Rosenberg method is affected on both its x-axis (IBS0) and its y-axis (IBS2), which means that the expected region of the plot for each relationship will be different for different data sets (top part of Figure 5). The method presented in Manichaikul *et al.* 2010 is affected by the SNP ascertainment mainly on its x-axis (IBS0, bottom part of Figure 5). Therefore, the ascertainment mainly affects the ability to distinguish between parent-offspring and full siblings, since the y-axis, which is only slightly affected by ascertainment, is the kinship coefficient, which can be used to distinguish between most close relationships except for parent-offspring and full siblings. The x-axis, IBS0, is included in part to help make the distinction between PO and FS (Manichaikul *et al.* 2010), but this ability is clearly affected by SNP ascertainment (bottom part of Figure 5).

To further explore the effect of SNP ascertainment on the new method, we also performed analyses of the previously mentioned simulated of data from population with a constant population size. This time we only analyzed SNPs with MAF above 5% and compared the results to the results for the full dataset. This confirmed the results from the real data analyses: SNP ascertainment does change the values a bit compared to

when all sites are analyzed, however the change is limited (Figure 2). This is well in line

with the fact that we got very similar results for the simulated data from three

populations with quite different population size histories and consequently different

allele frequency spectra. Indeed, the effect of population size decline is similar to that of

ascertaining for common SNPs, which makes sense because population decline is known

to lead to a skew in the allele frequency spectrum towards more common SNPs.

Assessing the effect of a limited number of sites

Genome-wide shotgun sequencing data, as is available for the 1000G individuals, is not

available for all species. Studies may instead have RADseq or similar data, covering only

a fraction of genomic sites. To assess to what extent the new method can be used to

analyze such data sets, we performed analyses of subsets of the 1000G data, constructed

to mimic RAD sequencing data. Specifically, we analyzed four subsets that consisted of

the sites from 10k 50k, 100k and 250k 200bp windows, respectively. For all but the

smallest data subset, the point estimates were similar to those obtained using the full

dataset, showing the method is robust to reducing the number of sites with ~4x

coverage (Figure 6, supplemental file 1). This suggests that even with the reduced

number of sites tested, there was sufficient data to characterize the genome-wide mean

IBD fractions for both closely related and unrelated pairs. The uncertainty in the

estimates, as estimated by a chromosome jackknife, increased with fewer sites, but the

effect was limited, suggesting the biological variation in IBD sharing across

chromosomes was larger than sampling variance across the examined sites.

Discussion

We have presented a simple new method for inferring if and how two individuals are

related based solely on genetic data from the two target individuals. This new method

ways in which it can also be applied directly to sequencing data via genotype likelihoods.

And importantly, we show that by using these, the method provides useful results when

applied to ~4x sequencing data as well as RADseq like subsets of such data. All of this

combined implies that – unlike previous methods - this new method can be used even if

all you have is low-depth sequencing data from a few individuals from a species without

a reference genome.

Comparison to similar methods

The new method is based on plotting two statistics, $R0$ and KING-robust kinship against

a third, new statistic $R1$. The $R0$ statistic similar to the test statistic proposed by Lee

(2003) to test for relatedness. The only differences are that the numerator and

denominator are flipped and that E, the proportion of sites where both individuals are

heterozygous, is included in the denominator in the statistic defined by Lee but absent

in $R0$. $R0$ is also similar to the pairwise population concordance (PPC) statistic in

PLINK (Purcell *et al.* 2007), a test of if the genotypes of a pair of individuals have more

IBS0 sites than two unrelated individuals with the same ancestry are expected to have,

signaling they have different ancestries. The KING-robust kinship estimator was

proposed by Manichaikul *et al.* (2010) and implemented for genotype data in the

program KING. Here we extend it to estimation directly from sequencing data. Notably,

our results suggest that this extension is vital for successful application to low-depth

sequencing data, because estimates based on genotypes called from low-depth

sequencing data are very poor (rightmost panels of Figure 4 and 5), which makes

programs like KING inappropriate to apply to such data. And importantly this extension

can also be used for similar statistics and thus makes existing methods based on such

statistics, like KING-robust kinship, more widely applicable.

However, this extension is not the only contribution of this study. Another key new

contribution is provide an alternative to the IBS0 statistic (the proportion of sites where

the two individuals share zero alleles IBS) that was utilized by Rosenberg (2006) and

Manichaikul *et al.* (2010). As we have shown, the fraction of sites that are IBS0 or IBS2

is very sensitive SNP ascertainment, meaning that results are only comparable within

each different ascertainment scheme. The method from Rosenberg (2006), where IBS0

is combined with IBS2, is difficult, if not impossible, to use for relatedness inference

because the fraction of sites that are IBS2 and IBS0 varies so wildly across different

ascertainment schemes, such as between SNP arrays and sequencing data, , , . On the

other hand, KING-robust kinship is still very useful, but it loses the ability to distinguish

between parent-offspring and full siblings, as IBS0 was used for this. Due to this

sensitivity to ascertainment, samples cannot be analyzed in isolation and must be placed

in the context of other samples with known relationships and the same ascertainment

scheme. This requirement makes it difficult to apply these previous methods to ancient

humans or other species with limited sample sizes.

In contrast, the ability to identify relatives based on expected values is maintained in the

new method, regardless of ascertainment scheme due to the use of R0, instead of IBS0,

which makes the new method robust to SNP ascertainment. Parent-offspring pairs tend

to have an R0 estimate extremely close to 0, making them particularly easy to identify

via the R1-R0 plot. The R1-KING-robust kinship plot, on the other hand, has the

appealing aspect that the kinship axis has a biological interpretation, defined as the

probability that two alleles sampled at random from two individuals are identical by

descent. Hence the two plots types, R1-R0 and R1-KING-robust kinship, each have their

advantages. Finally, it is worth noticing that the two plots types seem to work better

than a range of other plots constructed from similar ratio statistics that we explored

(Supplementary Figure S6).

Limitations and applications

While the new method provides substantial advantages over previous methods in

situations with limited data, it does have some limitations. First, like most other

relatedness inference methods, like PLINK, the proposed method assumes that the

individuals are not inbred and that they originate from the same homogenous,

population. And like many other relationship inference methods, it is not necessarily

robust to violations of these assumptions. Previous studies have shown the effect of

population structure and admixture on relatedness inference is complex and can

potentially lead to bias in either direction depending on the circumstances, and this is

true even for KING, which was developed to be robust to population structure

(Thornton *et al.* 2012; Conomos *et al.* 2016; Ramstetter *et al.* 2017). Specific methods

have been developed to correct for admixture when the allele frequencies in the

admixing populations are known (e.g. Thornton *et al.* 2012; Moltke and Albrechtsen

2014), or enough samples are available (Conomos *et al.* 2016; Dou *et al.* 2017). But since

these methods work by exploiting knowledge about allele frequencies or access to many

samples for their correction; the pairwise R0, R1 and KING-robust kinship statistics

cannot be corrected in a similar manner. However, we note that Lee (2003) showed that

the statistic he proposed for testing for relatedness can also be used to detect if two

unrelated samples are not from the same homogeneous population. If this is the case,

the Lee's statistic will be significantly smaller than 2/3; and equivalently R0 will be

significantly above 0.5, which may be useful when interpreting R0, R1, KING-kinship

plots in the presence of admixture or population structure more generally. Regarding

inbreeding, one potential way to assess if one of the individuals is inbred is to compare

heterozygosities across individuals; non-inbred and non-admixed individuals from the

same population should have similar heterozygosity, so marked heterozygosity

differences can be a warning signal.

A second limitation, which is shared with other relatedness estimation methods, is that

there is significant biological variation in the amount of IBD sharing between relatives

with the same pedigree relationship due to randomness inherent in the process of

recombination (Hill 1993; Rasmuson 1993). For humans, this means that a pair of

relatives, say first cousins, will sometimes share less of their genomes IBD than another

pair with a more distant pedigree relationship, say second cousins. This makes

classification into specific relationships difficult. The degree of biological variation in

IBD sharing between relatives varies across species and can even differ between sexes

due to sex-specific recombination patterns. This makes it difficult to provide general

guidance appropriate for all species. In general, species with more chromosomes and

more recombination will have less variation in IBD sharing for a defined pedigree

relationship, making it easier to distinguish among various potential relationship

categories. To quantify this uncertainty, we propose a chromosomal bootstrap

procedure that can be used if reads can be assigned to chromosomes.

Biological variation in IBD sharing is also related to the estimation and interpretation of

confidence intervals on statistics like R0, R1, and KING-robust kinship. Relatedness and

limited recombination also cause correlation between sites in the genome, due to shared

IBD segments and LD. This correlation between sites increases the variance in the

estimates of these statistics in a way that can be difficult to fully account for when

computing confidence intervals. For statistics that test for introgression such as the D-

statistic (Patterson et al. 2012), where the main concern is correlation due to LD, a block

jackknife, leaving out contiguous blocks (e.g. 5Mb) is a common approach. When

considering relatedness, we want to compare our estimates to the expectations of each

relationship category. Since shared IBD segments can be much longer than the range of

LD we propose a more appropriate chromosome-jackknife. In either case, a jackknife (or

bootstrap) over single sites will fail to provide a confidence interval that accounts for the

non-independence of the sites. For more discussion on this topic see Thompson (2013).

Unfortunately, this means that it is difficult to provide the most appropriate confidence

intervals when no information about genomic positions is available.

Despite these limitations, we believe that the results presented here suggest the new

method constitutes a helpful new tool for relatedness inference for studies with limited

data. Identifying related samples is a crucial step in nearly any genetic analysis and can

also reveal other problems such as duplicate samples or cross contamination of genetic

material. Removing the requirements to specify allele frequencies and to have accurate

genotypes has the potential allow the identification of relatives even in small studies of

non-model species or ancient samples. These types of studies do not currently have

many good options to address relatedness.

Data Accessibility

697 The IBS method is available at:

698 http://www.popgen.dk/angsd/index.php/Genotype_Distribution

699 The data sets used are publicly available.
700 The HGDP SNP array data are available at:
701 ftp.cephb.fr/hgdp_supp1
702
703 The 1000G phase 3 aligned sequencing data are available at:
704 ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/phase3/data
705
706 The 1000G phase 3 called genotypes are available at:
707 ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/
708

References

Abecasis, G. R., S. S. Cherny, W. Cookson and L. R. Cardon, 2001 GRR: graphical
representation of relationship errors. Bioinformatics 17: 742-743.

Albrechtsen, A., T. Sand Korneliussen, I. Moltke, T. van Overseem Hansen, F. C. Nielsen
et al., 2009 Relatedness mapping and tracts of relatedness for genome-wide data
in the presence of linkage disequilibrium. Genet Epidemiol 33: 266-274.

Baca, M., K. Doan, M. Sobczyk, A. Stankovic and P. Weglenski, 2012 Ancient DNA
reveals kinship burial patterns of a pre-Columbian Andean community. BMC
Genet 13: 30.

Balding, D. J., 2006 A tutorial on statistical methods for population association studies.
Nat Rev Genet 7: 781-791.

Busing, F. M. T. A., E. Meijer and R. Van Der Leeden, 1999 Delete-m jackknife for
unequal m. Statistics and Computing 9: 3-8.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al.*, 2015 Second-
generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience 4: 7.

Conomos, M. P., A. P. Reiner, B. S. Weir and T. A. Thornton, 2016 Model-free
Estimation of Recent Genetic Relatedness. Am J Hum Genet 98: 127-148.

Derrien, T., J. Estelle, S. Marco Sola, D. G. Knowles, E. Raineri *et al.*, 2012 Fast
computation and applications of genome mappability. PLoS One 7: e30377.

Dou, J., B. Sun, X. Sim, J. D. Hughes, D. F. Reilly *et al.*, 2017 Estimation of kinship
coefficient in structured and admixed populations using sparse sequencing data.
PLoS Genet 13: e1007021.

Gusev, A., J. K. Lowe, M. Stoffel, M. J. Daly, D. Altshuler *et al.*, 2009 Whole population,
genome-wide mapping of hidden relatedness. Genome research.

Hill, W. G., 1993 Variation in Genetic Identity within Kinships. Heredity 71: 652-653.

Hill, W. G., and B. S. Weir, 2011 Variation in actual relationship as a consequence of
Mendelian sampling and linkage. Genet Res (Camb) 93: 47-64.

Kardos, M., G. Luikart and F. W. Allendorf, 2015 Measuring individual inbreeding in the
age of genomics: marker-based measures are better than pedigrees. Heredity
(Edinb) 115: 63-72.

Kelleher, J., A. M. Etheridge and G. McVean, 2016 Efficient Coalescent Simulation and
Genealogical Analysis for Large Sample Sizes. PLoS Comput Biol 12: e1004842.

Korneliussen, T. S., A. Albrechtsen and R. Nielsen, 2014 ANGSD: Analysis of Next
Generation Sequencing Data. BMC Bioinformatics 15: 356.

Korneliussen, T. S., and I. Moltke, 2015 NgsRelate: a software tool for estimating
pairwise relatedness from next-generation sequencing data. Bioinformatics 31:
4009-4011.

Kuhn, J. M. M., M. Jakobsson and T. Gunther, 2018 Estimating genetic kin
relationships in prehistoric populations. PLoS One 13: e0195491.

Lee, W. C., 2003 Testing the genetic relation between two individuals using a panel of
frequency-unknown single nucleotide polymorphisms. Annals of Human
Genetics 67: 618-619.

Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data.
Bioinformatics 27: 2987-2993.

Manichaikul, A., J. C. Mychaleckyj, S. S. Rich, K. Daly, M. Sale *et al.*, 2010 Robust
relationship inference in genome-wide association studies. Bioinformatics 26:
2867-2873.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res 20: 1297-1303.

Moltke, I., and A. Albrechtsen, 2014 RelateAdmix: a software tool for estimating
relatedness between admixed individuals. Bioinformatics 30: 1027-1028.

Nielsen, R., T. Korneliussen, A. Albrechtsen, Y. Li and J. Wang, 2012 SNP calling,
genotype calling, and sample allele frequency estimation from New-Generation
Sequencing data. PLoS One 7: e37558.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira *et al.*, 2007 PLINK: a
tool set for whole-genome association and population-based linkage analyses. Am
J Hum Genet 81: 559-575.

Ramstetter, M. D., T. D. Dyer, D. M. Lehman, J. E. Curran, R. Duggirala *et al.*, 2017
Benchmarking Relatedness Inference Methods with Genome-Wide Data from
Thousands of Relatives. Genetics 207: 75-82.

Rasmuson, M., 1993 Variation in Genetic Identity within Kinships. Heredity 70: 266-
268.

Rosenberg, N. A., 2006 Standardized subsets of the HGDP-CEPH Human Genome
Diversity Cell Line Panel, accounting for atypical and duplicated samples and
pairs of close relatives. Ann Hum Genet 70: 841-847.

Sikora, M., A. Seguin-Orlando, V. C. Sousa, A. Albrechtsen, T. Korneliussen *et al.*, 2017
Ancient genomes show social and reproductive behavior of early Upper
Paleolithic foragers. Science 358: 659-662.

Stevens, E. L., G. Heckenberg, E. D. Roberson, J. D. Baugher, T. J. Downey *et al.*, 2011
Inference of relationships in population data using identity-by-descent and
identity-by-state. PLoS Genet 7: e1002287.

The Genomes Project, C., 2015 A global reference for human genetic variation. Nature
526: 68-74.

Thompson, E. A., 1975 The estimation of pairwise relationships. Ann Hum Genet 39:
173-188.

Thompson, E. A., 2000 Statistical inference from genetic data on pedigrees, pp. i-169 in
NSF-CBMS regional conference series in probability and statistics. JSTOR.

Thompson, E. A., 2013 Identity by descent: variation in meiosis, across genomes, and in
populations. Genetics 194: 301-326.

Thornton, T., H. Tang, T. J. Hoffmann, H. M. Ochs-Balcom, B. J. Caan *et al.*, 2012
Estimating kinship in admixed populations. Am J Hum Genet 91: 122-138.

Wiuf, C., 2006 Consistency of estimators of population scaled parameters using
composite likelihood. J Math Biol 53: 821-841.

Author Contributions

801 AA, IM, and RKW designed the research. RKW performed research, with input from IM

802 and AA. IM and RKM wrote the paper, with help from AA.

Figures for:

Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

Authors:

Ryan K. Waples
Anders Albrechtsen
Ida Moltke

(a)

(b)

Ratio 0 (R0):

$$\frac{C+G}{E}$$

(c)

Ratio 1 (R1):

$$\frac{E}{B+D+H+F+C+G}$$

(d)

KING-robust kinship:

$$\frac{E-2(C+G)}{B+D+H+F+2E}$$

Figure 1. Definitions of pairwise genotype categories A - I and the R0, R1, and KING-robust kinship statistics. **(a)** Definition of the pairwise genotype categories A - I. Here g_1 and g_2 denotes the genotype for each of the two diploid individuals, 1 and 2, respectively. These genotypes are defined as the number of copies of a certain allele carried by 1 and 2, respectively. We assume diallelic variants such that g_1 and g_2 each has 3 possible values: 0, 1, and 2. For a pair of individuals, there are nine possible genotype combinations. We organize them into a 3x3 matrix and denote them with the letters from A to I. The values A – I can equivalently be either counts or proportions. **(b)** Definition of the R0 statistic based on the notation illustrated in A. **(c)** Definition of the R1 statistic based on the notation illustrated in A. **(d)** Definition of the KING-robust kinship estimator (Manichaikul *et al.* 2010), formulated using the notation illustrated in (a).

Figure 2. Ranges of expected values and simulation results for R1-R0 and for R1-KING robust kinship for each of five relationship categories: parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1), and unrelated (UR). **Top)** The colored shaded areas (sometimes just lines) show theoretically derived ranges of the joint expectation for R1-R0 based on expected IBD sharing (i.e. values of k0, k1 and k2) for each relationship across all possible allele frequency spectra. For PO this range is a singular point and is shown as a shaded purple X. The colored symbols (triangle, circle, star) and black 'x's show values for each relationship obtained from data simulated under four different scenarios. Three of the scenarios are different demographic histories: 1) a 10-fold increase in Ne over 100 generations, 2) constant N_e, and 3) a 10-fold decrease in N_e over 100 generations. The fourth scenario is also constant N_e, but sites are ascertained to have allele frequency above 5%. Note that, while it is difficult to see due to overplotting, all simulated values for PO fall very close to (R1, R0) = (0.5, 0). **Bottom)** same top, but for (R1, KING robust kinship). Here all simulated values for PO fall very close to (R1, KING robust kinship) = (0.5, 0.25).

Figure 3. R1-R0 and R1-KING-robust kinship scatterplots for seven HGDP populations. Each colored point represents a pair of individuals and is colored according to the relationship category inferred using an allele frequency based approach. Colored shaded areas/lines show the theoretically derived range of expected values for specific relationship categories as in Figure 2. Black 'X's show the values for a pair of individuals simulated under a constant population size demography from Figure 2. Note that in addition to the relationship categories for Figure 2 there is an additional category here representing distantly related pairs (DR). **(a)** R1-R0 plot for all pairs of individuals within each population **(b)** R1-KING-robust kinship plot for all pairs of individuals within each population. (**c**) Scatterplot of the two relatedness coefficients k_1 and k_2 for all pairs of individuals within each population estimated using the allele frequency-based approach implemented in PLINK. Note that the black 'X's here show simulated values for k_1 and k_2 and are not inferred by PLINK, they approximately coincide with the expected values of k_1 and k_2 for each relationship category (Supplementary Table S1).

Figure 4. Relatedness plots for all pairs among five LWK individuals from the 1000G Project. **Top)** R1-R0 scatterplots for pairs of five LWK individuals for five different analysis methods: 1) IBS: reference-free estimation from (~4x) 1000G bam files, 2) realSFS: site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G sites: curated 1000G genotypes from the 1000G project, 4) HGDP sites: curated 1000G genotypes but only at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes: genotypes called de novo from (~4x) 1000G bam files.. Points are colored by their true relationship status, as reported by 1000G. Thin grey lines show confidence intervals (+/- 2 SE) estimated using a chromosome jackknife. Colored shaded areas/lines show the theoretically derived range of expected values for specific relationship categories from Figure 2. Black 'X's show the values for a pairs of related individuals simulated under a neutral demography, as in Figure 2. **Bottom)** R1-KING-robust kinship scatter plots for the same data sets, confidence intervals and expected ranges are constructed in the same way.

Figure 5. Plots of two alternate frequency-free methods for different subsets of SNP sites. **Top)** Results from applying the plotting approach from Rosenberg (2006) to pairs of the same five LWK individuals for five different analysis methods: 1) IBS: reference-free estimation from (~4x) 1000G bam files, 2) realSFS: site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G sites: curated 1000G genotypes from the 1000G project, 4) HGDP sites: curated 1000G genotypes at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes: genotype called de novo from (~4x) 1000G bam files. Pairs are colored by their true relationship status, as in Figure 3. Fraction IBS0/IBS2 are the overall fraction of sites that are IBS0/IBS2, respectively. Gray lines centered on each point show confidence intervals (+/- 2 SE) based on a chromosome jackknife. **Bottom)** Results from applying the KING-robust based approach to the same pairs of LWK individuals using the same five different analysis methods as in A. The horizontal black lines show the kinship thresholds used to distinguish unrelated (UR), first-cousins (C1) half siblings (HS), full siblings (FS) and (PO) following (Manichaikul *et al.* 2010) from bottom to top, respectively. Like in A, thin grey lines centered on each point show confidence intervals (+/- 2 SE) estimated using chromosome jackknife.

Figure 6. The effect on estimates of R0, R1, and KING-robust kinship from reducing the number of sites analyzed from the same five LWK individuals as in figures 4 and 5. Each point shows the point estimate and error bars show +/- 2SE estimated using chromosome jackknife. Each column shows results for different numbers of examined basepairs. Pairs of individuals are colored by their true relationship. These plots show the results of the IBS-based method, see supplemental for results from the SFS-based approach. Colored shaded areas/lines show the theoretically derived range of expected values for specific relationship categories from Figure 2 and black 'X's show the values for a pair of individuals simulated under a neutral demography from Figure 2.