Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

For submission to Genetics as an Investigation

Authors:

Ryan K Waples*, Anders Albrechtsen*,[1], Ida Moltke*,[1]

*Department of Biology, Section for Computational and RNA Biology, University of Copenhagen, 2200 Copenhagen N, Denmark,

[1]Corresponding authors: Department of Biology, University of Copenhagen, Ole Maaloes Vej 5, 2200 Copenhagen N, Denmark. E-mail: albrecht@binf.ku.dk; and ida@binf.ku.dk

Abstract

Knowledge of how individuals are related is important in many areas of research and numerous methods for inferring pairwise relatedness from genetic data have been developed. However, the majority of these methods were not developed for situations where data is limited. Specifically, most methods rely on the availability of population allele frequencies, the relative genomic position of variants, and genotype data. But in studies of non-model organisms or ancient human samples, such data is not always available. Motivated by this, we present a new method for pairwise relatedness inference, which requires neither allele frequency information nor information on genomic position. Furthermore, it can be applied to both genotype data and to low-depth sequencing data where genotypes cannot be accurately called. We evaluate it using data from SNP arrays and low-depth sequencing from a range of human populations and show that it can be used to infer close familial relationships with a similar accuracy as a widely used method that relies on population allele frequencies. Additionally, we show that our method is robust to SNP ascertainment, which is important for application to a diverse range of populations and species.

Introduction

The ability to infer the familial relationship between a pair of individuals from genetic data plays a key role in several research fields. In conservation biology, it is used to design breeding programs that minimize inbreeding (Kardos *et al.* 2015), in archeology it is helpful to understand burial patterns and other cultural traditions (Baca *et al.* 2012; Sikora *et al.* 2017), and in population and disease genetics it is often used to exclude relatives, because many analysis methods within those fields assume all sampled individuals are unrelated and violations of this assumption can lead to wrong conclusions (Balding 2006).

Numerous pairwise relatedness inference methods have been developed, e.g., Thompson (1975), Lee (2003), Purcell *et al.* (2007), Albrechtsen *et al.* (2009), Manichaikul *et al.* (2010), Stevens *et al.* (2011), Korneliussen and Moltke (2015), and many are available in popular software packages, e.g., PLINK (Purcell *et al.* 2007), and KING (Manichaikul *et al.* 2010). Most of these methods estimate either the three relatedness coefficients k_0, k_1 and k_2, or the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$ for each pair of individuals, where k_0, k_1 and k_2 are the proportions of the genome where a pair of individuals share 0, 1, or 2 alleles identical by descent (IBD) (Thompson 2000). By definition, alleles are IBD when they are identical due to *recent* shared ancestry, but because alleles can also be identical due to older shared ancestry or recurrent mutations, IBD status cannot be directly observed. Therefore, the pairwise relatedness coefficients and the kinship coefficient have to be estimated, which can be done from patterns of observed genetic identity (identity by state or IBS) together with population allele frequencies. Once estimated, these relatedness statistics, k_0, k_1, k_2, and the kinship coefficient can be used to infer familial relationships by comparison to the expectation of the statistics under different familial relationships (Hill and Weir 2011).

Although inference of relatedness is of wide interest, most existing methods are not immediately applicable in studies with limited data or genetic resources. First, most existing methods require that the allele frequencies of the source population are known. For most studies in modern humans this is not a problem. However, in studies of

ancient humans or other species, accurate estimates of allele frequencies are not always

available, due to limited number of samples.

Second, several existing methods examine loci in sliding windows or use hidden Markov

models to leverage the non-independence of allele sharing along the genome between

relatives, (e.g., Stevens *et al.* 2011; Monroy Kuhn *et al.* 2017). These methods are

powerful, but require information about the genomic position of variable sites, and for

non-model organisms high-quality reference genome assemblies are often not available.

Finally, nearly all existing methods --both frequency-based and others-- require

genotype data. However, in sequencing studies, samples are often only sequenced to low

depth due to cost and technical issues, which makes it infeasible to call genotypes

accurately (Nielsen *et al.* 2012), precluding the use of these methods. There are a few

methods that estimate relatedness from low depth data by utilizing genotype likelihoods

instead of genotypes (e.g., Korneliussen and Moltke 2015), however, these methods still

rely on accurate allele frequency estimates.

Hence, most existing methods to infer close familial relationships are not immediately

applicable in studies where data is limited, including many studies of non-model

organisms and ancient humans. One of the few exceptions is a simple but elegant test

for pairwise relatedness proposed in Lee (2003), that requires neither information on

allele frequencies nor the relative genomic position of the variable sites. This test relies

entirely on the relative frequency of different genotype combinations within the two

individuals and thus only requires genotype data from the two target individuals. While

useful, this test does not provide any means to distinguish between different types of

close familial relationships; it only provides a statistical test for relatedness vs the null

hypothesis of unrelated.

To our knowledge, there are only two published methods that can be used to distinguish

between different types of familial relationships when neither allele frequencies nor

information about the relative genomic position of sites is obtainable. The first of these

two methods is to plot the proportion of the genomic sites in which the two individuals

share both alleles IBS (which we denote IBS2) versus the proportion of the genomic

sites in which they share zero alleles IBS (which we denote IBS0). This method was used

in Rosenberg (2006), where it was applied to the Human Genome Diversity Project

(HGDP) dataset. In the resulting plot of the HGDP data (figure 1 of Rosenberg (2006)),

pairs of individuals with the same relationship category form distinct clusters in such a

way that it is possible to identify parent-offspring pairs, full sibling pairs, and to a lesser

extent more distant relationships such as half-siblings/avuncular/grandparent-

grandchild and first cousins.

The second existing method is based in part on the KING-robust kinship estimator

(Manichaikul *et al.* 2010), which was developed to overcome biases in kinship estimates

caused by cryptic population and family structure. More specifically, the KING-robust

estimator was developed to be robust to violations of the assumption that the genotypes

of the pair of individuals are sampled from a single homogenous population. It was not

developed with the aim of providing a kinship estimator that works when allele

frequencies within a single population are unknown. However, it is directly applicable to

this scenario because, like the test suggested in Lee (2003), it relies only on the genotype

combinations within the two target individuals and does not require specification of

allele frequencies. Importantly, Manichaikul *et al.* (2010) show it is possible to use SNP

array data to infer if a pair of individuals are parent-offspring, full sibling, half

siblings/avuncular/grandparent-grandchild, cousins, or unrelated by jointly considering

the KING-robust estimator and the fraction of sites IBS0. For example, see figure 3A in

Manichaikul *et al.* (2010), a scatterplot of KING-robust kinship vs the fraction of sites

IBS0.

Notably, both existing methods discussed above have two key limitations. First, like

most other pairwise relatedness estimation methods, they were developed for accurately

called genotypes. For example, the KING software (Manichaikul *et al.* 2010)

implementing the KING-robust kinship estimator requires genotype data as input,

which can be problematic for studies where only medium or low-depth sequencing data

is available. Second, both methods rely on estimates of the fraction of the genomic sites

in which the pair of individuals share zero alleles identical by state, IBS0. This can be

problematic because this fraction is highly sensitive to the SNP ascertainment

procedure. This means that the results of the methods are platform-dependent and are

likely to differ between different SNP arrays and especially between SNP array and
sequencing datasets.

Motivated by the outlined limitations to the existing methods, we present a method for
relationship inference among non-inbred individuals from the same homogeneous
population that relies neither on allele frequencies nor on information about the relative
position of the variant sites. The method is inspired by previous methods; it uses the
KING-robust kinship estimator and a statistic R_0, which is similar to the test statistic
from the test for relatedness suggested by Lee (2003). However, it also uses a third
statistic, R_1, which helps distinguish between full siblings and parent-offspring. Very
briefly, the method consists of using two combinations of these three statistics, R_1-R_0
and R_1-KING-robust kinship, to infer relationships and compared to previous similar
methods it has the advantage that it is robust to SNP ascertainment. Also, we present
two ways to estimate the three statistics directly from sequencing data in the absence of
called genotypes, which makes the method applicable not only to genotype data, like
previous similar methods, but also to low-depth sequencing data.

In the following we first fully describe the three statistics, R_0, R_1 and KING-robust
kinship, how they can be estimated and other methodological details. Next, using
publicly available SNP array data, we show that the new method provides similar
accuracy to the frequency-based method implemented in PLINK, without utilizing any
information on allele frequencies or genomic positions. Then using sequence data from
the 1000 Genomes, we show that these statistics can be estimated directly from
sequencing data of moderate depth (~4x) and that utilizing the sequencing data directly
is markedly better than calling genotypes and subsequently estimating R_0, R_1, and
kinship from the called genotypes. Also, using different subsets of the same data, we
show that this new method, unlike previous similar methods, is robust to SNP
ascertainment. Finally, we discuss some potential applications and limitations.

Methods and materials

The R0, R1, and KING-kinship statistics

The method for relationship inference we propose consists of first estimating three statistics called R0, R1 and KING-robust kinship from genetic data, then constructing two plots: (R1, R0) and (R1, KING-robust kinship), and finally interpreting these plots.

To define the three statistics, R0, R1 and the KING-robust kinship, we note that for a pair of diploid individuals, all diallelic sites fall into one of nine genotype combinations; all possible combinations of the two individuals carrying 0, 1, or 2 copies of a specific allele, e.g. the ancestral allele. For a pair of individuals, we can therefore fully characterize their genome-wide IBS-sharing pattern in nine counts or proportions denoted: A, B, C, D, E, F, G, H, and I (Figure 1A), similar to a two-dimensional site-frequency spectrum (SFS) across the two individuals. This is important to note because R0 and R1 statistics are defined as simple functions of a subset of these nine counts as shown in Figure 1B and C, and the KING-robust kinship statistic, originally defined by Manichaikul *et al.* (2010), can also be re-formulated as a function of these 9 values (Figure 1D).

The new method is motivated by several observations. First, the expected values of A-I vary depending on the familial relationship. Consequently, so do functions of these counts, including R0, R1 and KING-robust kinship. Specifically, for two non-inbred individuals from the same homogenous population [R1, R0] and [R1, KING-robust kinship], both have the property that the joint expectation ranges for the five relationship categories parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1) and unrelated (UR) do not overlap (Figure 2, for derivations see supplementary text) regardless of the underlying allele frequency spectrum. Second, this method requires neither population allele frequencies nor information about the relative position of the genomics sites analyzed. The only information that is required to estimate the three statistics is genetic data from the two individuals of interest. Third, we expect these statistics to be robust to

SNP ascertainment because they are ratios computed from sites that are variable within
the two samples and should thus be unaffected by the number of non-variable sites and
because the (unknown) underlying frequency spectrum should only have a limited effect
on these ratios. Finally, while the counts of the nine genotype combinations, A-I, and
thus R0, R1, and the KING-robust kinship estimator can be calculated from genotype
data, they can also be estimated directly from next-generation sequencing (NGS) data
based on the expected number of sites with each genotype combination. This makes the
approach appropriate even when the sequencing depth is too low for accurate genotype
calling (see below for methodological details).

Estimation from sequencing data

The nine genotype combinations, A-I, and thus R0, R1, and the KING-robust kinship for
a pair of individual, can be estimated directly from NGS data via the use of genotype
likelihoods calculated from the aligned sequencing reads. By using genotype likelihoods
it is possible to account for the genotype uncertainly inherent to low-depth NGS data.
We used two distinct, but related approaches to estimate these statistics from
sequencing data that both build on this idea.

The first approach, which we denote the IBS-based approach, considers all ten possible
genotypes at a diallelic site for each individual and consists of a maximum-likelihood
(ML) estimation of the counts of each of the 100 (10 × 10) possible genotype pairs (for
details see supplementary text). To perform the ML estimation we used an EM
algorithm, which we have added to the `ANGSD` software package under the name `IBS`.
After obtaining the estimate of the counts of all the 100 possible genotype pairs, we
convert them into estimates of the genotype combinations A-I, by summing over the
counts that correspond to each combination. For example, the genotype pairs: [AA/AA,
CC/CC, GG/GG, TT/TT] all contribute to cells A or I of Figure 1. Count corresponding to
genotype pairs with more than two alleles were discarded. The advantage of this IBS-
based approach is that it does not require specification of an allele at each site, and can
thus be applied to nearly any sequencing data set, even with low depth, without any
prior information on the alleles at each site.

The second approach, which we denote the SFS approach, consisted of performing ML
estimation of the two-dimensional site-frequency spectrum (SFS) (Nielsen *et al.* 2012).
To find the ML estimate of the SFS, we used an expectation-maximization (EM) method
`realSFS` from the `ANGSD` software package (Korneliussen *et al.* 2014). The SFS method
requires one allele to be specified for each site e.g., the ancestral, consensus or reference
allele. The model underlying this method assumes that all genotypes for each site have
the possibility of containing this specified allele and up to one other unspecified allele.
To specify the alleles that do exist at each site, we used the consensus sequences from
the highest depth individual (NA19042) and restricted our analysis to sites where the
depth in this individual was at least three.

For both these genotype likelihood-based methods, we analyzed each chromosome
separately due to computational constraints. We used the original `GATK` genotype
likelihood model (Mckenna *et al.* 2010) with independent errors, as implemented in
`ANGSD`, to calculate the likelihood of sequencing data for each possible genotype. We also
tested the `samtools` genotype likelihood model (Li 2011) with a more complicated error
structure, but found that it produced worse results in this case (data not shown). Both
the IBS approach and the `realSFS` produces the expected counts for each value A-I and
the values for each chromosome were summed over to produce a genome-wide estimate.
For example command lines for each analysis, see supplemental text.

Confidence intervals

All of the above estimation methods treat each site as independent. This assumption
should not affect our expectation of each statistic (Wiuf 2006), but statistical non-
independence (e.g. linkage disequilibrium and IBD) does affect standard estimates of
uncertainty. To quantify the uncertainty, we therefore estimated confidence intervals for
all statistics using a block-jackknife procedure. Confidence intervals were calculated by
leaving each chromosome out (chromosome jackknife), which takes both the IBD and
the LD correlation into account. The weighted block jackknife variance estimator of
Busing *et al.* (1999) was used to estimate the variance from the distribution of estimates

234 for each statistic. The square root of this variance was interpreted as the standard error
235 in our estimate.

Application to real datasets

237 To assess the utility of the new method, we applied it to two different publicly available
238 datasets: SNP array data from seven HGDP populations (Rosenberg 2006), and
239 sequencing data from five related individuals from the Luhya in Webuye, Kenya (LWK)
240 population of the 1000 Genomes Project phase 3 (The Genomes Project 2015).

HGDP SNP array

The HGDP SNP array dataset was downloaded from `ftp.cephb.fr/hgdp_supp1` (accessed
Jan. 13, 2017). Prior to any filters, the collated HGDP dataset had 1043 individuals,
644,258 variants, and a genotyping rate of 0.999.

Following Rosenberg (2006), we performed basic quality control to exclude error-prone
or duplicated samples. Specifically, based on the HGDP subsets described in:
`web.stanford.edu/group/rosenberglab/supplements/hgdprelSupp.pdf` (last accessed Jan. 13,
2017), we excluded subset 3: "found by (Rosenberg *et al.* 2002) to be extremely atypical
and potentially mislabeled". We also removed the duplicate samples listed in subsets 6
and 7 and retained subset 9 containing all remaining samples.

Next, we selected populations from the HGDP based on two criteria: 1) at least 15
unadmixed individuals per population, and 2) presence of several close familial
relationship among the individuals. Based on these criteria, we considered samples from
seven target populations: five non-African populations: Surui, Pima, Karitiana, Maya,
and Melanesian, and two African populations: Mbuti Pygmies and Biaka Pygmies.

We excluded individuals showing obvious signs of admixture or inbreeding from the
selected HGDP populations. To quantify the admixture proportions of each individual
we ran `ADMIXTURE` (Alexander *et al.* 2009) separately for each of the seven target
population, while also including samples from several reference populations. In the
analyses of each of the non-African target populations we included French, Han, and

Yoruban individuals and selected K=4, and in the analyses of each of the African target

populations we included French and Han individuals only and selected K=3. To select a

set of autosomal loci for ADMIXTURE, filters for genotyping rate (>99%), MAF (>5%),

HWE (p > 10^{-10} with midp adjustment), and LD (indep-pairwise 50 5 0.5) were applied

to each set of selected individuals. Twenty replicate runs of ADMIXTURE were performed

for each of the seven target populations; all replicates gave rise to likelihoods within 3

log likelihood units of each other, suggesting convergence was reached. The results from

the replicate with the highest likelihood were selected and individuals in the target

population with >5% contribution from more than one ancestry component were

excluded. Admixture analyses excluded 16 individuals: 1 Karitiana, 1 Surui, 5 Pima, and

9 Maya. The remaining individuals were split by population and we constructed a final

set of loci by retaining autosomal loci with genotyping rate (>99%), MAF (>5%), HWE

(p > 10^{-4}, with midp adjustment), from the original HGDP data (no LD filter). Finally,

estimates of the inbreeding coefficient (F) of each individual were calculated within each

population and individuals with F > 0.0625 were excluded, excluding two Karitiana.

This left us with a total of 142 remaining individuals across the six populations: Biaka

Pygmies N=31, Karitiana N=21, Maya N=16, Mbuti Pygmies N=15, Melanesian N=19,

Pima = 20, Surui N=20. This produced 2902 within-population pairs of individuals that

were evaluated for relationship status. The R0, R1, and KING-robust kinship statistics

for each pair were calculated from all sites where both individuals had non-missing

genotypes.

1000 Genomes sequencing data

We selected five individuals from two families in the Phase 3 1000 Genomes (1000G)

Luhya in Webuye, Kenya (LWK) population: NA19027, NA19042, NA19313, NA19331,

NA19334. Across the five individuals, there is one pair of half-sibs (NA19027 &

NA19042), and a separate trio of related individuals with a pair of full siblings (NA19331

& NA19334), one parent-offspring relationship (NA19313 & NA19331), and another

unspecified second-degree relationship (NA19313 & NA19334), possibly avuncular (The

Genomes Project 2015). These stated relationships leave six unrelated pairs among the
five individuals.

For each pair of the five LWK individuals we estimated the R0, R1 and KING-robust
kinship statistics in four different ways: 1) by simple genotype counting based on the
phased and high-quality curated genotypes provided in the hg37 1000G VCF files, 2&3)
by applying the two different sequencing-based approaches described above to the
1000G aligned sequence files (bam files) and 4) by re-calling genotypes from the same
1000G bam files and then simple counting from the called genotypes. In the fourth
estimation method, genotypes were called in a basic manner meant to mimic data from
a species with a reference genome but few other genetic resources. Specifically, we used
`samtools mpileup` (v1.3.1) to summarize the reads overlapping each position, and
`bcftools call` (v1.3.1) to assign the most likely genotype at each position. We used
mostly default settings; non-default flags to `samtools` specified skipping indel positions.
Non-default flags to `bcftools` specified using the consensus caller and requesting only
variant sites. For all sequence-based analyses we only considered reads with a minimum
phred-scaled quality score of 30 and bases with minimum phred-scaled quality score of
20 and we restricted our analyses to regions with a GEM 75mer mappability of 1
(Derrien *et al*. 2012).

Assessing the effect of SNP ascertainment

To evaluate the effect of SNP ascertainment, we created a subset of the high-quality
genotype data from the five 1000 Genomes individuals. We selected the sites that
overlap with the Illumina 650Y array that was used for the HGDP and estimated our
three relatedness statistics. We compared the results from this subset of HGDP sites to
results for the full genotype dataset and also to the sequence-based analyses. For an
additional comparison, we also performed the same comparison for the methods
presented in Rosenberg (2006) and Manichaikul *et al*. (2010) by constructing
scatterplots by their methods.

Comparison to other methods

To get an allele frequency-based categorization of relationships for the HGDP genotype dataset described above, we first applied the commonly used the allele frequency based relatedness inference program PLINK (v1.9) (Chang *et al.* 2015) to the individuals from each population separately to estimate the genome-wide IBD fractions k_0, k_1, k_2, (Z0, Z1 and Z2 in PLINK (Purcell *et al.* 2007). Next, we applied the relationship criteria proposed in Table 1 of Manichaikul *et al.* (2010) to the relatedness coefficient estimates: the estimated k values were combined into an estimate of the kinship coefficient $\theta = \frac{k_1}{4} + \frac{k_2}{2}$, and a relationship degree was assigned to each pair of individuals based on comparing the estimated kinship coefficient to the criteria in the table. Parent-offspring and full siblings were differentiated based on k_2 values. This provided us with a categorization into five categories: parent offspring (PO), full siblings (FS), half-siblings / avuncular / grandparent-grandchild (HS), first cousin (C1), and unrelated (UR). To achieve additional resolution, we further divided the last category (UR) into two: distantly related (DR) and unrelated (UR). We did this by simply extending the logic behind the criteria proposed above. Specifically, we set the kinship threshold between DR and UR to $1/2^{13/2}$, which corresponds to including 2nd to 5th degree relatives in the DR category.

To estimate the statistics for identifying related individuals proposed by Rosenberg (Rosenberg 2006) and KING (Manichaikul *et al.* 2010) we note that they can all be calculated directly from the same nine proportions, A-I, as R1, R2 and King-robust kinship:

KING-kingship = (E − 2(C+G))/ (B+D+H+F+2E)

Fraction IBS0 = (C+G)/(A+B+C+D+E+F+G+H+I)

Fraction IBS2 = (A+E+I)/(A+B+C+D+E+F+G+H+I)

We used these formulas, because they allow for estimation of the statistics directly from sequence data in the same manner as for R0, R1 and KING-robust kinship. This allowed us to use sequence data to compare all the methods on a level playing field.

Results

To assess to how the new method works in practice, we applied it to real data from different data generation platforms and compared the results to those of other methods.

Application to SNP array data

We first applied the method to SNP array data from the Human Genome Diversity Project (HGDP) to see how the method worked on this type of data. More specifically, we applied it to data from seven populations from the HGDP after removing samples with signs of admixture or inbreeding. This resulted in the R1-R0 and R1-KING-robust kinship plots shown in Figure 3 A and B (for population-specific plots see Figure S1 and S2). Because the true relationships for the pairs of individuals is not known, we colored each point not according to the truth, but according to the relationship category inferred using a common allele frequency-based method (Figure 3C, Figure S3, for further details see Methods). Since there are least 15 individuals in each of the seven selected HGDP populations, the allele frequency estimates for these populations should be reasonably accurate even with some relatedness. Thus, although this classification and coloring scheme does not necessarily reveal the true relationships for all pairs (Hill and Weir 2011) it should provide a fair picture of how well our method performs in comparison to a standard allele frequency-based method.

In general, points representing pairs of individuals that according to the allele frequency-based method have the same relationship, clearly cluster together on both the R1-R0 plot and the R1-KING-robust kinship plot (Figure 3). Also, they are located fairly close to their expected ranges. More specifically, almost all pairs identified as parent-offspring (PO) by the frequency-based method are easy to identify as such in both the R1-R0 plot and the R1-KING-robust kinship plot, which is not the case when only a single statistic is used (see also Figures S1 and S2). The same is true for full siblings (FS). Points classified as half-siblings/avuncular/grandparent-grandchild (HS) or first cousins (C1) by the frequency-based method have a minimal overlap with each other and with less-related pairs (see highlighted areas of Figure 3A and B). But notably the

372 few pairs of individuals that are difficult to classify are the same pairs as those that are
373 edge cases for the allele frequency-based method. This is apparent in an R1-R0 plot of
374 the HGDP data constructed excluding pairs that are closer than 0.01 to the kinship
375 coefficient thresholds that the frequency based method used when categorizing
376 relationships (Figure S4).

Application to sequencing data

To assess how well the new method works on sequencing data, we applied it to
sequencing data from five low-depth (~4x) human genomes from the 1000 Genomes
project. Among the five selected samples there is a parent-offspring pair, a pair of full
siblings, a pair of half siblings, an unspecified 2nd degree relationship (e.g. avuncular),
and the rest are unrelated. We estimated the three statistics for each pair in several
ways. First, by using an IBS-based approach that estimates the proportion all pairwise
combinations of the 10 possible genotypes (Figure 4). Second, we estimated the two-
dimensional site frequency spectrum (2D-SFS) of each pair with a bi-allelic model and
calculated R0, KING-robust kinship, and R1 based on this spectrum (Figure 4). Both
these approaches base their estimates on genotype likelihoods calculated from
sequencing read data, instead of on called genotypes, and take the uncertainty of the
underlying genotypes that is inherent to low-depth sequencing data into account. The
key difference between them is that the SFS-based approach requires specification of an
allele at each site, potentially excluding studies with low-depth data from non-model
species, whereas the IBS approach has no such requirement (for methodological details
see Methods and materials). The methods also differ in how they deal with sites with
more than two unique alleles. We include both methods here to compare their
performance on real data.

Additionally, we calculated the three statistics from the high-quality phased genotypes
for the same five individuals available from the 1000 Genomes Project Phase 3 (Figure
4) to see how well the two genotype-likelihood-based approaches applied to low-depth
sequencing data perform compared to direct calculations from high-quality genotype
data for the same samples. In this comparison, results obtained by using the genotype

likelihood-based approaches applied to low depth sequencing data are close to those obtained from the high-quality genotypes for many of the pairs (Figure 4). Chromosome block-jackknife estimates of uncertainty were generally small, and varied by relationship type, with the pair of full siblings having the most uncertainty in R0, R1, and KING-robust kinship. Notably, the IBS-based approach provided meaningful results, even though it did not make use of any external allele information. With the IBS method, unrelated individuals had a slight decrease in R0 and a slight increase in R1 and kinship, relative to the SFS method, making them appear slightly more related than expected for unrelated individuals from a homogenous population. However, despite this bias, it is still possible to correctly categorize of all the pairs to their relationship category within the set of close familial relationships [PO, FS, HS, or UR]. Suggesting that this method can be used when not enough information is available for the realSFS method.

Finally, we also made R1-R0 plot and the R1-KING-robust kinship plots based on genotypes that we obtained through a standard genotype calling procedure from the raw read data. We did this to investigate if the genotype likelihood-based approaches are necessary or one could just as well use genotypes called from the 4x data. Genotype calling had a large negative effect on the outcome; in the resulting R1-R0 and R1-KING-robust kinship plots the half siblings appear where first cousins are expected and both the parent-offspring and full sibling pairs appear in the expected range of half siblings (Figure 4). These results demonstrate the potential pitfalls of basing relationship inferences, including R1-R0 and R1-KING-robust kinship plots, on genotypes called from low-depth data. Notably, this is also the case for the methods presented in (Rosenberg 2006) and (Manichaikul *et al.* 2010) (Figure 5A). Hence, for 4x sequencing data, called genotypes are not a good alternative to genotype likelihood-based approaches.

Assessing the effect of SNP ascertainment scheme

To assess the effect of SNP ascertainment on the new method, and thus the effect of the data-generation platform or SNP filtering, we applied it to three different subsets of data from the five 1000 Genomes individuals. The results for each of the three ascertainment

schemes (Illumina 650Y SNP array sites, 1000 Genomes release sites and whole

genomes) are similar (Figure 4E and F versus Figure 4B and C, respectively), suggesting

SNP ascertainment does not have a large effect.

For comparison, we performed the same assessment of the methods presented in

Rosenberg (Rosenberg 2006) and (Manichaikul *et al.* 2010) by constructing scatterplots

of the same type as those shown in their papers (Figure 5). This shows that both these

other methods are much more affected by ascertainment than the method proposed

here. The Rosenberg method is affected on both its x-axis (IBS0) and its y-axis (IBS2),

which means that the expected region of the plot for each relationship will be different

for different data sets. The method presented in Manichaikul *et al.* 2010 is affected by

the SNP ascertainment mainly on its x-axis (IBS0). In this case, the ascertainment

mainly affects the ability to distinguish between parent-offspring and full siblings, since

the y-axis, which is only slightly affected by ascertainment, is the kinship coefficient,

which can be used to distinguish between most close relationships except for parent-

offspring and full siblings. The x-axis, IBS0, is included in part to help make the

distinction between PO and FS (Manichaikul *et al.* 2010), but this ability is clearly

affected by SNP ascertainment (Figure 5B).

Discussion

We have presented a simple new method for inferring if and how two individuals are

related based solely on genetic data from the two target individuals. This new method

has several advantages compared to most existing relatedness inference methods, which

are almost all dependent on additional information. First, it does not require any

information about allele frequencies. Second, it does not require information about the

relative genomic positions of the sites included in the analysis. Third, although the

method is straightforward when applied to genotype data (where it boils down to simple

counting, addition, and division), we demonstrate two ways to utilize sequencing data

via genotype likelihoods and we show that 4x sequencing data that is insufficient for

high-quality genotype calling can provide useful results. This implies that this new

method can be used even if all you have is low-depth sequencing data from a few

individuals from a species without a reference genome. This is not likely to be the typical

situation in studies of modern humans, however it is much more likely to occur in

studies of with non-model organisms or ancient samples. Fourth and finally, we have

shown that the new method is robust to the SNP ascertainment present in common SNP

array data sets.

Comparison to similar methods

The new method is based on plotting two statistics, $R0$ and KING-robust kinship against

a third, new statistic $R1$. The $R0$ statistic is derived from the test statistic proposed by

Lee (2003) to test for relatedness; the only differences are that the numerator and

denominator are flipped and that E, the proportion of sites where both individuals are

heterozygous, is included in the denominator in the statistic defined by Lee but absent

in $R0$. The KING-robust kinship estimator was proposed by Manichaikul *et al.* (2010),

here we simply extend it to estimation directly from sequencing data. Notably, our

results suggest that this extension is important for low-depth sequencing data, because

estimates based on genotypes called from low-depth sequencing data are not very good.

And importantly this extension can also be used for similar statistics and thus makes

existing methods based on such statistics, like KING-robust kinship, more widely

applicable.

However, this extension is not the only contribution of this study. Another key new

contribution of this study is the fact that $R0$ and KING-robust kinship are combined

with the new $R1$ statistic instead of with IBS0 (the proportion of sites where the two

individuals share zero alleles IBS) as was previously proposed by Rosenberg (2006) and

Manichaikul *et al.* (2010). As we have shown, the fraction of sites that are IBS0 or IBS2

is very sensitive SNP ascertainment, meaning that results are only comparable within

each different ascertainment scheme. Across different ascertainment schemes, such as

between SNP arrays and sequencing data, the method from Rosenberg (2006), where

IBS0 is combined with IBS2, is difficult, if not impossible, to use for relatedness

inference because the fraction of sites that are IBS2 and IBS0 varies so wildly. On the

other hand, KING-robust kinship is still very useful, but it loses its ability to distinguish

between parent-offspring and full siblings, as IBS0 was used for this. Due to this

sensitivity to ascertainment samples cannot be analyzed in isolation, and must be placed

in the context of other samples with known relationships and the same ascertainment

scheme. This requirement makes it difficult to apply these previous methods to ancient

humans or other species with limited sample sizes.

In contrast, the ability to identify relatives based on expected values is maintained in the

new method, regardless of ascertainment scheme due to the use of R1, instead of IBS0,

which makes the new method robust to SNP ascertainment. Here the R1-R0 plot is

especially useful, because parent-offspring pairs, tend to have an R0 estimate extremely

close to 0, making them particularly easy to identify. Additionally, it is worth noticing

that R1 not only makes it easier to distinguish between parent-offspring and full

siblings, both the new plot types, R1-R0 and R1-KING-robust kinship provide better

separation of several of the different relationship categories than KING-robust kinship

does alone (Figure S2). One appealing aspect of KING-robust kinship is that kinship

axis has a biological interpretation, defined as the probability that two alleles sampled at

random from two individuals are identical by descent. Hence the two plots types, R1-R0

and R1-KING-robust kinship, each have their advantages.

Finally, it is worth noticing that the two plots types seem to work better than a range of

other plots constructed from similar ration statistics that we explored (Figure S5).

Limitations and applications

While the new method provides significant advantages over previous methods in

situations with limited data, it does have some limitations. First, like most other

relatedness inference methods the proposed method assumes that the individuals are

not inbred and that they originate from the same homogenous, population. And like

many other relationship inference methods, it is not necessarily robust to violations of

these assumptions. Previous studies have shown the effect of admixture on relatedness

inference is complex and can potentially lead to bias in either direction depending on

the circumstances (Thornton *et al.* 2012; Moltke and Albrechtsen 2014; Conomos *et al.*

2016; Ramstetter *et al.* 2017). Specific methods have been developed to correct for

admixture when the allele frequencies in the admixing populations are known (e.g.

Thornton *et al.* 2012; Moltke and Albrechtsen 2014), or enough samples are available

(Conomos *et al.* 2016). However, since these methods work by exploiting knowledge

about allele frequencies or access to large sample sizes for their correction, the R_0, R_1

and KING-robust kinship statistics can not be corrected in a similar manner. Instead we

suggest trying to detect inbreeding and admixture before performing relatedness

analyses and to proceed with caution when it is present. One potential way to identify

admixture or inbreeding is to compare the heterozygosities for the pair of individuals;

two non-inbred and non-admixed individuals from the same population should have

similar heterozygosity, so heterozygosity differences can be a warning signal.

We can also flip these relatedness statistics around and ask if the pair of individuals

appear to be *less* related than we expect from two individuals sampled from a single

population. In this vein, Lee (2003) showed that the statistic he proposed can also be

used to detect if two unrelated samples are not from the same homogeneous population.

If this is the case, the Lee's statistic will be significantly larger than 2/3; equivalently, R_0

will be significantly above 0.5 and KING-robust kinship will be negative.

A second limitation is that the categorization of different relationship pairs is based on

visual inspection of plots and that in some cases such classification will be difficult

because the ranges of the different relationship in the plots are close to each other and

even overlap. However, this is to a large extent due to biological variation caused by the

randomness of recombination (Hill and Weir 2011). This limitation in classification

accuracy is therefore shared with the allele frequency based inference methods, as

illustrated by Figure S5.

Despite these limitations, we believe that the results presented here suggest that the new

method constitutes a helpful new tool for relatedness inference that is especially well-

suited for situations with limited data. Identifying related individuals is a crucial step in

nearly any genetic analysis and removing the requirements to specify allele frequencies

and to call genotypes has the potential allow the identification of relatives even in small

studies of non-model species or ancient samples. These types of studies do not currently

have many good options to address relatedness among their samples. An additional

appeal of the IBS-based approach introduced here, is that because it considers all

possible pairwise genotype categories, it can potentially help identify specific error types

that are expected to occur in DNA from ancient samples, by allowing researchers to

specifically identify transitions. Also, due to its non-reliance on allele frequencies the

method is also potentially useful when relatedness is pervasive among many samples,

complicating allele frequency estimation in the broader population (Mcpeek *et al.*

2004). Hence, the method will hopefully be useful to many.

Code availability:

The IBS method is available at:

http://www.popgen.dk/angsd/index.php/Genotype_Distribution

References

Albrechtsen, A., T. Sand Korneliussen, I. Moltke, T. van Overseem Hansen, F. C. Nielsen *et al.*, 2009 Relatedness mapping and tracts of relatedness for genome-wide data in the presence of linkage disequilibrium. Genet Epidemiol 33: 266-274.

Alexander, D. H., J. Novembre and K. Lange, 2009 Fast model-based estimation of ancestry in unrelated individuals. Genome Res 19: 1655-1664.

Baca, M., K. Doan, M. Sobczyk, A. Stankovic and P. Weglenski, 2012 Ancient DNA reveals kinship burial patterns of a pre-Columbian Andean community. BMC Genet 13: 30.

Balding, D. J., 2006 A tutorial on statistical methods for population association studies. Nat Rev Genet 7: 781-791.

Busing, F. M. T. A., E. Meijer and R. Van Der Leeden, 1999 Delete-m jackknife for unequal m. Statistics and Computing 9: 3-8.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al.*, 2015 Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4: 7.

Conomos, M. P., A. P. Reiner, B. S. Weir and T. A. Thornton, 2016 Model-free Estimation of Recent Genetic Relatedness. Am J Hum Genet 98: 127-148.

Derrien, T., J. Estelle, S. Marco Sola, D. G. Knowles, E. Raineri *et al.*, 2012 Fast computation and applications of genome mappability. PLoS One 7: e30377.

Hill, W. G., and B. S. Weir, 2011 Variation in actual relationship as a consequence of Mendelian sampling and linkage. Genet Res (Camb) 93: 47-64.

Kardos, M., G. Luikart and F. W. Allendorf, 2015 Measuring individual inbreeding in the age of genomics: marker-based measures are better than pedigrees. Heredity (Edinb) 115: 63-72.

Korneliussen, T. S., A. Albrechtsen and R. Nielsen, 2014 ANGSD: Analysis of Next Generation Sequencing Data. BMC Bioinformatics 15: 356.

Korneliussen, T. S., and I. Moltke, 2015 NgsRelate: a software tool for estimating pairwise relatedness from next-generation sequencing data. Bioinformatics 31: 4009-4011.

Lee, W. C., 2003 Testing the genetic relation between two individuals using a panel of frequency-unknown single nucleotide polymorphisms. Annals of Human Genetics 67: 618-619.

Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics 27: 2987-2993.

Manichaikul, A., J. C. Mychaleckyj, S. S. Rich, K. Daly, M. Sale *et al.*, 2010 Robust
relationship inference in genome-wide association studies. Bioinformatics 26:
2867-2873.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res 20: 1297-1303.

McPeek, M. S., X. Wu and C. Ober, 2004 Best linear unbiased allele-frequency
estimation in complex pedigrees. Biometrics 60: 359-367.

Moltke, I., and A. Albrechtsen, 2014 RelateAdmix: a software tool for estimating
relatedness between admixed individuals. Bioinformatics 30: 1027-1028.

Monroy Kuhn, J. M., M. Jakobsson and T. Günther, 2017 Estimating genetic kin
relationships in prehistoric populations. bioRxiv.

Nielsen, R., T. Korneliussen, A. Albrechtsen, Y. Li and J. Wang, 2012 SNP calling,
genotype calling, and sample allele frequency estimation from New-Generation
Sequencing data. PLoS One 7: e37558.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira *et al.*, 2007 PLINK: a
tool set for whole-genome association and population-based linkage analyses. Am
J Hum Genet 81: 559-575.

Ramstetter, M. D., T. D. Dyer, D. M. Lehman, J. E. Curran, R. Duggirala *et al.*, 2017
Benchmarking Relatedness Inference Methods with Genome-Wide Data from
Thousands of Relatives. Genetics 207: 75-82.

Rosenberg, N. A., 2006 Standardized subsets of the HGDP-CEPH Human Genome
Diversity Cell Line Panel, accounting for atypical and duplicated samples and
pairs of close relatives. Ann Hum Genet 70: 841-847.

Rosenberg, N. A., J. K. Pritchard, J. L. Weber, H. M. Cann, K. K. Kidd *et al.*, 2002
Genetic structure of human populations. Science 298: 2381-2385.

Sikora, M., A. Seguin-Orlando, V. C. Sousa, A. Albrechtsen, T. Korneliussen *et al.*, 2017
Ancient genomes show social and reproductive behavior of early Upper
Paleolithic foragers. Science 358: 659-662.

Stevens, E. L., G. Heckenberg, E. D. Roberson, J. D. Baugher, T. J. Downey *et al.*, 2011
Inference of relationships in population data using identity-by-descent and
identity-by-state. PLoS Genet 7: e1002287.

The Genomes Project, C., 2015 A global reference for human genetic variation. Nature
526: 68-74.

Thompson, E. A., 1975 The estimation of pairwise relationships. Ann Hum Genet 39:
173-188.

Thompson, E. A., 2000 Statistical inference from genetic data on pedigrees, pp. i-169 in
NSF-CBMS regional conference series in probability and statistics. JSTOR.

Thornton, T., H. Tang, T. J. Hoffmann, H. M. Ochs-Balcom, B. J. Caan *et al.*, 2012
Estimating kinship in admixed populations. Am J Hum Genet 91: 122-138.

Wiuf, C., 2006 Consistency of estimators of population scaled parameters using
composite likelihood. Journal of Mathematical Biology 53: 821-841.
638

Figures for:

Allele frequency-free inference of close familial relationships from genotypes or low depth sequencing data

For submission to Genetics

Authors:

Ryan K. Waples
Anders Albrechtsen
Ida Moltke

Figure 1. Overview of the R0, R1, and King-robust kinship statistics
A Definition of the pairwise genotype categories A - I. Here g_1 and g_2 denotes the genotype for each of the two diploid individuals, 1 and 2. These genotypes are defined as the number of copies of a certain allele carried by 1 and 2, respectively. We assume diallelic variants such that g_1 and g_2 each has 3 possible values: 0, 1, and 2. For a pair of individuals, there are nine possible genotype combinations. We organize them into a 3x3 matrix and denote them with the letters from A to I. The values $A - I$ can equivalently be either counts or proportions. **B** Definition of the R0 statistic based on the notation illustrated in A. **C** Definition of the R1 statistic based on the notation illustrated in A. **D** Definition of the KING-robust kinship estimator (Manichaikul *et al.* 2010), based on the notation illustrated in A.

Figure 2. Joint expectation ranges for R1-R0 and for R1-KING robust kinship.

A Joint expectation ranges for R0 - R1 for five different relationship categories: Parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1), and unrelated (UR). The expected value for PO is shown as a purple X. The expected value range for monozygotic twins (MZ) is not depicted because it is (R1=0, R0=∞).

B Joint expectation ranges for KING-robust kinship and R1 jointly for six different relationship categories: Parent-offspring (PO), full siblings (FS), half siblings/avuncular/grandparent-grandchild (HS), first cousins (C1), and unrelated (UR). The expected value for PO is shown as a purple X.

Figure 3.

R1-R0 and R1-KING-robust kinship scatterplots for seven HGDP populations. Each point represents a pair of individuals and is colored according to the relationship category inferred using an allele frequency based approach. Shaded areas and lines show the expected ranges for specific relationship categories, as in Figure 2. Note that in addition to the relationship categories for Figure 1 there an additional category here representing distantly related pairs (DR). **A** R1-R0 plot for all pairs of individuals within each population. **B** R1-KING-robust kinship plot for all pairs of individuals within each population. **C** Scatterplot of the two relatedness coefficients k1 and k2 estimated using the allele frequency-based approach implemented in PLINK. The coloring illustrates how these estimates were used to categorize the pairwise relationships, and thus determine the coloring in panel A-B. **D** Zoom into the outlined area in A. **E** Zoom into the outlined area in B.

Figure 4.

Relatedness plots for all pairs among 5 LWK individuals from the 1000G Project.

A R1-R0 plots for pairs of the same five LWK individuals for five different analysis methods: 1) IBS = reference-free estimation from (~4x) 1000G bam files, 2) realSFS = site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G genotypes from the 1000G project, 4) HGDP sites=1000G genotypes at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes= de novo genotype calling from 1000G bam files. Shaded areas show the joint expected ranges across the two statistics. Points are colored by their true relationship status, as in Figure 3. Thin grey lines show the confidence intervals (+/- 2 SE) with a chromosome jackknife.

B R1-KING-robust kinship plots for the same data sets, confidence intervals and expected ranges are constructed in the same way. The dashed gray line shows kinship of zero.

Figure 5. Plots of the two other frequency-free methods for different subsets of SNPs.
A Results from applying the plotting approach from Rosenberg (2006) to pairs of the same five LWK individuals for five different analysis methods: 1) IBS = reference-free estimation from (~4x) 1000G bam files, 2) realSFS = site-frequency spectrum based estimation from (~4x) 1000G bam files, 3) 1000G genotypes from the 1000G project, 4) HGDP sites=1000G genotypes at sites that overlap with the Illumina 650Y array used for the HGDP, and 5) called genotypes= de novo genotype calling from 1000G bam files. Pairs are colored by their true relationship status, as in Figure 3. Thin grey lines on each point show the confidence intervals (+/- 2 SE) with a chromosome jackknife. Fraction IBS0/IBS2 are the overall fraction of sites that are IBS0/IBS2, respectively.
B Results from applying the KING-robust based approach to different subsets of SNPs to pairs of the five LWK individuals.